V. An Account of a Journey into Africa from the Cape of Good-Hope, and a Description of a new Species of Cuckow. By Dr. Andreas Sparrman, of the Royal Academy of Stockholm, in a Letter to Dr. John Reinhold Forster, F. R. S.

DEAR SIR,

Gottenburg, Sept. 16, 1776.

Read Dec. 19, after an absence of five years from it, I will endeavour to give you a short account of my expedition into Africa, which I undertook soon after parting with you at the Cape of Good-Hope. The voyage round the world, of which I shared the perils and pleasures with you, had only made me more eager to continue my rambles in quest of new discoveries. I set out therefore from the neighbourhood of the Cape-town in the beginning of August 1775, with no other company than the son of the Dutch lieutenant EMELMAN, who had formerly accompanied my learned friend Dr. THUNBERG on a similar journey, and some Hottentots who took care of my oxen.

The

The first misfortune I met with was the loss of the thermometer which you had left me, and which broke before I had reached the hot-baths. This was only a prelude to greater diffresses. The drought was so violent this year, that the like had not been experienced in the colony within the memory of man, and it obliged the inhabitants to leave their country-feats. A great part of their cattle perished for want of grass and water, and I have frequently fuffered the most raging thirst in the hot defarts which I traversed; but I was too well seasoned during the voyage to dread the hardships of a scanty fubfistence, the fatigues of travelling, or the power of the climate. The most sensible misfortune which the dry feafon brought along with it, was the defolation of the vegetable kingdom. Far from being fo fortunate as Dr. THUNBERG, who has added above a thousand species to the Flora Capensis, I found every thing burnt up, and only in the thickest forests met with some perennial plants which were new to me, and which, upon a revifal of that gentleman's herbal, I believe are likewise unknown to him. Of these I propose to send you specimens as foon as I can find time to bring my collection into some kind of order. On the other hand, I have been fortunate with animals, and especially in the class of quadrupeds. I shall not speak of lions and other beafts

of prey, which I have frequently feen in broad day-light, and heard roaring about me at night, though they never ventured to attack our cattle. But it was chiefly among the antelopes and animals of that fort that I hunted. Mr. EMELMAN and myfelf, with nine hottentots, a waggon drawn by feveral pairs of oxen, and feveral hunting horfes, happily traverfed a defart of fifty miles, where we had greater fport than any German prince could ever boaft of. On that route I penetrated farther into the country than any of my predeceffors, having gone one hundred miles beyond the laft Christian's or Dutchman's hut, into the district of the Yellow or (as they are vulgarly called) Chinese Hottentots.

The great buffaloes which inhabit the wilds of Africa, do not appear to me to differ in any respect from the North American Bison, although I have seen great numbers of them. I have likewise found a species of pole-cat on that continent which LINNÆUS calls Viverra Putorius, contrary to M. DE BUFFON'S opinion, who seems to confine this animal and its species to America. By the sea-side I was fortunate enough to catch a Manatee alive, notwithstanding the difficulty which must attend the capture of such an unweildy animal. There I likewise saw some islands, on which I was told an English ship had been lost. These I suspected at first to be the Doddingtons; but afterwards

afterwards had reason to doubt it, those islands being supposed to lie in a more southerly latitude.

I have had opportunities of making many curious and valuable Observations relative to the different tribes of Hottentots, their oeconomy, hunting-matches, and other customs; an account of which, together with some remarks on the natural history of the elephant, the rhinoceros, and other animals, I intend to prepare for the press. I am possessed of an accurate map of that part of Africa which I have visited, containing all the hills, together with the smallest rivulets, as far as the Bay de la Goa, which, I think, will be a great addition to the work. I only regret that I was not able to draw the objects of natural history, and have an hundred times wished that your son had been with me for this purpose.

As I had been upwards of nine months on this journey, at my return to the Cape I found that my acquaintance had given up all hopes of feeing me again, having had no tidings of me for fo long a space of time. Notwithstanding the many dangers to which I had been exposed on this expedition, I assure you I was greatly tempted to stay another year, in hopes of being more successful in botanical discoveries. However, the prospect of securing the spoils which I had collected, prevailed on me to relinquish that scheme. Indeed I little thought at that time that the greatest Vol. LXVII.

danger awaited my collection in Sweden. A few days ago a great part of it has been damaged here by fire, which has been particularly fatal to my stuffed birds, having destroyed some which were not yet described.

As I am well acquainted with the pleasure which every new discovery in the history of nature gives you, I take this early opportunity of expressing the readiness with which I wish to contribute to your satisfaction, and have subjoined to this letter an account of a curious bird, a species of Cuckow, which I have saved out of the fire. I only beg that you will consider it as an earnest of more important communications, as soon as the hurry of my affairs will permit me to bring my papers into order. In the mean time if you should think that account, and the annexed drawing, worthy the attention of the Royal Society, I should be greatly flattered if you would do me the homour to lay it before that learned Body.

With the greatest esteem I remain, &c.

THE HISTORY OF THE HONEY-GUIDE, OR CUCULUS INDICATOR.

THIS curious species of Cuckow is found at a considerable diffance from the Cape of Good-Hope, in the interior parts of Africa, being entirely unknown at that fettlement. The first place where I heard of it was in a wood, called the Groot Vaader's Bosch, the Grand Father's Wood, fituated in a defart near the river which the Hottentots call T'kaut'kai. The Dutch fettlers thereabouts have given this bird the name of Honig-wyzer, or Honey-guide, from its quality of discovering wildhoney to travellers. Its colour has nothing striking or beautiful, as will appear from the description and drawing annexed; and its fize is confiderably fmaller than that of our Cuckow in Europe: but in return, the instinct which prompts it to feek its food in a fingular manner, is truly admirable. Not only the Dutch and Hottentots, but likewife a species of quadruped, which the Dutch name a Ratel(a), are frequently conducted to wild bee-hives by this bird, which as it were pilots them to the very fpot. The honey being its favourite food, its own interest prompts it to be inftrumental in robbing the hive, as

(a) Probably a new species of badger.

fome fcraps are commonly left for its fupport. The morning and evening are its times of feeding, and it is then heard calling in a shrill tone cherr, cherr, which the honeyhunters carefully attend to as the fummons to the chace. From time to time they answer with a foft whiftle, which the bird hearing, always continues its note. As foon as they are in fight of each other, the bird gradually flutters towards the place where the hive is fituated, continually repeating its former call of cherr, cherr: nay, if it should happen to have gained a considerable way before the men (who may eafily be hindered in the purfuit by bushes, rivers, and the like) it returns to them again, and redoubles its note, as if to reproach them with their inactivity. At last the bird is observed to hover for a few moments over a certain fpot, and then filently retiring to a neighbouring bush or other resting-place, the hunters are fure of finding the bees nest in that identical spot, whether it be in a tree, or in the crevice of a rock, or (as is most commonly the case) in the earth. Whilst the hunters are bufy in taking the honey, the bird is feen looking on attentively to what is going forward, and waiting for its share of the spoil. The bee-hunters never fail to leave a fmall portion for their conductor, but commonly take care not to leave fo much as would fatisfy its hunger. The bird's appetite being only whetted by this parfimony,

it is obliged to commit a fecond treason, by discovering another bees-nest, in hopes of a better falary. It is further observed, that the nearer the bird approaches the hidden hive, the more frequently it repeats its call, and feems more impatient.

I have had frequent opportunities of feeing this bird, and have been witness of the destruction of several republicks of bees, by means of its treachery. I had however but two opportunities of shooting it, which I did to the great indignation of my Hottentots. From those specimens (both of which are supposed to be females) I have made the subsequent description. The inhabitants in general accuse the same bird of sometimes conducting its followers where wild beasts and venomous serpents have their places of abode: this however I never had an opportunity of ascertaining myself; but am apt to believe such cases to be accidental, when dangerous animals happen to be in the neighbourhood of a bees-nest.

Whilst I staid in the interior parts of Africa, a nest was shewn to me, which some peasants assured me was the nest of a Honey-guide. It was woven of slender filaments or fibres of bark, in the form of a bottle. The neck and opening hung downwards, and a string in an arched

fhape

shape was suspended across the opening, fastened by the two ends, perhaps for the bird to perch upon.

DESCRIPTIO CUCULI INDICATORIS.

ROSTRUM craffiusculum, versus basin suscum, apice luteum.

Angulus oris usque infra oculos extensus.

Nares postremæ ad basin rostri, supremæ vicinæ ut carinula dorsali saltem separerentur, oblongæ, margine prominulo.

Pili aliquot ad bafin rostri, præcipuè in mandibula inferiore.

Lingua plana, subsagittata.

Oculorum irides ferrugineo-grifeæ.

Palpebræ nudæ, nigræ.

Pedes nigri, scansorii. Tibiæ breves; Ungues tenues, nigri.

Pileus læte grifeus e pennis brevibus latiusculis.


Gula, Jugulum, Pectus, fordidè alba, cum aliquo virore vix notabili in pectore.

Dorsum et Uropygium ferrugineo-grisea.

Abdomen, Crissumque alba.

Femora tecta pennis albis, macula longitudinali nigra notatis.

Alarum



CUCULUS INDICATOR.

ICATOR.

The Honey Guide.

Befor fo

Alarum tectrices superiores omnes griseo-susceptis fummis aliquot quæ flavis apicibus formant maculam flavam in humeris, exiguam, et a plumis scapularibus sæpe tectam.

Tectrices infra alam albidæ, harum fupremæ ex albido nigroque maculatæ.

Remiges omnes supra fusci, primarii octo, secundarii sex, subrus cinereo-fusci.

Alula griseo-fusca.


Cauda cuneiformis, rectricibus duodecim: liarum duæ intermediæ longiores angustiores, supra et infra æruginoso-fuscæ; proximæ duæ suliginosæ, margine interiore albicantes; duæ utrinque his proximæ, albæ, apice susce, et exterius ad basin macula nigra notatæ; extima utrinque reliquis brevior, alba, apice susce s

Ale complicatæ caudæ partem quartam attingunt.

Longitudo ab apice rostri ad extremum caudæ circiter septem uncias pedis Anglicani explet.

Rostrum a basi superiore ad apicem semunciale.





CUCULUS INDICATOR.

The Honey Guide.