

PHILOSOPHICAL TRANSACTIONS.

I. *Catalogue of Nebulæ and Clusters of Stars.*
By Sir JOHN FREDERICK WILLIAM HERSCHEL, Bart., F.R.S.

Received October 16,—Read November 19, 1863.

Introduction.

THE study of the Nebulæ has, within the last quarter of a century, attracted much more of the attention of observers than heretofore—as well on account of the singularity of the phenomena presented by many of these objects, as in consequence of the increased optical power of the telescopes which the skill and industry of modern inventors and artists have placed within their reach. The brighter nebulæ cannot be viewed to any advantage, and the fainter cannot be seen at all, except by the aid of telescopes of large aperture; and, thanks to the exertions of Lord ROSSE, Mr. LASSELL, Messrs. NASMYTH and DE LA RUE in England, and Messrs. STEINHEIL, FOUCAULT, and PORRO in Germany and France, as regards reflecting telescopes, and to those of FRAUNHOFER, MERZ, CAUCHOIX, CLARKE, COOK, SECRETAN, ROSS, and DALLMEYER as regards refractors; instruments of abundantly sufficient optical capacity not only to repeat and verify the earlier observations, but to disclose new and more interesting features in many cases, have now come into the hands of many observers, both professional astronomers and amateurs, and may be had by any one who is willing to incur a cost which may be considered moderate when it is remembered that instruments of similar dimensions and goodness could not be obtained fifty years ago at any price. In consequence we find a continually increasing attention directed to this department of astronomy. Not to insist on the observations of the Earl of ROSSE and Mr. LASSELL with their transcendent reflectors, we find a systematic examination and review of them undertaken by M. D'ARREST in the year 1855, by the aid of a refractor of 6-feet focal length and $4\frac{1}{2}$ inches aperture in the Leipzig Observatory, whose results, consisting in the carefully determined places, by repeated observations, of about 230 nebulæ, were published in 1856, in a work entitled “*Resultate aus Beobachtungen der Nebelflecken und Sternhaufen*” (Erste Reihe, Leipzig). This review has since been carried on by the same excellent astronomer, with the great refractor by MERZ of 11 inches in aperture and 16-feet focus, erected in the year 1861 at the Royal Observa-

tory of Copenhagen. Again, from the Observatory of the Collegio Romano, under the direction of Signor SECCHI, have emanated many valuable observations, and from that at Harvard College, Cambridge, U.S., under the late and present Professors BOND, some of the most striking pictorial representations of particular nebulae which we possess. Neither ought a short but very valuable memoir by the late E. MASON, printed in the 7th volume of the Memoirs of the American Academy of Arts and Sciences, to be passed in silence; containing as it does a very elaborate and minute examination, and some excellent delineations of several highly interesting nebulae, particularly those in the great nebulous region of Cygnus. To M. AUWERS also we owe many accurate and valuable observations, besides a Catalogue comprising the whole series of Sir WILLIAM HERSCHEL'S nebulae arranged in order of right ascension and reduced to a common epoch, of which more hereafter. Should the efforts which are now making to procure for the University of Melbourne in Australia a reflector of the first magnitude prove, as is to be hoped, successful, it is understood that one of the principal uses to which it will be devoted will be the examination and exact delineation of the numerous and wonderful objects of this class which the southern hemisphere presents.

These circumstances, but more especially the last-mentioned, render it extremely desirable to have presented in one work, without the necessity of turning over many volumes, a general catalogue of all the nebulae and clusters of stars actually known, both northern and southern, arranged in order of right ascension and reduced to a common and sufficiently advanced epoch which may serve as a general index to them, and enable an observer at once to turn his instrument on any one of them, as well as to put it in his power immediately to ascertain whether any object of this nature which he may encounter in his observations is new, or should be set down as one previously observed. For want of such a general catalogue, in fact, a great many nebulae have been, from time to time, in the 'Astronomische Nachrichten' and elsewhere, introduced to the world as new discoveries, which have since been identified with nebulae already described and well known. Many a supposed comet, too, would have been recognized at once as a nebula, had such a general catalogue been at hand, and much valuable time been thus saved to their observers in looking out for them again.

Besides these there are other considerations which have weighed with me in undertaking the task of compiling such a general catalogue. Having, in the course of my own observations, received the greatest possible assistance from the possession of a Manuscript Catalogue of all the nebulae and clusters discovered by my Father, brought to the common epoch 1800.0, and arranged in zones of 1° in breadth in polar distance, by his sister the late Miss CAROLINA L. HERSCHEL, it seemed to me nothing less than a debt of gratitude, not merely to acknowledge that assistance, but to avail myself still further of it to complete the list of his nebulae by supplying from that catalogue the places of all those nebulae among them which had escaped my own observation (a very numerous list), and by inserting from it all those places of nebulae observed by myself which were deficient in either element (of R.A. or P.D.), or in which I had reason to apprehend

greater errors than those which probably affected her results. This I have accordingly done. But to do it effectually, and at the same time to effect a thoroughly correct identification of the objects in my catalogues with those of the older series, involved, as a necessary preliminary step, the reduction to 1830·0 of the whole of her catalogue, an operation in which I received the assistance of my sons; the computations being executed for each nebula in duplicate and checked by myself, and which, taken leisurely, zone by zone, as time and circumstance permitted, proved less onerous and wearisome than might have been expected. The Catalogue thus reduced to the same epoch as my own, afforded the means of detecting and rectifying a great many errors of nomenclature in the latter. And it was in the course of this part of the inquiry, in which many cases of considerable intricacy and difficulty occurred (as will be evident on a perusal of the notes appended to this Catalogue), and in which it became necessary to recur both to the original sweeps and to a series of registered extracts from them (the nature of which will be more distinctly stated hereafter), that I learned fully to appreciate the skill, diligence, and accuracy which that indefatigable lady brought to bear on a task which only the most boundless devotion could have induced her to undertake or enabled her to accomplish.

Arrived at this stage—that is to say, the mean results of all the observations in my own Catalogues taken, and all the deficient or imperfectly observed nebulæ in my Father's list supplied, as above stated, and the whole arranged, not in zones, but in general order of right ascension,—it then became necessary, in order to produce a work available for future observation, to bring the whole up to a still more advanced epoch. The work required for this purpose, calling no longer for any discussion, or collation of the original observations or registers, but being one of simple arithmetical computation from a definite formula—the Royal Society, at my application, very liberally undertook to supply, from the funds at their annual disposal, the amount necessary to procure its execution by an experienced computer (Mr. KERSCHNER, one of the occasional computists for the Royal Observatory of Greenwich). This work the Astronomer Royal most obligingly offered to superintend, affording at the same time his advice as to the general principle on which the computation should be conducted. The plan suggested by him and adopted in effect was this. Each object in the Catalogue was first roughly brought up to the year 1880 by the application of approximate precessions in R.A. and P.D. The places so obtained were then employed to compute the exact precessions in both by the usual formulæ, with coefficients for the year 1880·0, viz.

$$\text{Precession in R.A.} = 3^{\text{s}}\cdot072 + 1^{\text{s}}\cdot337 \cdot \sin \text{R.A.} \cotan \text{P.D.}$$

$$\text{Precession in P.D.} = -20''\cdot06 \cos \text{R.A.}$$

And the precessions, so calculated, were then used to bring up the places from 1830 to 1860, the epoch of the Catalogue; so that, the places being given for 1860 and the precessions for twenty years in advance, the application of those precessions to those places shall give dependable places for any year up to the year 1930, at which time the small error in excess or defect of the true precession consequent on using the fifty years'

antecedent place of the object will be exactly compensated by the further change of place in the same direction in the *subsequent* fifty. Two cases of excessive proximity to the poles, northern and southern, viz. those of the nebulae Nos. 2043 and 1652 of the present Catalogue, are excepted, the precessions changing so rapidly, and with so much deviation from uniformity, that a rigorous computation, at least in R.A., will always be necessary. In the case of No. 2043, the effect of precession in the thirty years from 1830 to 1860 has been to change the R.A. from $2^{\text{h}} 32^{\text{m}}$ to $10^{\text{h}} 8^{\text{m}}$.

This computation was completed, and a fair copy of the resulting places, arranged *de novo* in their order of R.A. for 1860, forwarded to me on the 6th of February last (1863). The nomenclature of the objects having in the interim been settled satisfactorily by myself, and a description of each nebula, from a careful comparison of all the descriptions given, prepared, it remained only to fill in the columns left blank for these and the other necessary particulars, and to complete the Catalogue by the insertion in their proper places of the places and descriptions of all such other nebulae, non-observed by either my Father or myself, similarly reduced, of which I could gather any accounts. These will be found enumerated further on in the "Explanation and arrangement of the Catalogue."

On the 23rd of February last, while engaged in this work, I received, by the kindness of the Astronomer Royal, a copy of the important work of M. AUWERS before alluded to, entitled "WILLIAM HERSCHEL'S Verzeichnissen von Nebelflecken und Sternhaufen, bearbeitet von ARTHUR AUWERS. Königsberg, 1862," of whose existence this was my first notice. It contains a complete and most elaborate reduction to 1830, from the observed differences in R.A. and P.D. with known stars, recorded in the Philosophical Transactions, of all the nebulae and clusters in my Father's three Catalogues; together with a separate catalogue of all those collected by MESSIER from his own observations, or those of MECHAIN and others (101 in number), similarly reduced; another of LACAILLE'S southern nebulae, and one of 50 "new nebulae," comprising nearly all those observed by other astronomers (Lord ROSSE excepted) in this hemisphere—all brought up to the same epoch.

It may be readily supposed that I lost no time in comparing my own previous work with this of M. AUWERS; the places of which having been obtained by the aid of far better and more dependable catalogues of stars, to give the true positions of the zero-points or determining stars in the differential observations, as well as of more exact precessions, and doubtless, a much more systematic process of treatment, would be entitled, *observation for observation*, to be considered as representing the original sweeps more faithfully than could be expected from my own preparatory catalogue. On the other hand, however, the Zone Catalogue from which that was derived possessed the advantage of having been deduced, not from a single difference of R.A. and P.D. between each nebula and a single determining star, but from *all* the observations of each nebula; often in many different sweeps, and in the same sweep often from more than one star; thus eliminating, no doubt, a great deal of casual error. In that catalogue, too, as in my own catalogues of 1833 and that of the southern nebulae, the individual results of each observation, or, to speak more exactly, of each differential comparison, is separately

recorded, so that any suspiciously large deviation from the mean of all may be at once noticed and traced to its origin in the sweeping books. My reduction was of course based on the means of all these (rejecting such as were obviously and grossly faulty), and might therefore, *pro tanto*, be regarded as of superior authority. This consideration, joined to that before adduced, decided me to retain those places in the present Catalogue which had been derived from this source, except in a few instances (specified in the notes) when it proved, by careful examination of the causes of discordance, that actual mistakes had been committed. And I must not omit to add that the comparison so instituted with M. AUWERS'S results has led me to the detection of several grave errors in my own work which would certainly have otherwise escaped notice (and in some cases have caused the loss of future observations by missetting the telescope), and whose rectification has added materially to its value. On the other hand, as no human work is perfect, I have been led to notice some errors in M. AUWERS'S work itself, which are set down in a list of *errata* and *corrigenda* at the end of this Catalogue; and besides, a good many cases in which, owing to mistakes in the printed catalogue in the volumes of the Philosophical Transactions (many of which stood corrected in MS. in the margin of the copy of those Transactions in my possession, and many more have been silently detected and rectified by Miss C. H. in her subsequent computations), his calculations have been founded on erroneous data, and have therefore led him to assign erroneous places to the objects so affected. Thus on every account the result has been what may be considered a complete expurgation of both our catalogues.

It remains for me to say a few words on the way in which the reduction to 1860 and the calculation of the precessions have been performed by Mr. KERSCHNER, the computist employed by the Astronomer Royal for that purpose. The whole work has been executed on printed forms, which being preserved may at any time be referred to. Since error in computation, however practised the computer, and however checked, is always possible, and occasional error of copying, especially when the order of the entries has to be rearranged, is absolutely unavoidable, I considered it incumbent on me to recalculate, *seriatim* from my original MS. Catalogue for 1830, and taking for granted the precessions set down in the fair copy, for 1880, the places both in R.A. and P.D. of every object included in the Catalogue; keeping an eye meanwhile to the precessions themselves, and their signs, to seize the least indication of error in that quarter. It would have been too laborious to recompute these. As for the precessions in P.D., their regular progression of itself ensures their correctness, as far at least as the integer seconds and the first decimal place. A pretty considerable number of errors (most of them of little moment) was thus detected and corrected—not more, however, than might reasonably be expected in the work of the most expert computist in so extensive a work, consisting of between nine and ten thousand computed entries (taking both elements), and traceable moreover in many instances to obvious misreading, and in some to actual misentry on my part, of figures in the original MS., which but for this further examination would also have escaped notice altogether.

The correction of these and the other errors already spoken of necessitated, in a great

many instances, a change in the order of R.A., and a consequent erasure and interlineation in the MS. The introduction, too, of the other nebulæ (those of M. AUWERS'S catalogue of "novæ," those communicated to me for insertion by M. D'ARREST, and those noticed by Lord ROSSE in his memoir of 1861, amounting altogether to 433 objects) necessitated many more interlineations, often occurring very inconveniently, two or three together, in a way to disfigure the MS. considerably. Unfortunately, too, in the MS. itself the column headed "No. in the Catalogue," which I had intended to have been left blank till all the rest of the work was completed, had been filled in by the transcriber with a series of numbers in regular progression, from 1 to 4629, the actual number of lines of which it then consisted. This made it necessary to renumber the whole *ab initio* in red ink, striking out the former numbers, and thus producing a still more unsightly appearance. Under these circumstances, I debated whether or not to recopy the whole. But, to say nothing of the sacrifice of time (since I could have entrusted it to no other hand), I believe it impossible to copy so voluminous a mass of figures and abbreviated writing without numerous errors. And being satisfied, from the repeated and careful revision it has undergone, of its present correctness, and equally so that with ordinary care on the part of the compositor (should the Council of the Royal Society decide on printing it) no *mistake* can arise from any of the alterations and interlineations it contains, I have decided in favour of presenting it as it stands, with the exception of two sheets which it was absolutely necessary to recopy owing to the extreme closeness of the interlineations, the smallness of the writing, and the transpositions needed. These have each been twice carefully read with the original.

In presenting to the Royal Society this Catalogue, it will be accompanied by the following series of records and documents which it may become desirable hereafter to refer to in elucidation of any point which may arise respecting the history or reduction of such of the objects as occur in my Father's classes and numbers printed in the Philosophical Transactions, viz.—

1st. A series of "*register sheets*," in which are entered up *all* the observations of *each* nebula or cluster copied verbatim from the sweeps, the nebulæ, &c. being arranged in the order of their dates of discovery. These are the "register sheets" referred to in the notes on this Catalogue, and cited by their *general (i. e. current) number*, as H, 1; H, 2; ... H, 2508.

2nd. A similar set of register sheets of all the observations of each of MESSIER'S nebulæ, arranged according to MESSIER'S numbers.

3rd. A general index of the 2508 nebulæ in classes and numbers, to find the "general number" of each to facilitate reference to the register sheets. (This index was drawn up by myself.)

4th. An index list of the same nebulæ, &c. arranged according to the "general number," to find the class and number of each.

5th. A more complete ditto ditto, containing also the rough approximate R.A. and P.D. of each object for 1800, and the determining stars as in the Philosophical Transactions.

6th. A catalogue in zones of P.D. of all the said nebulæ and clusters arranged in each

zone in order of R.A., and reduced to the year 1800 by Miss C. L. HERSCHEL, exhibiting the reduced result of each separate observation of each nebula; together with the determining star or stars in each case, and the differences of R.A. and P.D. from such star, with references to the current number of the sweep in which the observation is contained.

7th. The original sweeps with the 20-foot reflector at Slough in which the nebulæ were observed, contained in three small quarto and four folio volumes of MS.

All these manuscripts, with exception of the index No. 3, are in the original hand-writings of my Father and his Sister, in most cases easily distinguishable, in some others not so readily. The Zone Catalogue No. 6 is entirely the autograph of the latter.

Explanation and arrangement of the Catalogue.

The Catalogue is arranged in twelve columns, of which the first contains the general or current number in order from 1 up to 5063, the total number of objects comprised, including six supplementary ones, whose insertion in their proper order in R.A. would have involved altering all the numbering both of the catalogue and the annotations, &c., and would have proved a source of confusion and unavoidable error. Nevertheless, to prevent their being overlooked by any observer who may consult the catalogue for the purpose of a general review of the nebulæ, or for the verification of a new one, their numbers are interpolated into the general series so as to catch the eye, and a reference made to the supplementary catalogue in each case in the column of descriptions.

Column 2 contains the numbers of those nebulæ of which observations are given in my two former catalogues, and those of the two nubeculæ; the numbers from 1 to 2307 inclusive being from that in Philosophical Transactions 1833, and from 2308 to 4021 from my Cape observations. Where a number in this column is enclosed in hooks thus [], it is taken from the Catalogue of Objects in the Nubecula minor in pp. 153 to 155 of that work. Where in parentheses thus (), from those in the Nubecula major, pp. 156 to 163.

Column 3 contains the classes and numbers of nebulæ as given by my Father in his three Catalogues in the Philosophical Transactions for 1786, 1789, and 1802. One only is omitted, viz. V. 35. It is an immense diffused nebosity, extending from $5^{\text{h}} 27^{\text{m}}$ to $5^{\text{h}} 42^{\text{m}}$ in R.A., and from $98^{\circ} 6'$ to $87^{\circ} 43'$ in P.D. A special list of these great diffusions of nebula is given by M. AUWERS in p. 42 of the work above cited.

Column 4 contains references to other authorities, and gives either the name of the first discoverer of the nebula, or a reference to the particular list or catalogue of nebula which has been taken as the authority for the place set down. The principal of these are—1st. The list of “new nebula” (Verzeichniss neuer Nebelflecke), in pp. 73 to 76 of the work of M. AUWERS already cited. These are referred to in the following form:—Auw. N. 1, Auw. N. 2, &c. 2ndly. Under the form D’Arr. 1, 2, &c., are given a series of objects contained in a MS. list of 125 nebula, kindly communicated to me by their discoverer, M. D’ARREST, Director of the Royal Observatory of Copenhagen, and reduced by him to the epoch (1860.0) of this Catalogue, with their precessions for 1880. 3rdly. A great number of nebula cited under the form “R. novæ,” whose places have been approxi-

mately obtained from the diagrams accompanied by micrometrical measures of position and distance, or from more loose and general indications contained in Lord Rosse's paper in the Philosophical Transactions for 1861, the comparisons being in all cases made with those nebulae in my Catalogue of 1833 whose numbers stand annexed to them in column 2, with an italic letter appended, thus:—

322, *a* R. nova; 319, *a* R. 3 novae.

In cases of which latter kind it is intended to express merely that nebulae to the number indicated, not otherwise identifiable, will be found on due search in the immediate neighbourhood of the place approximately set down. Lastly. The names of Professor G. P. BOND, Mr. S. COOLIDGE, and Mr. J. T. SAFFORD in this column of the supplementary list of nebulae refer to the places of nebulae and clusters in a list of objects of that description discovered at the Observatory of Harvard College, obligingly communicated to me by Professor BOND, Director of that establishment, too late for their introduction into the body of the Catalogue.

Besides these references, in which the places set down have been adopted from the catalogues above mentioned, column 4 also contains synonyms or identifications of objects observed by myself with those contained in MESSIER'S lists communicated to the French Academy, or to the Connaissance des Temps for 1783 and 1784. These are cited by the number they bear in MESSIER'S own list, thus, M. 1, M. 2, &c. They have, with very few exceptions, been observed and described by myself or my Father, and their places here set down are given as results from our observations. In the few excepted cases they are taken from M. AUWERS'S catalogue already spoken of. The nebulae also whose identity has been (sometimes satisfactorily, but for the most part very doubtfully) made out with objects in Mr. DUNLOP'S Catalogue of Southern Nebulae, are indicated by the letter Δ , thus, Δ . 169, &c. In a few cases, chiefly those of nebulous stars, planetary nebulae, or very star-like objects, which have been set down as stars in catalogues of authority; these are also referred to by name and number in column 4.

Many of Mr. DUNLOP'S nebulae are contained in LACAILLE'S catalogue, as also some of MESSIER'S, but of that catalogue two objects only, not so identifiable, viz. Nos. 38 and 40 of M. AUWERS'S catalogue of LACAILLE'S nebulae, have been considered as definitely enough described (*nébuleuses sans étoiles*) by that astronomer to be included in the present Catalogue.

Column 5 contains the Right Ascension in time for 1860.0 of each object in the Catalogue. When this is given to decimals of seconds, it is to be understood as having been brought up from the mean of the observations given in my former Catalogues, or from the mean of those (where not observed by myself) in Miss C. HERSCHEL'S Zone Catalogue above mentioned*. When the R.A. is given only to the nearest minute or degree, it will of course be understood that the place is too loosely determined to render

* In some cases a careful subsequent revision of the catalogued observations *seriatim* has necessitated altering these R.A.'s by a few decimals of a second (seldom more) *after the process of reduction to 1860*. In all such cases the alteration has been applied as a *correction* to Mr. KERSCHNER'S figures, so as not to disturb the amounts of precession allowed—a procedure perfectly legitimate and productive of no error. The same remark applies to col. 8.

further precision of statement other than illusory. This is the case with the greater part of those set down as “R. novæ.”

Column 6 contains the precession, in seconds and decimals, in R.A. for 1880·0.

Column 7 contains the number of observations in R.A. which have been actually used in concluding the R.A. for 1830, from which that for 1860 has been computed. In all cases (unless where the contrary is especially indicated in a note, or otherwise as by the letters B.A.C. or A.S.C., Au., &c. inserted in place of a number in the column itself—which indicate that the R.A. is that of a star in one of those catalogues, or rests upon that other authority), the observations used for all objects included in my former catalogues are brought up from the data there registered, to the exclusion of all others; and in such cases (the vast majority) no parenthesis or other distinctive mark is applied. When, however, no satisfactory R.A. is there recorded, or when the R.A. is there expressly stated to have been set down from the “working list,” the R.A. adopted is that brought up from the Zone Catalogue of C.H., and in such cases the number of observations used is enclosed in parentheses (). Dots attached (·) indicate some uncertainty in the R.A.; (::) a very considerable doubt, extending, perhaps, to a whole minute; ? and ?? express still wider limits of uncertainty. In those nebulæ of my Father’s catalogues which have no number corresponding in column 2 (indicating the absence of any observations of my own), the places set down both in R.A. and Declination are those brought up from the Zone Catalogue of Miss C. H., and the numbers of observations on which they rely are set down in the appropriate column without any parenthesis or distinctive marks, the absence of any number in column 2 being a sufficient indication. In the case of M. D’ARREST’S nebulæ, the numbers in column 6 enclosed thus [] indicate the number of his observations of the nebula employed by himself to give the place.

Columns 8, 9, and 10 contain, in like manner, the North Polar Distance for 1860, the precessions for 1880, and the numbers of observations used for P.D. in the case of each object; and the same remarks apply to these as to columns 5, 6, and 7.

In column 11 is given a short description of the nebula or cluster in abbreviated words, made out from an assemblage and comparison of *all* the descriptions of each object given in my Father’s *and* my own observations. As regards the former, recourse was had, not to the printed account in the Philosophical Transactions (which gives only a single description), but to a series of manuscript sheets in the nature of a REGISTER (and *as such cited* in the notes which follow this Catalogue), into which have been transcribed, *verbatim*, from the original sweeps, all the descriptive parts of each and every observation of each cluster or nebula in the order of their dates, and the data for computing their places, derived from the sweeps by applying the index and clock corrections pertaining to each. In this Register the nebulæ are entered, each with its class and number, and each on a separate sheet; the whole series being arranged, however, not in the order of their classes and numbers, but in the order of the dates of their discovery, from No. 1, corresponding to October 28, 1783, to No. 2508, corresponding to September 30, 1802. Of these, the first 2500 only are included in the catalogues com-

municated to the Royal Society; the other 8 are printed in the form of an Appendix to my Cape Catalogue, in p. 128 of the "Results of Observations," &c. A similar and separate Register in sheets has been kept for my Father's observations of MESSIER'S nebulae, and these have in like manner been collated with my own observations of the same objects in framing the ultimate, or, as it may be termed, the average description of each.

In making out these descriptions, it was found to a certain degree practicable, in the particulars of brightness, size, and extension, to make a kind of arithmetical approximation to a mean conclusion, by arranging the degrees of brightness, &c. in a progressive upward scale from 1 to 10, and taking a mean of these numbers in each case, as indicating the designating words to be finally adopted. Thus, taking the extreme degree of faintness when a nebula was declared to be "excessively faint," or "barely visible," or "hardly more than suspected" for 1, and "extremely" or "excessively bright" for 10, the intermediate degrees, such as *very faint*, *faint*, *considerably faint*, *pretty faint*, *pretty bright*, *considerably bright*, *Bright*, *very bright*, were denoted by the intermediate numbers 2, 3, 4, 5, 6, 7, 8, 9; and similarly for the scale of sizes, exchanging the words Small and Large for Faint and Bright. In the case of extension, the scale 1 to 10 was supposed arranged in the order, *Round*, *very little extended*, *elliptic or oval*, *considerably extended*, *pretty much extended*, *much extended*, *very much extended*, *extremely extended*, or *a long ray*. It is obvious that the qualifying words, such as "pretty" and "considerably," admit of a good deal of latitude of interpretation, and that, in reference to brightness or faintness, greatness or smallness, their meaning is rather relative than absolute; and especially, that as between *bright or faint*, and "considerably bright" or "considerably faint," for instance, there is so little real distinction of an absolute kind, that it is impossible to say which is to be accepted as indicating the superior degree. In the case of extension there is the same indistinctness as to precedence between the qualifying phrases "considerably" and "pretty much." Nicety, however, in this respect would be misplaced, when it is considered that when several descriptions of the same nebula, observed at different times, come to be compared, they can hardly ever be reconciled except by allowing to each qualification a latitude of meaning extending over several degrees of our arbitrary scale. In many instances, indeed, the discordance, or rather contradiction is so great, as to authorize a strong suspicion of variability in the object itself. In a few cases where, from the low altitude of the object in England, coupled with corresponding discordances of description, it was evident that it must have been seen to much greater advantage from the Cape station (as, for example, in that of h. 3375 = H. III. 754), additional weight has been attributed to the Cape observations.

In the descriptions, I have found it absolutely necessary to abstain from any specification of the estimated sizes of nebulae or clusters in angular measure. In comparing estimations of this kind I find the discordance so great, and (to speak only of my own practice) so little evidence of adherence to any definite standard of estimation, that nothing but confusion would have arisen from introducing such estimates. Nevertheless, as in the use of such a catalogue as the present some guide is necessary for the

observer, to advertise him of what sort of object he may expect to see, the following scale may be taken as conveying a general idea of the magnitudes intended by the conventional words used. Thus, a round nebula of 3" or 4" in diameter would be called *extremely small*;

one of 10" or 12", *very small*;
 20" or 30", *small*, or *considerably small*;
 50" or 60", *pretty small*, or *pretty large*;
 3' or 4', *considerably large*, or *large*;
 8' or 10', *very large*;
 20' and upwards, *extremely large*.

In estimating clusters of stars (that is to say, of well separated and scattered stars) a wider acceptation must be understood, so that, for instance, a cluster of only 1' in extent would be considered *extremely* or *very small*; one of 15' or 20' *large*, and one of 30' or 40' *very large*. This amplification of scale, however, must not be held applicable to those resolved or resolvable clusters of a "globular" character marked in the descriptions as \oplus , which must be understood as belonging to "nebulae" and not to "clusters," so far as the conventional terms used in the descriptions are concerned. I should observe also, that when in making out the average appropriate phrase in size I have found any extravagant discordance between the estimate in words and that in figures, as, for instance, where a nebula has been described in words as *very large*, and the diameter then set down as 2', a compromise has usually been made, and the word modified, as, for instance, to *large* or *considerably large*.

The abbreviations employed in the column of descriptions and elsewhere, in the notes, &c., are as follows:—

ab.	about.	ch.	chevelure.
alm.	almost.	com.	cometic.
am.	among.	cont.	in contact.
app.	appended.	C.	Compressed.
att.	attached.	Cl.	cluster.
Auw.	Auwers.	C.G.H.	"Results of observations, &c. at the Cape of Good Hope."
A.S.C.	Astronomical Society's Catalogue.	C.H.	Miss Carolina Herschel. When it occurs in column 4 it indicates that the object was discovered by her.
b.	brighter.	d.	diameter.
bet.	between.	dist.	distance.
biN.	binuclear.		distant.
bn.	brightest towards the north side.	dif.	diffused.
bs.	brightest towards the south side.	diffic.	difficult.
bp.	brightest towards the preceding side.	D.	double.
bf.	brightest towards the following side.	D'Arr.	D'Arrest.
B.	Bright.	Δ.	Dunlop.
Br.	Brisbane (Sir T.'s) Catalogue of Stars.	def.	defined.
Bo.	Bode.	e.	extremely.
B.A.C.	British Association Catalogue.		
c.	considerably.		
co.	coarse, coarsely.		

ee.	excessively.	rr.	partially resolved—some stars visible.
er.	easily resolvable.	rrr.	well resolved—clearly seen to consist of stars.
exc.	excentric.	R.	round.
E.	extended.	RR.	exactly round.
f.	following.	R. nova.	New nebula discovered by Lord Rosse.
F.	faint.	R. MS.	Manuscript notes furnished by His Lordship.
g.	gradually.	Ri.	Rich.
gr.	group.	R.	The Earl of Rosse.
H.	Sir William Herschel.	s.	suddenly.
h.	Sir John Herschel.	s.	south.
h.o.n.	list of omitted nebulae in C.G.H.	sp.	south preceding.
i.	irregular.	sf.	south following.
inv.	involved.	sc.	scattered.
	involving.	st.	stars.
iF.	irregular figure.	sev.	several
l.	little (adv.).	susp.	suspected.
	long (adj.).	sh.	shaped.
L.	Large.	stell.	stellar.
Lac.	Lacaille.	S.	small.
Lal.	Lalande.	sm.	smaller.
Lass.	Lassell.	sw.	sweep.
m.	much.	Σ.	Struve.
mm.	mixed magnitudes.	tri-N.	tri-nuclear.
mn.	milky nebulosity.	trap.	trapezium.
mon.	monograph.	v.	very.
M.	Middle, or in the middle.	vv.	an intensive of v.
M.	(in col. 3) Messier.	var.	variable.
Mess.	Messier.	W. H.	Sir W. Herschel.
n.	north.		Besides these abbreviations of words, the following arbitrary signs are used.
neb.	nebula.		* a star; *10 a star of the 10th magnitude.
np.	north preceding.		** a double star; *** a triple star.
nf.	north following.		! a remarkable object; !! very much so; !!! a magnificent or otherwise exceedingly interesting object.
nr.	near.		? doubtful; ?? very doubtful, either as to accuracy of place or reality of existence, according to the column in which it occurs.
N.	nucleus, or to a nucleus.		: , :: , see explanations already given.
o.	omitted.		△ a triangle. Forms a triangle with.
ON.	omitted nebula.		⊕ a globular cluster of stars.
p.	preceding.		○ a planetary nebula.
p.	pretty (before F, B, L, S, &c.).		⊙ an annular nebula.
pg.	pretty gradually.	st. 9	Stars from the ninth (or other) magnitude downwards.
pm.	pretty much.	st. 9 13	Stars from the ninth down to the 13th magnitude.
ps.	pretty suddenly.		
P.	poor.		
Pi.	Piazzi.		
P.T.	Philosophical Transactions.		
quad.	quadrilateral.		
quar.	quartile.		
r.	resolvable, barely (mottled as if with stars).		

As examples of the interpretation and expansion of these abbreviations some examples are subjoined.

Ex. 1. pB; vL; vg, vsmbMN 15"; pmE 162°·3; "pretty Bright; very Large; at first very gradually, then very suddenly much brighter in the middle to a nucleus 15" in diameter; pretty much extended—the position of the longer dimension micrometrically measured 162°·3 (*i. e.* reckoned from the north round to 162°·3 in the direction nfsp)."

The angles of position in all cases are to be understood as so reckoned. When decimals of degrees are annexed (or if integer, written decimally thus 151°·0), they have been micrometrically measured. If thus, E 0° or E 45°, E 90°, they mean only in or near the meridian, or parallel or oblique to the meridian from nf to sp, &c., as the case may be. If with a \pm annexed, the position is from a more or less careful estimation.

Ex. 2. R; psbM ill def O; pB*10 125°·4, 70"; "Round, pretty suddenly brighter in the middle to an ill-defined planetary disc; has a pretty Bright star of the 10th magnitude, whose position measured *from* the centre of the nebula is 125°·4, and whose distance also from the centre is 70" by estimation."

The relative situations of neighbouring stars or nebulae are *invariably* to be understood as thus reckoned, *i. e.* taking the centre of the nebula or other object described as a starting-point or origin of angle or distance. Thus S*s will mean that a small star is south of *the nebula*, *np nr that a star is *near the nebula* in a north preceding direction from it; * 4^sf, 3'n, that a double star follows the centre of the nebula 4 seconds of time, and is 3' to the north of it.

Ex. 3. Cl; pRi; pmC; L; st6, 10 ... 15. "A cluster; pretty rich; pretty much compressed; Large; consisting of stars one of which is of the 6th, and the rest from the 10th to the 15th magnitudes."

Attached or vicinary stars or small nebulae are always placed at the ends of the descriptions. Thus \oplus sf means that the nebula described "has a globular cluster following and to the southward of it." When, however, the description of a cluster ends abruptly thus, *_{*}, it is to be understood that "the place taken is that of a conspicuous double star."

The 12th column of the Catalogue contains the number of times that each nebula has been observed by both my Father and myself, whether its place were taken or not, comprising all the cases in which the object has been seen, and whether described or not. Since attention has been drawn to the real or supposed variability of nebulae, and since it can hardly be doubted that comets have occasionally been observed as nebulae, this enumeration is not without its importance. In this column the abbreviation "mon" occasionally occurs. In such cases the nebulae have been so often and diligently observed for the purpose of exact delineation or "monographing," that a special enumeration of the observations would be impossible or useless.

Finally, at the end of the line allotted to each nebula occur occasionally one or both of the marks * and †. The former refers to the notes appended to the Catalogue, the latter to the list of figured nebulae in which the publications wherein are contained figures of the nebulae are referred to by plate and figure—those at least which seem entitled, in the present state of astronomical instrument-making and pictorial representation, to be pointed out to the observer as conveying any idea of their appearance.

Notes on the Catalogue.

No.

- 12 h. 5. D'Arrest says, "h. II. positio certe erronea," but gives no indication of the correction required in R.A. or P.D.
- 29 h. 13; II. 241=II. 243. In P.T. the determining * is omitted, and in the statement of the places of these nebulae, as well as of II. 239, 240, 242, and III. 199, there is much confusion, for the correction of which see the list of errata subjoined. Auwers has threaded the intricacies of this maze with singular felicity, but has been misled in the case of II. 243 into assigning to it a totally erroneous place ($22^{\text{h}} 48^{\text{m}}$ R.A., $73^{\circ} 37'$ P.D. 1830), and, in consequence, has not perceived its identity with II. 241.
- 78 II. 3. Auwers makes the P.D. of this neb. (1830)= $99^{\circ} 32'$, from P.T., which places it $2^{\circ} \pm n$ of 17 Ceti. C.H. makes it $1^{\circ} 51' n$ of the same star, or for 1830, $99^{\circ} 42'$. In fact H. has two observations of it, neither of them more than eye-drafts with neighbouring stars, and the P.D. is concluded graphically by C.H. from these diagrams.
- 88 III. 876. The P.D. of Auwers ($81^{\circ} 16'$) is 1° wrong. The place given in P.T. is $1^{\circ} 43' n$ of 51 Piscium; so also in Register (H. 2296).
- 119 Auw. N. 4=D'Arr. 6. The place given is that brought up from D'Arrest's observations, the R.A. being set down only roughly in Auw.
- 132 h. 57=V. 20. Once looked for by Lord Rosse and not seen. Having been observed both by H. and h., there can be no doubt of its existence.
- 138 h. 61=h. 2345=V. 1. In h.'s sweep 733 the position reading is set down as $324^{\circ} \cdot 5$. This is in contradiction with a diagram made at the time, and is an obvious mistake for $234^{\circ} \cdot 5$, which = $180^{\circ} + 54^{\circ} \cdot 5$, agreeing well with the diagram and with 2 obs. of W.H., in both of which it is described as "*nf* to *sp*." There is also an erratum in the C.G.H. Catal., for $143^{\circ} \cdot 8$ read $144^{\circ} \cdot 5$, since $324 \cdot 5 - 180 = 144 \cdot 5$.
- 145 h. 64=II. 621=II. 703. Auwers remarks that A Ceti, the determining star of W.H., does not exist; but C.H. has perceived this, and by using another determining star (13 Ceti, sw. 756, W.H.), has fixed the place of the nebula II. 703 for 1800 at R.A. $0^{\text{h}} 37^{\text{m}} 47^{\text{s}}$, P.D. $93^{\circ} 53'$ (= $93^{\circ} 43'$, 1830), thereby identifying it with II. 621. Auwers, using a conjectural star, sets down the P.D. erroneously as $92^{\circ} 52'$ (1830).
- 165 h. 2356. This is the main body of the nubecula minor.
- 169 h. 2359. A complex object with several nuclei. There is an erratum in the R.A. set down in C.G.H. as resulting from sw. 488, for $46^{\text{m}} 12^{\text{s}} \cdot 1$ read $47^{\text{m}} 12^{\text{s}} \cdot 1$.
- 177 79, *a*, *b*. In Lord Rosse's diagram, α =h. 79, β =h. 78, γ =nova, accidentally omitted in the body of the Catalogue, but inserted as No. 5058 at the end. The whole Catalogue having been finally numbered before the omission was detected, it could not be inserted in its place. δ is a star; ϵ =h. 79, *a*.

- No.
 178 } h. 4007, 4008, 4012. In the Catalogue of C.G.H. these nebulae are placed
 179 } erroneously in the 23^h of R.A., owing to a mistake of a whole hour in
 196 } reducing.
 202 }
 203 } These constitute the group laid down by Lord Rosse as seen in and about the
 205 } places of h. 84, 85, 86, viz. his α , β , γ , γ' , δ , ϵ , ζ , θ . Of these, α is No. 202=h. 84;
 206 } β =No. 203=h. 85; γ =No. 206=h. 86; γ' =No. 205=86, a ; δ =No. 209=86, b ;
 207 } ϵ =No. 208=D'Arrest No. 10; ζ =No. 207=D'Arr. 9, and θ =86, c . In the MS.
 208 } notes furnished me by Lord R. it is stated that α =h. 84, β =h. 85, and θ =h. 86.
 209 } The latter identification, however, is incorrect.
 210 }
 214 h. 88=I. 54. This is *not* the I. 54 of the P.T., which proved to be one of
 Messier's nebulae, but another subsequently inserted by W.H., so as not to
 break the order of the numbers, as appears from a MS. correction in P.T., and
 from Register (H. 570).
 275 } These constitute Lord Rosse's group seen in or near the place of h. 103, and
 276 } marked in his diagram as A, β , δ , ϵ , and another unlettered (which call γ).
 277 } These I identify as follows:—A=No. 276=h. 103; β =No. 277=103, b ;
 280 } γ =No. 275=103, a ; δ =No. 280=103, c ; and ϵ =No. 290=103, d .
 290 }
 297 } In reference to M. Auwers's remark on the nebulae 170, 171, as also 167, 168
 311 } (H. class III.), after very careful examination of all the data, I can arrive at no
 317 } other conclusion than that embodied in the present Catalogue under these Nos.:
 319 } h. 118 is certainly not III. 171, neither is h. 120. Both places and descriptions
 325 } disagree.
 313 } h. 119 was taken for III. 556, but no R.A. was obtained, that set down being the
 314 } R.A. brought up from C.H. The descriptions differ so materially, especially in
 the particular of extension, that they are most probably distinct nebulae.
 330 h. 124=VII. 48. Auwers remarks in his 'Verbesserungen zu *h*,' that this cluster,
 h. 124, is not nova, but VII. 48. This is correct. Re-examining sweep 216, I
 find an error of 1° committed in reducing the P.D.
 358 This is not in M. D'Arrest's final list, communicated to me in MS.; but being set
 down by M. Auwers as No. 15 in his 'Verzeichniss neuer Nebelflecke,' I felt
 bound to retain it.
 418 h. 160=h. 2442=I. 62. This nebula, though set down by W.H. as of the 1st
 class (*i. e.* as a bright nebula), could not be seen by D'Arrest with the Leipzig
 Fraunhofer of 6-feet focus and 4½ inches aperture. It is marked in this Cata-
 logue, however, by a mean of 4 observations, only as "F."
 428 55 Andromedæ. Although this star has been eight times examined by Lord Rosse
 without perceiving any nebulous atmosphere, yet as my observation is corrobora-

No.

rative of Piazzi's designation of it as "Nebulosa," it is retained for occasional future examination.

442) h. 169, II. 221. The places agree almost exactly, but the descriptions are irre-
444) conciliable. One makes the nebula round, the other much extended. They
are therefore almost certainly distinct nebulae, and there is therefore probably
some error in the R.A. of II. 221. The neighbourhood is rich in nebulae (see
the next note, however).

442) }
444) In Lord Rosse's diagram of the group about h. 169, assuming α to be h. 169
445) = No. 444, the others will be β = No. 445 = 169, α ; γ = No. 446 = 169, b ;
446) δ = No. 447 = 169, c ; and ϵ = II. 221.
447) }

462 h. 179 = 50 Cassiopeiæ. Retained in the Catalogue for future occasional obser-
vation. Nothing can be more difficult than to verify or disprove the nebulosity
of a considerable star under ordinary atmospheric circumstances.

472 h. 184 = III. 583. Though Lord Rosse on one occasion did not find this nebula,
its existence cannot be doubted, having been found by h. nearly in the place
assigned by C.H.

487 h. 193 = I. 152. M. D'Arrest found this nebula too faint for observation with the
Leipzig refractor, though placed by W.H. in Class I., and standing in this Cata-
logue (from a mean of 3 observations) as a "bright" nebula.

501 h. 204 = III. 604. C.H. and Auwers make the R.A. 1^m less. Both H. and h. rely
on single observations. Sweep 188 h. examined and reduction found correct.

510 h. 206 = III. 457. Not found by Lord Rosse; once looked for. See notes on
Nos. 472 and 132.

516 h. 210 = II. 246. Singularly enough, h. and H. are at issue about the two adja-
cent stars. h. makes the stars south of the nebula; H., on the contrary, places
the nebula south of the stars, and says expressly that both this nebula and
III. 201, observed just previously, were similarly situated with regard to their
attendant stars. Now in h.'s obs. of III. 201 (No. 513) the attendant star *is*
stated to be *sf* the nebula, and in that of II. 246 the larger of the two stars is
south and only a very few degrees preceding. I believe the error to lie on the
side of the older observations, as I have a diagram of the small star nearer to
II. 246, *sf*, which shows that I made no mistake of *n* and *s*.

536 I. 153. Auwers makes the R.A. for 1830 $1^h 28^m 45^s$, whereas C.H. makes it
 $2^h 15^m 13^s$. The cause of the discordance lies in an erratum in P.T. (see list
of errata). In C.H.'s reductions the error is corrected, and I find the correc-
tion verified on reference both to the Register (H. 1488) and the original sweep
(sw. 596). The nebula *follows* (not precedes) the determining star.

549 h. 226 = I. 154. Auwers makes the R.A. of this for 1830, $2^h 23^m 8^s$; C.H.

No.

- 2^h 20^m 57^s.8, by the observations in different sweeps differing only 18^s in R.A. The latter is the more correct; so that M. Auwers's remarks on this nebula are not confirmed. The cause of the disagreement lies in a misprint in P.T. (See List of Errata.)
- 557 } In Lord Rosse's description of this group, α =No. 557=h. 231; β =No. 563=h. 234;
 558 } γ =No. 558=231, α ; δ =No. 559=231, β . The other nebula, "about 12' south
 559 } following," is probably No. 563=h. 234. No. 561=h. 233 seems to have escaped
 561 } notice.
 563 }
- 571 h. 240=II. 238=III. 198. C.H. has overlooked or omitted an obs. of W.H. of III. 198 in sw. 574, which, referred to, confirms Mr. Marth's surmise that the nebulae are identical.
- 573 II. 6. This was probably really a comet, as indicated by its description, having been subsequently looked for and not found.
- 574 h. 244=I. 102. M. D'Arrest found this nebula, when observed with the Leipzig refractor of 4½ inches aperture, inferior to a 1st class nebula. In this Catalogue, from a mean of 5 observations, it ranks as "considerably bright."
- 591 h. 258=I. 1. M. D'Arrest found this nebula, when examined with the Leipzig refractor, not entitled to rank above the 2nd class. With this our present Catalogue agrees, it being set down from a mean of 8 observations as "pretty faint."
- 614 This nebula of Bessel was also looked for and not found by D'Arrest, who therefore supposes it to have been a comet.
- 636 h. 280=II. 502. II. 502 is described by H. as eS; F; stellar. Either then the identity is doubtful, or some change must be suspected. The place, however, agrees well.
- 639 h. 281=IV. 43. Once looked for by Lord Rosse, but not found. (See notes on 134, 472, 510.)
- 646 h. 284=III. 578. The same remark. Twice looked for unsuccessfully by Lord Rosse. On one occasion clouds were passing.
- 654 } In Lord Rosse's diagram of this pair and the neighbouring stars γ and δ , the figure
 655 } is in contradiction with the measures. The position of $\alpha\gamma$, instead of 2°, should, I presume, have been stated thus, $\gamma\alpha=178^\circ$, or, which comes to the same thing, $\alpha\gamma=-2^\circ$. This has been assumed in deducing the place of No. 655=289, α from No. 654=h. 289.
- 656 h. 291=III. 591. H. makes this nebula to be the nf of two, but both those of h. the sf.
- 674 h. 293=II. 603. H.'s description is pB; stellar; a pc* with eS, vF chevelure. The place, however, agrees well with that of h. 293.
- 684 III. 195. Auwers makes the R.A. (1830)=3^h 11^m 50^s and C.H. 3^h 10^m 13^s; but

No.

- a misprint in P.T. (see List of Errata) accounts for the difference of the minute at least.
- 708) III. 959; I. 60. The catalogued places contradict the described position of and
709) np; but this is owing to the error in R.A. of I. 60, which D'Arrest makes less by 40^s, which would place I. 60 at 3^h 19^m 35^s (1860).
- 710 Au. N. 17. The discovery of this nebula is attributed by Au. to Schönfeld in 1858, but it seems to be identical with that described by Tuttle (*Astronom. Notices*, xix. p. 224). Auwers's place is preferred, Tuttle's being only approximate.
- 768 Au. N. 18. The celebrated variable nebula of Tempel, discovered Oct. 19, 1859.
- 774 II. 594. Auwers considers this as identical with II. 548, with 1° mistaken in P.D.
- 778 h. 309=I. 155. Auwers makes the R.A. of I. 155 for 1830=3^h 53^m 33^s, destroying the identity of these two nebulae. But his place is deduced from an erroneous entry in P.T. (see List of Errata). C.H., by 2 observations in sweeps 608, 638 agreeing to 3^s in R.A. and 2' in P.D., gives a place which, brought up to 1830, gives R.A. 3^h 37^m 58^s; P.D. 94° 29' 7".
810. h. 311=IV. 69. M. D'Arrest found the nebulous atmosphere around the central star of this nebula very conspicuous with the Leipzig 4½-inch refractor.
- 826 h. 2618=IV. 26. D'Arrest's R.A. is preferred, that of h. 2618 being clearly shown to be erroneous.
- 836 II. 464. The P.D. is given by W.H. as the same with that of 44 Eridani. C.H., using an erroneous place of this star, makes the P.D. 5' too small. This is here corrected, and the result agrees with Auwers.
- 839 Auw. N. 20. This is the remarkable variable nebula discovered by Mr. Hind on Oct. 11, 1852. M. D'Arrest testifies to its complete disappearance on the 3rd and 4th of Oct. 1861, "Hujus nebulae . . . ne umbram quidem detegere valeo." —"Caelo serenissimo regionem summâ curâ perlustravi adjuvante Dr. Schjellerup. *Nebula reverâ* deest." (In 1855 and 1856 it was found by M. D'Arrest within 2' of Mr. Hind's original place.) On Dec. 29, 1861, it was seen by M. Otto Struve with the great Pulkowa refractor, but so excessively faint as to be barely within the power of that instrument. On March 22, 1862, with the same telescope, it was again seen, but considerably brighter, so as to bear a faint illumination of the wires.
- 851 h. 314=III. 587. Not seen by Lord Rosse, once looked for, clouds passing. See notes on Nos. 639, 646, &c.
- 880 h. 322. The bright star preceding is ν Eridani.
- 908 h. 333=II. 547. Not seen by Lord Rosse, once looked for. See notes 132, 472, &c.
- 926 h. 335. Erroneously identified in my Catalogue of 1833 with III. 453 (No. 981). See the note on that nebula.

- No.
- 953 h. 341=D'Arrest 48. Observed by him as "nova," but since recognized as unquestionably =h. 341.
- 970 VIII. 43. Auwers makes the P.D. of this cluster for 1830 = $66^{\circ} 25'$, which is incorrect. The determining star is 109, *n*, Tauri, the cluster being $1^{\circ} 29'$ north of the star. This would give $66^{\circ} 39'$ for the P.D. for 1800, agreeing with C.H., and $66^{\circ} 36'$ for 1830.
- 975 h. 343. A very large diffused nebulosity, distributed in zigzags. This has been looked for seven times by Lord Rosse and not found. Its existence is therefore very doubtful.
- 979 h. 2709. The place graphically determined by measurement of a diagram, as compared with h. 2710.
- 981 III. 453. This was erroneously identified with h. 335 in my Catalogue of 1833. By an unlucky coincidence, its place per working list, roughly brought up from C.H., agreed so well with the latter nebula as taken in sw. 322 (h.), that it was unhesitatingly assumed to be the same. It appears, however, that in C.H.'s reduction an error of 10^m in R.A. has been committed, the star of comparison being 10 Orionis, and the nebula *following* the star by $5^m 7^s$ (as ascertained by reference both to the register sheet (H.1160) and the original sweep (sw. 462, H.)). M. Auwers, misled by my erroneous identification, has assumed that the nebula must have *preceded* the star, which would (nearly) account for the difference, and in consequence, his R.A. of this nebula is 10^m too small. C.H.'s error probably arose from misapplying in like manner the sign of the Δ . R.A.
- 998 III. 268. Auwers's R.A. ($4^h 57^m 23^s$, 1830) is adopted in preference to $5^h 0^m 28^s$, that brought up from C.H. to the same epoch. In the sweep 367 (H.) three stars of comparison are given, 58 Eridani, α Leporis, and 19 Leporis. The Δ . R.A. of α and 19 comes out correct, but that of 58 from each is wrong by $3^m 5^s$, so that the star must have been mistaken. C.H. has used 58 and α , and has rightly brought out the place of the nebula by the former (the wrong star), and wrongly by the right one; and by an odd coincidence the two results agree well, though both wrong.
- 1030 h. 349=VII. 4. Described by D'Arrest as "Ein Ausserordentlich reicher Hauf," an extraordinarily rich cluster.
- 1133 h. 356. Looked for four times by Lord Rosse, in two of which the sky was fancied to have a milky appearance.
- 1138 } h. 2841. Double nebula. In my Cape Catalogue, sweep 538, for "first" and
1139 } "second" read "larger" and "smaller." The smaller is sp. The position 260° is right. It is very remarkable that in sweeps 508, 522, 658, and 761 the smaller of the two was not noticed. Is it variable?
- 1167 III. 747. Auwers makes the P.D. $8' 20''$ greater. It is difficult to identify the determining star used by C.H.

- No.
- 1165 } h. 2866, 2867, 2868, 2869. $16^s.2$ added to all the R.A.'s of these nebulae in the
 1168 } Cape Catalogue to compensate an error detected in sw. 538. The correction is
 1171 } deduced from a comparison of the diagram fig. 20, Pl. VI. C.G.H. with the
 1174 } place of No. 1171.
- 1179 h. 360. $3^s.3$ added to h.'s P.D. to bring it to the place in B.A.C.
- 1180 V. 30. The place of V. 30 corrected by $+3^s.2$ in R.A. and $+25' 45''.4$ in P.D. to bring it to the place of ϵ' 42 Orionis in the B.A.C.
- 1183 h. 361=V. 31. h.'s place corrected by $+0^s.4$ in R.A. and $-0' 27''.2$ in P.D. to bring it to that of ν 44 Orionis in B.A.C.
- 1185 III. 1. ?? There are two observations by H. of III. 1, but they differ enormously. One agrees with M. 43. The place of M. 43 is corrected to agree with its place in the Catalogue of Stars, &c. in the great nebula in Orion, C.G.H. p. 28.
- 1191 Chacornac's recently discovered nebula. Place from Moigno's "les Mondes," No. 9, p. 241.
- 1196 III. 269. Auwers gives as the R.A. of this nebula for 1830 $6^h 27^m 57^s$, which is mistaken by 1^h . The Philosophical Transactions says that it precedes 19 Leporis by $32^m 23^s$, and that this is no misprint appears from C.H.'s reductions.
- 1226 IV. 24. Annular according to Lord Rosse.
- 1287 III. 270. Auwers places this nebula in R.A. $6^h 40^m 20^s$, or an hour too late. Its place is very distinctly settled by two determining stars, α Leporis and 19 Leporis, the former of which it followed by $15^m 4^s$, and preceded the latter by $20^m 0^s$.
- 1425 h. 393=IV. 3. Lord Rosse's account of this nebula is extremely remarkable. "This h. 393," he says, "is an enormous nebulosity which I have traced f and n of it to a great distance—*some degrees*. It narrows at times to a band across the finding eyepiece about $6'$ or $8'$."
- 1440 h. 401=V. 27=VIII. 5. Retained as a cluster, though but a poor one. Nine times examined by Lord Rosse for nebulosity, but none seen.
- 1452 III. 271. Auwers places this nebula in R.A. $8^h 3^m 35^s$, P.D. $76^\circ 21'$ (1830). There has been some mistake. III. 271 is stated to follow 8 (ν 3) Canis, $8^m 0^s$, and to be $4'$ n of that star, which gives a place agreeing with C.H. and with the present Catalogue.
- 1454 h. 441=M. 41. This nebula was also observed by Flamsteed.
- 1455 } In Lord Rosse's diagram of this group, α is No. 1457=h. 410; β =No. 1455=410, α ;
 1456 } γ =No. 1456=410, b ; δ =No. 1458=h. 409; and ϵ =No. 1460=410, c . But
 1457 } some suspicion seems to have arisen that the principal nebulae observed were
 1458 } not really h. 409, 410, but h. 406, 407. In that case the identification will
 1460 } stand as follows:—
- α =No. 1448=h. 406.
 β =h. 406— $5^s.2$ in R.A., and $-1' 25''$ in P.D.
 γ =No. 1449=h. 407.

No.

 $\delta = \text{h. } 406 + 1^{\text{s}}.6$ in R.A., and $-5' 6''$ in P.D. $\epsilon = \text{h. } 406 + 14^{\text{s}}.7$ in R.A., and $-5' 2''$ in P.D.

1480 h. 423. This nebula is entered by C.H. as VIII. 1. B, with a remark "not in print."

1508 h. 439=VI. 6. The R.A. is nearly 2^{m} in excess of C.H. and of Auwers. Examined sweep (h.) 393 in which it was observed. Found all clear and correctly reduced.

1527 } Compared with Lord Rosse's two diagrams of the nebulae composing this group.
 1528 } None of them are "novae." $\alpha = \text{h. } 449$; $\beta = \text{h. } 448$; $\gamma = \text{h. } 447$; $\delta = \beta$; $\epsilon = \gamma$;
 1530 } $\zeta = \text{h. } 446$.
 1531 }

1533 VIII. 44. Auwers's P.D. is 84° , instead of 82° , owing to an erratum in P.T. (See List of Errata.)

1578 h. 468=III. 479. No nebulosity seen by Lord Rosse in 5 observations. In H.'s single observation the nebula is "suspected," and in those of h. it is not positively ascertained. The object seems therefore to be merely a small resolved cluster of vFst.

1594. M. 47. Auwers assigns a R.A. greater by 4^{m} . The cluster has not since been observed. It is probably a very loose and poor one.1611 h. 480=VI. 37. h.'s P.D. corrected by $-10'$ as the presumed error of reading in the single observation obtained. Harding in 1827 (it appears) observed its P.D.= $100^{\circ} 10'$ (for 1830), and W.H.'s place for that epoch is $100^{\circ} 12'$, that of h. being $100^{\circ} 19' 4''$.

1615 } In Lord Rosse's diagram, $\alpha = \text{No. } 1617 = \text{h. } 483$; $\beta = \text{No. } 1616 = \text{D'Arr. } 51$;
 1616 } $\gamma = \text{No. } 1615 = 483$, α . D'Arrest's place for β is preferred to that which results
 1617 } from comparison with the diagram. h. 284 could not have been in the field,
 being almost a degree distant.

1633 h. 493=II. 719. h.'s R.A. in P.T. diminished by 1^{m} for an error of 1^{m} detected in the reduction of the observation. This brings it nearer to Auwers.1652 h. 3176. *Polarissima Australis*. This nebula is so near the south pole that its precession in R.A. varies from year to year with great rapidity, so that its R.A. cannot be computed correctly by the ordinary approximate method.

1666 } The four nebulae h. 508, 510; 510, α ; 510, b evidently include among them that
 1667 } third nebula referred to by Lord Rosse as the accompanying "nova" "forming
 1668 } a triangle with h. 507, 508—of the last degree of faintness." h. 507, however,
 is 30° distant in P.D., so that in the observation of Feb. 9, 1850, the P.D. of h. 507 must doubtless have been read as 36° instead of 66° , giving rise to a mistaken identity with one of the *two* really new nebulae at that time in view.

1696 III. 50. I find a memorandum to the effect that this nebula is lost, and was probably a comet; but I cannot recover my authority for the statement. It is described by H. as "of the last degree of faintness," and it is therefore no way

No.

surprising that it should not have been again perceived without some time and trouble bestowed, and in clear weather.

- 1707 h. 527=II. 48. M. Auwers, owing to an erratum in P.T. (see List of Errata), makes the R.A. of II. 48 two minutes too great, and is thus led to doubt its identity with h. 527. There still remains the rather considerable disagreement of 5' in P.D. D'Arrest found neither of these nebulae; but there can be no doubt of the existence of one at least, in or near the place here given. This is *not* the nebula seen by Lord Rosse "nearly in contact with h. 526." This latter (described already by h. as "bi-nuclear") was seen by R. as distinctly double.
- 1712 h. 531=M. 67. Discovered by Oriani.
- 1720 h. 535=II. 823. W.H. describes this nebula as "Round;" h. as "much extended," while Lord Rosse saw it as bi-nuclear, or a double nebula joined by faint nebulosity. Is it separating into two, like Biela's comet?
- 1735 } h. 542 and II. 557. The descriptions are irreconcilable, and they must be two
1736 } distinct nebulae. The R.A. of h. 542 was *not* observed, and its P.D. is set down as "hardly more than conjectural," having been looked for by working list as II. 557 and set down as such.
- 1742 h. 545=II. 834. Misprinted II. 844 by Auwers in the Catalogue, but the number is correct in his general list of the nebulae by numbers and classes.
- 1743 h. 546. Not seen by Lord Rosse in one observation. Examined sweep 21 (h.) and found all right.
- 1756 III. 291=D'Arr. 60. These are assuredly one and the same nebula. Auwers's declination of III. 291 ($+27^{\circ} 7'$) should be $+26^{\circ} 7'$.
- 1773 h. 565=III. 61. The P.D. according to H. is 70° .
- 1788 II. 708. Owing to an erratum in the determining star in Phil. Trans. (see List of Errata), Auwers has given the place of this nebula for 1830 R.A. $9^{\text{h}} 12^{\text{m}} 39^{\text{s}}$; P.D. $39^{\circ} 17'$, instead of $9^{\text{h}} 6^{\text{m}} 29^{\text{s}}$; $47^{\circ} 20'$.
- 1791 }
1794 } h. 577; h. 578. Not seen by Lord Rosse in one observation. (See next note.)
- 1792 D'Arrest 62. This nebula must surely be variable, as it is inconceivable else that it should not have been seen by h., when h. 578, to which it is almost close, was observed and its place taken. D'Arrest says, "Fugerat Herschelium necnon me anno 1862." Neither of the three (Nos. 1791, 1792, 1794) were seen by Lord Rosse. Sweep 59 (h.) and the reductions re-examined. Found all clearly written and all correct.
- 1804 } h. 581, 582; 581, *a, b, c, d*, 582, *a, b, c, d, e, f, g*; D'Arr. 63. Of this very complex
to }
1815 }
1817 }
1818 }
1821 } group of 15 nebulae or "knots" (as they are called by Lord Rosse), six have been determined from his diagram, and six more by the aid of notes subsequently furnished me from the records of the observatory at Birr Castle, containing differences of R.A. and P.D. from one or other of the former. These

No.

are indicated by the letters MS. attached in the column of descriptions. The others I identify as follows:—

α	(in Lord R.'s diagram)	is No. 1813=582, <i>c</i> .
β	”	” 1812=582, <i>b</i> .
γ	”	” 1811=h. 582.
δ	”	” 1806=h. 581.
ϵ	”	” 1815=582, <i>e</i> .
ζ	”	” 1821=582, <i>g</i> .

One of those for which no data are given must have been D'Arr. 63, and the two remaining ones are included under the entries Nos. 1817, 1818 as 582, *f*.

- 1832 h. 590. Not seen by Lord Rosse; once looked for. Re-examined the sweep and reductions. Found all correct.
- 1868 h. 3171. In the omitted observations of nebulae in the last page of the C.G.H. observations, *for* h. 3170 *read* h. 3171; and this observation, combined with the two in the body of the work, gives the mean result for 1830 employed to deduce the place in the present Catalogue.
- 1911 h. 3185=III. 289. In consequence of a misprint in P.T. (see List of Errata), the P.D. of Auwers is 5' too small. Corrected by this, his place agrees well with my observation.
- 1953 M. 81 ?? A nebula observed by W.H. as described, but differing most materially in place from M. 81. It would certainly be very extraordinary should *three* nebulae so extremely remarkable as M. 81 and 82 and this be found to lie so near together.
- 1959 } h. 3198, 3202 are distinct nebulae, and were observed consecutively in one and
1962 } the same sweep—sw. 561 (h.).
- 1960 } h. 3199 and 3201 are also distinct nebulae, and were observed consecutively in
1961 } sweep 562 (h.).
- 1974 III. 293. M. Auwers makes the place of this nebula $9^{\text{h}} 24^{\text{m}} 4^{\text{s}}$; $66^{\circ} 30'$ (1830), instead of $9^{\text{h}} 48^{\text{m}} 48^{\text{s}}$; $60^{\circ} 13'$. The cause of the error is an erratum (see List) in P.T., where the determining star is set down as 23 Leonis instead of 23 Leonis Minoris, another of the instances of confusion arising from the use of this silly and barbarous nomenclature.
- 2014 h. 669=III. 65. Not seen by Lord Rosse in one observation. It was found by h. in its place *per* working list.
- 2019 h. 672. Not seen by Lord Rosse in one observation. Examined the sweep and reductions, and found all correct.
- 2043 h. 250. This nebula is so very close to the North Pole, that its place cannot be calculated by a precession proportional to the time in the usual approximate mode, the R.A. changing from year to year with extreme rapidity.

- No.
 2054 }
 2055 } In Lord Rosse's diagram, α =No. 2058=h. 692=II. 44; β =No. 2061=h. 693
 2057 } =II. 45; γ =No. 2055=692, *b*; δ =No. 2054=D'Arr. 61; ϵ =No. 2057=692, *c*,
 2058 } not lettered in the diagram.
 2061 }
- 2088 } II. 28, 29. Both D'Arrest and Secchi agree in placing this double nebula more
 2089 } to the south than W.H. by $15' \pm$, and D'Arrest supposes the P.D. to have been
 misread to that extent. As so great a proper motion is most improbable, and
 the identity is indisputable, I have adopted this supposition and made the neces-
 sary correction.
- 2094 h. 706. Not seen by Lord Rosse in 6 observations. Re-examined the record of
 the original obs. Sweep 115 (h.), No. 68, and the reductions. The entries are
 all clear and perfectly legible. Reduction in P.D. correct; reduction in R.A.
 erroneous by $-0^m 26^s.6$. This, however, could not have caused its non-observ-
 ation by R. This then was a comet, or is a lost nebula. The error of reduction
 is corrected in the present Catalogue.
- 2111 III. 316. C.H.'s reduction of this nebula being affected with a considerable error,
 Auwers's R.A. is adopted, after verification.
- 2144 h. 3276. Place approximate, by equatoreal zone review.
- 2189 h. 745=V. 52. Not seen by Lord Rosse when once looked for (see note on
 No. 132, &c.).
- 2192 h. 3294. The minute in R.A. doubtful.
- 2197 h. 3295. The great nebula about η Argus. According to a letter from Mr. Eyre
 B. Powell of Madras, a most extraordinary change has taken place in this nebula
 since my figure of it was delineated. He states that the southern end of the
 curious oval vacuity close to the great star, which was *decidedly closed* when I
 depicted it, is now *decidedly open*. Should this be established, it will be the
 most extraordinary fact that has yet appeared in the history of a nebula.
- 2201 h. 754=II. 99. M. D'Arrest found this nebula in the Leipzig refractor, bright
 enough to be ranked in the first class. And it is marked as "very bright" in
 this Catalogue by a mean of 5 observations. It must have been ill seen in the
 earlier observation when classed as II.
- 2231 } IV. 6=II. 131 and h. 777=III. 88. I adopt, on due consideration, the opinion
 2234 } of Auwers, that III. 88 and II. 131 are not the same. Their having been
 successively observed in the same sweep is decisive. Also, that IV. 6 is not
 III. 88, but in reality identical with II. 131. The descriptions are made out
 in conformity with this.
- 2233 } I. 118 and h. 779. The degree of P.D. is probably mistaken in I. 118. Marth,
 2236 } according to Auw., suggests that the determining star 46 Ursæ (which though

No.

not so called in B.A.C., is doubtless No. 3741 of that catalogue) was mistaken, and should have been called 46 Leonis minoris. Consulting the original sweep (sw. 487, H.), I find this surmise *not* corroborated; for the nebula, when reduced by the star next preceding it (37 Leonis minoris), gives the same Polar distance, and, within a few seconds, the same R.A. But there is some faint indication of the figure 6 in the reading of the Polar distance piece $56^{\circ} 55'$ having been written over a 7, which would have thrown the nebula somewhat below the southern limit of the sweep, and might have caused a suspicion of error at the time. I found no nebula in the catalogued place in my sweep No. 337 (h.), so that the probability of an erroneous degree is strengthened. At the same time, it is not impossible that this nebula may be identical with No. 2236=h. 779, the mistake in the degree lying the other way.

2238 h. 780=I. 172. h., in Ph.Tr., suggests that this nebula may have moved. There is, however, no ground for this supposition, as its place agrees quite remarkably with that brought up from C.H. But query if the double star have not moved, since one of the observations places it "in the middle," and a subsequent one makes the southern extremity of the nebula touch the large star of the double star.

2276 h. 806=II. 101. Found to rank as a first-class nebula by M. D'Arrest with the $4\frac{1}{2}$ -in. Leipzig refractor. In this Catalogue it stands described as "very Bright," by a mean of 4 observations. See remark in note 2201.

2310 h. 823=III. 111. There is a strange amount of discordance between the observed and reduced places of this nebula. Auwers makes the P.D. for 1830= $84^{\circ} 29'$. C.H. has reduced the single observation of W.H. by two stars 84, τ Leonis and 349 Bode Leonis, and her results differ by $10'$; τ , which gives the greater, being stated to be "too far distant in P.D." The several results stand thus:—

P.D. 1830, by Auwers	$84^{\circ} 29'$
„ by τ Leonis (C.H.)	$84 20$
„ by h. obs.	$84 15$
„ by 349 B. Leonis (C.H.)	$84 9\frac{1}{2}$

My observed P.D. is nearly a mean between those of C.H.

2315 h. 828=II. 42. Not seen by Lord Rosse when once looked for (see notes on No. 132, &c.).

2319 h. 829=III. 351. The observations of this nebula, which are numerous, disagree so very remarkably in the particular of brightness, that a considerable suspicion of variability exists.

2373 h. 854=M. 65. There is a misprint, 45° for 75° np to sf, in the position of extension in my Catalogue of 1833. The diagram in the original sweep also corroborates this, as does also the figure (fig. 53) accompanying that Catalogue.

No.

- W.H. twice says mE in merid. (180°)—h.'s position 75° np to sf= 165° ; a mean of those of Winnecke and Auwers = 172° .
- 2377 h. 857, h. 875; M. 66. No doubt these are the same. fig. 54 P.T. 1833 corroborates their identity. The accompanying stars and their positions agree entirely. The R.A. of h. 875, however, requires to be corrected by -3^m , allowing the seconds and the P.D. observed in that observation their weight.
- 2382 II. 30. Auwers deduces his R.A. for 1830 ($11^h 12^m 21^s$) from the statement in P.T. "following 68, δ Leonis, $6^m 30^s$." C.H. from the same data concludes R.A. $11^h 11^m 31^s$ (also for 1830). The latter is (within 2^s) the correct result.
- 2388 h. 867=h. 861? These are very probably the same. But as, after all, the difference of the observed R.A.'s is sufficient to have allowed one to escape while observing the other, so that they *may* be different, and as moreover one is described as "Round," and the other as "extended," both are retained.
- 2405 h. 882=I. 20. This nebula would seem to have decreased in brightness. The bright * is 1341. A.S.C.
- 2411 h. 886=I. 131. Ranked by M. D'Arrest in the second class with the $4\frac{1}{2}$ -inch Leipzig telescope. In this Catalogue it stands as "pretty Bright" from a mean of three observations.
- 2417 III. 112. Auwers has reduced this nebula by the star given in P.T. ϕ , 74 Leonis. But I find a MS. note that this star was not dependable, and that Mayer's No. 510 is the proper determining star. The nebula was subsequently looked for and found, not in the place given by ϕ , but $8'$ from the P.D. concluded from Mayer 510. A mean of these two determinations is therefore used in this Catalogue.
- 2440 h. 907=III. 353. Auwers doubts the identity of these nebulae. But this is in consequence of a misprint in P.T. (see List of Errata), 53^m for 43^m . The error is found also in the Register Sheet (H. 937), but C.H. has avoided it and used 43^m in her reduction so as to give a R.A. agreeing within 35^s with that of h. 907.
- 2461 h. 918=II. 784. Lord Rosse, in his observation of this nebula, mentions "another brush-like, $20'$ np." This was no doubt II. 783=No. 2454.
- 2501 h. 945=I. 94. W.H. makes this nebula by one observation extended, n to s, by another nf to sp, while h. has two observations agreeing in making it extended in the parallel. Surely it does not rotate?
- 2540 h. 967. 1^m added to the R.A. It is evidently the first of the group of 4.
- 2577 III. 113. This nebula is reduced also in Auwers's catalogue by ϕ Leonis, the star set down in P.T. But C.H. remarks that ϕ was above the sweep, and otherwise observed under unfavourable circumstances, and Mayer's 510 zod. star. s. $0^\circ 31'$ is preferred, which gives a result differing by $+24'$ in P.D. and -48^s in R.A. The place adopted in the present Catalogue is in conformity with this remark. (See note on No. 2417.)

- No.
- 2591 h. 1000=III. 616. The star 6m, 5' n only noticed by W.H. The other 7m, f in the parallel only by h. Are there really two stars? and are they both variable?
- 2597 h. 1002=I. 203. Auwers, in consequence of an erratum in P.T. (see List of Errata), makes the R.A. of this nebula 7^m too small. The error is corrected in the Register (H. 1889) and in C.H.'s reduction.
- 2604 h. 1009=I. 202. The same misprint in P.T. mentioned in the last note on h. 2597 has also vitiated M. Auwers's R.A. of this nebula. It is corrected in the Register Sheet (H. 1886) and in C.H.
- 2608 h. 1013=III. 381. I adopt Mr. Marth's identification of these nebulae. The place of III. 381 in the catalogue of C.H., from which my working lists were made out, is vitiated by some great mistake. The P.D. is supposed to be derived from 1 Comæ, the neb. being 1° 12' south of the star. This, however, would give 68° 9' 29" for 1830 instead of 65° 45' 0", that brought up from C.H.
- 2650 h. 1039. This cannot be identical with h. 1036, and its brightness precludes its being accepted as III. 354. But there is extreme uncertainty as to its P.D. The degree may even be wrong.
- 2652 h. 1041=II. 733. According to W.H. the position of extension is "near the meridian." If *meridian* be not a mistake for *parallel* it has changed. h. has a measure 62° 3, and an estimation 65° in another observation.
- 2653 h. 1042. This cannot be III. 3, as C.H. has reduced two obs. of this latter well agreeing, and giving a R.A. 2^m exceeding that of h. 1042, which also rests on 2 obs. of h.
- 2668 h. 1050=I. 253. The difference of descriptions is extraordinary, so that they seem hardly to pertain to the same object; but the places agree.
- 2683
2684
2685
2686
2689 } h. 1062, 3, 4, 5, 7, 8, 1070, 1, 3, 5, III. 391, 2, 3, 4, 5, 6. The places set down
2690 } for the nebulae of this extensive group are made out by a most careful consider-
2693 } ation of all the observations and records in the sweeping books which seem
2694 } irreconcilable with a group of six nebulae only. The group, however, needs a
2697 } thorough re-examination.
2699
2701
2702
- 2730 II. 14. Owing to an erratum in P.T. (see List of Errata) Auwers gives quite an erroneous place for this nebula (11^h 39^m 27^s R.A., 81° 9' P.D. 1830).
- 2747 h. 1103=III. 814. Auwers suspects some error of the press, since his P.D. for 1830 comes out 36° 58', while that of h. 1103 is 35° 56'. There is, however,

No.

no error, either of printing, registry, or reduction in any part of the older work. The determining star is rightly set down as 5 Canum, whose P.D. for 1800 (the epoch of C.H.'s catalogue) is $37^{\circ} 19' 42''$, and III. 814 is declared to be $1^{\circ} 32'$ north of it, so that $35^{\circ} 48'$, the P.D. of C.H., is correct, and reduced to 1830 ($=35^{\circ} 58'$) agrees with my place within $2'$. Neither is there any error of the press or of reduction, or any apparent mistake of a clerical nature in all the process of h. 1103, and the nebula observed is set down *in* the sweeping book (of course from the working list) *as* III. 814. I consider their identity therefore as fully established.

- 2771 } h. 1211=II. 372. H. says, the most northerly of the pair II. 372, III. 360 the
2773 } largest: h., "by diagram," makes the following nebula, III. 360=No. 2773,
the larger of the two.
- 2814 II. 109. The reductions of the sweep 187 (H.) in which this occurs are somewhat precarious, and in C.H.'s revision of the sweep the Δ . P.D. from 6 Comæ is set down at $1^{\circ} 50'$, that in the P.T. at $1^{\circ} 54'$ (these changes are never made without good reason), and this accounts for $4'$ out of the $5'$ difference between her P.D. and that of M. Auwers.
- 2846 III. 535. In a sweep two years subsequent to the obs. of this nebula by H. it was looked for again but not found. ? if a comet.
- 2849 D'Arr. 89. M. D'Arrest makes mention in a letter which he has done me the honour to address to me, of a nebula having the same R.A. as this, but a P.D. (1860)= $83^{\circ} 46' 42''$. He does not include it in his final list. It should, however, be looked for.
- 2852 } h. 1183, 7, 9, 1190, 4; II. 568, 9, 570, 1, 2, 3. There cannot be a doubt that
2856 } II. 568, 569, 570, 571, are in 82° P.D., and II. 572, 3, in 83° . It is equally
2857 } certain that h. 1183, 1189, 1190, 1194 are in 83° . They were observed in two
2862 } distinct sweeps (sw. 111 and 238); I observed also II. 572 in sw. 238, and III.
2865 } 573 in sw. 250. There must be a set of nebulae, at least 8 in number, hereabouts.
2869 } N.B. W.H. makes II. 568, 569, 570, 571, $34'$ n. of 11 Virginis. If n. be a
mistake for s, these agree with h. 1187, 1189, 1190, 1194.
- 2855 h. 1186=I. 90=II. 322. Marth's conjecture is right (see Auwers's note on I. 90) as regards II. 322, but not so his conclusion that II. 322=II. 377.
- 2878 h. 1202=I. 139=M. 61. Discovered by Oriani. N.B. The first discoverers of the nebulae in Messier's list, when not Messier himself, are mentioned by M. Auwers in his catalogue of those nebulae (pp. 66-71), except in the cases of Oriani's nebulae, M. 14?, 18?, 35?, 61, 67.
- 2884 1202, *a*. Under h. 1196 and 1202, two nebulae, unidentifiable, are described as companions, but there must be some great error in Lord Rosse's account of them, as the place of one is referred to a scarlet star " $10'$ south of a scarlet star R.A. $12^{\text{h}} 25^{\text{m}}$." Now h. 1202 is in R.A. $12^{\text{h}} 14^{\text{m}}$. To afford a fair chance

No.

- of reobserving them, the companion $10'$ nf h. 1202 is entered here as 1202, *a*, and that south of the scarlet star, under No. 3060 as 1196, *a*.
- 2892 D'Arr. 90. "Reperta a me Mart. 4, 1862. Eandem reperit Schönfeldus, April 1, 1862. Vide Comptes Rendus, &c."
- 2951 II. 87. This *may* be h. 1240, but $7'$ in P.D. is a large error.
- 2961 h. 1253=M. 86. The nebula of Lord Rosse $14'$ sp this is no doubt II. 168.
- 2976 h. 1261=III. 492. III. 492 was looked for April 11, 1787, by W.H. in the place assigned to it, but was not seen. Auwers, however, makes it identical with h. 1261. Yet the descriptions are radically different, and after all there may be another nebula, the real III. 492, in the neighbourhood.
- 2992) R. novæ. 1274, *a*; 1275, *a*. Of the eleven "knots" seen by Lord Rosse in this
2995) place these two are the only really "novæ." The other 9 were h. 1237, 1244, 1250 (1 & 2), 1253, 1259, 1274, 1275, and Auw. N. 30, numbered in this Catalogue 2931, 2949, 2955, 2956, 2961, 2965, 2974, 2991, 2994. h. 1203, numbered by Lord Rosse as one of the group, seems too far remote in R.A. to have been seen on that occasion.
- 2999 h. 1279=II. 156. H. says "F;" h. "vB." The latter preferred, since F might arise from fog or haze.
- 3003 h. 1282. II. 56 and II. 90. Both II. 56 and II. 90 were seen in one sweep, March 1, 1784, at 1^m interval of time (by the same star, 25 Comæ), II. 56 being $1'$ more north, and II. 90 $3'$ more south than the star. This is a case of positive disappearance, for in sweep 334 (h.) the neighbourhood was carefully examined and only one nebula found.
- 3008 I. 23. By ρ Virginis, sw. 174 ; n. $1^\circ 31'$; \therefore P.D. (1830) $77^\circ 18' 29''$. By 34 Virginis in sw. 199 , s. $0^\circ 19'$, whence P.D.= $77^\circ 25' 33''$, mean $77^\circ 22'$. Auwers makes it $77^\circ 16'$. This nebula is placed in the 2nd class by M. D'Arrest as seen with the Leipzig refractor. In this Catalogue it is set down from a mean of two observations, as "pretty bright."
- 3011 h. 1289=II. 212=II. 750. The two nebulae so designated were not observed by H. in one sweep, and are, no doubt, identical.
- 3013 h. 1290=II. 122=II. 174. These two nebulae of the 2nd class were also not observed by H. in the same sweep, and are presumed to be identical, as the places agree.
- 3021 h. 1294=M. 49. Discovered by Oriani in 1771.
- 3026 h. 1295=II. 117=II. 629. The same remark applies as in the notes on Nos. 3011, 3013.
- 3029 II. 116. Not seen by D'Arrest.
- 3043 h. 1307=I. 83. Not found by Lord Rosse when once looked for. There can be no doubt, however, of its existence in or near this place.
- 3060 1196, *a*=R. nova. See note on No. 2884.

No.

- 3075 h. 1329=I. 31=I. 38. H. describes I. 31 as "between two bright stars." The places differ 15' in P.D.; h. describes I. 38 (the place well agreeing with that of H.) in one observation as having a large star *f*, and in two others as having a star 9m, *p*; that is, in effect, as lying between two bright stars. N.B. The star used for I. 31 is 31 *d* 1 Virginis, and for I. 38, 32 *d* 2 Virginis. The declination of 31 *d* 1 is 30' wrong in A.S.C. (No. 1469). In B.A.C. it is right. The P.D.'s of the two nebulae of H. differ, as already remarked, by 15'. The R.A.'s agree. They must be identical with a mistake of 15' in I. 31. D'Arrest says he is sure there are not two nebulae here.
- 3078 III. 26. Place as per C.H., 12^h 25^m 32^s, 68° 32' for 1830; as per Auwers, 12^h 25^m 40^s, 68° 47' (see List of Errata). The correction of the place in P.T. is not, properly speaking, an erratum, but the substitution of a good observation for a bad one. In the obs. sw. 177 (H.), where 20 Comæ was used as the determining star, the place is given only by description. In a sweep long subsequent (sw. 944) it was compared with 26 Comæ in the regular form of observation, and this is of course to be preferred. Auwers's place is deduced from the earlier, and that of C.H. from the later observation, rejecting the other.
- 3079 h. 1322=8 Canum. This very remarkable object occurs among the list of those observed by Lord Rosse in his paper in P.T. 1861, but without a word of remark or description; and it does *not* occur among his list of nebulosities looked for but not perceived. Surely it might be inferred from this that the nebulosity surrounding the star *was* seen, or its absence would have been noticed, as in the instance of 55 Andromedæ. Yet Mr. Lassell saw no nebulosity about 8 Canum.
- 3097 h. 1348=M. 89. Lord Rosse has h. 1343 and 1348, and in his account of them says, "two others, about 20' s. of 1348;" one of these must have been h. 1343, and the other h. 1349.
- 3103 h. 1353=I. 119. This nebula was barely perceptible, with straining the attention, by M. D'Arrest with the 4½-inch Leipzig refractor. It is described in this Catalogue as "considerably bright" by two observations.
- 3108 } h. 1358, 1359, 1363=IV. 8, 9. The obs. of 1363 in my Catalogue of 1833, in
3109 } which the R.A. is uncertain, undoubtedly refers to the same very remarkable double nebula, IV. 8, 9. D'Arrest is sure that there is no other double nebula in this neighbourhood.
- 3111 M. 90. The place is from two observations by W.H., as also the description.
- 3127 h. 1374=I. 273. The descriptions of H. differ so much that it is not impossible there may be another bright nebula near this place.
- 3138 h. 1379=II. 577. Two diagrams by h. in sweeps 141, 143, agreeing, represent this nebula as making a considerably acute-angled, nearly isosceles triangle with

No.

two following stars. H. says, "Between two Bright stars, making a triangle with them." No one now, looking at those diagrams, would call the situation of the nebula between the stars. A suspicion of proper motion arises in such a case.

3148 h. 1384=II. 148. In my Catalogue of 1833 this nebula is identified with II. 20, and in the Register Sheets (H. 320), under the head of II. 148, there is a memorandum, "Probably the same as II. 20 (H. 47)." But on examining all the observations of both nebulae, I arrive at the conclusion that they are different, II. 20 being nearly 2^m later in R.A.

3170 h. 1401. Query if not =II. 38, with one degree mistaken in P.D.

3174 See note on 3148, above.

3177 } h. 1406, 1407=II. 794 (1 & 2), III. 778; h. 1428, 1435=II. 795, 796. Auwers
 3179 } remarks, and justly, on the great apparent discordance of the observations of h.
 3206 } and his places of II. 794, 5, 6, and those of W.H. The fact is that the places
 3216 } of these in the P.T. all rest on comparisons with ϵ Ursæ in sweeps 921 and
 3224 } 1001 (H.); and the observation of that star has been erroneous or mistaken in
 sw. 921 by about 11' in P.D., as appears from an obs. of 73 Ursæ in the same
 sweep. The nebulae affected by this error are those here enumerated, and it
 requires very careful consideration to disentangle all the observations of each
 nebula by both stars, and to decide on their identities. My final conclusions
 are,—1st, that in these sweeps two distinct nebulae, II. 794, 1 and II. 794, 2,
 were observed, and confounded together under one number (=H. 2079 register).
 These are my h. 1406, 1407. 2ndly, that h. 1407 and III. 778, II. 795, 796
 are correctly determined in sw. 1001 (H.). 3rdly, that in sw. 921 (H.) the
 nebula set down as II. 794 was not the same as that called II. 794 in the
 reduction of sw. 1001; *i. e.* that it was in fact h. 1406, and that in this obser-
 vation there is also an error of 6' in P.D., or that, if not, there must be still
 another nebula in P.D. 33° 54' (1860). Finally, that the place of III. 778
 given in Phil. Tr., which is affected by the same *general* cause of error, requires
 a correction of +9' in P.D.

3180 h. 1405=III. 44. This is the companion of M. 60, and is placed by M. D'Arrest in the first class, even with the 4½-inch Leipzig refractor. Perhaps the very superior light of M. 60 may have led both H. and h. to under-estimate that of its, anyhow, much fainter companion.

3189 } h. 1414, 1415=I. 176, 177. These two, according to Lord Rosse, are connected
 3190 } by faint nebulosity.

3206 III. 778. See note on 3174.

3214 h. 1426=II. 181. Auwers points out a discordance of 19' in P.D. between my observation and that of II. 181. This is owing mainly, however, to a misprint in Phil. Trans. (See List of Errata.)

No.

- 3216 }
 3224 } II. 795, 796. See note on 3174.
- 3228 I. 8=III. 6. The later of these nebulae is expressly stated in the register (H. 38) to be of the 1st class, though set down (it does not appear why) in the 3rd.
- 3254 h. 1452=I. 41. The case of this nebula is a very odd one. H. has two observations of it. One on April 5, 1784, where it is described as a "L; B; r neb; sbM; iR Fig; Class I." Another on March 3, 1789, calls it "pB; cL; i Fig; er. Many of the st. visible." So that it may be called a cluster. Both the places of H. and that of h. agree so well, that the object in all must have been the same. Here seems evidence of change.
- 3256 h. 1453=II. 73. Contradictory descriptions, and possibly two nebulae differing 1^m in R.A.
- 3311 h. 1480=I. 141. Query if not changed. h.'s observations are positive as to the clearness of the sky. But query as to the state of the speculum.
- 3319 h. 1485=II. 384. Not seen by Lord Rosse in two observations (hazy).
- 3337 h. 1497=I. 68; II. 299; h. 1511=I. 69; h. 1536=II. 301; h. 1574=III. 382.
- 3338 }
 3358 } Auwers finds 5' Δ.P.D. between H. I. 68 and h. 1497. His place is from P.T.
 3420 } 53 Virginis n. 1° 4', whereas C.H. in her reductions uses n. 1° 11', and my
 3483 } observations of this and the other nebulae in this list justify the departure. I
 subjoin her note on this nebula (in zone 103° C.H.):—
- "I. 68, I. 69, III. 282 are each 7' more north than they are given in the "printed Catalogue. The disagreement is the result of the recalculation, and "is probably owing to my attempting more accuracy in valuing the 'numbers "to a degree,' &c. &c." (*i. e.* in the index reductions of the Polar distance readings which were parts of an arbitrary scale). And in the next zone (104° C.H.) occurs,
- "II. 299 and II. 301 require the same memorandum." In point of fact, comparing my own observations with those reduced by M. Auwers, the differences, as stated by him, run thus:
- | | | |
|--------------------|-------------|-----|
| I. 68 | Δ.P.D. H—h= | +5' |
| I. 69 | | +7' |
| III. 282 | | +7' |
| II. 299 | | — |
| II. 301 | | +6' |
- so that in each case, where I have observed the object, the alteration is justified. This is only one out of the innumerable instances of painstaking and laborious scrutiny bestowed by her upon these reductions which have occurred to me in the collation of her zone catalogue with the original observations and with my own results.
- 3356 h. 1509=I. 143. Auwers places this nebula 1° 13' too much to the south in consequence of an erratum in P.T. (see List of Errata).

No.

- 3358 See note on 3337.
- 3363 V. 3. Auwers makes the R.A. of this neb. for 1830 $13^{\text{h}} 2^{\text{m}} 31^{\text{s}}$, which is 10^{m} too great. The P.T., which in this instance is correct, makes it follow 75 Leonis $1^{\text{h}} 44^{\text{m}}$.
- 3393 h. 1527. This is not impossibly III. 937, but as both R.A.'s and P.D.'s differ very much, they *may* be different, and are therefore separately stated.
- 3415 h. 1535. Not seen by Lord Rosse in one observation; clouds passing. h. has two observations, both agreeing well.
- 3420 See note on No. 3337.
- 3421 II. 185. Auwers, misled by an error in P.T. (see List of Errata), makes the R.A. of this neb. too small by 10^{m} .
- 3426 Auw. N. 31. Not visible in the Königsberg Heliometer.
- 3483 See note on No. 3337.
- 3506 II. 22. P.D. extremely doubtful.
- 3512 II. 826. Place re-reduced by the star used by H. and A.S.C.
- 3527 h. 1597=II. 314. Auwers makes Δ .R.A. H.—h.= $+107^{\text{s}}$, and remarks that there is perhaps some error in P.T. This is the case (see List of Errata), and with the correction there indicated the agreement is satisfactory.
- 3550 D'Arr. 94. D'Arrest says "not found again, Feb. 19, 1863. Sky perfectly clear. Perhaps a comet."
- 3588 h. 1633=III. 926. H. says it is sp a considerable star. h. has "a $*9^{\text{m}}$ with a very dilute nebulous atmosphere." Has the star or the nebula moved?
- 3650 III. 946. Auwers makes the declination $+89^{\circ} 17'$, a misprint for $+80^{\circ} 17'$.
- 3662 h. 1674=I. 255. Evidently ill seen by h. The description of H. preferred.
- 3664 } h. 1676, 1679=III. 422, 423. Auwers makes the P.D. $12'$ too great by reason
3668 } of an erratum in P.T. (see List of Errata).
- 3728 h. 1720=III. 666. Auwers finding Δ .R.A. H.—h.= $+52^{\text{s}}$, supposes a mistake of 1^{m} . Examined sweep 146 (h.), and found all clearly written and right reduced.
- 3750 } h. 1734, 1735=II. 309, 310. H. says the second is the larger, h. the smaller of
3751 } the two.
- 3760
- 3762
- 3763 h. 1744=M. 101, and its attendants in more or less intimate nebulous connexion.
- 3764 Of those in Lord Rosse's woodcut, P.T. 1861, p. 729, N, the principal nucleus,
- 3766 is No. 3770=h. 1774; n_1 =No. 3774=1744, i; n_2 No. 3773=1744, h. The
- 3767 others are not lettered, and are made out from the joint evidence of this dia-
- 3770 gram and the measures of position and distance of the stars compared with the
- 3771 copper plate, fig. 35.—1744, α is not improbably=III. 787.
- 3773
- 3774

No.

- 3820 h. 1763=III. 804=III. 835. The identity of these nebulae rests on a memorandum in MS. in my copy of Ph. Tr., supported by the reductions of all the obs. by C.H. in 3 sweeps, each with two determining stars. Auwers makes them differ by 14' in P.D.
- 3836 III. 551. Place concluded from h. 1772=III. 552 from H.'s description, viz. that it precedes that nebula by 3' or 4' (3' 30'')=14^s of time.
- 3844 h. 1777=III. 347. Auwers makes Δ .P.D.=−59', but observes that there must be some misprint. Examining all, I find that such is the case (see List of Errata), which recognized, shows that 1° has been mistaken, and the identity is therefore proved.
- 3846 h. 1779=I. 144. Auwers makes the P.D. (1830)=86° 30', and H.—h.=1° 14'. The cause of the discordance is a misprint in P.T. (see List of Errata), in consequence of which the nebula is 1° 13' north of its printed place.
- 3858 } h. 1789, 1788, 1791=III. 416, 417. Lord Rosse says that of these three only
3859 } two were found. The obs. in sw. 28 re-examined—1789 and 1791 were both
3860 } observed. Moreover, in sw. 337, III. 417=h. 1791 and h. 1788 were both
observed, and 1791 is expressly stated to have been the sf of two seen in moonlight. Now the np of these could not be h. 1789, which is eF and not north, but south preceding, whereas h. 1788 by its place in sw. 338 is np. All three, therefore, really *existed* at the date of these observations. It was h. 1789 (eF) which escaped Lord Rosse's notice, though looked for with greater instrumental power. Perhaps it may have changed.
- 3863 III. 135. Auwers's P.D. for 1830 is 63° 0'. C.H. reduced to 1830 gives 62° 50' 20''. Auwers has used (P.T.) 1° 5' n. of *d*, 12 Bootis; C.H. 1° 16' n. of the same *. C.H. is to be preferred on every account to P.T. Her Δ .P.D.'s are grounded on a most complete and searching re-examination and recomputation (*according to the then existing star catalogues*) of all the data (in the earlier sweeps most obscure—*foliis sibyllinis obscuriora*) for determining the degrees and minutes of P.D. from the index numbers. In almost every case I find her corrections (or rather interpretations) to be justified; and I have no doubt that in this particular instance such will prove the case, though *here* I confess myself, after consulting the original sweep, unable to perceive the reason for the deviation.
- 3888 III. 319. Auwers, following P.T., which places the nebula 2° 26' north of β Ursæ min., makes the P.D. 1830 =12° 46', and so it stands in the Register sheet (H. 864). But it should be 2° 26' *south*. So C.H. has used it, and so it proves to be on reference to the original sweep, sw. 391 (H.), giving for the P.D. 17° 36' 12''.
- 3920 h. 1832=II. 695. Not seen by Lord Rosse in one observation. See note on No. 132.
- 3922 h. 3573= Δ . 342. In Auwers's list of Lacaille's nebulae, he sets down for the

No.

declination of this $-55^{\circ} 58'.8$. For $58'.8$ read $48'.8$, if it be the same object, but of that some doubt remains.

- 3967 VI. 8. Auwers, using χ Virginis, the determining star in P.T., places this cluster in R.A. $14^{\text{h}} 53^{\text{m}} 37^{\text{s}}$ (1830), $99^{\circ} 55'$ P.D. This, however, is declared by a subsequent MS. note to be a mistake for Mayer's 577 zod. star, whence the place in this Catalogue is accordingly derived. But this star, too, must have been mistaken, and on consulting the original sweep (sw. 209, H.) I find no star in the sweep whose identity can be satisfactorily ascertained. All that can be certainly affirmed is that, within a degree one way or the other in P.D., and from 5 to 10 minutes of time in R.A. of the place set down, there exists a fine cluster of the 6th class which should be looked for. Fortunately it is the only nebula observed in the sweep, a very short one.
- 3977 h. 1866=I. 184. Some suspicion of variability, inasmuch as one description calls it R, another E, and another mE, besides other indications in respect of brightness.
- 3998 III. 373. C.H., by three distinct observations in three different sweeps (400, 730, 917, H.) from the same determining star 11 Libræ (s. $0^{\circ} 13'$, s. $0^{\circ} 14'$, and s. $0^{\circ} 15'$), deduces a P.D., which reduced to 1830= $91^{\circ} 49' 39''$. Auwers, using the same star, s. $0^{\circ} 12'$ as *per* P.T., places it in P.D. $91^{\circ} 17'$, which, however, is probably a misprint for $91^{\circ} 47'$. Two of H.'s observations place the small star *south*, and one *north* of the nebula.
- 3999 h. 1881=II. 576. The binuclear character verified by R, who says that it is a close double nebula.
- 4016 h. 1892=III. 131. Query if not variable in brightness. H. in two observations calls it F and cB; h., in two others, vF and eF.
- 4025) II. 756=h. 1898?. In the two observations by H. of II. 756 it is described as
4029) cF; pL; iF; r;
pB; s; E;
- and no mention is made of a double star near it, so that though the places agree within the *possible* limits of discordance, they are most probably two distinct nebulae.
- 4043) 1901, a. Two of six seen by Lord Rosse. The others must have been h. 1901,
4044) h. 1902, II. 541 and III. 511.
- 4048) III. 886, 887. Auwers has made an error of $-12'$ in the declination, or $+12'$ in
4049) the P.D. of this double nebula as determined from P.T. ($20'$ n. of 7 Serpentis). The P.D. here set down is that correctly reduced, C.H. having on her part committed an error of $+2'$ in P.D.
- 4051 h. 1905=II. 751. In Auwers's declination, for $+20^{\circ} 44'$ read $+20^{\circ} 14'$, an evident misprint.
- 4065 II. 818. Owing to an erroneous designation of the determining star in P.T. (see

No.

- List of Errata), Auwers has given the place of this nebula (1830) as R.A. $14^{\text{h}} 41^{\text{m}} 3^{\text{s}}$; Decl. $+60^{\circ} 5'$.
- 4124 h. 1934, &c. In Lord Rosse's diagram of the group h. 1934, A, the most conspicuous, would naturally be selected as identical with that nebula, but in that case II. 766 would not be included in the group. On the other hand, if B be taken for h. 1934, the identifications will stand as follows:—A=No. 4131=II. 766; B=No. 4128=h. 1934; C=No. 4127=1934, *b*; D=No. 4124=1934, *a*. This, however, supposes an error of 45^{s} of R.A. in H.'s place of II. 766, which is not probable, while on the other hand it is difficult to account otherwise for its not having been noticed at all. All things considered, I have thought it best to enter A as a new nebula, No. 4133=1934, *c*, leaving 766 untouched.
- 4167 h. 1948=III. 74. Not seen by Lord Rosse, once looked for (see note on No. 132).
- 4173 h. 3624=M. 80. This is Pogson's globular cluster, with a variable star in the centre, for whose most singular history see the Monthly Notices of the R. Ast. Soc. xxi. pp. 32, 33, by Mr. Pogson. Mr. P. in that statement says that Sir J. Herschel (among others mentioned) had described it as either "cometary" or "nebulous." This is incorrect. In both my observations of this object it stands described as a globular cluster, *all completely resolved into stars*. (See C.G.H. h. 3624.)
- 4234 h. 1970= Σ . 5. D'Arrest calls this planetary nebula *blue*. The place used is a mean of his observations, that of h. (Catal. of 1833) being only Struve's roughly brought up. M. D'Arrest makes the diameter = $14''\cdot6$.
- 4247 III. 727. The comparison of the place here set down with that of Auwers is curious for the great number of perfectly accidental errors which have heaped themselves together. The place (C.H.) is rightly reduced by her from σ Herculis, $f 16^{\text{m}} 11^{\text{s}}$; $n 0' 14''$, which is that given in P.T., and which, reduced to 1830, gives for the R.A. $16^{\text{h}} 44^{\text{m}} 46^{\text{s}}\cdot8$ and for the P.D. $47^{\circ} 58' 16''$, differing $+8^{\text{s}}\cdot8$ and $+11''$ from the exact result. In M. Auwers's catalogue it is entered thus: III. 127; R.A. $16^{\text{h}} 14^{\text{m}} 47^{\text{s}}$; Decl. $+43^{\circ} 1'$ (corresponding to P.D. $46^{\circ} 59'$). That is to say, there is a misprint *in each of the three particulars*. This is not to be taken as a specimen of M. Auwers's work, which is an admirable example of painstaking devotion, and far beyond any eulogy in my power to offer. But it is a striking instance of the way in which, in the great run of chances, unlucky coincidences will happen.
- 4259 h. 1974. Doubtful whether a nebula or a very faint double or triple star.
- 4294 M. 92 (= also Lalande No. 31544). Not observed by h., but 8 times by H. Place from Wollaston's catalogue, which is almost identical with Auwers (Δ .R.A.= $0^{\text{s}}\cdot1$, Δ .P.D.= $0' 3''$).
- 4302 h. 1981=h. 3686=IV. 11. The annular form only perceived in the southern

No.

observations. Both H. and h., in their northern observations, describe it as of equable light throughout. It appears from Lord Rosse's observations that the annular form is much more common among these "planetary" nebulæ than H. or h. had any idea of.

4364 h. 3723=II. 200. On a ground astonishingly rich.

4368 V. 13. P.D. by Auwers = $113^{\circ} 36'$ (1830), owing to an error in P.T. (see List of Errata).

4372 h. 3726= Δ . 473. There is a singular statement respecting this cluster by Cacciatore in No. 113 of the *Astronomische Nachrichten*. He observed it as a nebula, he says, on the 19th of March, 1826 (of course, therefore, Dunlop has the priority in point of date). But *where* he saw it Lacaille, he says, noted his star 1483 (Cælum Australe). Also, Piazzzi in 1794 and 1801 in the same place saw only a star. Cacciatore in 1809 and 1810 observed the same star, but saw no nebula, only a star 9m following it (P. xvii. 341, 346). In looking for the comet of 1826, however, "fui colpito," he says, "da questa bella nebulosa." Unfortunately for this curious history, the place of Piazzzi's star referred to (and which he identifies with 1483 C.A.) differs by no less than 18' in P.D. from that of the nebula in question, which was therefore out of the field of view, both of his own and of Piazzzi's telescope, when observing the star.

4390 h. 2000. Σ . 6. Omitted by Auwers from his catalogue of new nebulæ, which contains many far less remarkable. Diameter, according to D'Arrest, = $7''\cdot05$. Bessel's place = h. + $0^{\circ}\cdot8$, - $0' 22''$.

4397 h. 2004=M. 24. H.'s two observations hardly consist with this description, and their deviation in R.A. of nearly 4^m from Messier's place makes it very doubtful whether he really saw this object.

4411 M. 69. Piazzzi, in a note on xviii. 122 of his catalogue, says that both M. 69 and M. 70 are 1° more to the south. But he is wrong.

4415 Auwers, N. 40. This is the nebula discovered by Tuttle on Sept. 1, 1859, and it would appear to be variable, for M. D'Arrest says (in a letter of May 8, 1863), "La nébuleuse de M. Tuttle (*Astron. Nachr.* No. 1337. p. 272) était, le 24 Sept. 1862, si brillante et si remarquable dans le chercheur (*grandis et præclara, ovalis, 2' longa, 80'' lata*), que je suis persuadé qu'elle n'a pas été telle du temps de Messier et de votre père, et de vos propres observations. Voici la position que j'ai obtenue. 1861.0 R.A. $275^{\circ} 55'\cdot6$, N.P.D. = $15^{\circ} 30'\cdot1$." The place given in the present Catalogue is that of M. Auwers, and differs somewhat, though not considerably, from this determination.

4428 M. 70. See the note on No. 4411.

4462 III. 742. This agrees too well with M. D'Arrest's place of his No. 113 not to be the same. His description is F; S; R; *10p $12'\cdot6$, s $2' 30''$.

4473 Auwers, N. 44. This is the nebula discovered by Mr. Hind on March 30, 1845.

No.

- It was observed in May 1852 as a nebula of the first class; subsequently as "pretty faint and diluted." M. Auwers found it "surprisingly faint," and of the 2nd class at the highest.
- 4487 h. 2037=III. 743. This was seen as a planetary nebula in the twilight by M. D'Arrest with the $4\frac{1}{2}$ -inch refractor, and can therefore hardly be ranked so low as Class III.
- 4536 h. 2062=III. 144. Not seen by Lord Rosse; once looked for. (See note on No. 132, &c.)
- 4570 h. 2073. Not seen by Lord Rosse; twice looked for. h. has three observations agreeing well. The object is an equivocal one.
- 4585 } h. 2081=I. 103. According to an observation of Olbers, cited by Auwers, this
4586 } is identical with No. 4585=I. 103, the place of the latter nebula, as assigned by H., being 20' wrong in P.D. This had escaped my notice until the nebulae in this Catalogue had been finally numbered and much other work accumulated on them; and it was considered better to let No. 4585 stand, though erroneous, than to hazard confusion by striking it out and altering all the subsequent numbering.
- 4618 h. 2093. In conformity with Mr. Mason's remarks on my observations of this nebula, and with his elaborate and excellent monograph of the great nebulous system of which it forms a part, I have diminished the P.D. in my Catalogue of 1833 by 1° . It is evident that the index reading must have been mistaken, 1° for 0° . Sweep 8 examined; the writing is clear and the reduction correct, but the conclusion from Mr. Mason's observations is irresistible.
- 4628 h. 2098=IV. 1. According to Lassell this is annular, an elliptic ring with a star in the centre.
- 4654 h. 2113. Not seen by Lord Rosse; twice looked for. Examined sw. 86 (h.), in which it was observed. All found apparently correct, the observation clearly written and right reduced: and it is added, "the double star" (h. 934 in my "3rd series of observations, &c. &c.," Mem. Ast. Soc. vol. iii.) "is a good guide." A diagram accompanying the observations, by indicating lines points out the relative situation of the double star and nebula.
- 4710 h. 2133. Not seen by Lord Rosse in four observations.
- 4714 h. 3897. Not found by Mr. Lassell within 30' all round the place.
- 4723 h. 2137=III. 920. Not seen by Lord Rosse in one observation.
- 4756 h. 2148. Not seen by Lord Rosse in three observations. In one a cloud passing.
- 4775 h. 2156=III. 932. H. says, "just sf a S* to which it seems almost to be attached, but is free from it." h. says, "has a * 13m at a distance from the edge = 1 diameter by diagram." Sw. 274 (h.). This sweep re-examined. The diagram makes the star north of the nebula. The description says, "Diagram certainly right."
- 4816 2172, a. In this group Lord Rosse has given only measures of relative position,

No.

- and none of distance; so that it is impossible to assign specific places to the individuals of which it consists. He speaks of five *near* to h. 2172. The diagram exhibits only four. One may possibly be III. 166.
- 4848 2184, α . In Lord Rosse's diagram of the group to which this belongs, α is h. 2183 = No. 4845; β = D'Arr. 117 = No. 4844; γ = h. 2184 = III. 217 = No. 4846; δ = D'Arr. 118 = No. 4847. That marked as 2184, α is not lettered in the diagram, and is "nova."
- 4892 h. 2205 = I. 55. Placed in the second class only by M. D'Arrest with the $4\frac{1}{2}$ -inch Leipzig refractor. In this Catalogue it is set down as only "pretty Bright," from a mean of seven observations.
- 4894 h. 3971 = h. 3972. These are assuredly identical; but the minute of R.A. being doubtful, that of the earlier 3971 is preferred. The mean of the seconds and the Polar distances is taken, blending the two, and also the descriptions.
- 4922 h. 2223 = III. 222. Three times called by h. "pretty Bright," and three times by h. and H., eF; vF; eF. Is this a case of variability?
- 4933 h. 2228 = h. 3982 = I. 104. Placed in the second class by M. D'Arrest. With this the present Catalogue agrees; making it "pretty Faint" by a mean of three observations.
- 4941 D'Arr. Not included by M. D'Arrest in his final list; but there are four observations of it recorded in his "Resultate," all agreeing well.
- 4964 h. 2241 = IV. 18. According to Mr. Lassell this superb "planetary nebula" is *bi-annular*, consisting of a nucleus and *two oval rings*.
- 4966 h. 2242 = III. 226. Called by h. in four observations, pB; pB; pB; pB, and in two by H. eF; vF.
- 4980 h. 2250 = III. 213. Not seen by Lord Rosse in 4 observations. In my observations of sweep 103, a very short sweep, using the quadrant instead of the index arc, and with no good zero star, both R.A. and P.D. may be a good deal wrong. My place, however, agrees pretty well with that of H. (Δ .R.A. = 5^s , Δ .P.D. = $4'$), and the existence of a nebula as described, *hereabouts*, is certain, but it should be looked for within somewhat wider limits.
- 4998 h. 2261 = I. 110. H. has two observations in which this nebula is called cB; h. has one where it is called eF; adding "sky quite clear."
- 5003 } h. 2263 = II. 208. These can hardly be the same. The R.A.'s differ by nearly
5004 } 2^m and the P.D.'s by $6'$. The descriptions also disagree. 255° , the position of the star 14m in h. 2263, is not np but sp, and the estimates of their magnitudes differ materially.
- 5015 h. 2271 = III. 854. A very problematic object, and in which there is great difficulty in making out its nature. Stars and nebula oddly mixed.
- 5020 } h. 2274 = II. 230; 2274, α ; h. 2275 = II. 231. In Lord Rosse's diagram of this
5021 } group, α = h. 2274; β = h. 2275; γ = nova = 2274, α . h. sweep 91 makes II.
5022 } 230 the np of two, and II. 231 "to have II. 230, 45° sp." This is contradicted

No.

by the diagram. There is some confusion among the observations as to whether the two nebulae II. 230, 231 really lie np or sp from each other, and it might be suspected that the P.D.'s had been read crossways, the R.A.'s being rightly set down; but Lord Rosse's diagram and measures decide the point in favour of the relative situation being here correctly given.

5051 h. 2302. Not seen by Lord Rosse in two observations. Examined the original observation, all clear and apparently correct. The nebula certainly exists in or very near the place here set down.

5061 2849, *a*. A nebula mentioned by M. D'Arrest, but not included in his MS. list of well-determined nebulae. Should, however, be looked for.

References to Figures of Nebulae in various works.

In the following list of figured nebulae, the first column contains the current number of the nebula or cluster in the present Catalogue; the second the number attached to it in my Catalogues in P.T. 1833 and C.G.H.; or if not found in either of these, the class and number in my Father's Catalogues or other sufficient designation. The third contains an abbreviated reference to the publication in which the figure will be found, viz.—

P.T. 33. The volume of the Philosophical Transactions of the Royal Society for A.D. 1833.

P.T. 44. Ditto, Ditto, for 1844 }
 P.T. 50. Ditto, Ditto, for 1850 } Lord Rosse's papers.
 P.T. 61. Ditto, Ditto, for 1861 }

C.G.H. Results of astronomical observations at the Cape of Good Hope by J.F.W.H.

R. di. The woodcut diagrams in Lord Rosse's paper, Philosophical Transactions, 1861; such only being referred to as express some distinct peculiarity not elsewhere figured.

B.A.A. Professor Bond's Memoirs in vol. iii. N.S. of the Transactions of the American Academy of Arts and Sciences.

M.A.A. Mr. Mason's Memoirs in vol. vii. of the Transactions of the American Academy.

D'Arr. M. D'Arrest's Inaugural dissertation and description of the Copenhagen Equatoreal, 1861.

Lam. Dr. Lamont's "Oeffentliche Vorlesung über die Nebelflecken." München 1837.

Lass. Mr. Lassell's Memoirs in vol. xxiii. of the Transactions of the Royal Astronomical Society.

Column 4 contains the number of the Plate in the volume referred to where the figure will be found, and column 5 the number of the figure in that Plate.

The figures annexed to Mr. Dunlop's catalogue are not included, as for the main part they offer no resemblance to the objects figured (when identifiable), and would serve only

to mislead. The same remark applies to most of the older figures of nebulae scattered through the volumes of the Histoire de l'Académie Française, and other collections. Of the older figures of the nebula in Orion, however, for curiosity's sake, a list is sub-joined. The figures accompanying my Father's memoir in Philosophical Transactions, 1811, are also omitted. They do not profess to be resemblances, and are given rather as types of certain classes of objects into which he there considers the nebulae to be distributable. At least they are made from very rude diagrams.

References to published figures of Nebulae.

No. in Catalogue.	h. &c.	Work cited.	No. of plate.	No. of fig.	No. in Catalogue.	h. &c.	Work cited.	No. of plate.	No. of fig.
27	2315	C.G.H.	iv.	8	1157	357	P.T. 33	viii.	81
31	15	P.T. 61	xxv.	1			P.T. 44	xix.	81
52	2322	C.G.H.	iii.	1			R. di.		
67	2327	C.G.H.	vi.	19			D'Arr.	ii.	4
105	44	B.A.A.	opp. p. 86		1163	2864	Lass.	ii.	1
106	45	B.A.A.	Ditto.		1164	2865	C.G.H.	iv.	7
116	50	B.A.A.	Ditto.		1165	2866	C.G.H.	vi.	20
117	51	B.A.A.	Ditto.		1168	2867	C.G.H.	vi.	20
138	61	P.T. 33	vi.	52	1171	Δ. 136	C.G.H.	vi.	20
169	2359	C.G.H.	v.	10	1171	2868	C.G.H.	vi.	20
187	2370	C.G.H.	iv.	6	1174	2872	C.G.H.	iv.	7
298	112	P.T. 33	v.	38	1175	2869	C.G.H.	vi.	20
303	116	R. di.			1176	2875	C.G.H.	iv.	7
352	131	P.T. 50	xxxvi.	5	1177	2876	C.G.H.	iv.	7
		P.T. 61	xxvi.	10	1179	360	C.G.H.	viii.	1
		R. di.					B.A.A.	opp. p. 96	
372	142	R. di.					Lass.	i.	1
400	151	P.T. 33	vi.	58			*	see note	
412	156	P.T. 61	xxv.	2	1180	V. 30	C.G.H.	ii.	3
527	218	P.T. 33	ii.	28	1183	361	C.G.H.	ii.	3
		R. di.					P.T. 50	xxxviii.	6
544	223	D'Arr.	ii.	7			Lass.	ii.	3
560	232	P.T. 61	xxv.	3	1185	M. 43	C.G.H.	viii.	1
567	2487	C.G.H.	vi.	14			B.A.A.	opp. p. 96	
572	241	P.T. 61	xxv.	4			Lass.	i.	1
575	242	P.T. 33	vi.	56			*	see note	
		P.T. 61	xxv.	5	1225	365	D'Arr.	ii.	2
600	262	P.T. 61	xxv.	6			Lass.	ii.	2
705	2534	C.G.H.	vi.	7	1226	iv. 24	D'Arr.	ii.	10
731	2552	C.G.H.	iv.	1	1233	2910	C.G.H.	iii.	5
810	311	P.T. 33	ii.	31	1235	2913	C.G.H.	iii.	5
		P.T. 61	xxv.	17	1238	2916	C.G.H.	iii.	5
822	2620	C.G.H.	v.	11	1243	2918	C.G.H.	iii.	5
823	2621	C.G.H.	v.	11	1248	2923	C.G.H.	iv.	9
826	2618	D'Arr.	ii.	9	1249	2925	C.G.H.	iv.	9
		Lass.	ii.	4	1258	2935	C.G.H.	iv.	9
853	315	P.T. 61	xxv.	8	1259	2933	C.G.H.	iv.	9
888	327	P.T. 61	xxv.	9	1260	2936	C.G.H.	iv.	9
979	2709	C.G.H.	iii.	3	1265	2938	C.G.H.	iv.	9
980	2710	C.G.H.	iii.	3	1266	2939	C.G.H.	iv.	9
981	2711	C.G.H.	iii.	3	1267	368	P.T. 33	iv.	36
987	2716	C.G.H.	iii.	3			R. di.		
1057	2775	C.G.H.	vi.	1	1269	2941	C.G.H.	ii.	4
1082	2802	C.G.H.	iii.	6	1276	2948	C.G.H.	iii.	4
1084	2803	C.G.H.	iii.	6	1277	2949	C.G.H.	iii.	4
1085	2804	C.G.H.	iii.	6	1278	2950	C.G.H.	iii.	4
1086	2805	C.G.H.	iii.	6	1279	2951	C.G.H.	iii.	4
1089	2808	C.G.H.	iii.	6	1281	2952	C.G.H.	iii.	4
1090	2810	C.G.H.	iii.	6	1282	2953	C.G.H.	iii.	4
1135	2840	C.G.H.	iii.	2	1283	2954	C.G.H.	iii.	4
1137	355	P.T. 33	v.	49	1419	390	R. di.		
1140	2842	C.G.H.	iii.	2	1425	393	P.T. 61	xxvii.	11
1141	2843	C.G.H.	iii.	2	1437	399	P.T. 33	vi.	64
1142	2844	C.G.H.	iii.	2			P.T. 50	xxxvii.	10
1143	2845	C.G.H.	iii.	2			Lass.	ii.	8
1156	2859	C.G.H.	iv.	7					

TABLE (continued).

No. in Catalogue.	h. &c.	Work cited.	No. of plate.	No. of fig.		No. in Catalogue.	h. &c.	Work cited.	No. of plate.	No. of fig.
1467	415	P.T. 33	viii.	91		2841	1175	P.T. 33	vi.	55
1477	421	P.T. 61	xxvii.	12		2870	1196	P.T. 61	xxvii.	21
1511	3075	C.G.H.	iv.	4		2878	1202	P.T. 33	vii.	69
1519	444	P.T. 33	vii.	72		2884	1196, <i>a</i>	P.T. 61	xxvii.	21
		P.T. 50	xxxvii.	6		2910	1225	P.T. 33	vi.	57
1520	445	Lass.	ii.	9		2950	1245	P.T. 61	xxvii.	22
1521						2958	1252	P.T. 33	vii.	68
1532	450	P.T. 50	xxxviii.	15		2962	1252	P.T. 33	vii.	68
		Lass.	ii.	6		2972	1258	R. di.		
1565	{ 464 }	P.T. 50	xxxviii.	12		3041	1306	P.T. 61	xxvii.	23
	{ 3093 }	Lass.	ii.	5		3042	1308	P.T. 61	xxvii.	23
1567	3095	Lass.	ii.	7		3085	1337	P.T. 33	iv.	37
1677	3131	C.G.H.	vi.	12				P.T. 61	xxviii.	24
1721	536	P.T. 33	vi.	61		3101	1352	P.T. 33	viii.	83
		Lam.	i.	8		3106	1357	P.T. 50	xxxvii.	9
1728	537	P.T. 33	v.	65		3108	1358 1363	P.T. 33	vii.	78
1745	3145	C.G.H.	vi.	12						
1801	3154	C.G.H.	v.	8		3109	{ 1359 1365 }	P.T. 33	vii.	78
1861	604	P.T. 33	vii.	70		3113	1362	P.T. 33	vi.	66
1863			P.T. 50	xxxvi.	3		3132	1376	P.T. 33	vi.
1911	639	P.T. 61	xxvii.	13		3151	1385	P.T. 61	xxviii.	25
2003	3221	C.G.H.	v.	9		3152				
2017	3228	C.G.H.	vi.	9		3165	1397	P.T. 33	vii.	76
		Lass.	ii.	10				P.T. 50	xxxvii.	9
2058	692	P.T. 61	xxvii.	14		3180	1405	P.T. 33	vii.	74
2063	3241	C.G.H.	vi.	2		3182	1408	P.T. 33	vii.	74
2067	3239	C.G.H.	iv.	3		3189	1414	P.T. 33	vii.	75
2102	3248	C.G.H.	vi.	5		3190	1415	P.T. 61	xxviii.	26
		Lass.	ii.	11		3240	1441	P.T. 61	xxviii.	27
2158	731	P.T. 33	v.	40		3249	1451	R. di.		
2197	3295	C.G.H.	ix.	1	<i>n</i> Arguts.	3258	1456	P.T. 33	v.	41
2216	765	P.T. 61	xxvii.	15		3275	3435	C.G.H.	i.	2
2217	766	P.T. 61	xxvii.	15		3278	1466	P.T. 33	viii.	84
2333	3324	C.G.H.	iv.	10		3321	1486	P.T. 33	ii.	27
2336	3325	C.G.H.	iv.	10		3340	1499	P.T. 33	vi.	62
2337	3326	C.G.H.	iv.	10		3356	1509	P.T. 33	vi.	67
2338	3327	C.G.H.	iv.	10		3511	1589	P.T. 61	xxviii.	28
2340	3329	C.G.H.	iv.	10		3525	3501	C.G.H.	iv.	2
2342	3330	C.G.H.	iv.	10		3531	3504	C.G.H.	v.	7
2343	838	P.T. 33	ii.	32		3570	3514	C.G.H.	vi.	1
		P.T. 50	xxxvii.	11		3572	1622	P.T. 33	ii.	25
2373	854	P.T. 33	vi.	53				P.T. 50	xxxv.	1
		P.T. 50	xxxvii.	7				R. di.		
		Lam.	i.	6		3606	3523	C.G.H.	iv.	5
2377	{ 857 }	P.T. 33	vi.	54		3614	1649	P.T. 33	v.	39
	{ 875 }	P.T. 61	xxvi.	16		3615	1650	P.T. 61	xxviii.	29
2378	859	P.T. 33	vi.	51		3651	3541	C.G.H.	vi.	15
2379	858	R. di.				3706	3548	C.G.H.	vi.	10
2445	910	R. di.				3717	1713	P.T. 61	xxviii.	30
2486	{ 934 }	P.T. 33	vii.	79		3750	1734	R. di.		
	{ 3355 }					3766	III. 787	P.T. 61	xxix.	35
2488	{ 936 }	P.T. 33	vii.	79		3770	1744	P.T. 61	xxix.	35
	{ 3356 }					3778	III. 788	P.T. 61	xxix.	35
2559	982	R. di.				3779	III. 789	P.T. 61	xxix.	35
2597	1002	R. di.				4051	1905	P.T. 33	vii.	77
2606	1011	P.T. 61	xxvi.	17		4052				
2652	1041	P.T. 50	xxxvii.	7		4058	1909	P.T. 61	xxviii.	8
2670	1052	P.T. 61	xxvi.	16		4066	3594	C.G.H.	vi.	8
2671	1053	P.T. 61	xxvi.	16		4083	1916	P.T. 33	viii.	87
2680	1061	P.T. 61	xxvii.	19		4087	1917	R. di.		
2733	1092	R. di.				4118	1929	P.T. 33	viii.	89
2756	1111	P.T. 61	xxvii.	20		4125	3610	C.G.H.	vi.	7
2760	1113	P.T. 61	xxvii.	20		4160	1946	P.T. 61	xxviii.	32
2804	1146	P.T. 33	vii.	71		4224	3641	C.G.H.	v.	4
2806	1148	P.T. 33	vi.	59		4229	3644	C.G.H.	v.	6
2807	1149	P.T. 50	xxxvii.	8		4230	1968	P.T. 33	viii.	86
2838	1173	P.T. 50	xxxv.	2		4234	1970	P.T. 61	xxviii.	33
								Lam.	i.	1

TABLE (continued).

No. in Catalogue.	h. &c.	Work cited.	No. of plate.	No. of fig.		No. in Catalogue.	h. &c.	Work cited.	No. of plate.	No. of fig.
4261	3661	C.G.H.	vi.	13	Milky Way.	4572	2075	P.T. 33	v.	47
4284	3675	C.G.H.	vi.	6		4594	2084	P.T. 61	xxviii.	34
4290	3680	C.G.H.	vi.	3		4600	2088	Lam.	i.	5
	3680, 2	C.G.H.	v.	3		4616	2092	P.T. 61	xxx.	36
4302	{ 1891 3686 }	C.G.H.	vi.	4		4618	2093	P.T. 33	iii.	33
4305	3688	C.G.H.	vi.	18				M.A.A.	vii.	1
	3702, 2	C.G.H.	v.	1				P.T. 33	viii.	82
4335	3707	C.G.H.	v.	5				M.A.A.	vii.	1
4342	3713, 2	C.G.H.	v.	2		4627	2099	P.T. 61	xxx.	37
4343	1989	P.T. 33	v.	42		4628	2098	P.T. 33	v.	44
4355	{ 1991 3718 }	P.T. 33	viii.	80			P.T. 50	xxxviii.	14	
		C.G.H.	ii.	2			D'Arr.	ii.	1	
		M.A.A.	iv.	1	4678	2125	Lam.	i.	4	
4361	3722	C.G.H.	i.	1			P.T. 33	viii.	88	
4375	3727	C.G.H.	vi.	16			P.T. 44	xviii.	88	
4395	2002	P.T. 33	ii.	30	4687	{ 2128 3878 }	P.T. 33	viii.	90	
4403	2008	P.T. 33	iv.	35	4729	3908	C.G.H.	iv.	11	
		C.G.H.	ii.	1	4730	3909	C.G.H.	iv.	11	
		Lam.	i.	10	4731	3910	C.G.H.	iv.	11	
		M.A.A.	vi.	1	4733	3911	C.G.H.	iv.	11	
4437	2019	Lam.	i.	9	4734	2139	P.T. 61	xxx.	38	
4447	2023	P.T. 33	ii.	29	4815	2172	P.T. 61	xxx.	39	
		P.T. 44	xix.	29	4876	2197	P.T. 33	vii.	73	
		D'Arr.	ii.	5	4877	2198	P.T. 33	vii.	73	
4487	2037	Lam.	i.	7	4892	2205	P.T. 33	vi.	63	
4510	2047	P.T. 33	v.	46			P.T. 50	xxxvi.	4	
		D'Arr.	ii.	3			D'Arr.	ii.	6	
		Lam.	i.	2	4950	2236	P.T. 33	vi.	60	
4514	2050	P.T. 33	v.	43	4964	2241	P.T. 33	v.	45	
4532	2060	P.T. 33	ii.	26			P.T. 50	xxxviii.	13	
		P.T. 44	xix.	26			P.T. 61	xxx.	40	
		P.T. 50	xxxviii.	17			Lam.	i.	3	
		P.T. 61	xxx.	43	4971	2245	P.T. 33	viii.	85	
		D'Arr.	ii.	8			P.T. 61	xxx.	41	
4565	2072	P.T. 33	v.	48	5046	2297	P.T. 61	xxx.	42	

* No. 1179=h. 360. Other figures of the great nebula in Orion will be found in Huyghens's *Systema Saturnium*, 1659; ditto, copied by Le Gentil in *Mém. Acad. Sci. Par.* 1759, pl. 21, fig. 1; Le Gentil's own figure in *do. do.* fig. 2; by Picard, *do. do.* fig. 5; another by Le Gentil, *do. do.* fig. 6. See also:—

Mairan, "Sur la Lumière Zodiacale," copied in Lalande's 'Astronomy.' These older representations, however, are mere curiosities, and present no points of exact resemblance.

Messier, *Hist. de l'Acad. Sci. Par.* 1771, p. 435...461. Plate 8 is a careful and (for the time) elaborate figure. J. F. W. Herschel, *Mem. Astron. Soc.* ii. 1826.

De Vico, *Memoria intorno ad alcune osservazioni fatte nel Collegio Romano nel corrente anno 1838*, nebula d'Orione osservata al Telescopio di Cauchoix. 1839.

Bond. A very fine engraving—not yet published.

† No. 4447. P.T. 44. xix. fig. 29. There is an erratum in this figure. For Decl. 32° 49' n read 22° 49' n.

The following nebulae have been indicated by Lord Rosse as being either "of spiral structure (S), having in them dark spaces (D), as knotted (K), or as in the form of rays (*i. e.* much elongated forms) with splits or clefts (R).

No. in Catalogue.	h. &c.		No. in Catalogue.	h. &c.		No. in Catalogue.	h. &c.		No. in Catalogue.	h. &c.	
202	84	K	2158	731	D	2717	1085	S	3249	1451	S
372	142	S	2194	749	S	2733	1092	S	3258	1456	S
594	257	K	2248	788	D	2749	1107	D	3474	1570	S
600	262	S	2373	854	S	2807	1149	R	3572	1622	S
604	264	D	2377	857	D	2870	1196	S	3750	1734	S
888	327	S	2379	858	S	2878	1202	S	3843	1776	S
895	329	K	2413	887	D	2890	1211	S	4045	1901	K
1267	368	D	2445	910	S	2910	1225	D	4058	1909	D
1458	409	K	2499	943	S	2991	1274	K	4087	1917	R
1527	446	K	2559	982	S	3049	1312	S	4572	2075	S
1676	514	D	2597	1002	S	3050			4815	2172	S
1806	581	K	2652	1041	R	3106	1357	R	4964	2241	D
2058	692	D	2670	1052	S	3121	1368	S	4971	2245	S
2066	695	S	2680	1061	S						

List of Errata and Corrigenda in Sir William Herschel's Catalogue of 2500 Nebulae in the Philosophical Transactions.

Class.	No.	No. in Catalogue.	Error and Correction.	
I.	6	3702	for f. 3 ^m 56 ^s read f. 33 ^m 56 ^s	
	54	214	for f. 12 ^m 44 ^s ; s. 2° 50' read f. 18 ^m 36 ^s ; s. 1° 26'	
	87	2274	for f. 9 ^m 30 ^s read f. 10 ^m 30 ^s	
	137	1837	for 42 Lyncis read 41 Lyncis	
	143	3356	for s. 2° 7' read s. 0° 54'	
	144	3846	for n. 0° 24' read n. 2° 7'	
	153	536	for p. 23 ^m 16 ^s read f. 23 ^m 16 ^s	
	154	549	for f. 1 ^m 23 ^s read p. 1 ^m 23 ^s	
	155	778	for f. 7 ^m 49 ^s read p. 7 ^m 49 ^s	
	202	2604	for f. 0 ^m 47 ^s read f. 7 ^m 47 ^s	
	203	2597	for f. 7 ^m 42 ^s read f. 14 ^m 42 ^s	
	II.	1	4738	for p. 15 ^m ::; s. 3° ::; read p. 11 ^m 45 ^s , n. 0° 17'
		11	2824	for f. 1 ^m 24 ^s , n. 0° 24' read f. 1 ^m 13 ^s , n. 0° 30'
		14	2730	for 3 (ν) Virginis f. 2 ^m 20 ^s , n. 1° 22' read 59 ε Virginis p. 69 ^m 0 ^s , n. 0° 11'
48		1707	for f. 56 ^m 45 ^s read f. 54 ^m 45 ^s	
181		3214	for s. 0° 48' read s. 1° 15'	
185		3421	for p. 11 ^m 0 ^s read p. 1 ^m 0 ^s	
239		634	for 27 (x) Persei p. 8 ^m 20 ^s , n. 0° 2' read 30 Persei p. 14 ^m 41 ^s , n. 0° 51'	
240		5046	for read 39 Pisc. p. 2 ^m 24 ^s , n. 1° 0'	
241		29	for read 39 Pisc. p. 14 ^m 24 ^s , s. 0° 11'	
242		4973	for 48 (μ) Pegasi read 87 (u) Pegasi	
264		1335	for 47 (δ) Cancri read 25 (δ) Canis	
265		1384	for 1 χ Can. read ξ 1 Can.	
286		654	for p. read 13 (ξ) Eridani p.	
314		3528	for f. 17 ^m 57 ^s read f. 15 ^m 57 ^s	
372		2771	for p. 74 ^m 24 ^s read p. 14 ^m 24 ^s	
658		1718	for 44 Lyncis read 43 Lyncis	
708		1788	for 37 Lyncis read 36 Lyncis	
794		{ 3177 3179 }	for s. 0° 49' read s. 1° 0' (see note on this No. in Catal.)	
795		3216	for s. 1° 13' read s. 1° 24' (see note on this No. in Catal.)	
796		3224	for s. 1° 25' read s. 1° 36' (see note on this No. in Catal.)	
818	4056	for 12 Draconis read 12 Draconis Hevelii		
853	14	for p. 25 ^m 38 ^s read p. 25 ^m 48 ^s		
II .	6	3228	for 59 (ε) Virginis p. 28 ^m 11 ^s read d Virginis f. 2 ^m 42 ^s , n. 0° 57'. The obs. belongs to I. 8	
	26	3078	for 20 Comæ f. 4 ^m 30 ^s , s. 0° 37' read 26 Comæ p. 5 ^m 5 ^s , s. 0° 32'	
	112	2417	for 74 φ Leonis f. 10 ^m 6 ^s , s. 1° 52' read Mayer 510. z. p. 61 ^m 48 ^s , s. 1° 10'	
	113	2577	for φ Leonis f. 34 ^m 18 ^s , s. 1° 3' read Mayer 510. z. p. 37 ^m 36 ^s , s. 0° 31'	
	178	631	for 17 (γ) Persei f. 9 ^m 6 ^s read 17 (r) Persei f. 10 ^m 0 ^s	
	192	419	for 72 Ceti read 62 Ceti	

TABLE (continued).

Class.	No.	No. in Catalogue.	Error and Correction.
III.	195	684	for f. 42 ^m 42 ^s read f. 41 ^m 6 ^s
	199	628	for 27 (κ) Persei p. 8 ^m 27 ^s , n. 0° 2' read 30 Persei p. 14 ^m 44 ^s , n. 0° 55'
	256	1641	for s. 0° 48' read s. 0° 58'
	289	1911	for s. 0° 25' read s. 0° 31'
	293	1974	for 23 Leonis read 23 Leonis minoris
	319	3888	for n. 2° 26' read s. 2° 26'
	347	3844	for s. 1° 17' read s. 0° 17'
	353	2440	for f. 53 ^m 4 ^s read f. 43 ^m 4 ^s
	369	3618	for — 25 ^m 41 ^s read f. 25 ^m 41 ^s
	422	3664	} for n. 0° 44' read n. 0° 36'
	423	3668	
	511	4042	
	607	1645	for f. 3 ^m 5 ^s read f. 3 ^m 11 ^s
	627	1820	for p. 12 ^m 33 ^s read p. 12 ^m 23 ^s
	739	4149	for 43 Lynceis read 42 Lynceis
	751	1897	for p. 32 ^m 30 ^s read p. 32 ^m 47 ^s
	778	3206	for 39 Lynceis read 38 Lynceis for s. 1° 4' read s. 1° 15' (see note on this No. in Catal.)
	IV.	29	2255
31		4802	for — 0° 37' read s. 0° 37'
V.	13	4368	for n. 0° 39' read s. 0° 38' ::
VI.	8	3967	for 26 χ Virginis f. 23 ^m 44 ^s , s. 0° 6' read Mayer 577. z. f. 1 ^m 48 ^s , n. 1° 26'
VII.	6	1509	for 50 Geminorum read 51 Geminorum
VIII.	11	1534	for 50 Geminorum read 51 Geminorum
	28	1229	for (1λ) Orionis read (1st χ) Orionis
	44	1533	for 5 (η) Can. min. read 4 (γ) Can. min.

The following nebulae are declared in MS. notes to be identical.

II. 6=I. 1; II. 119=II. 94; II. 148=II. 20; II. 176=II. 70; II. 243=II. 241; II. 703=II. 621;
III. 6=I. 8; III. 198=II. 238; III. 835=III. 804.

Errata and Corrigenda in M. Auwers's Catalogue.

Page.		For	Read	Page.		For	Read
19	II. 844	II. 834	37	III. 946 in Decl.	89°	80°
20	III. 291 in Decl.	27°	26°	39	III. 347 in Decl.	24°	25°
24	II. 30 in R.A.	11 ^h 12 ^m 21 ^s	11 ^h 11 ^m 33 ^s	40	II. 751 in Decl.	20° 44'	20° 14'
25	IV. 59 under Δα	—5	—31	40	I. 282	I. 182
26	III. 385 in R.A.	16 ^h	11 ^h	42	III. 127.....	III. 127	III. 727
26	III. 388 in R.A.	10 ^h	11 ^h	42	Do. in R.A.	14 ^m	44 ^m
26	II. 342 in R.A.	10 ^h	11 ^h	42	Do. in Decl.	43°	42°
28	III. 814 in Decl.	53°	54°	72	No. 27 Decl.	58'8	48'8
32	III. 858	III. 850	77	List of Errata, II. 341...	16 ^m & 11 ^m	16 ^u & 11 ^u
33	III. 778 in R.A.	13 ^h 37 ^m	12 ^h 37 ^m	77	Ditto, III. 680...	26 ^m & 16 ^m	26 ^u & 16 ^u

M. Auwers has given a list of errata and corrigenda required in my two previous Catalogues. They are very numerous, but relate almost exclusively to errors of identification with my Father's classes and numbers. They had been, with hardly an exception, detected and rectified during the process of preparing and arranging the present Catalogue, which being therefore expurgated of them, it is unnecessary to annex a list of them here.

One very important erratum, however, must be noticed, not having been set down by M. Auwers. In p. 494, explanation of plates, Phil. Trans. 1833, figs. 13...18, for pmbM; vbM; vmbM read psbM; sbM; vsbM.

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1	h.	H.	D'Arrest, 1	h m s	s	[4]	63 4 0	-20°05	[4]	F; S; R; bet*11 and *14 ...	0
2	4014	0 1 13.8	3.065	3	120 41 11.5	20°05	3	eF; cL; mE; vglbM.....	3
3	4015	0 1 28.3	3.062	2	124 38 54.5	20°05	2	F; cL; vLE; glbM.....	2
4	1	III. 868	0 1 34.2	3.073	2	86 7 38.5	20°05	2	eF; pL; vglbM.....	3
5	2	III. 866	0 1 35.7	3.084	2	57 20 43.5	20°05	2	vF; vS; Sst+neb.....	3
6	2, a	R. nova	0 1	57 20	0
7	3	II. 591	0 1 37.0	3.076	3	74 57 52.5	20°05	3	vF; pS; R; glbM.....	4
8	4	0 1 52.1	3.083	1	63 3 27.5	20°05	1	pB; S; R; bM.....	1
9	III. 147	0 2 33.4	3.081	1	64 49 57.5	20°05	1	3Sst+neb.....	1
10	2308	III. 461	0 2 47.2	3.061	1	115 44 58.5	20°05	1	vF; cL; mE; gbM.....	2
11	2309	0 2 57.1	3.053	2	147 48 14.5	20°05	2	vF; S; R.....	2
12	5	IV. 15	0 3 14.6	3.085	(1)	63 4 58.5	20°05	1	vF; vS; stell.....	2*
13	2310	0 3 25.2	3.033	1	147 46 24.5	20°05	1	eF; p of 2.....	1
14	6	II. 853	0 3 31.8	3.089	2	57 25 28.5	20°05	2	pB; pL; E 0°±.....	3
15	2311	0 3 39.0	3.020	2	147 46 21.5	20°05	2	eeF; S; R; f of 2.....	2
16	Auw. N. 1	0 3 41.7	3.078	...	71 59 0.7	20°05	...	F (Schmidt 1861, Oct. 10) ...	0
17	2312	0 4 22.0	3.023	1	147 43 45.5	20°05	1	eF; S; R.....	1
18	7	III. 861	0 5 7.5	3.093	2	59 44 3.5	20°05	3	vF; pS; R.....	4
19	III. 456	0 5 11.3	3.076	1	84 21 57.5	20°05	1	vF; pS; iF.....	1
20	8	IV. 58	0 5 31.4	3.188	3	18 15 26.5	20°05	3	vF; vS; R; vsmbM*10; *12 241°4; 25°.	4
21	9	0 5 45.3	3.095	1	59 51 46.5	20°05	1	eF; *12, 45'', 325°.....	1
22	10	0 5 59.4	3.096	1	59 29 20.5	20°05	1	eF; vS.....	1
23	2313	0 6 47.8	3.052	1	113 57 14.8	20°04	1	eF; L; vglbM; L*cont, f ...	1
24	Auw. N. 2	0 6 59.2	3.074	...	84 47 28.8	20°04	...	A nebula (Markree Cat. 1852, Oct. 22).	0
25	11	III. 183	0 7 47.8	3.089	1	72 14 19.8	20°04	1	vF; S; E.....	2
26	2314	0 7 53.1	2.978	1	151 6 13.8	20°04	1	eF; S; R; bM.....	1
27	2315	Δ. 507	0 8 0.4	3.028	3	129 59 33.8	20°04	3	vB; vL; vmiE; tri-N.....	3†
28	12	0 8 8.9	3.083	1	78 19 58.8	20°04	1::	eL; eF; diff.....	1
29	13	II. 241 = II. 243	0 8 18.0	3.088	1	73 26 48.8	20°04	1	F; S; R; sbM.....	4*
30	14	III. 248	0 9 14.6	3.065	1	97 5 46.8	20°04	1	vF; S; iR; psvlbM.....	3
31	15	V. 16	0 11 6.2	3.112	1	60 42 9.1	20°03	1	eF; L; 3 or 4st + neb.....	2†
32	15, a	R. 6 novæ	0 11	60 42	Nos. 32...37 incl.....	0
38	16	0 13 54.8	3.106	1	68 24 45.7	20°01	1	F; S; R; psbM.....	1
39	17	0 14 5.3	3.107	1	68 19 15.7	20°01	1	E; bi-N; 3Bst near.....	1
40
41	17, a	R. 3? novæ	0 14±	68 19±	{ Several F, S (3 novæ at least presumed). }	0
42
43	2316	0 14 18.5	2.968	2	139 24 41.7	20°01	2	eF; S; R; gbM; 1st of 4.....	2
44	2317	0 14 28.2	2.967	2	139 25 13.7	20°01	2	eF; vS; R; 2nd of 4.....	2
45	2318	0 14 30.0	2.967	2	139 26 36.7	20°01	2	vF; S; R; gbM; 3rd of 4.....	2
46	2319	0 14 36.9	2.966	2	139 24 21.7	20°01	2	F; S; R; gbM; 4th of 4.....	2
47	19	II. 257	0 15 9.8	3.088	1	80 17 48.7	20°01	1	F; pL; R; gbM.....	3
48	18	0 15 10.5	3.124	1	61 1 29.7	20°01	1	F; vS; R; gbM.....	1
49	2320	0 15 55.7	2.970	1	136 3 13.0	20°00	1	vF; pS; R; bM; r.....	1
50	2321	0 16 56.0	3.004	2	123 19 22.3	19°99	2	pB; pL; IE; *14, f.....	2
51	20	0 17 38.2	3.266	1	29 26 39.3	19°99	1	Cl; pS; pC; st11...18.....	1
52	2322	Δ. 18 = 47 Toucani	0 17 47.4	2.721	4	162 51 33.3	19°99	4	⊕; !l; vB; vL; vmCM.....	4†
53	21	III. 148	0 18 39.6	3.134	2	61 33 49.6	19°98	2	pF; pL; R; pslbM.....	3
54	D'Arrest, 2	0 18 49	3.117	[1]	68 58 20	19°98	[1]	vF; S; 3 st near, making quadr.	0
55	22	0 19 18.9	3.411	1	19 23 2.6	19°98	1	Cl; pR; IC; st9...12.....	1
56	2323	0 19 51.3	2.989	2	124 27 21.9	19°97	2	vF; pL; IE; D*2', np.....	2
57	2324	0 20 17.3	2.877	2	147 45 22.9	19°97	2	pB; S; R; mbM.....	2
58	2325	0 20 25.1	2.685	1	162 18 23.9	19°97	1	pB; pS; IE; vglbM.....	1
59	23	III. 869	0 21 41.4	3.123	2	87 56 3.2	19°96	2	vF; S; bM; D*vnr; p of 2...	3
60	23, a	R. nova	0 22 0.7	+3.123	...	87 57 38.5	-19°96	...	No descr (MS).....	0

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
61	h. 23, b	H.	R. nova	h m s	s	...	87° 54' 8.5"	-19.95	...	No descr (MS)	0
62	25	II. 854	0 22 4.7	3.077	2	87 54 38.5	19.95	2	pB; vS; IE, 0°±; bM; f of 2	3
63	24	VIII. 79	0 22 6.2	3.302	1	30 33 4.5	19.95	1	Cl; vL; pR; lC; st 9...13 ...	2
64	23, c	R. nova	0 22 8.7	3.123	...	87 54 8.5	19.95	...	No descr (MS)	0
65	2326	0 22 42.6	2.979	2	124 2 2.5	19.95	2	F; pL; pmE; vgbm; p of 2...	2
66	26	II. 855	0 23 0.2	3.076	3	88 41 12.5	19.95	3	pF; cL; R; vglbm; r	6
67	2327	0 23 27.6	2.976	2	124 1 45.8	19.94	2	vB; L; vmE, 47°±5; psbM; f of 2; *10 327°±9 45''	2†
68	VI. 35	0 23 48.3	3.332	1	29 16 1.8	19.94	1	⊕; vF; S; eC	1
69	II. 471	0 24 3.5	3.096	1	80 34 1.8	19.94	1	F; iF; lbM	1
70	{ 27 = 2328 }	0 24 38.0	3.057	2	95 55 33.1	19.93	2	{ F; pL; vIE; vgbM; } *8.9, 75°±; 5'.	2
71	28	0 25 12.7	3.367	1	27 29 19.4	19.92	1	Cl; pL; lC; st 11...12; D*...	1
72	29	0 25 30.6	3.243	1	42 16 32.4	19.92	1	vF; vL; iR; g; smbM*11 ...	1
73	2329	0 26 22.3	2.971	1	122 33 57.7	19.91	1	vB; S; IE 90°; smbM*11 ...	2
74	{ 30 = 2330 }	II. 478	0 26 57.8	3.042	2	100 28 30.7	19.91	2	pF; pL; IE 90°; vglbM	4
75	2331	0 27 3.7	2.513	1	163 53 11.0	19.90	1	vF; L; R; vglbM	1
76	31	III. 467	0 27 14.0	3.033	1	103 25 59.0	19.90	1	eF; vS; R	2
77	2332	0 28 2.9	2.818	2	146 33 23.3	19.89	2	vF; pS; R; glbM; 3stf	2
78	II. 3	0 28 32.5	3.044	2	99 32 4.3	19.89	2	F; L; mE; bet 2cBst	2*
79	32	III. 476	0 28 43.0	3.147	1	66 48 49.3	19.89	1	vF; vS; stellar; *7, 15°, 5' ...	2
80	32, a	R. nova	0 28	66 48	Makes Dneb with h. 32	0
81	III. 954	0 29 20.1	3.039	1	100 51 4.6	19.88	1	eeF; S	1
82	D'Arrest, 3	0 29 33	3.15	[1]	66 46 18	19.87	[1]	F; pL; R; *6, 3½ dist	0
83	III. 223??	0 29 41.2	3.008	1	109 44 5.6	19.88	1	vF; pL; IE; 2pBst sf	1
84	33	III. 871	0 30 1.5	3.076	2	88 48 44.9	19.87	2	vF; S; R; vgbM; *11, 225°±; 30''.	3
85	2333	0 30 2.4	2.968	3	120 14 30.9	19.87	3	eF; S; vIE; amBst	3
86	2334	III. 223	0 30 20.6	3.004	1	110 42 9.9	19.87	1	pB; pL; E; gbM; r	1
87	2335	0 30 21.5	2.446	2	163 56 28.9	19.87	2	eF; S; vIE; r; *8 near	2
88	III. 876	0 30 46.6	3.098	1	82 6 6.2	19.86	1	vF; pL; iR; *np inv	1*
89	III. 870	0 31 0.5	3.079	1	88 1 7.2	19.86	1	vF; S; iR; vgbM	1
90	35	II. 707	0 31 11.4	3.278	1	42 26 0.2	19.86	1	pB; vL; iR; vgbM; r	2
91	D'Arrest, 4	0 31 15	3.08	[2]	87 36 6	19.85	[2]	F; S; R; lbM	0
92	34	0 31 40.6	5.151	1	5 26 27.6	19.84	1	Cl; vL; R; 150...200st 10 ... 18.	1
93	36	0 31 40.7	3.407	1	29 42 31.5	19.85	1	Cl; pL; R; st 11...15	1
94	37	0 31 49.4	3.081	1::	87 25 4.5	19.85	1	vF; L; p of 2; st 15 close ...	1
95	38	II. 479	0 31 53.4	3.039	1	99 46 25.5	19.85	1	pB; pL; iE 0°±	3
96	39	III. 872	0 32 2.3	3.072	5	89 54 28.5	19.85	5	F; pS; pmE; bM; 1st of 3...	6
97	II. 857	0 32 4.5	3.079	1	87 57 7.5	19.85	1	F; S; vgbM	1
98	40	II. 856	0 32 5.6	3.080	1	87 42 54.8	19.84	1	pB; S; R; vgbM	2
99	40, a	R. nova	0 32	87 42	(?=h. 37 or II. 857)	0
100	41	III. 595	0 32 6.8	3.072	1	89 51 23.8	19.84	1	F; pS; R; psmbM; f of 2 ...	2
101	42	II. 860	0 32 14.4	3.081	1::	87 27 4.8	19.84	1::	pB; pS; R; vgbM; 2nd of 3	5
102	43	III. 873	0 32 21.4	3.072	2	89 55 7.8	19.84	2	vF; cL; E; vglbM; f of 3 ...	3
103	D'Arrest, 5	0 32 22	3.08	[1]	87 37 24	19.84	[1]	F; vS; *8, p 27°7, ls	0
104	II. 858	0 32 22.5	3.079	1	87 52 7.8	19.84	1	pB; S; vgbM	1
105	44	V. 18	C.H.	0 32 45.4	3.243	1	49 4 49.8	19.84	1	vB; vL; mE 165°; vgvmbM	6†
106	45	V. 36	0 32 47.8	3.237	1	50 1 34.8	19.84	1	vF; vL; mE 0°	4†
107	46	II. 452	0 33 32.4	3.020	3	104 38 20.1	19.83	3	B; pS; R; psbM; r; *90''±	4
108	46, a	R. nova	0 33 32	104 38	E 0°±	0
109	III. 200	0 33 48.8	3.129	2	74 17 8.4	19.82	2	F; S; bet 2 Sst	2
110	2336	0 33 49.9	2.763	2	146 56 14.4	19.82	2	vF; S; R; p of 2	2
111	47	II. 209	0 34 5.4	3.166	1	65 16 6.4	19.82	2	pF; pS; vglbM; r	4
112	2337	0 34 25.7	2.757	2	146 58 27.7	19.81	2	F; S; R; amst; f of 2	2
113	49	III. 244	0 34 28.9	2.990	1	111 48 59.7	19.81	1	eF; vS; IE 0°...90°	2
114	48	II. 480	0 34 30.2	3.033	1	100 47 1.7	19.81	1	F; S; IE 90°±; glbM	3
5058	0 35 1.0	89 50 6.6	See No. 5058.	
115	2338	0 35 2.1	+ 2.344	3	164 9 52.7	- 19.81	3	F; iR; vgbM; 1st of several	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
116	50	M. 31	0 35 3.9	+3.252	1	49 29 45.7	-19.81	1	{!!!eeB; eL; vmE; (An- drom. Gt. Neb.) Bifid (Bond)	13†
117	51	M. 32	0 35 5.3	3.250	1	49 54 12.7	19.81	1	!vvB; L; R; psmbMN	8†
118	2339	0 35 5.4	2.338	1	164 14 1.7	19.81	1	vF; R; 2nd of several.....	1
119	{D'Arrest, 6= Auw.N.4	0 35 6	3.07	[3]	89 55 0	19.81	[3]	vF; pL; R (Bond, Jan. 1853)	0*
120	52	VIII. 78	C.H.	0 35 19.2	3.457	1	28 58 43.0	19.80	1	Cl; L; IC; st 9...10	3
121	53	0 35 24.0	3.204	1	58 10 55.0	19.80	1	eF; S; R; *13, 20" 180° ...	1
122	II. 444	0 35 33.8	3.064	1	92 18 11.0	19.80	1	F; pL; lbM.....	1
123	2340	Δ. 2??	0 35 50.6	2.329	1	164 7 16.6	19.78	1	i train of st and neb	1
124	54	III. 149	0 36 5.2	3.196	2	60 10 58.3	19.79	2	F; vS; R; lbM	3
125	II. 245	0 36 10.9	3.124	4	76 27 54.3	19.79	4	F; pS; iLE; bM	4
126	2341	0 36 53.3	2.803	1	140 56 42.6	19.78	1	eF; pl; R; gvlbM	1
127	2342	0 38 13.4	2.278	1	164 11 39.2	19.76	1	vF; R	1
128	2343	0 38 18.6	2.275	3	164 12 17.2	19.76	3	vF; S; bi-N.....	3
129	55	III. 485	0 38 45.1	3.005	1	106 20 58.5	19.75	1	vF; S; iR; r; *10, 5' s	3
130	II. 445	0 39 3.7	3.063	1	92 29 13.5	19.75	1	F; pS; iF; er	1
131	56	V. 25	0 40 1.4	3.019	2	102 38 24.1	19.73	2	vF; L; 4st in diff n.....	3
132	57	V. 20	0 40 4.2	2.979	1	111 30 57.1	19.73	1	F; vL; vmE 172°	2*
133	2344	0 40 13.4	2.240	2	164 8 36.1	19.73	2	F; S; E or bi-N; vglbM.....	2
134	2346	Δ. 19? 21?	0 40 17.9	2.255	3	163 51 1.1	19.73	3	F; pL; vLE; r	3
135	58	III. 204?	0 40 22.9	3.154	(2)	71 9 48.1	19.73	1	vF; S; R; lbM; 2vSstf; *inv	3
136	59	II. 609	0 40 32.9	3.195	5	63 8 35.1	19.73	5	pB; S; R; pmbM; r; * p ...	6
137	59, a	R. nova	0 40	63 8	One of R.'s novæ; the other =h. 60.	0
138	{61= 2345}	V. 1	C.H.	0 40 37.6	2.954	3	116 3 40.4	19.72	3	{!!; vvB; vvL; vmE } 54°5 gbM; 4st.	9*†
139	60	0 40 39.3	3.195	1	63 8 14.4	19.72	1	pF; R; bM	1
140	2347	0 40 40.7	2.920	3	122 11 28.4	19.72	3	vB; pS; lE; smbM; *8, 5'nf	3
141	62	II. 472	0 40 43.5	3.020	1	102 14 39.4	19.72	1	F; pS; R; gbM	2
142	2348	0 40 44.3	2.223	4	164 16 44.4	19.72	4	F; S; R; gbM; *9, 40"nf ...	4
143	II. 863	0 40 47.1	3.105	1	82 27 14.4	19.72	1	pL; lE; gbM; r	1
144	63	0 40 55.7	3.057	1	93 37 22.4	19.72	1	vF; Δ 2st & neb	1
145	64	{II. 703 = II. 621	0 40 56.1	3.058	1	93 32 40.4	19.72	1	F; S; E 135°±; lbM.....	3*
146	2349	Δ. 3, 4, 21?	0 41 23.8	2.233	3	163 52 6.7	19.71	3	F; pL; R; gbM*13	3
147	2350	0 41 41.6	2.872	1	129 0 20.7	19.71	1	F; S; R; vsymbM*13	1
148	2351	0 42 3.8	2.198	4	164 15 4.0	19.70	4	F; pS; R;	4
149	65	III. 153	0 42 13.6	3.226	4	58 29 23.0	19.70	4	pL; pS; lE; psbM; r; *8sf4',	5
150	2352	0 42 49.8	2.195	1	164 2 37.3	19.69	1	Cl; F; pL; stvS	1
151	66	III. 463	0 43 1.3	3.046	2	95 57 59.3	19.69	2	vF; pS; iLE; r.....	4
152	2353	0 43 20.8	2.169	1	164 18 1.6	19.68	1	vF; S; R.....	1
153	68	III. 955	0 43 28.7	3.030	(1)	99 25 39.6	19.68	2	pF; vS; iR; pgbM.....	3
154	67	II. 446	0 43 34.3	3.061	1	92 39 59.6	19.68	1	pF; S; lE; psbM; *8 f5*5...	3
155	III. 430	0 43 47.0	3.038	1	97 39 17.9	19.67	1	vF; vS.....	1
156	69	III. 429	0 43 56.8	3.037	1	97 49 22.9	19.67	1	pB; pS; smbM; sp of Dneb .	3
157	70	0 43 59.1	3.037	1	97 49 2.9	19.67	1	vF; S; R; nf of Dneb	2
158	71	I. 159	0 44 10.2	3.352	1	43 12 15.9	19.67	1	cB; pL; R; 2st10nr	4
159	73	III. 439	0 44 59.5	3.059	2	92 59 10.5	19.65	2	vF; S; iR; bM; stellar	4
160	72	III. 477	0 45 0.3	3.189	1	66 25 23.5	19.65	1	eF; S; R; *15, f30"	3
161	75	0 45 52.1	3.241	1	58 16 59.8	19.64	1	eF; S; R	1
162	{74= 2354}	VI. 20	0 45 52.1	2.932	2	117 20 41.8	19.64	2	⊕; B; L; lE; st 12...16.....	3
163	2355	0 45 57.6	2.903	3	121 57 53.1	19.63	3	vB; L; pmE; glbM; *11np	4
164	2357	0 46 16.9	2.135	1	163 54 48.1	19.63	1	eF	1
165	2356	0 46 33.5	2.115	2	164 6 33.1	19.63	2	Cl; F; eeL; R; st 12...18 ...	2*
166	2358	Δ. 5, 6?	0 47 14.8	2.102	1	164 8 28.7	19.61	1	vF; pL; R; vglbM; r.....	2
167	II. 214	0 47 51.9	+3.242	1	59 11 21.0	-19.60	1	F; E; aB*f, vnr	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
168	h. 2360	H.	h m s	s	4	162° 57' 22.3	- 19.59	4	pB; vS; R; gvlbM; r.....	4
169	2359	0 48 23.0	+ 2.151	3	128 27 8.3	19.59	4	pB; vL; vmiE; vgpmbM ...	5*+
170	76	0 48 24.5	2.844	1	78 40 51.6	19.58	1	Cl; S; sest	1
171	2361	0 48 59.6	3.131	1	163 0 0.9	19.57	1	F; vS	1
172	77	0 49 15.8	2.132	1	92 31 27.9	19.57	1	pF; S; E	1
173	78	0 49 25.0	3.060	1	60 28 21.2	19.56	2	pF; vS; R; gbM.....	2
174	2363	0 49 55.7	3.240	2	143 32 6.2	19.56	2	F; S; R; *12 f 90°.....	2
175	2362	0 50 3.5	2.675	2	122 43 13.2	19.56	2	eF; vS; R; pB* f 2'	2
176	79	II. 210	0 50 10.7	2.883	2	60 24 25.2	19.56	3	pB; pL; R; gbM; *9, 3' 135°	4
177	79, a, b	R. 2 novæ	0 50 11.5	3.241	3	60 21 31.2	19.56	{ F; S; R (ε of Lord R.). } For b, see No. 5059. }	0*
178				4007					
178	4007	0 50 29.3	2.780	1	134 35 56.5	19.55	1	eF; vS; R; lbM	1*
179	4008	0 50 36.2	2.780	1	134 30 1.5	19.55	1	vF; vS; R; lbM; 3stp	1*
180	2365	0 50 36.7	2.668	2	143 43 58.5	19.55	2	pF; S; R; bM; p of 2	2
181	2364	0 50 40.4	2.810	1	131 12 36.5	19.55	1	(?)F; S; stellar	1
182	2366	0 50 51.5	2.667	1	143 39 58.8	19.54	1	vF; iE; vgbM; f of 2	1
183	2367	Δ. 23	0 51 27.2	2.078	5	163 13 35.1	19.53	5	⊕; vB; S; iE; st 13...15 ...	5
184	2368	0 52 10.5	2.850	2	125 53 21.4	19.52	2	vF; S; R; glbM; 2st11s.....	2
185	80	II. 433	0 52 46.1	3.027	...	98 19 51.0	19.50	...	pF; L; E 0°±; glbM; *10, f 20*5.	2
186	2369	0 53 13.3	1.902	1	165 12 27.0	19.50	1	F; L; R; vgbM	1
187	2370	Δ. 25	0 54 19.9	2.043	5	162 56 9.3	19.49	5	B; L; viF; mbMD*; r	5+
188	2371	0 54 45.4	2.632	2	143 59 49.6	19.48	2	eF; S; R	2
189	81	III. 191	0 55 5.2	3.044	2	94 59 49.5	19.45	2	pF; S; iE; *8f97 ^s	4
190	82	II. 434	0 56 16.4	3.032	1	97 5 49.1	19.43	3	F; S; iR; sbM; *14nf 20''...	4
191	2372	0 57 26.8	2.308	1	156 21 30.0	19.40	1	eF; vmiE 145°4; vlbM	1
192	2374	Δ. 55 ??	0 57 28.7	2.022	2	162 22 33.0	19.40	2	vvF; pL; viE; vgbM	2
193	2375	Δ. 62	0 57 28.8	2.069	3	161 35 59.0	19.40	3	⊕; vB; vL; vC; vmbM; st 13...14.	4
194	2373	0 57 44.3	2.828	2	125 53 46.0	19.40	2	F; S; R; glbM	2
195	83	0 57 45.8	3.700	1	28 34 5.0	19.40	1	Cl; S	1
196	4012	0 57 57.6	2.745	1	134 1 50.3	19.39	1	eF; vS; *7.8 sp 3'	1*
197	D'Arrest, 7	0 58 53	3.28	[1]	58 20 18	19.37	[1]	vF; *13, s 15''; m diff	0
198	2376	Δ. 31 ??	0 58 56.4	1.969	5	162 48 27.9	19.37	5	Cl; F; L; R; pC; st 14...16.	5
199	D'Arrest, 8	0 59 21	3.29	[1]	57 57 18	19.36	...	F; S; bet 2 st 15.....	0
200	2378	Δ. 36 ??	0 59 27.3	1.904	2	163 34 23.2	19.36	2	⊕; B; S; R	2
201	2377	0 59 31.7	2.864	1	120 55 53.2	19.36	1	vF; S; R; gbM	1
202	84	II. 215	0 59 32.3	3.289	3	58 14 9.2	19.36	3	pF; S; R; bM; 1st of 3	4*
203	85	II. 216	0 59 34.3	3.289	3	58 16 41.2	19.36	3	pF; S; R; sbM; 2nd of 3 ...	4*
204	VIII. 64	C.H.	0 59 37.4	3.703	1	29 9 36.2	19.36	1	Cl; pC	1
205	86, a	R. nova	0 59 39.6	3.289	...	58 20 48.2	19.36	...	γ' in Lord R.'s diagram	0*
206	86	II. 217	0 59 40.5	3.289	3	58 20 16.2	19.36	3	pF; pL; R; gbM; 3rd of 3...	4*
207	D'Arrest, 9	0 59 42	3.29	[3]	58 26 0	19.36	[3]	p of Dneb; vF; pS; { Δ.R.A.=0 } { Δ.P.D.=93'' }	0*
208	D'Arrest, 10	0 59 42	3.29	[3]	58 25 18	19.36	[3]	vF; R; pS; f of D neb	0*
209	86, b	R. nova	0 59 46.3	3.288	...	58 23 20.2	19.35	...	δ in Lord R.'s diagram	0*
210	86, c	R. nova	1 0 2.3	3.288	...	58 27 16.2	19.34	...	θ in Lord R.'s diagram	0*
211	Auw. N. 9	1 0 15.1	3.071	...	89 48 55.8	19.34	...	F nebula (Bond, Jan. 1853).	0*
212	87	II. 218	1 0 39.6	3.299	1	57 37 10.1	19.33	1	F; vS; R; mbM; bet 2 st ...	2
213	87, a	R. nova	1 0	59 37	makes a D neb with h. 87 ...	0
214	88	I. 54	1 0 43.4	3.361	2	51 5 25.1	19.33	2	F; vS; viE; gbM; 4Sstnr ...	4*
215	D'Arrest, 11	1 0 45	3.30	[1]	57 36 18	19.33	[1]	F; S; pos from h. 87=40°; dist 47''.	0
216	2379	1 0 45.8	1.940	1	162 44 38.1	19.33	1	vF; pL; R; glbM	1
217	D'Arrest, 12	1 1 29	3.30	[1]	57 59 48	19.31	[1]	vF; S; R; *10f1 ^s .8, s 80'' ...	0
218	89	II. 224	1 1 39.0	3.326	2	55 2 13.7	19.31	2	pB; cL; R; gbM; β Androm. nr	3
219	2380	1 2 5.5	2.679	1	137 25 34.0	19.30	1	eS; stellar; =*7m	1
220	2381	1 2 39.6	2.041	1	160 37 28.3	19.29	1	F; vL; R; vglbM	1
221	II. 219	1 2 44.2	3.306	(1)	57 34 0.3	19.29	(1)	eS; F; p of D neb	1
222	II. 220	1 2 44.2	+ 3.306	(1)	57 34 0.3	- 19.29	(1)	pL; f of D neb	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
223	2382	1 3 9.2	+2.800	1	126 31 25.9	-19.27	1	eF; S; R; vS*nr	1
224	2384	1 3 27.6	1.908	1	162 30 42.9	19.27	1	eF; pL; R; gvlbM	2
225	2383	1 3 37.2	2.801	2	126 14 35.2	19.26	2	vF; S; R; glbM	2
226	2386	1 3 39.2	1.861	4	163 6 23.2	19.26	4	F; pS; R; gbM	4
227	2385	1 3 57.5	2.849	3	120 57 48.5	19.25	3	F; pL; R; vglbM; p of 2	3
228	2387	Δ. 36?	1 4 4.2	1.814	1	163 37 59.5	19.25	1	vF; vS; R	4
229	90	III. 15	1 4 25.7	3.303	1	58 37 12.8	19.24	1	F; pS; R; bM	2
230	III. 15 ⁴	1 4 26.1	3.303	1::	58 36 22.8	19.24	1::	eF; vS	1
231	[162]	1 4 28.3	1.890	1	162 30 54.8	19.24	1	vF	1
232	2388	1 4 41.6	2.855	2	119 58 44.8	19.24	2	eF; S; E; glbM; f of 2	2
233	2389	1 4 57.6	2.768	1	128 49 39.1	19.23	1	vF; S; R; glbM	1
234	91	III. 592	1 5 41.2	3.066	1	91 3 7.7	19.21	1	vF; vS; R	2
235	91, a	R. nova	1 5	91 3	Forms a Δ with h. 91 & 92	0
236	2390	1 5 43.8	2.826	2	122 49 44.7	19.21	2	2 vSt+ F neb	2
237	92	III. 593	1 5 46.4	3.065	1	91 4 22.7	19.21	1	eF; eS	2
238	II. 622	1 5 49.2	3.074	1	89 45 44.0	19.20	1	F; L; R; bM; er	1
239	93	II. 447	1 5 51.7	3.066	1	90 59 22.7	19.21	1	F; vS; R; vsbM*	3
240	95	1 6 16.1	3.324	1	57 1 50.0	19.20	1	F; S; vsbM	1
241	2391	1 6 16.3	2.340	1	152 20 39.0	19.20	1	F; S; R; gbM; *12 f	2
242	94	1 6 27.6	3.732	1	30 36 52.3	19.19	1	Cl; S; lC	1
243	2392	1 6 36.9	2.430	2	148 59 52.3	19.19	2	B; S; R; psbM	2
244	VII. 45	1 6 53.4	3.703	2	31 56 6.6	19.18	2	Cl; S; iF; pC	2
245	2393	1 7 7.0	2.761	2	128 38 59.9	19.17	2	pF; S; R; glbM	2
246	2394	1 7 12.4	2.823	2	122 29 25.9	19.17	2	pB; S; R; gbM	2
247	2396	1 7 13.1	2.423	2	149 1 44.9	19.17	2	F; vS; R	2
248	2395	1 7 16.6	2.823	2	122 32 55.9	19.17	2	pF; S; R; gbM	2
249	D'Arrest, 13	1 7 21	3.32	[1]	57 31 18	19.16	[1]	F; S; R; *15 p, 8 ^s .3, 270°	0
250	D'Arrest, 14	1 7 51	3.32	[2]	57 40 42	19.15	[2]	F; pL; bM; *11 nr	0
251	96, a	R. nova	1 8	59 43	vF; mE135°±; lbM; nph.96	0
252	96	1 8 29.3	3.306	1	59 42 41.8	19.14	1	vF; E; *9np; S*nf, vnr	1
253	2397	1 8 35.1	2.480	1	146 8 37.8	19.14	1	vF; S; R; bM	1
254	III. 440	1 8 48.8	3.061	1	91 35 19.1	19.13	1	vF; vL	1
255	2399	Δ. 7, 10?	1 9 50.6	1.668	3	164 2 15.0	19.10	4	pF; pL; iR; r; 1st of sev, neb and st.	3
256	97	VII. 42	1 10 24.8	3.722	1	32 24 35.3	19.09	1	Cl; B; L; pR; st 7, 8, 10	3
257	2401	Δ. 60?	1 10 40.8	1.799	2	162 17 13.6	19.08	2	pF; L; R; vgbM	2
258	III. 205	1 10 41.4	3.198	1	73 4 51.6	19.08	1	eF	1
259	2402	Δ. 8, 10?	1 10 48.6	1.650	4	164 2 41.6	19.08	4	F; pL; iR; gbM; r; 2nd of sev.	4
260	2400	1 10 54.4	2.794	1	124 6 11.6	19.08	1	pB; R; glbM; ? 1° in P.D.	1
261	2404	Δ. 9?	1 11 42.3	1.629	1	164 4 36.5	19.05	1	pB; pL; iF; 3rd of sev.	1
262	2403	1 11 43.3	2.365	1	149 38 46.5	19.05	1	vF; R; gbM; 30	1
263	99	III. 250	1 11 56.6	3.091	1	87 25 16.5	19.05	1	pB; pL; R; gmbM; p of 2	4
264	I. 108	1 12 1.0	3.091	1	87 30 53.5	19.05	1	eB; vL; iR; pB*f	1
265	98	1 12 2.9	3.335	1	58 1 55.5	19.05	1	vF; eS; stellar	1
266	D'Arrest, 15	1 12 41	3.33	[2]	58 2 0	19.03	[2]	eF; vS; *9p14 ^s ; v diffc.	0
267	III. 206	1 12 41.2	3.192	1	74 14 55.1	19.03	1	eF; S	1
268	100	III. 577	1 13 19.9	3.428	1	50 14 18.7	19.01	1	vF; pS; vIE; vglbM	2
269	III. 251	1 13 21.4	3.092	3:	87 22 9.7	19.01	3:	pB; S; smbM; f of 2	3
270	101	1 13 47.2	3.119	1::	83 43 2	19.00	1	eF; pL; R; red *7.8, 225°	1
271	2405	1 14 3.5	2.691	1	131 42 52.3	18.99	1	eF; IE	1
272	102	III. 156	1 14 6.0	3.350	1	57 17 30.3	18.99	1::	vF; eS; 1st of 3	3
273	2406	1 14 7.3	2.353	2	149 15 33.3	18.99	2	vB; S; IE; psmbM	2
274	D'Arrest, 16	1 14 29	3.14	[1]	81 31 48	18.97	[1]	pB; S; E	0
275	103, a	R. nova	1 14 29	3.107	...	85 24 26.6	18.98	...	γ in Lord R.'s diagram	0
276	103	III. 252	1 14 31.2	3.107	2	85 28 25.6	18.98	2	pB; L; R; svmbM; *7 f 1 ^m	3
277	103, b	R. nova	1 14 46.4	3.107	...	85 21 47.6	18.98	...	β in Lord R.'s diagram	0
278	III. 157	1 14 51.3	3.351	2	57 15 28.9	18.97	2	vF; S; 2nd of 3	2
279	2407	1 14 53.5	2.772	2	124 48 15.9	18.97	2	B; S; vIE; bM; vS*nr	2
280	103, c	R. nova	1 14 57.1	3.107	...	85 18 37.6	18.97	...	δ in Lord R.'s diagram	0
281	105	III. 594	1 14 59.5	3.075	1	89 47 7.2	18.96	1	vF; L; mE 60°±; lbM	2
282	104	1 15 4.4	+3.352	1	57 33 18.2	-18.96	1	vF; E; *s	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. O.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. O.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
.....	h.	H.		h m s	s		° ' "	"			
288	104, a	R. 5 novæ	1 15	+	57 33	-18.95	...	No descr Nos. 283...287 incl	0
289	106, a	R. nova	1 15	57 16	18.95	...	No description	0
290	106	III. 158	1 15 19.6	3.354	2	57 16 46.5	18.95	2	pB; pL; R; 3rd of 3	4
291	103, d	R. nova	1 15 23.2	3.107	...	85 20 17.6	18.95	...	ε in Lord R.'s diagram	0
292	D'Arrest, 17	1 15 35	3.35	[3]	57 31 18	18.94	[3]	vF; S; obs. with H. 157, 158, 159, 160.	0
293	107	1 15 36.6	3.352	1:	57 36 31.8	18.94	1:	No description	1
294	D'Arrest, 18	1 15 38	3.13	[2]	81 41 6	18.94	[2]	cB; S; R; bMN	0
295	108	III. 159	1 15 48.0	3.354	2	57 28 37.8	18.94	2	vF; pL; R; bM; p of 2	3
296	109	III. 160	1 15 49.0	3.354	1	57 26 52.8	18.94	1	vF; S; f of 2	2
297	110	1 16 7.8	3.362	1	56 49 53.1	18.93	1	vF; vS	1
298	111	III. 169	1 16 34.4	3.362	1	56 56 28.4	18.92	1	F; S; stellar	2
299	112	II. 252	1 16 35.8	3.169	2	77 49 10.7	18.91	2	F; L; IE; vglbM; * f.	3+
300	113	III. 167	1 16 45.3	3.360	1	57 16 37.7	18.91	1	Stellar; p of 2	2
301	D'Arrest, 19	1 16 48	3.14	[2]	81 10 6	18.91	[2]	eF; S; v diff; I 151 f41s ...	0
302	114	III. 168	1 16 48.8	3.360	1	57 18 37.7	18.91	1	pB; R; stellar; f of 2	2
303	114, a	R. nova	1 16	57 18	S; R; bM	0
304	116	III. 253	1 17 19.4	3.097	1	86 55 16.3	18.89	1	F; cL; E 135° ±	3+
305	115	II. 461	1 17 19.8	3.078	2	89 0 21.3	18.89	2	F; pL; R; gbM	5
306	D'Arrest, 20	1 17 22	3.14	[2]	80 44 24	18.89	[2]	eF; pL; iF; ?Cl + neb.	0
307	D'Arrest, 21	1 17 25	3.37	[1]	56 42 18	18.89	[1]	D neb; vF; 90° pos	0
308	117	I. 151	1 17 26.8	3.142	1	81 11 50.3	18.89	1	vB; pL; mbM; 4S st nr	2
309	D'Arrest, 22	1 17 31	3.14	[2]	81 1 12	18.88	[2]	vF; vS; * 11.12 p 5s	0
310	2408	1 17 35.1	2.750	3	125 48 8.3	18.89	3	F; S; IE; p of 2	3
311	2409	1 17 37.5	2.748	3	125 51 0.3	18.89	3	F; S; IE; bM; f of 2	3
312	118	1 17 44.5	3.377	1	56 1 31.3	18.89	1	pB; vS; sbM; p of 2	1*
313	118, a	R. nova	1 17	56 1	One of 4 neb nr h. 120	0
314	III. 556	1 17 51.6	3.140	1	81 28 43.6	18.88	1	vF; pL; mE 15° ±	1*
315	119	1 17 51.6	3.140	(1)?	81 29 30.6	18.88	1	Not vF; L; R; bM	2*
316	121	II. 462	1 18 19.5	3.081	2	88 58 4.9	18.87	2	pB; pL; R; gmbM	3
317	2410	1 18 22.9	2.708	2	128 52 11.9	18.87	2	eeF; S; R; vgbM; 1st of 4... ..	2
318	120	1 18 24.6	3.379	1	56 2 19	18.87	1	pB; pL; gbM; f of 2	1*
319	120, a	R. nova	1 18	56 2	One of 4, see h. 118, 120 ...	0
320	III. 170	1 18 26.6	3.373	1	56 39 5.2	18.86	1	Stellar	1*
321	2411	1 18 51.2	2.707	2	128 48 57.5	18.85	2	eeF; S; R; vgbM; 2nd of 4... ..	2
322	2412	1 18 52.4	2.707	2	128 47 17.5	18.85	2	eeF; S; R; vgbM; 3rd of 4... ..	2
323	II. 448	1 18 55.2	3.056	1	92 3 54.5	18.85	1	Stellar; p of D neb	1
324	II. 449	1 18 55.2	3.056	1	92 3 54.5	18.85	1	Stellar; f of D neb	1
325	2413	1 19 6.8	2.707	1	128 44 40.8	18.84	1	eeF; S; R; vgbM; 4th of 4... ..	1
326	III. 171	1 19 27.2	3.383	1	56 4 7.1	18.83	1	Stellar	1*
327	122	II. 463	1 19 33.1	3.083	2	88 42 53.1	18.83	2	F; S; E90°; bM; r.	5
328	123	III. 560	1 19 42.9	3.415	1	53 32 28.1	18.83	1	vF; S; E; vglbM; *13 nr ...	2
329	III. 172	1 19 56.5	3.371	1	57 16 7.4	18.82	1	vS; stellar; p of 2	1
330	III. 173	1 19 56.5	3.371	1	57 16 7.4	18.82	1	vS; stellar; f of 2	1
331	124	VII. 48	1 20 10.3	3.969	1	27 25 54.7	18.81	1	Cl; B; pL; pRi; st mm	2*
332	D'Arrest, 23	1 20 21	3.38	[1]	56 25 18	18.80	[1]	eF; pL; R	0
333	III. 441	1 20 34.8	3.051	1	92 40 8.0	18.80	1	vF; vS; iE; p of 2	1
334	III. 442	1 20 39.9	3.052	1	92 37 8.0	18.80	1	vF; vS; iF; f of 2	1
335	125	1 21 10.0	3.362	1	58 23 57.6	18.78	1	vF; S; R	1
336	2414	1 21 37.4	2.726	1	126 27 1.2	18.76	1	vF; S; R	1
337	2415	1 22 20.5	2.675	1	130 2 23.8	18.74	1	eF; S; att to S*; B*nr	1
338	2416	1 22 43.5	2.723	1	126 19 27.1	18.73	2	vS; *pos 225° inv	3
339	2417	1 23 15.2	2.453	1	142 18 52.7	18.71	1	F; S; R; bM; am st 11	1
340	2418	1 23 43.9	2.864	1	113 23 44.0	18.70	1	B; L; pmE; gpmbM	1
341	127	1 23 47.8	3.390	1	57 6 32.0	18.70	1	vF; pL; gbM	1
342	126	Σ.131 = M.103	1 23 59.8	3.916	2	30 2 9.3	18.69	2	Cl; B; R; Ri; pL; st 10...11 ..	5
343	128	I. 100	1 24 18.7	3.008	3	97 35 25.6	18.68	3	vB; pL; R; mbM; p of 2 ...	4
344	128, a	R. nova	1 24	97 35	No description	0
345	III. 431	1 24 22.2	3.008	(1)	97 36 39.6	18.68	(1)	eF; S; f of 2	1
346	129	1 24 33.4	3.059	1	91 38 46.6	18.68	1	vF; S; R; bM	1
347	130	1 24 35.7	+ 3.008	*1	97 36 47.9	-18.67	1	vF; vS; R	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
347	h.	H.	D'Arrest, 24	h m s	s	[1]	55° 26' 18"	-18°66'	[1]	vvF; S; ?rr	0
348	D'Arrest, 25	1 25 0	3·35	[2]	60 5 0	18·65	[2]	F; M. 33, f65 ^s ; another f28 ^s ..	0
349	D'Arrest, 26	1 25 28	3·35	[2]	60 5 0	18·64	[2]	F; pL, f of 2.....	0
350	2419	1 25 53·4	2·737	2	124 13 32·1	18·63	2	F; S; R; bM	2
351	132	II. 4	1 25 54·1	3·006	1	97 44 13·1	18·63	1	pB; R; bM; r; *6, f47 ^s ·5 ...	7
352	131	V. 17	M. 33	1 25 56·3	3·358	1	60 3 50·1	18·63	1	!; eB; eL; R; vRi; vgbMN; rr.	14†
353	II. 473	1 26 1·4	2·960	1	102 53 18·1	18·63	1	F; S; iF; er.....	1
354	III. 432	1 26 15·0	3·003	1	98 2 38·4	18·62	1	eeF	1
355	133	III. 150	1 26 17·0	3·361	1	59 57 47·4	18·62	1	vS; R; vvlbM	5
356	2421	Δ. 17?	1 26 24·6	1·327	2	164 16 37·4	18·62	2	B; S; R; psbM*; r.....	2
357	R. nova	1 26 33·5	3·355	::	60 31 53·9	18·61	::	S, a neb or Cl with 3stin v...	0
358	D'Arr. = Auw.N.15	1 27 18·1	3·002	...	98 5 42·6	18·58	...	{ Nebulous *11, m (D'Arr. Resultate). }	0*
359	134	1 27 23·6	3·399	1	57 4 11·6	18·58	1	vF; psbM; stellar	1
360	2423	1 27 44·5	2·690	1	127 13 26·9	18·57	1	F; vS; R; *12, p	1
361	{ 139 = 2422 }	I. 281	1 27 47·4	2·780	3	120 7 37·9	18·57	3	{ vB; vL; vmE, 118°·3; } sbmM; *34°·5, 6 ^s ·5. }	5
362	135	III. 174	1 27 53·1	3·401	1	57 2 16·9	18·57	1	pF; psbM; stellar	2
363	137	II. 282	1 28 18·0	3·001	1:	98 2 43·5	18·55	1	pB; pL; iE; gmbM; r; *8, np10'.	4
364	136	1 28 19·5	3·399	1	57 19 26·5	18·55	1	pB; pL; bM; *f, 2 ^m 51 ^s	1
365	2424	1 28 37·9	2·687	1	127 12 19·8	18·54	1	eeeF; vS; R; p of 2.....	1
366	138	III. 454	1 28 48·9	3·072	1	90 2 25·1	18·53	2	eF; pL; not bM	2
367	2425	1 28 50·4	2·686	1	127 12 40·1	18·53	1	F; S; R; f of 2	1
368	140	III. 471	1 28 55·4	2·976	1	100 43 34·1	18·53	1	eF; S; R; am vSst	2
369	2426	Δ. 479	1 28 57·2	2·612	1	132 9 23·1	18·53	2	B; pL; mE; gpmbM	3
370	2427	1 29 2·6	2·647	1	129 51 50·1	18·53	1	pF; S; R; bM.....	1
371	141	1 29 9·2	3·405	1?	57 7 44·4	18·52	1::	vF; R; f of 2	1
372	142	M. 74	1 29 11·1	3·211	2	74 55 46·4	18·52	2	⊕; F; vL; R; vg, psmbM; rr	11†
373	Auw. N. 16	1 29 14·4	4·681	...	17 49 45·0	18·50	...	3st+neb (Struve, Σ. 2)...	0
374	2428	1 29 27·8	2·642	1	130 3 42·7	18·51	1	iF; S; R; bM.....	1
375	143	1 29 59·6	3·119	1	84 50 14·0	18·50	1	pB; S; R; psbM.....	1
376	2429	1 30 9·9	2·669	2	128 2 13·0	18·50	2	pB; S; R; gbM; *np.....	2
377	144	II. 283	1 32 7·6	2·997	2	98 13 12·4	18·42	2	pB; vS; R; mbM; r	4
378	VII. 49	1 32 8·2	4·132	1	26 40 32·4	18·42	1	Cl; pS; L & vSst.....	1
379	2430	1 32 32·0	2·759	3	120 38 10·7	18·41	3	vF; vS; p of 2.....	3
380	2432	1 32 38·7	2·575	2	133 14 25·0	18·40	3	F; S; R; gpmbM; p of 2 ...	3
381	2431	1 32 39·3	2·759	3	120 37 15·0	18·40	3	vF; pS; R; gbM; *f, nr.....	3
382	2435	1 32 40·7	0·929	2	166 16 4·0	18·40	2	vF; pS; R; vglbM	2
383	2433	1 32 50·9	2·573	2	133 18 22·0	18·40	2	F; S; vLE; glbM; f of 2	2
384	2434	1 32 55·1	1·907	1	155 36 37·3	18·39	1	vF; iR; vglbM	1
385	M. 76	1 33 28·5	3·334	1	39 8 52·4	18·37	1	vB; p of D neb	2
386	I. 193	1 33 37·5	3·734	1	39 7 27·4	18·37	1	vB; f of D neb.....	1
387	145	VII. 46	1 34 28·0	4·062	2	28 49 19·8	18·34	2	Cl; iF; Ri; one * 6·7; st 11 ...14.	4
388	146	1 34 41·6	3·856	1	34 49 52·1	18·33	1	Cl; pRi; st 12, m.....	1
389	VIII. 65	C.H.	1 34 45·0	4·018	1	30 0 37·1	18·33	1	Cl; S; iRi; stL	1
390	II. 253	1 35 35·6	3·199	1	77 4 36·0	18·30	1	pB; pL; E; bM; r	1
391	147	II. 610	1 36 23·8	3·366	1	62 0 24·9	18·27	1	F; S; R; bM; r	3
392	VI. 31	1 36 29·2	4·055	1	29 27 40·9	18·27	1	Cl; B; L; eR; st pL	1
393	148	1 36 30·8	3·107	1	86 28 36·9	18·27	1	vF; S; R	1
394	II. 588	1 37 15·0	3·165	2	80 17 10·8	18·24	2	F; S; iE; bM; r	2
395	149	II. 611	1 39 33·1	3·365	2	62 48 58·2	18·16	2	F; S; iE 0°+	3
396	150	I. 157	1 40 0·8	3·360	2	63 16 25·8	18·14	3	F; E 0° ...90°; bet 2 st.....	4
397	II. 589	1 40 35·6	3·183	2	79 10 46·7	18·11	2	F; pL; E; lbM; *nf, 2'	2
398	D'Arrest, 27	1 41 29	3·30	[1]	68 21 3	18·08	[1]	pB; vmE; *14, f8 ^s	0
399	II. 228	1 41 34·5	3·299	2	68 42 50·6	18·08	2	pB; S; iR; mbM; 1st of 2 ...	2
400	151	IV. 42	1 41 37·9	3·126	2	84 47 12·6	18·08	2	vF; vmE, 165°±; sbM*9 ...	3†
401	III. 175	1 41 51·1	+3·482	1	54 53 51·9	-18·07	1	F; stellar	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
402	h.	H.		h m s	s		° ' "	"			
403	152	II. 229	1 41 57.7	+3.299	2	68 44 49.2	-18.06	2	pB; S; iR; mbM; 2nd of 2...	2
404	2436	II. 612	1 42 3.5	3.369	(1)	63 3 33.2	18.06	1	F; vLE, 90°; *15, nr	2
405	153	II. 481	1 42 17.0	2.959	1	101 7 2.5	18.05	1	pF; cL; R; glbM; S*p, 90''..	2
406	154	1 42 17.6	3.186	1::	79 0 17.5	18.05	1	eF	1
407	2438	II. 501	1 42 18.3	2.910	1	105 39 54.5	18.05	1	cF; S; R; gvlbMN.....	2
408	{ 155 = 2437 }	1 42 20.7	2.290	1	143 29 3.5	18.05	1	F; vL; R; gvlbM	1
409	III. 459	1 42 21.3	2.808	1	114 29 31.5	18.05	1	vF; vS; R; gbM; er; 2str ..	4
410	III. 561	1 42 25.7	3.492	1	54 21 52.8	18.04	1	vF; stellar	1
411	2439	II. 617	1 42 55.7	3.299	1	68 56 54.1	18.03	1	F; cL; vglbM.....	1
412	156	1 43 2.1	2.393	2	139 19 59.1	18.02	2	B; S; R; gbM	2
413	II. 859	1 43 10.9	3.129	1	84 33 6.4	18.02	1	pF; S; E90°; vglbM; *10, nf	2†
414	D'Arrest, 28	1 43 13	3.30	[1]	68 42 18	18.01	[1]	F; S; R; bet 2 st 15.....	0
415	2440	II. 618	1 43 29.0	3.309	1	68 7 54.7	18.01	1	vS; stellar	1
416	1 43 32.7	2.652	1	125 38 52.7	18.01	1	F; S; R	1
417	2441	III. 179	1 43 39.3	3.307	2	68 20 25.0	18.00	2	F; cL; E; mbM	2
418	{ 160 = 2442 }	1 43 42.0	2.653	1	125 34 5.0	18.00	1	eF; S	1
419	158	I. 62	1 44 8.0	2.964	1	100 23 45.6	17.98	2	F; pL; E; vglbM; r.....	4*
420	III. 192	1 44 17.0	3.024	2	94 44 57.9	17.97	3	eF; vLE 0° ±, *13, s, 90''.....	4
421	III. 564	1 44 25.9	3.496	1	54 35 56.9	17.97	1	Stellar; 3rd of 4	1
422	III. 565	1 44 25.9	3.496	1	54 35 56.9	17.97	1	Stellar; last of 4	1
423	157	III. 562	1 44 27.3	3.497	1	54 31 58.9	17.97	1	vF; stellar; 1st of 4.....	2
424	{ 157, a }	R. 3 novæ	1 44	54 32	Near h. 157, 159	0
425
426	161	II. 596	1 44 30.7	3.131	1	84 24 17.9	17.97	1	F; S; bM; *13 1', n	3
427	159	III. 563	1 44 34.3	3.497	1	54 33 1.9	17.97	1	F; pL; bM; 2nd of 4	2
428	162	55 Androm.	1 44 55.9	3.575	1	49 57 41.5	17.95	1	Fine nebulous * with strong atm.	1*
429	163	1 44 56.2	3.509	1	53 52 9.5	17.95	1	vF; R; am pBst	1
430	164	II. 270	1 45 56.9	3.110	2	86 29 23.7	17.91	2	pB; S; iR; psmbM.....	3
431	{ 165 = 2443 }	I. 105	1 46 12.4	2.918	2	104 25 39.0	17.90	2	cB; pL; lE; psmbM	3
432	D'Arrest, 29	1 46 19	3.28	[1]	70 50 18	17.89	[1]	eF; R; *19, f	0
433	D'Arrest, 30	1 46 32	3.56	[1]	51 18 3	17.88	[1]	eF; pL.....	0
434	D'Arrest, 31	1 47 5	3.29	[3]	69 59 18	17.86	[3]	vF; vS; R; β Arietis in field	0
435	{ 166 = 2444 }	III. 460	1 47 13.9	2.797	2	114 26 46.2	17.86	2	pF; vS; R; vgbM	3
436	167	1 47 14.6	2.795	1	114 33 21.2	17.86	1	vF; pL; R; gbM; S*195° ...	1
437	2445	1 47 43.1	2.621	1	126 34 24.8	17.84	1	F; S; R; bM	1
438	168	1 47 47.9	3.110	1	86 29 14.8	17.84	1	Suspected neb.....	1
439	2446	1 47 50.1	2.621	1	126 32 0.8	17.84	1	eeeF; S; R	1
440	III. 266	1 48 6.0	2.969	1	99 38 5.4	17.82	1	eF; stellar	1
441	{ 177 = 2447 }	III. 265	1 48 17.0	2.968	1	99 44 29.4	17.82	1	vF; pS; vLE	3
442	II. 221	1 48 26.0	3.463	1	57 38 16.7	17.81	1	F; pL; mE; r	1*
443	III. 176	1 48 33.4	3.483	1	56 23 6.7	17.81	1	eeeF; stellar	1
444	169	1 48 33.9	3.463	3	57 38 53.7	17.81	3	pB; R; bM; *13, np ..	3*
445	169, a	R. nova	1 48 34.5	3.463	...	57 38 25.7	17.81	...	{ β, γ, δ of Lord R.'s diag. ε=II. 221 H.	{ 0* 0* 0*
446	169, b	R. nova	1 48 38.4	3.463	...	57 38 0.7	17.81	...		
447	169, c	R. nova	1 48 47.9	3.463	...	57 40 30.7	17.81	...		
448	172	II. 271	1 49 2.8	3.126	1	85 3 29.3	17.79	1	pF; S; R; p of 2; pos=102°.4	1
449	173	II. 272	1 49 3.2	3.126	1	85 3 41.3	17.79	1	vF; vS; R; sbM; f of 2	4
450	170	1 49 6.3	4.127	1	30 30 49.6	17.78	1	Cl; not Ri; *	1
451	171	1 49 7.6	+3.953	1	35 13 13.6	-17.78	1	Cl; pL; pRi; iF; st 11...13..	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
452	h. 2449	H.	h m s 1 49 12.4	s +2.112	1	147 22 44.6	-17.78	1	pB; S; R; gbM	1
453	176	III. 193	1 49 20.4	3.017	2	95 9 13.6	17.78	2	eF; *9, 315° ±	3
454	2448	1 49 21.5	2.707	3	120 36 23.6	17.78	3	pB; S; E; 1M	3
455	175	II. 222	1 49 23.6	3.468	1	57 29 41.6	17.78	1	F; pL; mE; r; f of 2	3
456	175, a	R. nova	1 49	57 29	nf h. 175	0
457	174	VII. 32	1 49 25.8	3.541	1	53 1 14.9	17.77	1	Cl; vvL; Ri; st L and se ...	5
458	2450	1 49 27.7	2.107	1	147 26 49.9	17.77	1	vF; S; R; bM	1
459	{ 178 = } 2451	III. 464	1 49 57.4	3.006	3	96 5 19.5	17.75	3	vF; S; lE; vglbM	4
460	180	1 51 17.8	3.158	1	82 20 13.3	17.69	1	vF; S; R; *10, 2' 285°	1
461	D'Arrest, 32	1 51 35	3.28	[2]	71 42 48	17.68	[2]	vF; S; R; nr I. 112 H.	0
462	179	50 Cassiop.	1 51 36.4	5.005	1	18 15 36.6	17.68	1	Suspected nebulous *	1*
463	181	112	1 51 39.0	3.283	2	71 40 12.6	17.68	2	B; cL; R; gbM; r	3
464	181, a	R. nova	1 51 39	3.283	::	71 45 ±	::	5 or 6' s of h. 181	0
465	2452	III. 468	1 52 0.8	2.936	1	102 10 50.9	17.67	1	cF; pL; E 0° ±; glbM	2
466	III. 214	1 52 2.0	3.223	1	76 41 15.9	17.67	1	vF; stellar	1
467	2453	1 52 3.7	2.751	2	116 58 27.9	17.67	2	pF; S; R; glbM	1
468	D'Arrest, 33	1 52 5	3.34	[1]	67 2 18	17.66	[1]	F; pL	0
469	182	II. 223	1 52 10.0	3.451	1	59 15 3.2	17.66	1	pB; pL; R; glbM	2
470	183	I. 101	1 52 40.6	2.999	3	96 38 24.8	17.64	3	cB; L; mE 163° 0; mbM ...	6
471	III. 215	1 52 43.3	3.209	1	77 59 17.8	17.64	1	eF; stellar	1
472	184	III. 583	1 52 52.3	3.406	1	62 26 29.1	17.63	1	vF; vS; E; 3 stp; *250° ...	2*
473	2454	1 52 57.3	2.039	1	148 28 9.1	17.63	1	pB; pL; lE; *12 att	1
474	185	II. 435	1 54 2.1	2.989	(1)	97 30 22.6	17.58	1	pF; pS; R; bM	2
475	186	III. 433	1 54 20.6	3.005	1	96 3 38.9	17.57	1	cF; cS; R; bM	3
476	D'Arrest, 34	1 54 25	3.16	[1]	82 10 18	17.56	[1]	vF; S; *14 f 90°; 11.65 ...	0
477	187	1 54 39.4	3.247	1	74 57 35.2	17.56	1	eF; S; R; *11 75°	1
478	188	III. 207	1 54 49.2	3.280	1	72 17 54.5	17.55	1	vF; cS; R; stellar	2
479	2455	1 54 55.4	2.100	1	146 30 26.8	17.54	1	pF; S; R; 2 st 11, nr	1
480	2456	1 55 1.3	0.699	3	164 54 18.8	17.54	3	eF; vS; R; *12, 25" 315° ...	3
481	189	III. 566	1 55 8.5	3.571	1	52 33 52.1	17.53	1	vF; S; iR; sbM; *nr	3
482	2457	1 55 46.1	1.428	1	158 32 40.7	17.51	1	eeF; vS; R; *13 p 100" ...	1
483	190	III. 208	1 56 8.7	3.253	1	74 38 1.3	17.49	1	vF; S; iR; glbM; *10 p 3s.5	2
484	191	III. 151	1 56 54.9	3.428	1	61 40 47.2	17.46	1	vF; vS; iR; bet2stn and sp..	3
485	{ 192 = } 2458	1 57 28.5	2.780	2	113 58 18.1	17.43	2	vF; pS; vLE	2
486	2459	1 58 26.4	1.343	1	159 7 19.3	17.39	1	pF; S; R; gbM	1
487	193	I. 152	1 59 51.2	3.196	1	79 40 47.1	17.33	1	B; vS; vLE; svmbM; *10, 55" 320°.	3*
488	194	II. 604	2 0 19.0	3.604	1	51 54 21.7	17.31	1	pB; cL; lE; mbM	2
489	195	2 0 49.6	3.238	1	76 18 58.6	17.28	1	F; R; vS; bM	1
490	2461	2 0 55.4	2.468	2	131 49 14.6	17.28	2	cF; vS; R; sbM; r	2
491	{ 196 = } 2640	2 0 56.9	2.741	2	116 7 18.6	17.28	2	vF; vF * inv	2
492	2462	2 0 58.2	2.561	1	127 9 11.6	17.28	2	F; S; R; vsymbM*13	2
493	198	III. 227	2 1 34.0	3.161	1	82 41 44.2	17.26	1	vF; S; R; bM; am st	2
494	197	II. 605	2 1 40.3	3.617	1	51 28 53.5	17.25	1	pB; S; iR; * f 15s	2
495	{ 199 = } 2463	II. 482	2 2 30.1	2.942	3	100 47 38.7	17.21	3	F; S; R; 1st of 4	4
496	III. 567	2 2 32.5	3.591	1	53 0 41.7	17.21	1	vF; S; lE	1
497	{ 200 = } 2464	II. 483	2 2 32.8	2.942	3	100 47 50.7	17.21	3	F; S; R; 2nd of 4	4
498	{ 201 = } 2465	II. 484	2 2 47.1	+2.942	3	100 48 31.0	-17.20	3	vF; vS; R; 3rd of 4	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
499	{ h. 202 = 2466 }	II. 485	h m s 2 2 51.4	s +2.941	2	° ' " 100 51 11.0	" -17.20	3	vF; pS; R; 4th of 4	4
500	203	2 2 55.0	2.971	1	98 25 7.3	17.19	1	vF; vS; R; psbM	1
501	204	III. 604	2 3 56.7	3.593	1	53 10 55.5	17.15	1	vF; iF; stellar	2*
502	2467	2 4 15.8	1.992	1	147 23 34.1	17.13	1	pF; pS; R; glbM; r	1
503	III. 259	2 4 27.5	3.047	1	92 9 47.4	17.12	1	eF; eS; iF	1
504	II. 486	2 4 45.7	2.951	1	99 58 5.7	17.11	1	F; S; E	1
505	2468	2 5 36.7	2.555	3	126 30 33.9	17.07	4	cF; pS; lE 0°; gbM	4
506	II. 613	2 5 51.5	3.435	1	62 46 51.2	17.06	1	F; S; lE 90°; bM	1
507	2469	2 6 30.8	2.623	2	122 36 11.1	17.03	2	cB; S; E; psmbM	2
508	2470	2 7 24.0	2.420	2	132 41 12.3	16.99	2	F; vS; svmbM	2
509	205	III. 260	2 7 24.4	3.056	1	91 24 41.3	16.99	1	vF; R; bM; stellar	2
510	206	III. 457	2 8 6.4	3.140	1	84 39 32.5	16.95	1	eF; cL; R; gbM; *12 sf att.	2*
511	III. 2	2 8 41.3	3.078	1	89 36 2.1	16.93	1::	eF; vS; R; bM	1
512	207	VI. 33	2 9 15.3	4.166	2	33 29 55.0	16.90	2	!; Cl; vL; vRi; st7...14 ...	5
513	208	III. 201	2 9 33.1	3.252	1	76 5 52.3	16.89	1	vF; vS; E; *10 sf 4'	2
514	208, a	R. nova	2 9	76 5+	neb s of h. 208	0
515	{ 209 = 2471 }	II. 474	2 9 43.2	2.920	5	101 59 58.6	16.88	5	F; pL; R; vglbM	7
516	210	II. 246	2 10 22.6	3.253	1	76 6 11.5	16.85	1	pF; pL; lE; pgbM; { *9, 185° ± 5' S* sf 1'	2*
517	210, a	R. nova	2 10	76 6	neb s of h. 210	0
518	211	II. 436	2 11 14.5	2.980	1	97 17 24.7	16.81	1	F; pS; E; bM; 2 or 3 st nr...	2
519	213	2 11 59.3	3.272	1	74 48 57.9	16.77	1	eF; R; gbM; *16 nr	1
520	215	II. 437	2 12 8.7	2.978	2	97 25 44.2	16.75	2	pF; pS; vLE; bM; * nr	3
521	212	VI. 34	2 12 34.2	4.188	2	33 32 49.5	16.75	2	!; Cl; vL; vRi; ruby * M ...	5
522	214	2 12 46.6	4.536	1	26 52 20.1	16.73	1	Cl; L; lC; sc st 9...13	1
523	216	III. 486	2 12 54.3	2.852	1	106 42 4.1	16.73	1	F; S; iR; pgbM	2
524	2473	2 13 20.1	1.774	1	150 30 8.7	16.71	1	vF; S; R; 2 or 3 vSst nr ...	1
525	2472	2 13 30.8	2.400	2	132 23 1.0	16.70	2	eF; vS; R; bM; *7 sf and 6 more.	2
526	217	II. 225	2 13 41.2	3.548	4	57 22 48.3	16.69	4	B; S; R; bM; 3S st sp	5
527	218	V. 19	2 13 50.4	3.737	1	48 16 47.6	16.68	1	!; B; vL; vmE 22° 3'	5+
528	2474	2 14 21.4	2.404	2	132 2 58.2	16.66	2	pF; pS; R; lbM; *8 90°, 4'	2
529	219	II. 438	2 14 35.7	2.993	1	96 10 8.8	16.64	2	F; vL; iR; gbM	4
530	219, a	R. nova	2 14	96 10	E; F; bM; makes D neb with h. 219; both E.	0
531	III. 695	2 15 1.0	4.448	1	28 40 47.1	16.63	1	eF; pL; iF	1
532	2475	2 15 9.4	2.563	1	124 21 30.4	16.62	1	pB; S; R; psbM; *10f 90° 0 35''.	1
533	III. 570	2 15 14.0	3.732	2	48 42 17.4	16.62	1	eF; vS; lE	1
534	2476	2 15 26.4	2.779	1	111 27 12.0	16.60	1	pB; S; gbM; r; *p	1
535	2477	III. 224	2 16 34.0	2.778	1	111 20 33.5	16.55	1	F; S; E 90°; gbM	3
536	I. 153	2 16 36.1	2.770	1	111 52 23.5	16.55	1	cB; vL; E 0°...90°	1*
537	III. 571	2 16 44.2	3.736	1	48 49 21.5	16.55	1	eF; stellar	1
538	221	2 17 49.4	3.319	1	72 7 46.3	16.49	1	pF; L; R; *10 sf 3'	1
539	220	2 17 50.6	3.543	1	58 23 33.3	16.49	1	vF; S; R; 4 st nr	1
540	2478	III. 239	2 18 44.7	2.710	1	115 26 13.5	16.45	1	eF; pL; R; gpmbM	2
541	III. 474	2 18 50.8	3.350	1	70 7 27.8	16.44	1	eF; vS; iR	1
542	222	III. 177	2 18 55.7	3.571	2	57 3 25.1	16.43	2	cF; cL; E; vgbM; 2st13np..	3
543	II. 489	2 19 58.5	3.350	1	70 15 30.6	16.38	1	F; S; lE; 3 st inv	1
544	223	IV. 23	2 20 29.1	3.049	1	91 46 18.5	16.35	1	vB; vL; R; mbMN	2+
545	2479	2 20 59.9	2.295	2	135 4 17.1	16.33	2	vvF; S; R; gvlbM	2
546	224	III. 261	2 21 21.5	3.049	1	91 47 36.7	16.31	1	vF; cL; R; f of 2	2
547	{ 225 = 2480 }	II. 487	2 21 48.4	2.919	2	101 10 10.3	16.29	2	vF; L; iR; glbM	3
548	2481	2 22 1.5	+2.795	1	109 40 3.6	-16.28	1	pB; E; gbM	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
549	h. 226	H. I. 154	h m s	s	1	53 29 51.2	-16.26	1	cB; L; E; vgbM	3*
550	2482	2 22 18.6	+3.654	4	132 1 30.7	16.21	4	vF; pL; lE; gbM; *8sf 3'...	4
551	229	II. 278	2 23 20.3	2.366	1	91 43 1.7	16.21	1	pB; S; E; psbM.....	3
552	228	2 23 24.6	3.050	1	45 59 37.0	16.20	1	Cl; pRi; st 9...15	1
553	227	2 23 27.5	3.837	1	33 5 46.0	16.20	1	Cl; pL; pRi; st 13...15	1
554	230	II. 237?	2 23 28.9	4.288	1	93 33 56.3	16.19	1	pF; iE 0°±; bM	2
555	2483	2 25 38.8	3.024	2	126 39 0.0	16.10	4	pB; pS; mE 215°.7	4
556	2484	2 23 21.5	2.482	2	107 49 36.8	16.09	2	F; S; iR; gbM	2
557	231	2 25 38.1	2.817	1	57 40 44.3	16.09	1	S; R; psbM; 1st of 3	1*
558	231, a	R. nova	2 25 44.2	3.580	...	57 38 59.3	16.09	...	} γ and δ of Lord R.'s diag.	{ 0*
559	231, b	R. nova	2 25 47.7	3.580	...	57 38 50.3	16.09	...		
560	232	II. 211	2 25 52.0	3.580	(2)	61 17 57.6	16.08	(2)	pB; cL; lE 0...90°; gmbM; 3st s.	3†
561	233	2 25 52.4	3.512	1	57 40 14.9	16.07	1	vF; R; bM; 2nd of 3	1*
562	2485	III. 472	2 26 0.7	3.580	1	101 22 37.5	16.05	1	eF; pS; R; vlbM; amscst...	2
563	234	2 26 16.8	2.912	1	57 46 23.5	16.05	1	pB; R; 3rd of 3	1*
564	2486	2 26 21.2	3.580	1	135 8 21.8	16.04	2	F; S; R; bet 2 st in par.....	2
565	235	III. 572	2 26 26.9	2.270	1:	49 47 23.8	16.04	1:	vF; pS; p of 2; 210''; 157°...	2
566	236	III. 573	2 26 33.1	3.753	1	49 44 28.1	16.03	1	F; S; f of 2; 210''; 337° ...	2
567	2487	Δ. 519??	2 26 37.4	3.754	2	129 39 17.2	15.96	2	pB; L; pmE; smbM; bi-N...	2†
568	237	III. 161	2 28 0.1	2.404	2	57 17 16.5	15.95	2	F; S; vIE; bM; r; 2 st 14 np	4
569	238	III. 557	2 28 23.1	3.595	1	78 58 34.1	15.93	1	F; S; vIE; psbM; r	2
570	239	III. 434	2 28 43.8	3.232	1	97 46 30.4	15.92	1	vF; cL; iF; vlbM	2
571	240	II. 238 = III. 198	2 28 46.2	2.963	1	49 43 53.8	15.84	1	pF; L; E 90°±; mbM; r ...	4*
572	241	III. 152	2 30 24.9	3.761	1:	60 27 46.0	15.80	1	F; pS; iR; bM; st inv	3†
573	II. 6	2 30 56.6	3.540	1	89 44 6.9	15.77	1	S; cometic	1*
574	244	I. 102	2 31 35.3	3.077	3	97 17 18.9	15.77	3	cB; pL; vR; mbM.....	5*
575	242	I. 156	2 31 36.0	2.968	1	51 32 45.9	15.77	1	vB; vL; vmE; vvmB	3†
576	243	II. 592	2 31 38.3	3.731	1	79 45 46.9	15.77	1	pF; S; lE; bM; *11, 25' 50°	2
577	2488	2 31 38.5	3.222	1	145 28 45.9	15.77	1	eF; S; R; p of 2.....	1
578	VIII. 66	2 31 43.4	1.877	2	29 3 23.5	15.75	2	Cl; L; sc st, one 10.....	2
579	245	III. 581	2 32 1.1	4.572	1:	72 34 8.5	15.75	1:	vF; iE.....	2
580	2490	2 32 2.2	3.333	1	145 28 16.8	15.74	1	F; S; R; gbM; *11, s 2' ...	1
581	246	II. 5	2 32 8.3	1.874	2	89 30 24.8	15.74	2	pB; S; vIE 0...90°; bm; 3st trap.	10
582	{ 249 = 2489 }	II. 284	2 32 11.6	3.080	2	98 44 27.7	15.71	2	pF; L; mE; r; *17, att sf...	4
583	247	III. 475	2 32 36.5	2.945	(1)	71 19 38.7	15.71	1	F; S; R; lbM.....	2
584	248	M. 34	2 32 42.1	3.354	2	47 49 25.0	15.70	2	Cl; B; vL; lC; sc st 9	8
585	251	III. 228	2 33 2.3	3.829	1	81 52 7.2	15.66	1	vF; vS; p of 2; *10 p	2
586	{ 253 = 2491 }	II. 488	2 33 36.3	3.193	2	101 53 28.2	15.66	2	F; S; R; bM	3
587	252	III. 229	2 33 42.9	2.898	1	81 53 22.2	15.66	1	eF; vS; f of 2.....	2
588	2492	2 33 43.8	3.192	1	124 52 20.8	15.64	1	pB; S; R; stellar	1
589	{ 254 = 2493 }	I. 63	2 33 57.9	2.491	3	98 51 19.1	15.63	3	B; pL; R; mbM*12	4
590	256	III. 584	2 34 12.1	2.943	1:	62 1 17.0	15.60	1	F; S; R; psbM	2
591	258	I. 1	2 34 35.0	3.520	2	90 9 14.0	15.60	2	pF; cL; iE 80°; bM; pB*nr	8*
592	255	II. 633	2 34 35.1	3.072	1	53 16 21.0	15.60	1	pF; cL; R; glbM	3
593	259	2 34 39.9	3.701	1	72 35 27.0	15.60	1	eF; ?	1
594	257	III. 162	2 34 44.5	3.337	2	58 10 18.3	15.59	2	F; pL; R; lbM; *7.8 p 43°.5	4
595	} 257, a	R. 3 novæ	2 34 49.1	3.596	...	58 10	{ 6 seen (including .: h.) } { 257, 260, 261). }	0
596		2 34
597	260	III. 163	2 35 23.6	3.598	1	58 7 34.9	15.57	1	vF; pL; R; lbM; sp of 2 ...	2
598	261	2 35 26.2	+3.599	1	58 4 35.2	-15.56	1	eF; S; nf of 2.....	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
600	h. 262	H.	M. 77	h m s 2 35 30.2	s +3.065	1	90° 35' 56".2	-15.56	1	vB; pL; iR; sbMrrN; *130°, 2'	13†
601	263	II. 273	2 36 0.3	3.138	1	85 37 19.1	15.53	1	pF; S; iR; gbM	2
602	III. 455	2 36 33.3	3.085	2	89 13 22.0	15.50	2	vF; L; lbM; er	2
603	2494	2 37 43.7	2.589	1	119 35 49.8	15.44	1	B; pL; pmE; sbM	1
604	264	I. 64	2 39 6.9	2.950	1	98 9 52.5	15.35	1	vB; pL; E; gpmbM	3
605	265	II. 466	2 39 15.4	3.057	1	91 5 53.8	15.34	1	pB; cL; lE; mbM	4
606	266	II. 465	2 39 21.1	3.061	1	90 49 5.8	15.34	1	vF; pL; iR; bM	4
607	III. 582	2 39 22.7	3.313	1	74 25 31.8	15.34	1	vF; S; iF	1
608	267	III. 462	2 40 18.4	3.061	1	90 50 57.3	15.29	1	F; S; R; 2Sst p	2
609	2496	2 40 20.8	1.552	1	150 30 14.3	15.29	1	F; pS; R; glbM	1
610	2495	V. 48	2 40 21.5	2.557	2	120 51 30.3	15.29	2	vB; L; vmE 151°.1; vbMN	3
611	{ 269 = 2497 }	III. 449	2 42 35.2	2.796	2	107 34 23.5	15.15	2	pF; pL; pmE; glbM	3
612	268	2 42 39.0	3.839	1	48 55 4.5	15.15	1	neb or vScl of vSst	1
613	270	II. 601	2 43 44.5	3.858	1	48 22 21.3	15.09	1	cF; S; iR; vgbM; r	2
614	Bessel	2 44 6.5	3.738	...	53 4 14.9	15.07	...	? a comet	0*
615	272	III. 450	2 45 8.9	2.799	1	107 12 50.7	15.01	1	vF; S; lE; gbM	3
616	271	II. 602	2 45 25.7	3.847	1	49 0 2.3	14.99	1	cF; pS; iR; vglbM	2
617	{ 271, a }	R. 2 novæ	2 45	49	h. 271 is D; another near	0.
619	273	2 45 44.4	3.044	1	91 51 24.6	14.98	1	eF; pL; gbM; *8f	1
620	II. 254	2 46 0.9	3.269	1	77 34 54.2	14.96	1	F; S; iR; r	1
621	2498	2 46 35.8	1.778	1	145 32 30.4	14.92	1	F; R; gbM	1
622	2499	2 46 44.5	1.772	1	145 38 31.4	14.92	1	F; R; gbM	1
623	274	III. 580	2 47 29.7	3.897	1	47 31 21.6	14.88	1	vF; vS; R; gbM; 2Sst Δ	2
624	{ 275 = 2500 }	II. 470	2 47 46.2	2.905	2	100 36 9.2	14.86	2	vF; S; R; stellar	4
625	2501	2 48 7.7	2.760	1	109 13 6.8	14.84	1	F; pL; vmE; 2Sst f	1
626	276	II. 274	2 50 52.8	3.118	1	87 11 7.6	14.68	1	F; vS; iE; sbM; er	2
627	II. 619	2 51 35.0	3.493	1	65 21 16.1	14.63	1	pB; cL; pmE 0°; r; *n 1'	1
628	277	III. 199	2 51 54.7	3.949	1	45 40 24.7	14.61	1	F; pS; lE; SbM; *p 6°.5	4
629	277, a	R. nova	2 52 +	45 40 ±	R (nisi = H. II. 239)	0
630	2502	III. 469	2 52 16.6	2.863	1	102 57 33.3	14.59	1	F; R; gbM; stellar	2
631	III. 178	2 52 47.8	3.711	1	55 18 19.2	14.56	1	vF; pL; R; spmbM	1
632	278	2 52 51.3	3.908	1	47 58 42.2	14.56	1	eF; vS	1
633	2503	2 52 56.7	2.484	2	122 39 22.5	14.55	2	vF; pL; E; vlbM	2
634	II. 239?	2 53 0.1	3.977	2	45 35 20.5	14.55	2	pB; pL; iR; mbM	3
635	279	II. 620	2 53 28.9	4.025	(2)	44 10 52.4	14.52	(2)	pF; pS; iF; sbM	2
636	280	II. 502	2 55 0.2	2.818	1	105 23 8.1	14.43	1	pF; pL; R; psbM	2*
637	II. 607	2 55 21.2	3.906	1	48 13 26.7	14.41	1	F; cL; E	1
638	II. 704	2 55 44.9	8.608	1	9 42 40.3	14.39	1	F; pL; mE 90°...180°	2
639	281	IV. 43	2 56 20.1	3.925	1	47 43 32.5	14.35	1	F; mE; smbMS*	3*
640	2504	III. 245	2 56 25.0	2.669	1	113 25 4.8	14.34	1	pF; cL; pmE; gbM*16; r...	2
641	II. 608	2 56 32.2	3.975	1	46 10 29.0	14.40	1	F; cL; er	1
642	2505	2 56 55.1	2.864	1	102 38 19.0	14.30	1	vF; sp of 2	2
643	282	II. 503	2 57 5.8	2.802	1	106 8 59.3	14.29	1	cB; pS; iR; smbM	2
644	2506	II. 475	2 57 14.1	2.865	1	102 33 6.3	14.29	1	pF; cL; iR; bM; nf of 2	3
645	283	I. 109	2 58 3.7	2.603	1	116 35 45.8	14.24	1	cB; pS; vIE 0°; r; S*nr	4
646	284	III. 578	2 59 19.7	3.810	1	52 9 24.2	14.16	1	cF; vS; R; psbM	2*
647	{ 285 = 2507 }	II. 285	2 59 25.2	2.905	3	100 5 8.5	14.15	3	pB; S; lE 80°±; lbM	7
648	286	II. 504	2 59 30.2	2.799	1	106 8 39.5	14.15	1	B; S; cE; psbM	2
649	2508	3 0 48.0	2.288	1	129 34 26.9	14.07	1	pF; S; R; psbM	2
650	287	3 1 29.2	4.337	1	37 11 56.1	14.03	1	Cl; vS of Sst	1
651	2509	II. 258	3 3 27.9	2.702	3	111 7 0.0	13.90	3	pB; cL; R; gbM; r	7
652	288	III. 262	3 4 20.1	3.022	(1)	93 3 4.5	13.85	1	:: stellar; difficult	2
653	III. 164	3 4 23.3	+3.634	1	59 57 41.8	-13.84	1	eF; vS; ? vSst	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
654	{ 289 = 2510 }	II. 286	3 4 26.8	+2.912	3	99 27 28.8	-13.84	3	F; pL; R; vglbM; *9np ...	5
655	289, a	R. nova	3 4 31.4	2.912	...	99 26 25.8	13.84	...	No description.....	0*
656	{ 291 = 2511 }	III. 591	3 4 36.2	2.912	2	99 29 11.6	13.78	2	eF; vS; R; stell; sf of 2.....	3*
657	2512	Δ. 205??	3 4 41.9	0.771	2	157 18 47.1	13.83	2	F; S; pmE; gbM	3
658	290	VI. 25	3 5 6.3	4.106	1	43 17 22.3	13.79	1	Cl; pL; Ri; C; iR; st 12...15	3
659	2513	3 5 18.2	0.747	2	157 29 13.6	13.78	2	pF; S; R; glbM	2
660	II. 900	3 5 38.6	2.884	1	101 0 7.2	13.76	1	F; pL; E 80°±	1
661	292	III. 443	3 5 49.8	2.975	3	95 45 4.5	13.75	3	cF; S; lE; bM; *9, n 5' ...	4
662	2514	3 5 57.7	1.775	1	143 52 15.8	13.74	1	B; L; vmE 80°; vgbM	2
5060	3 6 55.2	89 4 27.5	See No. 5060.	
663	2515	3 7 7.5	1.474	1	148 40 18.9	13.67	1	Cl of 18 or 20 st.....	1
664	IV. 17	3 7 7.6	3.016	1	93 27 10.9	13.67	1	* with neb att 90° 1	1
665	2516	3 7 46.9	2.667	1	112 30 46.1	13.63	1	F; S; E; alm stell; *8, np...	1
666	2517	Δ. 337	3 8 25.7	1.635	2	145 44 50.6	13.58	2	⊕; B; L; R; rr	2
667	III. 194	3 8 43.9	3.020	1	93 8 18.9	13.57	1	eF; eS	1
668	D'Arrest, 35	3 9 31	3.93	[2]	49 2 30	13.49	[2]	F; vS; R; stellar; 1st of 7...	0
669	D'Arrest, 36	3 9 31	3.93	[2]	49 1 42	13.49	[2]	eF; S; lE; cometary; 2d of 7	0
670	2518	3 9 41.5	2.198	1	131 36 25.7	13.51	1	vB; R; gmbM	1
671	D'Arrest, 37	3 9 45	3.93	[2]	49 1 54	13.48	[2]	vF; S; R; 3rd of 7.....	0
672	D'Arrest, 38	3 10 8	3.93	[2]	49 1 36	13.46	[2]	F; S; R; 4th of 7	0
673	D'Arrest, 39	3 10 11	3.94	[2]	48 58 30	13.45	[2]	vF; vS; 5th of 7.....	0
674	293	II. 603	3 10 32.7	3.937	1	49 0 5.5	13.45	1	pB; pS; R; bM; 6th of 7 ...	2*
675	D'Arrest, 40	3 10 35	3.97	[2]	49 0 3	13.42	[2]	F; S; *17; 7th of 7	0
676	2519	III. 956	3 11 2.0	2.884	1	100 48 41.4	13.42	1	eF; vS; 2 st 2' or 3' s.....	2
677
678
679	{ 293, a }	R. 6 novæ	3 11 ±	49 ±	6 of 15 (including probably h. 294, 295)	0
680											
681											
682
683	2520	3 11 34.2	2.425	1	123 5 43.3	13.39	1	vF; L; R; vglbM	1
684	III. 195	3 11 43.7	3.017	1	93 14 30.6	13.38	1	eeF; eS	1*
685	2521	Δ. 487	3 12 15.0	2.189	2	131 36 47.1	13.33	2	⊕; vB; pL; R; mbM; er ...	2
686	294	III. 574	3 12 23.9	3.938	1	49 11 42.1	13.33	1	vF; R; bM; p & sm of 2 ...	2
687	295	III. 575	3 12 25.0	3.938	1	49 10 4.4	13.32	1	vF; R; bM; f of 2; 100'', 352°4.	2
688	296	II. 287	3 13 15.0	2.954	4	96 45 57.9	13.27	4	vF; S; vLE; gbM; er	7
689	2522	3 13 22.5	2.710	2	110 55 8.9	13.27	2	cB; vL; vmE; psvmbM.....	2
690	III. 444	3 14 7.5	2.983	1	95 8 11.7	13.21	1	eF; vS	1
691	III. 568	3 15 25.7	3.016	1	93 15 44.1	13.13	1	eF; S; iF; am 3 or 4 st	1
692	2523	I. 106	3 15 36.2	2.786	1	105 53 51.7	13.11	1	pB; cL; iR; gbM; *7, f 7 ^s .5, 211°0.	3
693	2524	3 15 41.5	2.295	3	127 38 38.7	13.11	3	⊕; vF; pL; R; vgvlbM.....	3
694	2525	3 16 0.3	1.749	1	142 41 0.3	13.09	1	F; pL; mE 37°3; gbm	1
5061	3 16 29.0	89 19 3.0	See No. 5061.	
695	2528	Δ. 206	3 16 34.1	0.698	1	156 59 37.2	13.06	1	pB; L; lE; vgbM; r	1
696	2526	3 16 54.8	2.666	1	111 52 19.1	13.03	1	pB; S; R; gbM	1
697	2527	Δ. 548	3 17 20.4	2.288	2	127 43 40.0	13.00	2	vB; cL; vLE; vsvmbMN ...	2
698	2529	Δ. 547	3 17 23.1	2.291	2	127 36 24.0	13.00	2	pB; pS; psbM.....	2
699	2533	3 17 45.6	2.662	1	112 1 0.9	12.97	1	F; S; R; bM; p of 2	1
700	{ 298 = 2530 }	III. 197	3 17 48.3	3.011	1	93 31 34.9	12.97	1	vF; S; R; bM; 1st of 3	3
701	{ 297 = 2531 }	III. 196	3 17 48.8	3.011	1	93 30 34.9	12.97	1	vF; vS; E; ? neb *; 2nd of 3	3
702	2532	3 17 50.9	+3.012	1	93 25 54.9	-12.97	1	F; vS; R; bM; 3rd of 3.....	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
703	h. 297, a	H.	R. nova	h m s 3 17 53·6	s + 3·011	...	93 16 41·9	-12·97	0
704	299	III. 445	3 18 4·9	2·962	2	96 14 11·2	12·96	2	vF; pS; pmE	3
705	2534	IV. 77	3 18 18·0	2·661	2	112 1 49·8	12·94	2	F; mE; 239°·1; com; *9, 10 att.	3+
706	2535	3 18 37·6	2·305	1	126 58 12·7	12·91	1	O? pS; vsymbMN	1
707	2536	3 19 39·5	2·739	1	108 4 59·5	12·85	1	F; pS; R; glbM	1
708	III. 959	3 19 53·5	2·662	2	111 51 1·8	12·84	2	vF; vS; sf of 2	1*
709	I. 60	3 20 15·0	2·662	2	111 50 36·0	12·80	2	vB; S; E90°...180°; symbMN; np of 2.	2*
710	Auw. N. 17	3 20 41·7	3·690	...	59 6 31·2	12·76	...	F; L; *10f 4 ^s ; n 2'·5 (Schönfeld, 1858).	0*
711	D'Arrest, 41	3 20 47	3·98	[1]	48 39 6	12·72	[1]	eF; pL; lbM	0
712	2537	3 21 10·0	2·319	3	126 12 33·5	12·75	3	vF; S; vLE; gbM	3
713	2538	3 22 29·8	2·407	2	122 46 15·5	12·65	2	cB; pS; R; psbM; *p	2
714	2542	I. 257	3 22 38·1	2·437	(1)	121 34 40·8	12·64	1::	cB; pL; iR; vgbM	2
715	2539	3 22 38·5	2·441	1	121 23 12·8	12·64	1	vB; pS; lE; psbM	1
716	2540	3 22 39·6	2·275	1	127 38 11·8	12·64	1	F; S; R; *12, sf	1
717	301	VIII. 88	3 22 41·0	3·852	1	53 9 36·8	12·64	1	Cl; vL; ab 60 st	3
718	300	III. 694	3 22 54·4	6·312	2	17 54 18·1	12·63	2	vF; vS; iR; gbM; *vnr	3
719	2541	3 23 10·3	2·732	1	108 16 10·7	12·61	1	F; S; R; psbM	1
720	VIII. 84	3 23 42·3	4·361	1	39 3 16·6	12·58	1	Cl; lRi; stL	1
721	2545	Δ. 591	3 24 11·0	2·365	1?	124 12 43·8	12·54	1::	B; L; mE; vmbMRN	1
722	2544	3 25 8·3	2·332	1	125 20 9·9	12·47	1	pB; pS; R; psbM	1
723	2543	3 25 16·8	2·698	1	109 45 38·2	12·46	1	eF; psbM; v diff *8, sf	1
724	2546	III. 246	3 25 52·1	2·665	2	111 17 46·1	12·43	2	pB; cL; iE; mbM	5
725	2547	III. 487	3 26 2·0	2·778	1	105 41 32·4	12·42	1	vF; S; lE; glbM	2
726	2548	II. 290	3 26 44·0	2·808	2	104 9 24·9	12·37	2	pF; pL; R; lbM; pL*nf 5'	5
727	302	III. 446	3 26 44·3	2·971	2	95 33 27·9	12·37	2	vF; S; bet 2st	3
728	2549	3 26 30·7	1·789	2	140 45 56·6	12·38	2	vF; pL; iR; gbM; *nr	2
729	2550	3 27 28·6	2·690	1	109 58 38·4	12·32	1	F; L; R; vglbM	1
730	2551	III. 960	3 27 38·5	2·674	2	110 46 3·7	12·31	2	vF; S; R	3
731	2552	3 28 18·0	2·289	2	126 36 31·9	12·27	2	!! vB; vL; mE; rN in vLE Halo.	2+
732	2553	III. 857	3 28 18·2	2·421	(1)	121 40 37·9	12·27	(1)	vF; S; iF; lbM	2
733	2554	III. 559	3 29 0·5	2·670	3	110 50 45·4	12·22	3	vF; S; R; bet 2st 14	4
734	2555	II. 262	3 29 1·6	2·570	1	115 24 21·4	12·22	1	pB; pL; vLE; psbM	2
735	2556	3 29 50·6	2·310	1	125 42 24·2	12·16	1	eF; vS; p of 3	1
736	2557	3 30 0·6	2·311	1	125 41 4·5	12·15	1	vB; pL; lE; gmbM; 2nd of 3	1
737	2558	3 30 0·6	2·309	1	125 43 44·5	12·15	1	B; S; lE; pmbM; 3rd of 3...	1
738	303	II. 288	3 30 7·0	2·971	2	95 30 0·1	12·13	2	eF; pL; iR; bM; r	5
739	2559	Δ. 574	3 30 9·5	2·316	1?	125 28 56·8	12·14	1::	vB; L; R; psbM	1
740	2560	III. 961	3 30 28·7	2·657	2	111 21 44·4	12·12	2	F; S; R; gbM	3
741	2561	3 30 42·7	2·307	1	125 55 1·7	12·11	1	⊕; B; pL; R; gpmbM	2
742	2562	3 31 19·8	2·710	1	108 48 18·5	12·05	1	pF; S; R; psbM	1
743	2563	II. 263	3 31 26·9	2·576	1	114 58 4·5	12·05	1	pB; pS; R; gpmbM	2
744	2564	3 31 34·7	2·298	1	125 58 33·5	12·05	1	⊕; vB; pL; R; gmbM	2
745	2565	III. 451	3 32 19·8	2·706	1	108 53 34·6	11·98	1	F; S; R; glbM	2
746	2566	I. 58	3 32 24·1	2·608	2	113 28 55·6	11·98	2	B; pS; E; psbM	4
747	2567	II. 593	3 33 11·0	2·700	1	109 9 29·4	11·92	1	cB; pS; R; psbM	2
748	2569	3 33 11·6	2·296	2	125 54 53·1	11·93	2	⊕; vB; pL; psbM; rr	3
749	2568	III. 247	3 33 14·1	2·613	1	113 10 47·4	11·92	1	vF; vS; R	2
750	2571	3 33 43·1	2·291	1	126 2 48·0	11·90	1	vB; pL; R; psbM	2
751	2572	3 33 49·8	2·407	1	121 46 25·6	11·88	1	F; cL; vmE; vglbM; *7np	1
752	2570	I. 107	3 33 55·4	2·702	1	109 2 21·9	11·87	1	vB; L; R; vsymbMN	3
753	304	III. 263	3 34 4·2	3·040	(2)	91 45 34·5	11·85	(2)	eF; stellar or lE	2
754	304, a	R. nova	3 34	91 45	makes D neb with h. 304	0
755	2573	3 34 5·2	2·014	2	134 33 18·9	11·87	2	B; pS; R; symbM	2
756	305	III. 569	3 34 32·3	2·977	1	95 7 5·4	11·82	1::	eF; lE; er; 1st of 3	2
757	2574	3 34 36·5	2·531	1	116 40 8·4	11·82	1	F; S; E; gmbM; *sf 2'	1
758	306	II. 455	3 34 41·5	2·976	1	95 8 51·7	11·81	1	pF; pL; lE; lbM; *sf; 2d of 3	3
759	2575	II. 267	3 34 51·1	+ 2·615	3	113 0 43·7	-11·81	3	pB; S; lE; pglbM; *sf 2' ...	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
760	h. 307	H. II. 456	h m s 3 35 2.3	s + 2.976	3	95° 9' 45.6"	-11.78	2	vF; S; E; B* 135°, 1'; 3d of 3	4
761	2576	3 35 29.9	2.229	2	127 58 34.9	11.77	2	pF; pS; R; psbM	2
762	2577	II. 291	3 35 56.9	2.804	2	103 56 56.1	11.73	2	F; cL; mE 0° ±; r.....	4
763	307, a	R. nova	3 35 58.6	2.976	::	95 9 45.6	11.73	::	No description	0
764	II. 852	3 36 30.1	2.438	1	120 21 11.3	11.69	1	F; pL; iR; gbM	1
765	2578	III. 248	3 36 42.1	2.622	2	112 33 21.6	11.68	2	pF; S; lE; bM	3
766	2579	3 37 0.0	2.288	1	125 50 50.2	11.66	1	pF; S; R; psmbM	1
767	2580	Δ. 426	3 37 37.8	1.879	2	137 40 33.4	11.62	2	vB; L; pmE; vsvmbM*10...	2
768	Auw. N. 18	3 37 52.3	3.542	...	66 40 12.9	11.57	...	!!!B; vL; iF; VAR. (Tempel)	0*
769	2581	Δ. 562	3 38 7.9	2.264	1?	126 34 12.6	11.58	1?	⊕; vB; pmE; pgbM	2
770	2582	3 38 20.0	2.272	1	126 18 1.9	11.57	1	F; vL; R; gbM	1
771	2584	III. 249	3 38 43.4	2.623	3	112 21 40.1	11.53	3	F; pS; gpmbM	5
772	II. 597	3 38 44.4	2.986	2	94 31 51.1	11.53	2	vF; S; iE; *nr	1
773	2583	II. 458	3 38 44.4	2.703	2	108 43 32.1	11.53	2	pB; pS; R; smbM*13	3
774	II. 594	3 38 46.3	2.682	2	109 41 52.1	11.53	2	pB; vS; bM	1*
775	308	VIII. 80	3 38 52.0	4.490	2	37 46 18.6	11.48	2	Cl of ab 30st 12...14	3
776	2585	3 39 4.9	1.977	1	135 5 20.4	11.52	1	pB; L; vmE 221°6	1
777	II. 459	3 39 12.4	2.696	1	109 0 25.3	11.49	1	F; R; lbM	1
778	309	I. 155	3 39 32.4	2.989	1	94 24 37.9	11.47	1	pB; S; R; *17M	3*
779	2586	3 39 56.6	1.974	3	135 5 22.2	11.46	3	pF; pL; eE 42°3; vgpmbM	3
780	2587	3 40 59.9	2.241	1	127 7 51.9	11.37	1	F; S; R; *att	1
781	2588	II. 460	3 42 24.8	2.739	1	106 49 22.2	11.26	1	pB; S; lE; mbMN!	2
782	2589	3 43 32.5	+1.139	2	150 14 27.4	11.22	2	cF; S; R; glbM; am 7Bst	2
783	2590	3 45 15.1	-0.360	1	162 6 36.1	11.13	1	pF; pS; iR; glbM; *7f	2
784	2592	3 46 48.5	+0.224	2	158 38 24.7	11.01	2	cF; pL; R; gvlbM	2
785	2591	3 47 34.5	1.955	2	134 56 47.0	10.90	2	cF; S; E 90°; gbM	2
786	2593	3 48 18.8	2.644	1	110 52 29.8	10.84	1	eF; S; R; 2Bstf; p of 2	1
787	2594	III. 962	3 48 28.0	2.643	3	110 54 59.4	10.82	2	F; S; vLE; 2st 10nr; f of 2	4
788	2595	Δ. 427? 428?	3 48 31.1	1.830	2	137 53 54.8	10.84	2	cF; pL; R; vglbM	2
789	2596	3 49 9.1	2.212	1	127 24 18.6	10.78	1	vF; L; E; vglbM	1
790	2597	Δ. 480	3 51 4.2	2.029	3	132 46 40.8	10.64	3	pb; pL; R; gbM; 2st Δ	3
791	Auw. N. 19	3 52 1.3	3.444	...	71 49 55.8	10.54	...	*12 inv in neb (Markree Cat. Nov. 24, 1854).	0
792	2599	3 52 22.9	0.478	2	156 25 42.3	10.59	2	pB; S; vLE; pmbM	2
793	I. 258	3 52 32.8	4.482	1	39 1 19.4	10.52	1	vB; S; iF; bM; r; *inv	1
794	2598	3 52 59.7	2.251	1	125 51 46.3	10.49	1	vF; vS; R	1
795	2600	Δ. 438	3 53 5.8	1.870	2	136 36 53.0	10.50	2	F; cL; R; vglbM	2
796	2601	3 53 36.3	1.748	2	139 18 42.9	10.47	2	F; L; R; vglbM; 3st n	2
797	2602	3 53 47.2	1.721	3	134 52 57.5	10.45	3	eF; S; lE 90°; vglbM	3
798	310	3 53 47.4	4.551	1	37 45 45.6	10.38	1	Cl; segment of a ring	1
799	VII. 3	3 53 52.2	2.821	1	102 25 27.4	10.42	1	Cl; S; C	1
800	2603	Δ. 369?	3 54 33.6	1.573	1	142 43 35.3	10.39	1	F; vS; R; pmbM; *8 np	1
801	IV. 53	3 54 57.0	5.109	2	29 27 30.6	10.28	2	O; pB; pS; vLE; 1' diam	2
802	VII. 47	3 55 9.9	5.232	2	28 3 51.2	10.26	2	Cl; pRi; cC; iF	2
803	2604	3 55 25.3	0.456	1	156 25 44.2	10.36	1	eF; pS; R; *10 np	1
804	2605	3 56 44.9	1.552	1	142 58 5.1	10.23	2	eeF; S; R; bet 2st 12 & 13.	2
805	II. 279	3 57 19.5	3.021	2	92 35 1.5	10.15	2	vF; pL; mE; vlbM; er	2
806	2606	3 59 0.1	1.965	2	133 47 35.5	10.05	2	F; pL; R; vgmB	2
807	2608	3 59 3.1	0.218	3	158 1 23.7	10.09	3	pB; pS; mE 121°5; gbM	3
808	2607	Δ. 466	3 59 20.3	1.966	3	133 44 23.1	10.03	3	⊕; B; cL; R; bM; rr	3
809	VII. 60	3 59 35.9	4.416	1	40 51 49.5	9.95	1	Cl; L; vRi; pC; st vL	1
810	311	IV. 69	4 0 28.6	3.755	2	59 35 29.2	9.96	2	*8m in neb 3' diam	3*+
811	2609	Δ. 348	4 0 39.4	1.448	2	144 29 25.8	9.94	2	B; L; vmE 10°; bM	2
812	2610	III. 499	4 1 24.0	2.884	2	99 12 21.5	9.85	2	eeF; S; E; psmbM; er	3
813	2611	4 1 44.0	+2.613	1	111 33 0.1	9.83	1	B; L; pmE; gbM; *8 sp	1
814	2615	4 2 8.1	-2.026	1	167 13 0.4	9.92	1	Cl; pL; lRi; st 9...10	1
815	2612	4 2 10.3	+2.616	1	111 25 36.0	9.80	1	pB; R; bM	1
816	2613	4 2 35.2	1.525	2	143 2 40.3	9.79	2	eF; vS; R; vlbM	2
817	2614	4 2 48.1	1.440	1	144 28 35.9	9.77	1	vF; R	1
818	2617	4 4 8.8	0.425	2	156 12 37.0	9.70	2	eF; vS; R; glbM	2
819	2616	4 4 17.7	+1.761	2	138 15 52.5	- 9.65	2	pB; pS; E 77°; vsmbMRN	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
820	h.	H.		h m s	s		° ' "	"			
820	VII. 61	4 4 37.1	+4.520	1	39 7 14.2	-9.56	1	Cl; B; vRi; cC	1
821	2619	4 5 33.4	0.746	1	153 16 3.3	9.59	1	vF; S; R; gbM	1
822	2620	4 6 34.6	2.302	3	123 12 36.2	9.46	3	pB; pL; R; bM; np of 2 ...	3+
823	2621	Δ. 600	4 6 39.6	2.301	3	123 14 8.2	9.46	3	B; vL; vmE 32°2; psmbM...	3+
824	2622	4 6 51.0	1.297	2	146 29 15.9	9.47	2	vB; vL; R; smbM; 2st *10nf	2
825	2623	4 6 59.8	0.749	2	153 9 21.9	9.47	2	F; S; R; vS *3d sf	2
826	2618	IV. 26	4 7 50.8	2.792	A	103 5 32.2	9.36	1	⊕; vB; S; R; ps, vsbM; r...	5*+
827	2625	4 8 4.3	1.267	1	146 50 24.6	9.38	1	vF; R; pL; vlbM	1
828	2624	4 8 17.9	2.337	1	121 54 41.8	9.34	1	vB; pS; IE; psymbM	1
829	2626	4 9 32.7	2.420	2	118 50 16.1	9.23	2	vF; vS; E; gvlbM; r	2
830	2627	4 9 58.6	1.170	1	148 5 33.1	9.23	1	B; pL; E; smbMN=*11 ...	1
831	VIII. 85	4 10 25.3	4.489	1	40 5 39.7	9.11	1	Cl; pRi; IC; stL	1
832	2628	4 11 37.5	1.282	2	146 24 57.7	9.11	2	pB; IE; gbMEN; *p	2
833	312	4 11 45.8	3.959	1	53 25 50.1	9.03	1	Cl; vL; IRi; IC; st10...12 ...	1
834	2629	4 12 9.4	1.313	1:	145 55 46.2	9.06	1	B; pS; R	1
835	D'Arrest, 42	4 12 21	3.11	[1]	87 56 12	8.99	[1]	vF; S; R; *13nr	0
836	II. 464	4 12 22.5	3.097	1	88 55 34.0	9.00	1	F; vS; R	1*
837	313	III. 490	4 13 10.4	3.052	2	91 2 17.1	8.93	2	cF; pS; IE; vgbM; *11sf ...	3
838	2630	4 13 11.1	1.295	1	146 7 50.6	8.98	2	vB; pS; R; gmbM; am 3st...	2
839	Auw. N. 20	4 13 47.7	3.488	...	70 48 46.0	8.87	...	!!!; vF; S; variable (Hind)...	0*
840	2631	4 13 56.5	+1.622	2	140 30 9.7	8.91	2	cF; S; R; vglbM	2
841	2633	4 14 0.8	-0.349	1	160 46 26.2	8.96	1	Cl; vIC; ab 20 sc st	1
842	2632	4 14 56.3	+1.858	1	135 21 58.1	8.83	1	pF; S; E; gbM	2
843	2634	4 15 56.3	0.704	2	153 7 47.6	8.78	2	vB; vL; mE; vgpmbM; *14 att n.	2
844	2635	Δ. 338??	4 16 52.1	1.337	2	145 16 34.3	8.69	2	B; vL; vg, svmbM; 15 ^s d in R.A.	2
845	2636	4 17 9.1	1.707	1	138 35 23.2	8.66	1	F; S; R; bM	2
846	2637	4 17 42.3	1.916	1	133 47 30.7	8.61	1	F; S; R; gbM	1
847	II. 768	4 17 43.5	5.621	1	25 27 13.7	8.51	1	pB; S; IE; bNM; pB*n ...	1
848	2638	4 17 45.5	1.910	1	133 57 0.7	8.61	1	vF; S; R; gbM; *nf	1
849	2639	4 17 59.3	2.026	1	130 55 4.6	8.58	1	pF; S; R; *13 nf 1'	1
850	2640	4 19 13.4	1.190	1	147 17 51.7	8.51	1	pB; S; R; pgbM; 2S st sf ...	1
851	314	III. 587	4 19 24.3	2.989	1	93 56 58.5	8.45	1	eF; bM; bet 2 st	2*
852	2641	4 20 12.4	1.526	3	141 55 9.4	8.42	3	pF; S; R; bM	3
853	315	I. 217	4 21 2.8	3.925	3	55 2 25.3	8.29	3	pB; vL; iR; mbM; *8, 350°, 2'.	6+
854	2642	4 21 37.2	1.321	2	145 15 47.7	8.31	2	F; S; E; glbM	2
855	VIII. 70	4 22 9.3	4.233	1	46 27 35.0	8.20	1	Cl; vL; pRi; IC; stL	1
856	2643	4 22 58.1	1.957	1	132 27 41.3	8.19	1	pF; S; R; gbM; *12, 287°8	1
857	D'Arrest, 43	4 23 5	3.05	[1]	90 50 6	8.15	[1]	vF; iF; vlbM; bet * & *14.	0
858	316	II. 8	4 23 29.1	3.081	1	89 39 14.4	8.12	1	F; pS; R; r; p of D neb	5
859	317	II. 9	4 23 32.6	3.081	1	89 38 36.7	8.11	1	F; vS; R; r; f of D neb	5
860	318	II. 7	4 23 33.5	3.085	1	89 26 33.7	8.11	1	F; pL; IE 132°; *42°, 80'' ...	5
861	2644	4 23 48.7	2.450	3	117 0 49.0	8.10	3	pF; pS; R; gbM	3
862	2645	4 23 52.7	2.443	1	117 15 57.0	8.10	1:	vF; vS	1
863	2646	4 24 24.8	1.710	1	138 7 6.6	8.08	1	vF; S; R; bM	1
864	2648	4 24 35.0	1.305	2	145 20 17.6	8.08	2	B; pL; mE 15°0; smbM; p of 2.	2
865	2647	4 24 36.3	1.711	1	138 5 22.2	8.06	1	F; S; R; bM	1
866	319	I. 158	4 24 45.0	2.958	4	95 23 3.4	8.02	4	pB; pL; R; gmbM	6
867	} 319, a	4 24 ±	95 23 ±	0
868		R. 3 novæ		
869
870	2649	4 24 48.1	1.303	2	145 22 1.2	8.06	2	eF; pL; IE; f of 2	2
871	VI. 26	4 24 59.0	4.298	1	45 3 48.9	7.97	1	Cl; vF; pS; C; steS	1
872	III. 585	4 25 45.2	2.973	1	94 38 56.8	7.94	1	Susp in hazy weather	1
873	III. 586	4 26 10.8	2.975	2	94 33 58.0	7.90	2	eF; S; E90°+	2
874	2650	4 28 20.2	1.883	3	134 0 32.2	7.76	3	F; S; E; vglbM	3
875	2651	Δ. 339??	4 28 33.6	+1.321	2	144 54 14.2	-7.76	2	B; L; mE 105°8; } vg, vsmbMN5''.	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	'Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
876	h. 320	H. II. 524	4 29 6.8	+2.999	1	93 26 37.9	-7.67	1	F; S; iF; lbM; 2st sf	2
877	321	II. 514	4 29 26.0	3.064	2	90 25 50.1	7.63	2	vF; pL; mE0°...90°; B *nf.	3
878	320, a	D'Arrest, 44	4 29 34	3.00	[2]	93 28 48	7.63	[2]	vF; S; *20, 270°, 5°; II. 524 p (R).	0
879	V. 49	4 29 47.5	4.558	1	39 50 13.9	7.57	1	F; cL; iF; 6 or 7 st + neb ...	1
880	322	4 30 7.9	2.995	1	93 35 45.6	7.58	1	vF; E 90°...180°; sbM; B*p 40°.	1*
881	322, a	R. nova	4 30 19.9	+2.995	::	90 30 45.6	7.58	::	MS	0
882	2653	4 31 14.0	-0.756	1	162 8 7.3	7.59	1	vF; pL; R; glbM	1
883	2652	4 32 20.0	+2.598	1	110 55 34.4	7.42	1	Neb. No description	1
884	323	III. 952	4 32 36.3	3.227	2	82 56 7.6	7.38	2	eF; S; R; *8 sp; p of D neb	3
885	324	III. 953	4 32 36.5	3.226	1	82 57 2.6	7.38	1	eF; vS; f of D neb	2
886	325	II. 515	4 33 1.4	3.055	1	90 49 38.5	7.35	1	F; S; R; bM; *9 nf 12° 5 ...	2
887	{ 326 = 2654 }	II. 522	4 33 57.4	2.878	2	98 52 48.9	7.27	2	vF; pS; R; vgbM; r; *nf 1/.	4
888	327	I. 122	4 34 26.6	3.005	2	93 8 53.1	7.23	2	cB; L; R; vgbM; er	3†
889	2655	4 34 36.9	2.686	1	107 16 3.1	7.23	1	eF; vS; R; bet 2 st	1
890	II. 525	4 35 29.3	3.028	1	92 2 39.5	7.15	1	F; pL; iE	1
891	2656	4 35 30.1	0.266	1	156 4 46.4	7.22	1	Cl; pL; pRi; pmC; st 11...16	1
892	D'Arrest, 45	4 35 45	3.08	[1]	89 39 1	7.12	[1]	F; R; cometary; Δ with 2 st 18, f.	0
893	328	III. 588	4 36 51.6	2.950	1	95 35 42.8	7.04	1	eF; vS; iR; bM	2
894	2657	4 37 7.2	0.206	3	156 27 51.3	7.09	3	F; S; R; gbM	3
895	329	II. 523	4 37 41.4	2.879	1	98 47 13.9	6.97	1	F; vS; iR; bM; *7 np	2
896	VIII. 8	4 37 54.5	+3.500	2	71 11 21.8	6.94	2	Cl; vL; stL, sc	2
897	2660	4 38 37.3	-0.201	1	159 5 7.6	6.98	1	F; pS; R; gbM	1
898	2662	4 38 42.3	-0.531	3	160 51 32.6	6.98	3	pF; L; vIE; vglbM	3
899	2661	4 38 43.9	-0.177	3	158 56 26.9	6.97	3	vF; S; R; glbM	3
900	II. 526	4 38 49.7	+3.015	1	92 38 59.9	6.87	1	F; cS; R; lbM	1
901	330	4 39 0.9	2.954	1	95 24 18.2	6.86	1	eF; iF; ?	1
902	2658	4 39 28.8	1.952	1	131 45 5.5	6.85	1	F; pS; pmE; glbM	1
903	331	III. 589	4 39 36.7	2.962	1	95 2 37.7	6.81	1	pF; pS; iE90°±; bM	2
904	2659	4 39 37.6	1.950	1	131 46 55.1	6.83	1	vF; S; iE; glbM	1
905	332	VII. 1	4 40 44.9	3.310	1	79 19 18.7	6.71	1	Cl of L & S sc st	3
906	VIII. 7	4 40 45.1	3.361	2	77 6 7.7	6.71	2	Cl; iRi; st L & S	3
907	VIII. 59	4 41 2.4	4.268	1	46 33 17.5	6.65	1	Cl; iRi; iC; pL	1
908	333	II. 547	4 41 23.7	2.947	1	95 41 12.2	6.66	1	eF; pL; R; lbM	3*
909	2663	4 41 59.7	1.811	1	135 2 6.8	6.64	1	eF; R; att to *14	1
910	2664	4 42 20.4	0.236	1	156 4 5.2	6.66	1	eF; S; R	1
911	III. 501	4 42 46.2	3.007	1	93 0 18.8	6.54	1	vF; vS	1
912	2665	Δ. 296 ??	4 43 32.0	+0.930	2	149 30 3.8	6.54	2	B; L; smbMN	2
913	2667	4 43 34.7	-0.404	1	160 4 2.6	6.58	1	vF; S; att to *10	1
914	2669	4 44 19.2	-0.228	1	159 4 32.7	6.51	1	vF; pL; iR; r	1
915	III. 502	4 44 35.4	+3.010	1	92 51 28.0	6.40	1	vF; S	1
916	2666	4 44 39.1	2.276	2	122 12 45.4	6.42	2	vB; L; iR; 4st inv	2
917	2668	4 44 41.5	1.665	3	138 3 31.4	6.42	3	vF; S; R; r or st inv	3
918	334, a	R. nova	4 45 16.3	2.999	::	93 14 46.4	6.32	::	R, MS	0
919	D'Arrest, 46	4 45 22	3.00	[2]	93 21 0	6.33	[2]	vF; vS; II. 527, f 12°±	0
920	334	II. 527	4 45 32.3	2.999	2	93 20 46.4	6.32	2	pF; S; R; bM; *7, 225°± ...	4
921	334, b	R. nova	4 45 44.3	2.999	::	93 20 46.4	6.32	::	MS } No description	0
922	334, c	R. nova	4 45 44.3	2.999	::	93 8 46.4	6.32	::	MS }	
923	2670	4 46 9.1	2.212	2	124 10 25.0	6.30	2	vF; S; R; vglbM	2
924	II. 528	4 46 12.6	2.999	1	93 18 35.2	6.26	1	F; S; lbM	1
925	2671	4 46 18.6	0.875	3	150 2 6.7	6.31	3	pB; pL; iR; pgmbM	3
926	335	4 47 6.7	+3.104	1	88 35 49.3	6.19	1	vF; vS; am vSst; L* sp	1*
927	2672	4 48 23.2	-0.339	2	159 35 14.9	6.17	2	F; S; R	2
928	2673	4 48 30.2	0.345	2	159 37 5.2	6.16	2	F; S; R	2
929	2674	4 48 47.0	0.146	1	158 27 26.8	6.14	1	vF; E; vlbM	1
930	2675	4 49 1.9	0.203	5	158 47 31.4	6.12	5	⊕; pB; L; R; rr	5
931	2677	4 49 42.0	-0.302	1	159 20 44.2	-6.06	1	pB; pS; R; glbM	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
932	h. 336	H. IV. 32	h m s	s	1	95° 5' 43.5"	-5.95	1	eB; S; mbM*.....	3
933	2676	4 50 25.3	+2.333	3	120 6 11.1	5.93	3	F; S; vIE; glbM; *10, 75''.	3
934	2680	Δ. 73?	4 50 27.0	-0.443	1	160 5 32.0	6.00	1	Cl; vF; S	1
935	2678	4 50 48.7	+0.866	2	149 58 0.1	5.93	2	F; L; R; vglbM; * att.....	2
936	2683	4 50 53.7	-0.427	2	159 59 14.9	5.97	2	F; pS; IE; r	2
937	2679	4 51 8.5	+1.342	3	143 35 14.3	5.89	3	pF; S; R; pmbM	3
938	2682	4 51 15.3	0.544	1	153 13 27.7	5.91	1	F; pS; R; vglbM	1
939	338	4 51 15.6	3.253	1	81 58 39.1	5.83	1	S; R; rrr.....	1
940	337	4 51 32.4	+4.753	1	37 19 51.9	5.77	1	Cl; vL; pRi; lC; stL and S.	1
941	2684	Δ. 76?	4 51 44.5	-0.473	2	160 13 0.0	5.90	2	⊕; B; S; iR; rrr; st14	2
942	2685	4 51 46.1	-0.370	1	159 39 49.3	5.89	1	Cl; pB; S	1
943	339	II. 516	4 51 47.1	+3.057	2	90 42 45.3	5.79	2	F; S; R; bM; p of 2	2
944	339, a	R. nova	4 51 ±	90 42 ±	No description	0
945	2686	4 52 0.4	0.043	5	157 8 55.2	5.86	5	vB; S; Eor bi-N; bM; sp of 2	5
946	2687	4 52 2.6	0.046	3	157 7 48.2	5.86	3	vF; S; R; sbM; 2stnr; nfof 2	3
947	2681	4 52 16.4	2.592	1	110 34 42.9	5.77	1	pF; pL; R; glbM	1
948	340	4 52 18.9	+3.063	1::	90 28 7.5	5.75	1::	pF; S; iR; pslbM	2
949	340, a	R. nova	4 52 ±	90 28 ±	No description	1
950	2688	4 52 20.4	+0.022	3	157 16 42.8	5.84	3	F; pS; R; vglbM	3
951	D'Arrest, 47	4 52 39	+2.89	[2]	98 4 0	5.73	[2]	pF; pL; lbM; h. 341 nr.....	0
952	2689	4 52 41.7	-0.366	3	159 37 11.4	5.82	3	Cl; pF; S; R; 2nd of 3	3
953	341	4 52 57.2	+2.893	1	97 58 11.7	5.71	1	F; R; *13, s	1*
954	2690	4 53 0.9	-0.358	3	159 34 7.3	5.79	3	Cl; pB; pS; pmE; st12	3
955	III. 503	4 53 21.2	+2.994	1	93 32 9.9	5.67	1	vF; pL; 2B st v nr.....	1
956	2691	4 53 22.6	+0.038	3	157 8 36.5	5.75	3	Cl; pL; lRi; lC; st10...15...	3
957	2694	4 53 43.2	-0.238	2	158 52 23.4	5.72	2	S; R; close * in M	2
958	2693	4 53 49.2	+0.073	1	156 53 44.7	5.71	1	eF; pS; R; gbM.....	1
959	2695	4 54 8.4	-0.259	1	158 59 17.0	5.70	1	pB; L; R; gmbM	1
960	2696	4 54 18.5	+0.009	1	157 18 56.9	5.67	2	pF; pS; R; 2st att	2
961	2697	4 54 19.0	-0.143	1	158 17 20.9	5.67	1	B; R; r	1
962	2698	4 54 31.9	-0.335	1::	159 24 10.2	5.66	1::	vF; S; 1st of 4	1
963	2699	Δ. 114	4 54 49.4	-0.339	2	159 25 17.8	5.64	2	B; pL; R; gbM; r; 2nd of 4	2
964	2692	4 54 51.1	+2.440	1	116 14 38.2	5.56	1	F; vL; vmE; vglbM	1
965	342	4 54 58.2	+2.994	1	93 30 26.1	5.53	1	eF; vS; *12, sf	1
966	2702	4 55 7.9	-0.335	1::	159 23 11.7	5.61	1::	F; S; 3rd of 4.....	1
967	2701	4 55 12.5	-0.006	2	157 23 32.0	5.60	2	Cl; pS; lRi; stvS	2
968	2704	4 55 13.8	-0.340	1	159 25 6.0	5.60	1	pB; vS; R; 4th of 4	1
969	2703	4 55 16.5	-0.166	1	158 24 26.3	5.59	1	vF; R; p of 2	1
970	VIII. 43	4 55 18.9	+3.630	1	66 33 18.3	5.49	1	Cl; stL, vcsc	1*
971	2705	4 55 19.4	-0.456	1	160 1 59.0	5.60	1	eF; pL; iR	1
972	D'Arrest, 49	4 55 26	+2.88	[2]	98 26 48	5.49	2	F; pL; pmE; 2 or 3st11nf...	0
973	2708	4 55 36.9	-0.581	1	160 39 49.9	5.57	1	F; S; R; *13att, 135°	1
974	2707	4 55 37.8	-0.350	1	159 27 21.9	5.57	1	vF; S; R.....	1
975	343	4 55 45.0	+2.962	1	94 55 35.9	5.47	1	vLdiff neb in zigzags??	1*
976	2706	Δ. 167	4 55 49.5	0.266	1	155 25 43.1	5.53	1	vB; pL; R; gbM; f of 2	1
977	VII. 21	4 55 55.7	3.634	1	66 25 21.1	5.43	1	Cl; pC; stL and S	1
978	2700	4 56 5.5	2.037	1	128 55 2.2	5.46	1	vF; pL; vglbM	1
979	2709	4 56 15.6	0.087	1::	156 43 53.0	5.50	1::	vF; S; 3vSst inv.....	1*†
980	2710	4 56 19.0	0.093	3	156 41 5.0	5.50	3	Cl; L; mC; * 9	3†
981	III. 453	4 56 24.5	3.104	1	88 34 31.3	5.39	1	vS; vF.....	1*
982	2711	4 56 31.9	+0.102	5	156 37 13.3	5.49	5	vB; vL; vimE.....	5†
983	2713	4 56 42.0	-0.092	1	157 54 38.9	5.47	1	vF; S; R.....	1
984	2712	4 56 51.7	+0.631	2	152 14 11.8	5.44	2	cF; S; R; glbM	2
985	2717	4 57 15.1	-0.543	2	160 26 16.8	5.44	2	cF; S; gbM	2
5062	4 57 18.4	159 36 32.1	See No. 5062	1
986	2718	4 57 22.2	-0.185	1	158 28 4.4	5.42	1	F; S; R; gbM	1
987	2716	4 57 29.6	+0.091	5	156 39 56.0	5.40	5	B; L; iR; vsmbM * 10.....	5†
988	2715	Δ. 169	4 57 39.7	-0.213	2	158 37 42.3	5.39	2	Cl+neb; pL; pRi; st11...18	2
989	2720	4 57 48.0	+0.509	1	153 20 44.2	5.36	1	vF; mE; glbM; *7, 8np	1
990	2722	4 57 48.9	-0.416	3	159 46 2.3	5.39	3	pB; pS; iR; rr	3
991	2721	4 57 50.3	+0.105	1	156 33 59.9	-5.37	1	pF; pL; iR; 2 or 3Bst nr	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ′ ″	″			
992	2723	4 58 11.0	-0.027	3	157 27 6.5	-5.35	3	B; S; R; smbM; * + neb ...	3
993	2725	4 58 18.6	-0.590	1	160 38 26.5	5.35	1	eF; pL; iR	3
994	2724	4 58 23.2	+0.093	1	156 38 1.4	5.32	1	vF; S; R; gbM	1
995	2728	4 58 28.2	-1.562	1	164 29 44.2	5.36	1	eF; E; * 9att, f	2
996	344	VIII. 61	4 58 35.7	+4.041	1	53 8 20.3	5.19	1	Cl; pC; lRi; iF; stL	2
997	{ 345 = 2714 }	III. 500	4 58 37.6	+2.860	3	99 20 30.1	5.23	3	pB; S; R; gpmbM	4
998	III. 268	4 58 42.9	+2.643	1	108 22 53.0	5.31	1	eF; vS; stellar	1*
999	2727	4 58 43.5	-0.390	2	159 36 16.7	5.31	2	⊕; pB; S; R; pmbM; rr ...	2
1000	2726	4 58 44.0	+0.153	2	156 11 37.6	5.28	2	cB; L; R; vgpmbM; r	2
1001	2719	4 58 56.6	+2.797	1	102 4 3.7	5.21	1	pB; pL; vLE; vgbM; am st ..	1
1002	(147)	4 59 12.2	-0.294	1	159 3 36.9	5.27	1	No description	1
1003	2729	4 59 20.7	-0.109	3	157 57 2.5	5.25	3	vB; pS; lE; vsymbM*9 ...	3
1004	2731	4 59 52.1	+0.170	1	156 2 39.3	5.19	1	Cl; vL; pRi	1
1005	V. 32	4 59 55.9	+2.993	2	93 32 59.7	5.11	2	B; cL; R; bM * 15; * 10, 318°.	4
1006	2733	4 59 57.5	-0.926	1	162 5 37.4	5.22	1	vF; pS; R; vglbM	1
1007	346	5 0 12.6	+4.723	1	38 7 8.8	5.04	1	Cl group of 8 or 9 st10	1
1008	2734	5 0 20.2	-0.542	1	160 21 49.6	5.18	1	eF; S; R	1
1009	2730	Δ. 531?	5 0 23.9	+2.056	2	128 11 33.0	5.10	3	vB; vL; mE314°; glbM; rr.	3
1010	2736	5 0 36.4	-0.426	2	159 45 22.5	5.15	2	F; S; R; glbM	2
1011	2738	Δ. 81	5 0 58.7	-0.472	1	159 59 38.4	5.12	1	F; pL; lE	1
1012	2735	5 1 1.0	+0.710	2	151 19 31.6	5.08	2	pF; pS; pmE; vglbM	2
1013	2732	5 1 10.9	+2.260	1	122 8 40.1	5.03	1	pB; pmE; gpmbM; *13f ...	1
1014	2739	5 1 34.1	-0.440	4	159 48 27.9	5.07	4	F; pL; R; vglbM; p of 2 ...	4
1015	VIII. 41	5 1 44.4	+3.647	1	66 4 49.8	4.94	1	Cl; st c se	1
1016	2737	5 1 51.5	+1.544	1	139 45 48.3	4.99	1	F; S; R; vglbM; *11sf; ? neb	1
1017	2742	5 1 51.9	-0.344	2	159 16 55.8	5.04	2	F; S; R; bM	2
1018	2741	Δ. 233?	5 2 1.9	+0.128	5	156 18 4.4	5.02	5	B; vS; vsymbM; st + neb	5
1019	2745	5 2 31.5	-0.159	1	158 11 17.6	4.98	1	pB; L; gbM	1
1020	348	5 2 37.5	+3.453	1	73 39 38.9	4.87	1	Cl; pRi; stL and S	1
1021	2740	Δ. 549	5 2 50.1	+2.071	2	127 41 47.0	4.90	2	B; L; E; psbM	2
1022	2747	5 3 4.7	-0.448	1	159 49 11.8	4.94	1	pF; S; R; gbM; 2nd of 2 ...	1
1023	2746	Δ. 235	5 3 9.1	+0.086	5	156 34 15.4	4.92	5	cF; S; R; lbM; ⊕f	5
1024	2743	5 3 16.9	+2.340	2	119 27 56.5	4.85	2	cF; S; lE; p of 2	2
1025	2744	5 3 27.3	+2.341	2	119 26 7.1	4.83	2	F; S; R; glbM; f of 2	2
1026	2752	5 3 54.8	-0.586	1	160 30 43.6	4.88	1	vF; S; R; r	1
1027	2748	5 3 55.4	-0.054	2	157 29 41.2	4.86	2	vF; R; s of 2 in Cl	2
1028	2753	5 3 56.0	-0.648	2	160 48 33.6	4.88	2	F; vS; R; vlbM; am st	2
1029	2750	5 3 58.7	-0.048	1	157 27 2.5	4.85	1	vF; R; 2nd neb in Cl	1
1030	349	VII. 4	5 3 59.1	+3.458	2	73 28 42.5	4.75	2	Cl; L; Ri; lC; st11...14 ...	5*
1031	2749	Δ. 236	5 4 2.0	+0.076	6	156 37 3.5	4.85	6	⊕; vB; pL; R; vmC; rr ...	6
1032	2754	5 4 22.1	-0.049	1	157 26 48.4	4.82	1	Cl; pL; Ri; C; iF	1
1033	2756	5 4 47.9	+0.104	1	156 23 46.6	4.78	1	vF; S; p of 2	1
1034	2758	5 4 51.8	-0.595	1	160 31 53.0	4.80	1	Cl; pF; L; iF; st12...15 ...	1
1035	2755	5 4 53.3	+0.834	2	149 54 12.5	4.75	2	vF; pL; vmE162°0	2
1036	(199)	5 5 2.2	-0.328	1::	159 7 51.0	4.80	1::	No description	1
1037	2757	5 5 10.7	+0.101	1	156 24 57.5	4.75	1	vF; S; f of 2	1
1038	2751	5 5 12.8	+2.088	1	127 9 17.3	4.69	1	vF; vmE; long ray; *11inv.	1
1039	2761	5 5 16.9	-0.411	4	159 34 33.2	4.76	4	F; S; R; 1st of 3	4
1040	2760	5 5 22.7	-0.177	1	158 14 14.8	4.74	1	F; pL; R; r	1
1041	2762	5 5 33.9	-0.403	4	159 31 45.1	4.73	4	F; pS; R; 2nd of 3	4
1042	2759	Δ. 246	5 5 35.8	+0.275	2	155 6 40.7	4.71	2	B; L; R; glbM; r	2
1043	II. 292	5 5 52.9	+2.702	1	105 53 13.4	4.62	1	pB; iR; mbM; cSnf1'	1
1044	2765	5 5 56.2	-0.679	1	160 55 5.7	4.71	1	vF; pL; 1st of sev	1
1045	2764	5 6 3.3	0.381	1	159 23 56.5	4.68	1	⊙? B; eS; lE	1
1046	2763	5 6 3.7	0.415	4	159 35 14.6	4.68	4	cB; S; R; gmbM; 3rd of 3..	4
1047	2766	5 6 12.5	0.278	2	158 48 40.9	4.67	2	st + neb; 1st of sev	2
1048	2769	5 6 28.0	0.677	2	160 53 56.2	4.66	2	Cl; L; Ri; st sc	2
1049	2767	Br. 895	5 6 32.8	-0.246	1	158 37 28.8	-4.64	1	Cl; L; vlc	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1050	h. 2768	H.	Δ. 170?	h m s 5 6 38.4	s -0.278	2	158° 48' 37.5	-4.65	2	st + neb; pB; iF; 2nd of sev	2
1051	2771	5 6 46.2	0.909	1	161 56 6.8	4.64	1	F; R; bM; r(? min. of R.A.)	1
1052	2788	5 7 1.6	9.684	1	174 12 38.6	4.88	1	pF; L; iR; vsbM; r	1
1053	2772	5 7 20.8	0.060	2	157 27 15.9	4.57	2	vvF; R; p of 2	2
1054	2773	5 7 36.6	0.068	4	157 29 56.8	4.54	5	pF; pL; R; gbm; f of 2	5
1055	2770	5 7 38.3	0.651	2	160 44 41.2	4.56	2	Cl; vICM; st 9, 11...16	2
1056	2774	5 7 45.1	0.089	3	157 37 57.8	4.54	4	pB; cL; R; vglbM; r	4
1057	2775	5 7 56.4	0.343	2	159 8 54.1	4.53	2	B; S; iE; * in M	2+
1058	2776	5 8 5.8	-0.785	1	161 21 52.1	4.53	1	Cl; vIC; st 9, ...	1
1059	2778	5 9 20.8	+0.078	1	156 28 44.3	4.39	1	vF; S; iE; glbM	1
1060	2780	Δ. 170?	5 9 25.2	-0.309	6	158 55 47.0	4.40	6	⊕! vB; L; iE; vmCM; rr...	6
1061	2777	Δ. 508	5 9 29.8	+1.970	2	130 12 20.1	4.33	2	⊕! vB; vL; R; vsvbm; rrr.	3
1062	2781	5 9 41.5	-0.145	4	157 56 58.9	4.37	4	F; pL; R; vglbM	4
1063	2779	5 9 51.6	+1.019	3	147 33 40.1	4.33	3	F; S; mE 45°; vglbM; *11 nf.	3
1064	2782	5 10 2.3	-0.327	5	159 1 12.5	4.35	5	⊕; cB; S; R; gbm; 2d of 3	5
1065	2783	5 10 8.1	0.326	1	159 0 38.8	4.34	1	Cl; vB; L; R; st12	1
1066	2784	5 10 20.3	-0.379	3	159 17 59.1	4.33	3	B; pL; R; gbm; 12° diam R.A.	3
1067	350	VII. 33	5 10 26.0	+4.138	1	50 48 30.6	4.18	1	Cl; pRi; pC; st 7, ...	2
1068	2785	5 10 37.2	-0.337	7	159 3 44.0	4.30	7	B; L; iE; biN; Cl + neb	7
1069	2786	5 10 59.4	+0.220	1	155 24 46.2	4.26	1	F; S; R; vgbM; * 7 nf 6'	1
1070	2787	Δ. 172?	5 11 21.6	-0.316	1	158 55 38.1	4.23	1	F; pL; R; vgbM	1
1071	2790	5 12 2.3	-0.708	1	160 56 56.6	4.18	1	eF; pL; R; vglbM	1
1072	2789	5 12 6.1	+0.094	1	156 18 45.2	4.16	1	pF; L; iR; vgbM; r	1
1073	2791	Δ. 173?	5 12 19.1	-0.312	5	158 53 22.5	4.15	5	vB; vS; R; r or stellar	5
1074	2792	5 12 56.3	0.130	2	157 47 15.6	4.08	2	F; pS; iR; bM; r or stellar	2
1075	2794	Δ. 173??	5 13 6.0	-0.321	3	158 55 37.6	4.08	3	vF; pL; R; vglbM	3
1076	2793	Δ. 247? 248?	5 13 11.3	+0.185	1	155 37 35.2	4.06	1	vB; L; R; vgbm; r	1
1077	2795	5 13 27.1	+0.071	1	156 27 4.8	4.04	1	eF; pL; R	1
1078	2796	5 13 41.5	+0.376	2	154 6 36.7	4.01	2	pB; pL; R; vglbM	2
1079	2798	Δ. 210	5 13 58.4	-0.092	3	157 32 7.0	4.00	2	Cl; L; pRi; st sc	2
1080	2799	5 13 59.0	0.386	2	159 16 29.7	4.01	2	B; S; R; gbm	2
1081	2800	5 14 5.5	0.105	2	157 36 46.0	4.00	2	Cl; lRi; 2nd of sev	2
1082	2802	5 14 7.8	0.423	4	159 28 12.0	4.00	4	pB; R; gbm; 1st of group	4+
1083	2801	5 14 8.8	0.086	1	157 29 23.3	3.99	1	Cl; 3rd of sev	1
1084	2803	5 14 9.6	0.434	1	159 31 55.0	4.00	1	neb & Cl; biN	1+
1085	2804	5 14 16.2	0.432	4	159 31 1.6	3.98	4	pB; iR; biN; 2nd in group	4+
1086	2805	5 14 19.0	0.434	1	159 31 50.6	3.98	1	vF; 3rd of group in Cl	1+
1087	2807	5 14 26.2	-0.651	1	160 37 52.6	3.98	1	vF; iE; vglbM; r	1
1088	2797	5 14 37.2	+2.244	1	122 17 45.6	3.88	1	vF; L; R; vglbM; *12p	1
1089	2808	5 14 37.9	-0.435	1	159 31 47.2	3.96	1	4th N. of neb in Cl	1+
1090	2810	5 15 3.1	-0.421	1	159 27 2.4	3.92	1	vF; * p	1+
1091	2809	5 15 15.7	+0.091	1	156 16 34.6	3.88	1	pF; R; vglbM; r	1
1092	VII. 34	5 15 28.7	4.452	1	43 35 54.8	3.74	1	Cl; vF; pRi; pC; iF	1
1093	2812	5 15 45.0	0.084	1	156 18 45.8	3.84	1	eF; pL	1
1094	2814	5 15 47.4	0.362	1	159 7 24.2	3.86	1	pB; vS; R; bM	1
1095	2813	5 15 53.7	0.061	1	156 28 16.1	3.83	1	vF; vS; R; *p25"	1
1096	{ 352 = 2806 }	II. 289	5 16 3.7	+2.802	2	101 38 3.8	3.74	2	pB; pL; R; r	2
1097	352, a	R. nova	5 16	101 38	Makes a close D neb with h. 352.	0
1098	2816	5 16 5.8	-1.017	1	162 13 55.5	3.85	1	vF; S; R; gbm	1
1099	2811	5 16 20.8	+2.124	1	125 51 33.8	3.74	1	Cl; L; Sc; * taken	1
1100	2815	5 16 31.9	+0.245	2	155 6 43.9	3.77	2	eF; pL; E 90° ±; vglbM	2
1101	351	5 16 32.3	+3.935	1	56 44 33.9	3.67	1	Cl; L; Ri; iC	1
1102	2818	5 16 56.7	-0.457	1	159 36 42.2	3.76	1	F; pL; R; sbM; r; st inv	1
1103	2817	5 17 3.0	-0.092	4	157 28 36.8	3.74	4	pF; pL; R; vglbM	4
1104	353	VIII. 4	5 17 8.1	+3.802	1::	69 58 11.1	-3.63	1::	Cl; vLri; vIC; st 9...12	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
1105	2820	5 17 44.0	-0.112	1	157 35 20.6	-3.68	1	eF; S; R.....	1
1106	2822	5 17 51.1	-0.495	1	159 47 44.6	3.68	1	F; pS; R.....	1
1107	2821	5 17 58.4	-0.180	1	158 1 7.2	3.66	1	F; pS; R; vglbM; 3st10p ...	1
1108	2819	5 18 1.2	+0.475	2	153 10 14.8	3.64	2	F; pL; lE; vglbM; *7np...	2
1109	2824	5 18 7.8	-0.302	1	158 44 6.2	3.66	1	Cl; BM; lRi; st7	1
1110	2823	5 18 14.1	+0.013	5	156 46 1.1	3.63	5	⊕; pB; pL; R; pmbM; rr...	5
1111	2825	5 18 22.1	-0.436	5	159 28 39.8	3.64	5	vB; S; R; gmbM	5
1112	M. 79	5 18 25.6	+2.469	A	114 39 39.5	3.55	A	⊕; pL; eRi; eC; rrr	4
1113	4016	5 18 36.3	-0.087	1	157 25 24.7	3.61	1	F; S; R; r	1
1114	354	VII. 39	5 18 42.6	+4.001	3	54 48 28.6	3.48	3	Cl; pRi; pC; R; st9...12 ...	5
1115	V. 33	5 18 52.3	3.011	1	92 39 22.0	3.30	1	v diffused neb susp	1
1116	V. 38	5 19 10.4	+2.881	1	98 15 15.9	3.47	1	eL; strongly susp (2° in P.D.).	1
1117	2827	Δ. 129	5 19 10.4	-0.416	4	159 21 30.9	3.57	4	Cl; L; pRi; iR; st11...16 ...	4
1118	2826	5 19 13.2	-0.010	1	156 54 4.5	3.55	1	F; R; gbM; am st	1
1119	M. 38	5 19 17.0	+4.020	(2)	54 17 36.1	3.43	(2)	Cl; B; vL; vRi; iF; st L & S	7
1120	(356)	5 19 33.6	-0.480	1::	159 41 31.8	3.54	1::	No description.....	1
1121	2830	5 19 35.7	0.833	2	161 23 42.8	3.54	2	F; L; iE	2
1122	2828	5 19 36.7	0.016	1	156 56 22.4	3.52	1	eF; pL.....	1
1123	2829	5 19 39.1	0.453	3	159 32 47.1	3.53	3	B; S; R; vgvmbM; r.....	3
1124	2831	5 19 53.3	0.379	1	159 8 15.0	3.50	1	vF; L; R; vglbM	1
1125	(369)	5 20 10.1	0.496	1::	159 46 33.6	3.48	1::	No description	1
1126	2832	5 20 15.1	0.030	1	157 0 57.2	3.46	1	Cl; eF; L; iR; mC; rr	1
1127	2833	5 20 33.0	0.014	7	156 54 44.8	3.44	7	pB; pL; R; vglbM	7
1128	2834	5 20 33.9	0.527	1	159 55 8.5	3.45	1	vF; pS; lE; r	1
5063	5 20 52.9	-0.469	1::	159 36 35.4	3.42	1::	(See No. 5063)	1
1129	2835	5 21 6.5	+0.168	1	155 36 58.6	3.38	1	vF; pS; R	1
1130	III. 447	5 21 8.1	2.947	2	95 26 56.0	3.30	2	pF; pL; iR; st nr	2
1131	2837	5 21 22.2	+0.113	1	156 0 17.2	3.36	1	Cl; vRi; lC; st10	1
1132	2838	5 21 40.7	-0.480	3	159 39 51.5	3.35	3	pB; pL; iR; r; in diff n.....	3
1133	356	5 22 0.7	+2.875	1	98 29 41.4	3.22	1	Diffused nebulosity	1*
1134	2839	Δ. 131	5 22 1.5	-0.473	2	159 36 55.4	3.32	2	pF; pL; R; gbM	2
1135	2840	5 22 1.9	-0.198	1	158 3 33.7	3.31	1	F; p of group	1†
1136	2836	5 22 3.0	+1.666	2	136 51 6.2	3.26	2	pF; S; R; bM; 4B st p	2
1137	355	I. 261	5 22 10.0	3.968	3	55 52 3.6	3.18	3	vB; L; R; b ** in M	4†
1138	2841	5 22 14.9	+0.074	5	156 16 28.3	3.29	5	{ pB; S; R; smbM } D neb	} 5*
1139											
1140	2842	5 22 21.3	-0.197	1	158 2 52.6	3.28	1	2nd neb of group	1†
1141	2843	5 22 24.9	0.204	4	158 5 19.6	3.28	4	pF; S; R; 3rd of group	4†
1142	2844	Δ. 175	5 22 41.6	0.207	6	158 6 22.5	3.25	6	!; pB; S; R; 4th of group ...	6†
1143	2845	5 22 43.6	0.195	1	158 1 49.5	3.25	1	vF; pL; follows a group.....	1†
1144	2848	Δ. 89?	5 22 45.6	0.564	4	160 4 39.2	3.26	4	{ pB; pS; R; glbM } D neb	} 4
1145											
1146	2847	5 22 54.5	-0.082	2	157 18 47.1	3.23	2	pB; vS; R; bM; 2st 9 & 10 f	2
1147	2846	5 22 55.3	+0.039	1	156 30 33.4	3.22	1	vS; neb+st.....	1
1148	2849	5 23 51.4	+0.358	2	154 4 25.8	3.14	2::	eF; stell; *14+ neb	2
1149	2850	Δ. 90	5 23 53.2	-0.608	3	160 16 48.9	3.17	3	pF; pS; iR; vglbM; *15, 190°-6, 60'	3
1150	2852	5 24 26.6	-1.138	1	162 36 15.8	3.14	1	pB; pL; R; bM	1
1151	2851	5 24 48.3	+0.025	1	156 34 43.9	3.07	1	eeef; vL; irr diff	1
1152	2854	Δ. 237?	5 25 8.5	0.034	1	156 30 29.8	3.04	1	pF; R; gbM; r	1
1153	2855	5 25 37.5	0.255	1	154 52 39.3	2.99	1	pB; L; R; glbM; *9np	1
1154	2856	5 25 40.3	+0.051	1	156 23 6.3	2.99	1	Cl; cL; Ri; st13.....	1
1155	2857	5 25 47.4	-0.296	1	158 35 32.3	2.99	1	pB; S; R; psbM.....	1
1156	2859	5 25 50.5	-0.561	2	160 1 23.3	2.99	2	The 1st of a group of 7!.....	2†
1157	357	M. 1	5 26 3.9	+3.605	1	68 5 10.5	2.85	1	vB; vL; E135°±; vglbM; r	12†
1158	2858	5 26 4.1	+0.003	2	156 42 31.2	2.96	2	B; lE; sbM * 10 & 11	2
1159	2862	5 26 16.7	-0.362	2	158 57 2.5	2.95	2	pB; S; R; glbM	2
1160	2853	III. 590	5 26 24.5	+2.738	1	104 9 58.5	2.85	1	vF; S; R; smbM	2
1161	2863	Δ. 211	5 26 31.0	-0.137	1	157 37 22.4	2.92	1	Cl; Ri; 2nd of sev	1
1162	2874	5 26 39.9	-3.085	2	167 51 34.3	-2.99	2	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of used. Obs.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1163	h. 2864	H.	h m s 5 26 46.8	s -0.550	3	159 57 22.7	-2.91	3	F; pL; iR; vgbM; 2nd of group!	3†
1164	2865	5 26 50.2	-0.569	1	160 2 51.7	2.91	1	F; vL; vgbM; 3rd of group!	1†
1165	2866	Δ. 136?	5 27 15.3	-0.363	1::	158 57 7.3	2.89	1	vF; pL; R; 1st of 4!	1*†
1166	358	M. 36	5 27 3.1	+3.966	3	55 57 28.2	2.76	3	Cl; B; vL; vRi; IC; st 9...11sc	9
1167	III. 747	5 27 11.2	+6.681	1	20 35 59.9	2.67	1	cF; pL; iF; mbM; er; * inv (? P.D.).	1*
1168	2867	Δ. 136?	5 27 32.7	-0.358	2::	158 55 26.2	2.86	2::	F; S; 2nd of 4!	2*†
1169	2861	5 27 17.2	+2.094	1	126 28 48.3	2.79	1	Cl; st 8...11	1
1170	2860	IV. 21	5 27 26.0	+2.536	1	112 2 39.9	2.77	1	F; vS; R; vsymbM*12; 3st inv.	2
1171	2868	Δ. 136	5 27 37.6	-0.361	4	158 56 0.7	2.84	4	{ Cl; pL; iF; 1st 9; } { + group of 4n nebpB; } { R; psibM; 3rd of 4 }	3*† 1
1172	(456)	5 27 43.2	0.411	2:	159 12 33.1	2.83	2:	No description.....	2
1173	2870	5 27 44.2	0.132	1	157 34 9.4	2.82	1	Cl; Ri; 3rd of sev	1
1174	2872	5 27 46.7	0.553	1	159 57 36.7	2.82	1	F; S; 4th of gr of 7	1†
1175	2869	5 27 50.4	0.361	1::	158 56 34.5	2.84	1::	4th of 4	1*†
1176	2875	5 28 1.0	0.556	1	159 58 24.7	2.81	1	5th of gr of 7	1†
1177	2876	5 28 2.0	0.555	1	159 57 34.7	2.81	1	6th of gr of 7! D; a vS neb np	1†
1178	2877	Δ. 213	5 28 16.4	-0.129	1	157 32 34.9	2.77	1	Cl; L; irr.....	1
1179	360	{ M. 42= θ ¹ Orionis }	5 28 24.0	+2.945	B.A.C.	95 29 10.9	2.68	B.A.C.	!!!; θ ¹ Orionis & the great neb	Mon.*†
1180	V. 30	42, c ¹ Orionis	5 28 29.2	+2.958	B.A.C.	94 56 2.3	2.67	B.A.C.	!!; c ¹ 42 Orionis & neb.....	2
1181	2878	Δ. 238??	5 28 32.1	-0.019	3	156 20 18.8	2.74	3	vB; vL; iE; vgpmbM	3
1182	III. 240	5 28 32.9	+2.498	1	113 26 9.6	2.68	1	vF; vS; stellar.....	1
1183	361	V. 31	44, ι Orionis	5 28 35.3	2.933	B.A.C.	96 0 21.0	2.66	B.A.C.	vF; vL; ι 44 Orionis inv ...	3
1184	362	5 28 36.5	2.969	1	94 26 50.5	2.65	1	Cl; vB; iRi; stL, sc	1
1185	III. 1??	{ M. 43= 144 Bo. Orionis }	5 28 38.4	+2.948	...	95 21 48.7	2.65	...	{ !vB; vL; R, with tail; } { mbM*8.9 }	Mon.*†
1186	2881	5 28 39.5	-0.393	1	159 5 49.5	2.75	1	Cl; vL; pRi; iF	1
1187	2882	5 28 39.8	-0.419	1	159 14 12.5	2.75	1	Cl; place of *	2
1188	359	III. 865	5 28 41.1	+3.897	1	58 6 38.4	2.62	1	cF; S; R; psbM	2
1189	2883	5 28 57.6	-0.581	4	160 4 57.4	2.72	4	B; pL; R; gbM	4
1190	2885	5 28 59.3	-0.739	2	160 50 53.1	2.73	2	F; L; iR; 3stp	2
1191	Chacornac	5 29 4.0	+3.581	...	68 52 20.4	!!!; variable (Chacornac) ...	0*
1192	2871	5 29 5.5	2.979	3	120 53 59.1	2.63	3	vF; S; R; lbM; st nr	3
1193	363	V. 34	ε Orionis	5 29 6.6	+3.042	1	91 17 44.7	2.61	1	!!!; eL; ε Orionis inv	2
1194	2884	5 29 12.9	-0.130	1	157 32 21.3	2.69	1	Cl; 4th of sev	1
1195	2873	5 29 13.9	+2.276	1	120 59 31.4	2.62	1	eeF; vS	1
1196	III. 269	5 29 20.4	+2.644	1	107 55 16.0	2.60	1	eF; vS; stellar.....	1*
1197	2887	5 29 21.2	-0.423	2	159 15 1.3	2.69	2	Cl; eS; st 11...16	2
1198	2879	5 29 24.4	+1.559	1	138 46 56.4	2.62	1	eeF; R; bM; diffic; p of 2...	1
1199	VIII. 42	5 29 32.9	3.711	2	64 15 45.5	2.55	2	Cl; L; IC; iRi	2
1200	2886	5 29 35.3	0.437	1	153 18 40.5	2.65	1	eF; cS; R	1
1201	2880	5 29 37.1	1.559	2	138 47 31.0	2.60	2	vF; R; gbM; st s; f of 2 ...	2
1202	IV. 33	5 29 37.6	+2.914	3	96 48 42.9	2.57	4	B* inv in N.....	4
1203	2889	5 29 45.0	-0.997	1	161 58 34.9	2.67	1	F; pL; R; vlbM	1
1204	2888	Δ. 178??	5 29 51.1	0.350	2	158 50 55.5	2.65	2	Cl; st 13m	2
1205	2890	Δ. 214?	5 30 25.2	-0.048	2	156 59 13.3	2.59	2	vB; S; R; * + neb in vLCl...	2
1206	2981	5 30 49.7	+0.014	1	156 34 8.5	2.55	1	B; S; stellar; r	1
1207	2893	Δ. 215	5 30 54.5	-0.110	6	157 23 18.5	2.55	6	⊕; B; pL; pRi; C; st 12 ...	6
1208	(509)	5 31 22.8	0.541	1::	159 51 17.4	2.52	1::	No description.....	1
1209	2895	5 31 26.5	-0.061	2	157 4 8.3	2.49	2	Cl; eL; R; vBvSNM	2
1210	2892	5 31 36.8	+1.433	1	141 1 12.1	2.43	1	eF; pL; R	1
1211	2894	5 31 44.8	+1.431	1	141 2 52.1	2.43	1	eF; pL; R; vlbM	1
1212	2897	5 31 59.0	-0.434	4	159 16 53.2	2.46	4	pF; pS; R; glbM; in Cl.....	4
1213	2898	5 32 15.9	0.764	1	160 55 58.8	2.44	1	F; cL; R; vglbM	1
1214	2899	5 32 40.9	0.149	1	157 37 1.3	2.39	1	vB; S; R; psmbM	1
1215	2907	5 32 41.8	-4.411	1	169 57 24.4	-2.52	1	vF; S; iE; bM; 2 st 9nf	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1216	h. 364	H.	h m s	s	1	24 16 50.7	-2.21	1	Cl; vRi; st 11.....	1
1217	2900	5 32 43.1	+6.02	3	157 47 24.3	2.39	3	Cl; pL; pC; iF; st 9...15 ...	3
1218	2901	5 32 48.0	-0.177	1	159 21 16.3	2.39	1	Cl; vL; Ri; vIC	1
1219	2902	5 32 52.0	0.448	1	160 3 5.6	2.38	1	F; vL; iR; gbM	1
1220	2896	5 32 57.5	-0.583	1	107 55 45.9	2.27	1	Cl of Lst	1
1221	2904	5 33 7.7	+2.643	1	161 10 19.2	2.36	1	pB; pL; R; pglbM; *10pinv	1
1222	2905	Δ. 98	5 33 16.8	-0.819	2	160 15 3.1	2.33	2	B; pL; gbM	2
1223	2903	Δ. 218	5 33 29.3	0.624	2	157 48 16.7	2.31	2	F; vL; vIE; vglbM	2
1224	2906	5 33 31.3	0.180	2	157 32 58.0	2.30	1	vF; S; R; in pLCl	1
1225	365	IV. 34	5 33 46.9	-0.139	1	80 59 3.8	2.14	2	O; pB; vS; vIE; r?	4†
1226	IV. 24	5 34 26.4	+3.283	2	92 18 43.1	2.13	1	B* in M of L, IE neb	1*†
1227	V. 28	5 34 40.0	3.019	1	91 55 43.7	2.11	2	lir; B; vvL; black sp incl ...	2
1228	2909	5 34 47.2	+3.028	2	161 47 23.1	2.23	1	vB; vS; IE; gmbM; r	1
1229	VIII. 28	5 34 47.8	-0.964	1	69 57 43.0	2.10	1	Cl; iRi; lC; st pL	1
1230	2908	Δ. 241	5 34 48.5	+3.557	1	157 0 29.6	2.18	3	Cl; vL; Ri; st 9...11	3
1231	2912	Δ. 100?	5 35 4.3	-0.058	3	160 2 26.6	2.18	1	vF	1
1232	2911	Δ. 240	5 35 11.2	0.585	1	157 38 43.5	2.15	2	pB; pL; R; gbM; in cLCl...	2
1233	2910	5 35 20.3	-0.158	2	156 6 50.8	2.14	2	pB; L; iR; gbM; 1st of 3 ...	2†
1234	2915	5 35 26.6	+0.072	2	161 5 22.5	2.15	2	⊕; B; pL; R; gbM; rr	2
1235	2913	Δ. 219?	5 35 33.9	-0.804	2	157 39 47.1	2.13	5	B; L; E; 2nd of 3	5†
1236	(579)	5 35 39.8	0.162	5	159 52 28.8	2.14	1::	Cl; no description	1
1237	2914	5 35 45.4	0.553	1::	156 58 42.7	2.11	1	Cl; vL; Ri	1
1238	2916	Δ. 220	5 35 51.8	0.055	1	157 40 20.7	2.11	3	B; L; R; bM; 3rd of 3	3†
1239	2917	5 35 52.8	0.163	3	159 8 38.7	2.11	1	vF; pL; R; gbM	1
1240	(593)	5 35 58.0	0.416	1	159 51 30.6	2.08	1::	Cl; no description	1
1241	2920	5 36 18.4	0.553	1::	160 38 22.6	2.08	3	pB; S; R; gbM; *9, np 5'...	3
1242	366	5 36 23.5	-0.710	3	81 25 37.9	1.97	1	Cl; vL; iRi; lC	1
1243	2918	5 36 25.7	+3.273	1	157 39 3.2	2.06	5	F; L; iR; glbM; r	5†
1244	2919	5 36 27.9	-0.160	5	157 4 23.8	2.04	4	B; S; R; vglbM	4
1245	2922	5 36 35.9	0.070	4	159 0 17.4	2.02	1	Cl; vL; Ri; st 12...15	1
1246	(608)	5 37 1.6	0.391	1	159 17 32.7	2.01	1::	Cl; no description	1
1247	367	Lal. 10842	5 37 7.7	-0.444	1::	77 10 43.0	1.90	1	*8, 9, with Fneb	1
1248	2923	5 37 8.3	+3.375	1	160 18 59.7	2.01	1	vF; R; gbM; 1st of 7	1†
1249	2925	5 37 11.1	-0.644	1	160 16 15.3	1.99	1	F; S; IE; 2nd of 7	1†
1250	2926	5 37 27.0	0.635	1	159 41 40.9	1.97	1	vF; L; pmE	1
1251	2921	5 37 35.8	-0.522	1	120 8 42.2	1.86	2	vF; S; R; bM	2
1252	2928	5 37 53.2	+2.300	2	159 28 11.8	1.94	1	Cl + neb; mC; iF; st vS.....	1
1253	2930	5 37 54.6	-0.479	1	161 5 21.8	1.94	1	pB; S; R; gbM	1
1254	2929	5 38 2.8	0.810	1	159 52 11.8	1.94	1	eF; vvS; vglbM	1
1255	2927	5 38 3.3	0.557	1:	157 30 8.4	1.92	1	F; pL; IE; gbM	1
1256	2931	5 38 3.9	0.139	1	159 29 59.7	1.91	1	Cl; vL; Ri; st 10...15	1
1257	2932	5 38 14.7	0.485	1	160 45 15.4	1.92	1	pB; R; bM; p of 2; *9 bet..	1
1258	2935	5 38 17.9	0.755	1	160 20 40.7	1.91	2	pF; S; R; gbM; 4th of 7 ...	2†
1259	2933	Δ. 102	5 38 24.3	0.652	1	160 14 33.7	1.91	2	vB; pL; R; gbM; 3rd of 7...	2†
1260	2936	5 38 24.6	0.631	2	160 12 26.7	1.91	1	vF; 5th of 7	2†
1261	(642)	5 38 27.3	0.624	1::	159 14 36.6	1.88	1	neb; no description.....	1
1262	2924	5 38 42.9	-0.438	1:	124 1 5.3	1.79	1	Cl; L; lC; st13	1
1263	2924	5 38 49.3	+2.174	1	156 56 50.8	1.84	1	vF; pS; E; gbM; 2st 10, s...	1
1264	2937	5 38 57.3	-0.055	1	81 16 35.1	1.73	2	Cl; poor; S sc st	3
1265	VIII. 2	5 39 6.6	+3.276	1	160 18 26.8	1.84	2	B; R; 6th of 7	2†
1265	2938	Δ. 103?	5 39 10.9	-0.646	2	160 15 5.4	1.82	1::	vF; vS; E; 7th of 7	1†
1266	2939	5 39 10.9	-0.646	2	90 0 15.7	1.61	2	B; L; wisp-sh; vgmbN; 3st	8†
1267	368	M. 78	5 39 21.0	-0.634	1::				inv; r.	
1268	2940	Δ. 143	5 39 27.3	+3.072	2	159 4 43.3	1.79	1	F; L; E	1
1269	2941	Δ. 142	5 39 34.1	-0.409	1	159 10 18.3	1.79	8	!!!vB; vL; looped	8†
1270	IV. 36	5 39 40.7	-0.423	8	89 46 49.6	1.68	3	* with vF, Lchev	3
1271	2934	III. 241	5 39 49.8	+3.077	3	112 3 59.6	1.68	1	eF; vS; R; gbM	2
1272	2942	5 39 59.5	+2.532	1	159 33 35.2	1.76	1	pB; pL; mE; 5st inv	1
1273	2943	5 40 1.0	-0.499	1	160 45 20.2	1.76	1	B; R; bM; rr; f of 2	1
1274	III. 267	5 40 9.0	-0.740	1	106 48 12.4	1.62	1	vF; pS; iE; bM	1
1275	2947	5 40 35.0	+2.670	1	159 43 48.3	-1.69	1	F; R; p of D neb	1
				5 40 49.6	-0.534	1					

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
1276	2948	5 40 53.3	-0.550	1	159 48 52.3	-1.69	1	neb; np of gr of 4	1†
1277	2949	Δ. 152??	5 40 54.2	0.556	1	159 50 42.3	1.69	1	neb; sp of gr of 4	1†
1278	2950	5 40 56.7	0.532	1	159 43 22.6	1.68	1	B; R; f of D neb	1†
1279	2951	5 41 10.5	-0.484	1	159 28 20.2	1.66	1	Cl; vF; mC; st+neb	1†
1280	2945	5 41 11.3	+0.297	3	154 21 38.1	1.63	3	pF; L; R; glbM	3
1281	2952	5 41 11.8	-0.549	1:	159 48 24.2	1.66	1:	neb; nf of gr of 7	1†
1282	2953	5 41 15.6	0.554	1:	159 49 51.5	1.65	1:	neb; st of gr of 7	1†
1283	2954	5 41 18.8	0.537	1:	159 44 57.5	1.65	1:	vF; R; *10vnr	1†
1284	2956	5 41 33.6	-0.536	1:	159 44 30.1	1.63	1:	B; pS; R; lbM; *10, p	1
1285	2946	5 41 38.9	+1.127	1	145 35 32.9	1.57	1	eF; pS; R; vlbM	1
1286	2955	5 41 39.6	-0.314	2	158 31 43.4	1.62	2	vF; S; R	2
1287	III. 270	5 41 43.7	+2.648	1	107 39 20.4	1.52	1	vF; eS; stellar	1*
1288	2944	Δ. 594	5 41 56.8	+2.163	2	124 18 21.4	1.52	2	⊕; B; pL; iR; gbM	2
1289	2957	5 42 5.1	-0.495	1	159 31 7.6	1.58	1	vF; S; mE; glbM; ?D	1
1290	2962	5 42 38.5	0.450	1	159 16 34.1	1.53	1	vF; pL; R; rr	1
1291	2963	Δ. 184??	5 42 43.0	0.397	1	158 59 18.4	1.52	1	vF; S; R	1
1292	2959	5 42 45.6	0.298	1	158 25 39.4	1.52	2	vF; S; R	1
1293	2961	5 42 47.9	0.128	2	157 23 26.7	1.51	2	Cl; F; cS; irr	2
1294	(725)	5 43 2.4	-0.319	1::	158 32 44.3	1.49	1::	neb; no description	1
1295	369	M. 37	5 43 7.5	+3.922	3	57 29 38.3	1.49	3	Cl; Ri; pCM; st L & S	8
1296	2960	5 43 6.1	+0.474	3	152 50 34.2	1.46	3	vF; pS; iR; pslbM*16	3
1297	2965	Δ. 185??	5 43 7.3	-0.282	1	158 20 10.3	1.49	2	⊕; B; S; rr	2
1298	2966	{ Δ. 147? 151? 154? }	5 43 9.6	-0.450	5	159 16 26.3	1.49	5	⊕; B; pL; irrR; rr	5
1299	2958	5 43 9.8	+1.358	1	142 8 17.1	1.43	1	eF; pS; R; 3st10 sf	1
1300	(730)	5 43 24.9	-0.503	1::	159 33 29.2	1.46	1::	neb; no description	1
1301	2968	5 43 44.7	-0.888	1	161 23 47.8	1.44	1	pB; L; pmE; gbM*13	1
1302	2964	5 43 49.9	+1.390	1	141 36 16.9	1.37	1	pB; pS; R; glbM	1
1303	2969	5 44 23.6	-0.064	1	156 58 0.9	1.37	2	F; pS; R; gbM	2
1304	2967	5 44 49.2	+2.544	1	111 36 16.5	1.25	1	vF; S; vLE; gbM	1
1305	2971	5 44 54.6	-0.736	2	160 42 13.8	1.34	2	pB; pS; R; gbM	2
1306	2970	Δ. 153?	5 44 58.2	0.444	1	159 14 0.1	1.33	1	eF; pL; IE	1
1307	2972	5 45 6.7	-0.331	3	158 35 56.7	1.31	3	F; pS; R; vglbM	3
1308	370	{ III. 448 = III. 510 }	5 45 30.3	+2.897	(4)	97 30 3.2	1.16	1	eF; cS; IE; pslbM; er	5
1309	2973	5 46 27.5	-0.817	2	161 2 46.7	1.21	2	vF; S; R; gbM	2
1310	371	VII. 24	5 46 37.1	+3.080	1	89 38 56.6	1.08	1	Cl; pL; lRi; pC; stS	3
1311	2975	5 46 40.1	-0.560	4	159 49 44.6	1.18	5	Cl; F; S; iF; vIC; rr	5
1312	2974	5 46 44.7	-0.246	1	158 5 35.9	1.17	1	eF; pL; iR	1
1313	2976	5 47 56.7	+1.446	1	140 37 4.7	1.01	1	eeF; vS; 3st10 sp	1
1314	2977	5 47 57.0	-0.326	1	158 33 11.2	1.06	1	F; S; R; *11p	1
1315	2978	5 48 7.4	0.149	5	157 29 33.8	1.04	5	F; pL; iR; vlbM; rrr	5
1316	2979	5 48 39.2	-0.438	2	159 10 22.7	1.01	2	⊕; vB; vS; vsmbM; rr	2
1317	2980	5 49 41.3	+0.369	2	153 42 41.3	0.89	2	eF; pL; R; vglbM	2
1318	2982	5 50 15.5	-0.922	1	161 30 44.6	0.88	1	vF; eL; vgbM	1
1319	2981	Δ. 106	5 50 16.8	-0.619	5	160 6 17.6	0.88	5	Cl; pB; iF; gvmCM; st15	5
1320	2983	5 51 17.8	+0.162	3	155 20 50.2	0.76	3	pB; vS; R; gbM	3
1321	III. 225	5 51 46.8	+2.584	1	110 3 13.8	0.64	1	eeF; pS; E; r	1
1322	2985	5 51 59.6	-0.505	1	159 31 16.4	0.72	1	vF; pS; R; gbM	1
1323	VIII. 68	5 52 7.3	+4.659	1	40 6 11.5	0.55	1	Cl, not Ri; 1*7m	1
1324	2986	5 52 28.5	-0.481	3	159 23 33.9	0.67	3	pB; vS; R; gmbM	3
1325	372	VIII. 26	5 52 34.6	+3.647	1	66 42 23.8	0.54	1	Cl; pL; 40 or 50 st 8...15	2
1326	2987	5 52 41.1	-0.131	7	157 21 35.8	0.64	7	F; pS; R; glbM	7
1327	2984	5 53 11.9	+2.401	1	116 39 52.1	0.53	1	vF; pS; R; gbM	1
1328	2988	5 53 22.1	+0.765	1	149 55 53.5	0.55	1	Cl; vIC; st L & S	1
1329	2989	5 53 26.1	-0.854	2	161 12 17.0	0.60	2	F; pL; R; gpmbM	2
1330	2991	5 53 53.9	0.837	4	161 7 52.5	0.55	4	⊕; B; pL; R; gmbM; r	4
1331	2990	5 53 54.3	0.146	5	157 27 0.8	0.54	5	F; pS; R; r; am st	5
1332	2992	Δ. 160	5 54 7.4	-0.504	4	159 30 51.1	-0.53	4	⊕; pB; R; gmbM; rr; st 14...16.	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1333	h. 2994	H.	h m s 5 54 23.3	s -0.503	2	159° 30' 24.0"	-0.50	2	vF; S; R; f of 2	2
1334	2993	5 54 34.6	+0.090	1	155 51 30.9	0.47	1	eF; S; R	1
1335	II. 264	5 54 55.9	+2.482	1	113 49 28.9	0.37	1	F; S	1
1336	2995	5 55 1.4	-0.342	3	158 37 0.8	0.44	3	pF; pS; iR; bM	3
1337	373	5 55 15.5	+2.822	1	100 36 14.8	0.34	1	* (3 Monoc) inv in pL, F, n.	1
1338	374	5 55 39.0	+3.206	1	84 16 44.3	0.29	1	Cl; L; pRi; vIC; st10,	1
1339	3009	5 56 0.0	-6.638	2	172 9 27.8	0.54	2	F; pS; iR; bM	2
1340	2998	5 56 13.4	0.798	1	160 56 36.5	0.35	1	F; iE; r	1
1341	2997	5 56 22.5	-0.274	2	158 13 12.4	0.32	2	eF; S; R; bM	2
1342	2996	5 56 50.7	+0.836	1	149 7 34.5	0.25	1	eF; S; R; *12 vnr	1
1343	3000	5 57 0.8	-0.519	1	159 34 54.6	0.28	1	F; vS; R; vsmBM; stellar	1
1344	3001	5 57 13.7	-0.416	1	159 2 0.5	0.25	1	F; pS; R; bM	1
1345	2999	5 57 34.0	+1.437	2	140 44 5.6	0.18	2	eeF; R; *15 att	2
1346	3002	5 57 46.1	+0.010	1	156 24 55.3	0.19	1	eeF; iE; *16 att	1
1347	3003	5 57 52.5	-0.119	4	157 16 2.3	0.19	4	F; pL; R; vglbM	4
1348	3004	5 58 9.8	+0.141	3	155 28 41.2	0.16	3	F; pL; R; vglbM	3
1349	3005	Δ. 196	5 58 32.0	-0.316	5	158 28 3.8	0.14	5	pB; S; R; gbM; 1st of 3	5
1350	3006	Δ. 161?	5 58 35.9	-0.447	4	159 11 59.1	-0.13	4	⊕; vB; S; R; vgvmbM; rr.	4
1351	375	VI. 17	5 58 49.3	+3.670	1	65 53 46.0	0.00	1	Cl; pS; mC; vRi; nrΔ; steS.	3
1352	3007	Δ. 193	5 58 50.0	-0.345	4	158 38 5.4	-0.12	4	pF; S; R; gbM; *15 att nf.	4
1353	3008	5 58 50.7	0.287	1	158 17 48.7	0.11	1	pF; pS; R; gbM	1
1354	3013	5 59 33.7	-1.702	1	164 21 22.3	0.09	1	F; pL; R; gpmBM	1
1355	3010	5 59 36.3	+0.365	3	153 43 19.4	0.02	3	F; pL; R; vglbM	3
1356	3011	Δ 194	5 59 39.1	-0.325	5	158 31 0.8	-0.04	5	⊕; vB; R; mCM; rr	5
1357	376	5 59 59.5	+4.765	1	38 17 55.2	+0.14	1	Cl; pL; poor; st11	1
1358	3012	Δ. 223?	6 0 4.7	-0.229	1	157 56 54.0	0.00	1	F; S; R; gbM	1
1359	378	IV. 44	6 0 8.7	+2.927	1	96 11 46.0	+0.10	1	Nebulous *7; am 3 st	2
1360	377	M. 35	6 0 12.5	3.677	1	65 39 16.9	0.13	1	Cl; vL; cRi; pC; st 9...16...	8
1361	379	VIII. 24	6 0 33.4	3.405	1	76 1 38.5	0.15	1	Cl; S; IRi; pmC; *Σ. 848...	3
1362	IV. 19	6 0 44.0	+2.923	3	96 22 39.5	0.15	3	*9 in vF, pLneb; E 170°	3
1363	3016	6 0 44.6	-0.750	1	160 43 11.2	0.04	1	eF; L; R; glbM	1
1364	3015	6 0 52.1	0.348	2	158 38 48.1	0.07	2	F; cL; R; lbM	2
1365	3018	6 0 56.3	-1.295	1	162 58 41.2	0.04	1	pF; pL; R; gmbM	1
1366	Auw. N. 21	6 1 19.1	+3.569	...	69 29 42.5	0.01	...	*8m in neb (Bruhns)	0
1367	3017	6 1 27.2	-0.056	1	156 51 19.9	0.13	1	eeF; pL; R; gbM	1
1368	3020	6 1 42.0	-0.192	2	157 43 40.5	0.15	2	F; vS; iR; lbM; r	2
1369	3019	6 1 56.1	+0.359	1	153 45 54.4	0.18	1	eF; vS; R	1
1370	3014	6 2 6.2	2.539	1	111 43 53.8	0.26	1	F; pS; vmE; glbM	1
1371	380	VIII. 6	6 2 12.0	3.182	1	85 15 56.7	0.29	1	Cl; pRi; iC; st L & S	4
1372	3021	6 2 28.9	0.171	1	155 15 14.6	0.22	1	vF; S; R	1
1373	381	IV. 38	6 2 41.6	2.924	1	96 18 57.6	0.32	1	pB *; L*neb; E 90°±	3
1374	382	6 3 59.9	2.990	1	93 30 5.2	0.44	1	Cl; L; vR	1
1375	383	IV. 20	6 4 17.9	2.927	2	96 12 6.1	0.47	2	*11&4 S st in vF, L neb	5
1376	384	VII. 25	6 4 39.8	+3.200	1	84 31 53.0	0.50	1	Cl; pL; pRi; pC; st L & S	2
1377	3025	6 5 3.8	-0.516	3	159 33 49.6	0.42	3	{ pB; pS; R; gbM } D neb; { vF; R; glbM } 12°.5	3
1378	3022	6 5 5.6	+2.168	1	124 4 30.3	0.51	1	pF; pL; vmE; gvlbM	1
5064	6 5 7	88 50 39	See No. 5064.	...
1379	3027	6 5 7.1	-1.817	1	164 42 26.7	0.39	1	vF; pL; R; glbM	1
1380	3023	6 5 16.5	+1.331	1	142 29 20.0	0.50	1	pB; vS; E; vsbM; *9 p 5 ^s ...	1
1381	VII. 57	6 5 25.2	4.189	1	50 6 21.0	0.60	1	Cl; cL; C; iF; st vS	1
1382	3026	6 5 47.6	0.196	1	155 4 9.3	0.51	1	F; iF; glbM; 2 or 3 st inv	1
1383	VI. 5	6 6 2.3	3.377	2	77 9 55.9	0.63	2	Cl; L; Ri; gvmCM	2
1384	3024	II. 265	6 6 14.4	+2.539	4	111 46 26.6	0.62	4	pF; pS; vIE; pmbM; st nr	5
1385	3028	6 6 21.3	-0.087	2	157 4 22.5	+0.55	2	vF; pS; R; gbM	2
5065	6 6 40.9	88 58 11.2	See 5065.	...
1386	3031	6 7 57.9	-1.404	1	163 22 10.8	+0.66	1	F; vS; R; bM	1
1387	3029	6 9 5.0	+1.799	2	133 37 23.5	0.85	2	eF; pS; R; vlbM; ?134° PD.	2
1388	3030	6 9 18.6	1.798	2	133 39 27.1	0.87	2	eF; S; R; pslbM; ?134° PD.	2
1389	385	Σ. 885	6 9 23.0	+3.213	1	83 58 10.6	0.92	1	* Chief of Cl	1
1390	3035	6 9 26.6	-2.061	1	165 24 21.8	+0.76	1	pB; pL; iR; vgpmbM; r	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1391	h.	H.		h m s	s		108° 36' 47.3	+0.91	1	Cl; L; pRi; IC	1
1392	3034	VII. 13	6 9 33.3	+2.622	1	152 29 50.0	0.90	1	pF; S; R; bM	1
1393	3032	6 10 13.2	0.506	1	111 19 46.4	0.98	1	pB; pL; mE, 87°; psIbMRN	1
1394	3033	6 10 22.8	2.551	1	116 43 35.4	0.98	2	F; pS; vIE; psIbM	3
1395	3037	6 10 23.9	+2.399	2	163 47 55.3	1.01	1	vF; cL; R; gvlbM	1
1396	3036	6 12 5.0	-1.526	1	159 5 2.4	1.08	3	vB; pL; R; mbM; r	3
1397	3038	6 12 33.1	0.423	3	161 29 30.7	1.09	1	vF; S; R; gIbM; * p	1
1398	3039	6 12 45.1	0.917	1	158 13 12.4	1.18	2	B; pS; IE; gbM; rrr	2
1399	386	Δ. 201	6 13 34.4	-0.270	2	97 14 22.3	1.31	2	Cl; cL; pRi; pC; st 11...15.	5
1400	3040	VII. 20	6 14 5.0	+2.902	2	112 0 58.2	1.44	1	vF; pL; R; vglbM	1
1401	3041	6 15 36.9	2.532	1	117 10 38.1	1.47	2	vB; S; R; psmbM; r	2
1402	Auw. N. 22	6 16 4.8	2.387	2	70 35 28.2	1.54	...	F Cl (Markree Obs. Jan. 13, 1853).	0
1402	6 16 25.2	3.539	...	94 37 19.9	1.53	1	Cl; P; vIC; st 6, 11...12	1
1403	387	6 16 28.9	2.964	1	134 41 54.5	1.55	1	Cl; B; P; st 8, ...	1
1404	3042	6 17 5.4	1.753	1	147 29 35.0	1.60	1	vF; IE; vgbM; p of 2	1
1405	3044	6 18 1.5	0.980	1	147 27 30.0	1.60	1	vF; IE; vglbM; f of 2	1
1406	3045	6 18 1.9	0.983	1	112 46 8.0	1.70	3	F; pL; R; vglbM; 2st inv...	3
1407	3043	6 18 43.9	2.513	3	77 16 45.6	1.82	1	Cl; pC; with neb?	1
1408	VII. 35	6 19 44.5	3.372	1	99 34 30.9	1.83	1	Cl; P; ICM; st 12...15	2
1409	388	VII. 26	6 19 57.6	2.847	1	111 55 24.9	1.83	1	eF; R; * p 270°, 90'	1
1410	3046	6 20 1.4	2.536	1	154 23 21.3	1.81	1	F; S; R; gIbM	1
1411	3047	6 20 33.0	0.293	1	154 52 49.6	1.82	2	eF; vS; R; 1st of 3	2
1412	3048	6 20 49.4	0.231	1	154 55 25.2	1.84	1	eF; S; IE; 2nd of 3	1
1413	3049	6 20 55.3	+0.226	1	157 27 7.2	1.84	3	F; pL; R; gvlbM; * f	3
1414	3050	6 21 0.8	-0.136	3	94 40 44.6	1.92	1	B* (10 Monoc) + Cl	1
1415	VIII. 25	6 21 1.6	+2.963	1	154 58 9.5	1.85	1	eF; S; 3rd of 3	1
1416	3051	6 21 11.0	0.220	1	73 13 33.5	1.95	1	Cl; eL; pRi; IC; st L & S...	2
1417	389	VIII. 9	6 21 13.5	3.473	(1)	154 51 45.3	1.91	1	vF; S; R; * 12 nr	1
1418	3052	6 21 49.5	0.235	1	83 4 23.2	2.04	1	Cl; pRi; pC; st 10, 12...15...	2+
1419	390	VII. 5	6 22 11.0	3.233	1	84 57 32.2	2.14	1	* 8 in L; P; BCl	4
1420	392	6 23 28.8	3.189	1	54 42 17.1	2.17	2	Cl; pL; P; vIC; st 7, 10...15	3
1421	391	VIII. 49	6 23 31.7	+4.013	2	158 50 33.8	2.06	3	vF; pL; R; gIbM	3
1422	3054	6 23 42.4	-0.366	3	121 11 28.4	2.18	2	pB; cL; R; vglbM; 4'	2
1423	3053	Δ. 616?	6 24 13.8	+2.267	2	85 2 14.5	2.27	B.A.C.	Cl; beautiful; st sc	
1424	VII. 2	12 Monoc. B.A.C.	6 24 53.4	3.189	B.A.C.	79 44 43.1	2.27	2	pL; com; mbNsf alm*; *7.8 nf.	7*+
1425	393	IV. 3	6 24 58.0	3.312	2	63 35 16.9	2.36	...	Small cluster (Markree Obs. Dec. 23, 1853).	0
1426	Auw. N. 23	6 25 51.9	+3.729	...	158 50 0.3	2.31	4	pB; pL; R; vgbM; * p	4
1427	3055	6 26 38.1	-0.360	4	94 57 53.2	2.44	1	Cl; pRi; IC; iF; st 8, 12...14	1
1428	394	6 27 5.0	+2.956	1	81 32 16.8	2.46	(2)	Cl; vL; E; Ri; IC	4
1429	395	VIII. 3	6 27 8.4	3.269	(2)	84 32 14.7	2.49	1::	Cl; vL; pRi; IC; st S	3
1430	396	VIII. 50	6 27 23.7	3.199	(1)	24 2 15.6	2.62	1	vF; st eS	1
1431	VII. 54	6 27 56.0	6.043	1	82 13 44.4	2.58	1	Cl; S; pC; iF; st 11...15	2
1432	397	VII. 22	6 28 25.1	3.253	1	124 42 52.7	2.59	2	eF; S; IE; vIbM	2
1433	3056	6 28 59.1	2.153	2	154 13 23.4	2.58	3	F; cL; R; vglbM; r; 17 ^s .0d	3
1434	3057	6 29 25.9	0.325	3	79 0 33.4	2.78	1	Cl; cRi; eC; iF; st eS	1
1435	VI. 28	6 30 47.9	3.329	1	91 20 42.7	2.79	1	Cl; vL; P; vIC; st L & S	2
1436	398	VIII. 48	6 31 2.3	3.041	1	81 8 20.5	2.85	3	B; vmE 330°; Ncom = *11	7+
1437	399	IV. 2	6 31 31.4	3.278	3	88 43 47.6	2.92	1	Cl; vC; iR; bM; st eS	2
1438	400	VII. 37	6 32 24.0	3.101	1	114 43 55.6	2.92	1	pF; IE; bet 2 vS st; psIbM...	1
1439	3058	6 32 41.8	2.462	1	79 59 24.7	2.99	1	15 Monoc; Cl; *; ? neb	6*
1440	401	{ V. 27 VIII. 5 }	15 Monoc.	6 33 16.0	3.305	1	77 57 3.2	3.04	1	Cl; P; 30 or 40 st 12...13	1
1441	402	6 33 40.5	3.354	1	62 53 35.6	3.12	1	Cl; pS; eC; Ri; st 11...15	2
1442	403	VI. 21	6 34 32.6	3.748	1	122 20 49.8	3.16	1	pB; S; R; 2 or 3 st v nr	1
1443	3059	6 35 31.5	2.236	1	85 17 41.1	3.27	1	Cl; vmC; not Ri; st vS	2
1444	404	VI. 3	6 36 26.7	3.180	(1)	86 24 51.4	3.28	1	Cl; IC; not Ri	2
1445	405	VII. 36	6 36 35.5	3.154	1	113 20 15.0	3.30	3	pF; S; R; gbM; am st	3
1446	3060	6 37 3.1	2.503	3	117 19 37.3	+3.31	3	pF; pS; vIE; bM; r	3
1447	3061	6 37 7.0	+2.390	3					

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1448	h. 406	H. II. 615	h m s 6 38 2.9	s +3.950	2	56 17 42.9	+3.43	2	F; S; bM.....	4
1449	407	II. 614	6 38 2.9	3.951	2	56 15 56.9	3.43	2	eF; vS.....	4
1450	3062	6 39 16.6	2.386	1	117 30 7.7	3.49	1	pF; pL; lE; gbM.....	1
1451	VIII. 71	6 39 30.5	4.223	1	48 47 17.8	3.56	1	Cl; pRi; vIC; st pL.....	1
1452	III. 271	6 39 46.2	2.642	1	108 3 23.8	3.56	1	3 or 4 S st + neb.....	1*
1453	408	VIII. 31	6 40 39.4	3.002	1	93 1 23.9	3.63	1	Cl; L; C; ab 100 st 9...15...	3
1454	411	M. 41	6 41 0.3	2.578	1:	110 36 2.2	3.64	1	vL; B; lC; st 8,...	3*
1455	410, a	R. nova	6 41 34.1	3.944	...	56 23 51.2	3.74	...	No desc.; β of Lord R.'s diag.	0
1456	410, b	R. nova	6 41 35.9	3.944	...	56 22 58.2	3.74	...	No desc.; γ of Lord R.'s diag.	0*
1457	410	III. 898	6 41 39.3	3.944	1:	56 25 16.2	3.74	1	eF; vS.....	2
1458	409	III. 897	6 41 39.4	3.946	1:	56 21 16.2	3.74	1	eF; vS.....	2
1459	3063	6 41 44.2	2.414	1	116 35 49.0	3.70	1	{pB; R; gbM} D neb; am st {eF; R; gbM}	1
1460	410, c	R. nova	6 41 54.0	3.944	...	56 19 55.2	3.74	...	No desc.; ε of Lord R.'s diag.	0*
1461	3064	6 42 3.3	2.414	1	116 34 19.9	3.73	1	eF; S; R; bet st; D neb p ...	1
1462	3066	6 43 23.6	0.428	1	153 34 16.4	3.78	1	vF; S; R; vglbM.....	1
1463	3065	Δ. 578	6 44 1.7	2.124	4	125 50 56.4	3.88	4	⊕; B; pL; iR; gbM; rr.....	4
1464	412	6 44 17.3	2.915	1	96 49 35.2	3.94	1	Cl of 30 or 40 st.....	1
1465	413	VI. 27	6 44 35.1	3.086	2	89 22 44.1	3.97	3	Cl; Ri; L; iF; st L & S.....	5
1466	414	VIII. 39	6 45 3.1	2.912	3	96 55 13.0	4.00	3	Cl; L; P; lC.....	6
1467	415	VI. 2	6 46 55.4	3.501	2	71 49 12.7	4.09	2	Cl; pL; Ri; mC; st vS.....	5+
1468	3067	6 47 45.6	0.375	1	154 6 43.8	4.16	1	vF; vS; R; 2 st Δ.....	1
1469	416	VIII. 51	6 47 47.1	2.910	1	97 1 26.2	4.24	1	Cl; P; vIC.....	4
1470	3068	6 47 58.3	0.369	2	154 9 52.4	4.18	2	vF; pS; vIE 90°.....	2
1471	417	VI. 18	6 49 15.4	2.911	3	97 1 24.8	4.36	3	Cl; pL; pRi; mC; st 13... ..	7
1472	3069	6 49 23.0	1.949	2	130 41 29.2	4.34	2	pB; pL; Ri; vme E 44°8; pslbM..	2
1473	418	VIII. 60	6 50 51.9	2.971	1	94 24 15.0	4.50	1	Cl; lC; not Ri.....	2
1474	419	6 51 6.4	3.311	1	79 33 25.9	4.53	1	Cl; P.....	1
1475	D'Arrest, 50	6 51 23	2.89	[3]	97 45 24	4.51	[3]	F; vS; R.....	0
1476	420	6 51 49.9	4.665	2	39 12 51.9	4.63	2	eF.....	2
1477	421	II. 304	6 52 55.6	2.898	3	97 35 39.1	4.67	3	pF; S; R; r; S st inv.....	6+
1478	421, a	R. nova	6 52	97 35	Makes a close D neb with h. 421.	0
1479	{ 422 = 3070 }	VII. 14	6 53 4.4	2.759	2	103 30 44.4	4.68	2	Cl; L; Sc; st 8...9.....	3
1480	423	VIII. 1B	6 53 48.2	3.145	2	86 44 53.5	4.75	2	Cl of sc st; st 8, 9,	3*
1481	III. 874	6 54 46.7	4.658	1	39 15 39.4	4.88	1	vF; vS; lE.....	1
1482	424	II. 861	6 54 57.4	4.661	1	39 13 10.3	4.81	1	pB; S; iR; gbM; *8, 120°...	2
1483	425	M. 50	6 56 12.5	2.886	4	98 8 46.5	4.95	4	l Cl; vl; Ri; pC; E; st 12...16	8
1484	427	VII. 38	6 56 57.0	3.100	2	88 44 31.6	5.02	3	Cl; L; Ri; cC; st 12...16 ...	5
1485	3071	6 57 7.4	2.368	1	118 30 13.3	5.01	1	pB; pL; lE; gbM.....	1
1486	426	II. 734	6 57 28.6	4.663	1	39 5 50.6	5.02	1	vF; pL; iR; psmbM; st p ...	2
1487	428	IV. 25	6 57 33.2	2.817	1	101 6 47.8	5.06	1	pB* inv in S, vF, neb.....	3
1488	3072	6 58 6.8	1.912	2	131 51 55.4	5.08	2	vF; S; vIE; bM; am st.....	2
1489	429 { II. 735 = III. 875 }	6 58 33.7	4.547	1	41 10 42.0	5.20	1	vF; vS; stellar.....	4
1490	432	VIII. 40	6 58 47.5	3.741	2	62 35 40.7	5.19	2	Cl; L; vIC; ScI inv.....	3
1491	430	II. 862	6 58 49.2	4.631	1	39 36 2.6	5.22	1	F; S; R; psbM.....	2
1492	430, a	R. nova	6 58 ±	39 36 ±	Several near h. 430 (? 426, 433 & 1 nov).	0
1493	431	III. 899	6 58 54.3	3.986	(1)	54 39 36.3	5.21	1:	vF; S; R; bM.....	2
1494	VIII. 32	6 59 54.6	2.847	1	99 52 4.8	5.26	1	Cl; L; lC.....	1
1495	435	7 0 6.6	2.949	1	95 24 37.7	5.29	1	Cl; vIC.....	1
1496	434	II. 769	7 0 9.7	3.515	1	71 0 17.0	5.30	1	pB; pL; R; glbM.....	2
1497	433	II. 736	7 0 27.9	4.627	2	39 36 26.8	5.36	2	pF; S; R; glbM; r.....	4
1498	VIII. 33	7 1 35.7	2.834	1	100 26 14.0	5.40	1	Cl; cL; P; lC.....	1
1499	3073	7 1 52.0	2.775	1	102 57 4.9	5.43	1	Cl; pL; pRi; gbM; st 10...14	1
1500	IV. 65	7 2 14.0	3.055	2	90 30 8.1	5.47	1	*9 aff with S, vF, neb.....	1
1501	III. 746	7 2 47.5	+5.827	1	24 57 49.7	+5.59	1	vF; S; R; lbM.....	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1502	h. 3074	H.	h m s 7 3 7.1	s +0.009	1	157° 11' 8.8"	+5.46	1	Cl; P; IC; 30 st ±	1
1503	436	VII. 27	C. H.	7 3 16.3	2.881	1	98 23 59.5	5.55	1	Cl; cL; P; cC	3
1504	437	7 6 55.9	2.817	1	101 15 9.5	5.85	1	Cl; IC; * taken	1
1505	VII. 15	7 7 48.2	2.506	1	113 51 10.6	5.92	1	Cl; pRi; pC	1
1506	VIII. 34	7 7 50.5	2.845	1	100 3 44.9	5.93	1	Cl; L; IC; one vB*	1
1507	438	VII. 16	7 8 28.2	2.463	1	115 29 29.4	5.98	1	Cl; cRi; IC	2
1508	439	VI. 6	7 9 3.8	3.390	1	75 58 53.5	6.05	1	Cl; pS; pRi; mC; st 15...16	2*
1509	VII. 6	7 9 12.4	3.394	1	75 46 51.8	6.06	1	Cl; IC	1
1510	VIII. 45	7 10 31.8	2.687	2	106 47 59.8	6.16	2	Cl; P; IC	1
1511	3075	V. 21	7 11 3.0	2.778	1	102 57 54.0	6.20	1	!!; vF; vL; viF	3†
1512	{ 440 = 3076 441 = 3077 }	VII. 12	C. H.	7 11 23.4	2.721	2	105 23 19.9	6.23	2	Cl; vL; Ri; pC; st 9...12 ...	5
1513	{ 442 = 3077 }	VII. 17	A.S.C. 905	7 12 54.6	2.488	2	114 42 14.2	6.34	2	Cl; pL; Ri	4
1514	442	7 13 59.7	2.908	1	97 18 9.5	6.05	1	Cl; pC; st pL; bifid	1
1515	III. 748	7 14 2.3	6.410	1	20 42 16.2	6.54	1	vB; pL; R; mbM; r; vS* inv.	1
1516	VIII. 27	7 14 8.9	2.568	1	111 40 18.5	6.45	1	Cl; S; P; IC	1
1517	443	7 14 20.5	2.845	1	100 7 38.4	6.48	1	Cl; S; pRi; st 15	1
1518	3078	7 15 0.5	0.679	2	152 5 45.1	6.47	2	pB; pL; iE; glbM	2
1519	444	II. 316	7 16 42.9	3.792	1	60 14 41.0	6.70	1	B; S; R; bMN; p of D neb, 45°, 60°.	3†
1520	445	II. 317	7 16 44.7	3.792	1	60 14 21.0	6.70	1	pB; S; R; bMN; f of D neb	3†
1521	3080	VIII. 35	7 17 31.6	2.781	1	102 59 33.2	6.74	1	Cl; vL; pRi; IC; st L	5
1522	3079	7 18 18.9	2.423	1	157 15 54.7	6.79	1	pF; pS; R; vsmbM; am st ...	1
1523	3084	7 18 27.0	0.617	1	112 48 48.5	6.75	1	vF; vS; R; am st	1
1524	3082	7 18 32.8	2.427	1?	117 6 24.0	6.80	1.:	pF; S; R; bM	1
1525	3081	7 18 39.7	2.597	1	110 39 57.6	6.82	1	Cl; pS; pmC; st 12 ...	2
1526	3083	7 18 59.7	2.595	2	110 44 57.5	6.85	2	Cl; IC; bifid; *	2
1527	446	7 19 17.6	3.920	1?	55 54 27.3	6.91	1.:	neb; 1st of 4	1*
1528	447	III. 703	7 19 23.1	3.921	1	55 53 27.3	6.91	1.:	vF; vS; R; bM	3*
1529	II. 820	7 19 44.8	4.017	1	52 58 44.5	6.95	1	pB; S; stellar	1
1530	448	III. 900	7 19 45.1	3.920	1	55 55 8.2	6.95	1	vF; S; R; bM	2*
1531	449	III. 901	7 19 57.6	3.921	1	55 51 47.1	6.97	1	vF; S; R; psbM	2*
1532	450	IV. 45	7 20 54.4	3.557	1	68 48 33.2	7.04	1	B; S; R; *8 M	4†
1533	VIII. 44	7 21 2.5	3.232	1	82 41 1.2	7.04	1	Cl; L; P; vIC; st L	1*
1534	VIII. 11	7 21 11.7	3.384	1	75 56 51.5	7.05	1	Cl; pRi; C	1
1535	451	VIII. 36	7 21 38.5	+2.818	1?	101 27 32.1	7.07	1?	Cl; vL; vIC	2
1536	3085	7 21 42.1	-0.151	2	158 43 56.0	7.00	2	pB; cL; cE 117°, lbM	2
1537	Auw. N. 24	7 22 32.1	+3.070	::	89 55 49.5	7.15	...	} Two B neb (Bond, Feb. 1853).	0
1538	Auw. N. 25	7 22 32.1	3.070	::	89 55 49.5	7.15	...		0
1539	454	VII. 65	7 22 57.2	2.767	1	103 41 23.1	7.17	1	Cl; S; cRi; cC; st vS	2
1540	453	III. 19	7 23 6.9	3.250	2	80 3 25.0	7.20	3	eF; S; R; lbM; * inv	4
1541	V. 44	7 23 18.6	5.864	1	24 0 3.0	7.30	1	!!; cB; eL; vmE; vgbMN 7'	2
1542	452	7 24 35.7	6.898	1	18 1 37.9	7.43	1	Cl; vIC	1
1543	3086	7 25 21.7	2.693	1	106 54 16.1	7.37	1	Cl; S but B; st 8...10	1
1544	VIII. 52	7 26 49.5	2.789	1	102 47 49.7	7.49	1	Cl; vL; P; vIC	1
1545	455	VIII. 37	7 26 49.5	2.735	1	105 8 46.7	7.49	1	Cl; P; IC; st 9, &c	3
1546	456	II. 821	7 27 43.4	3.954	1	54 28 22.7	7.59	2	pB; cS; R; vglbM; r; alm○	3
1547	3087	7 28 25.2	0.743	1	151 57 47.1	7.57	1	vF; L; R; gbM; r	1
1548	457	I. 218	7 28 37.8	4.077	1	50 48 47.4	7.68	1	pB; pL; iE 90°; vgbM; *7, 8, 19°.	3
1549	458	VI. 1	7 30 5.7	3.566	1	68 7 10.4	7.78	1	Cl; cL; Ri; C; st 11...18 ...	10
1550	3089	VII. 67	7 30 10.0	2.615	1	110 18 0.8	7.76	1	Cl; L; cRi; st 11...13	3
1551	{ 459 = 3088 }	VIII. 38	7 30 10.8	2.760	2	104 10 31.8	7.76	2	Cl; B; vL; pRi; st L & S ...	4
1552	3090	VII. 28	7 30 38.0	2.774	1	103 32 55.0	7.80	1	Cl; vL; Ri; pC; st vS	2
1553	VIII. 87	7 31 51.7	+2.751	1	104 35 14.0	+7.90	1	Cl; P; S; st vS	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1554	h. 460	H. II. 822	7 32 26.8	+4.676	1	37° 20' 50.0"	+8.00	1	eF; R; vgbM; r; *8 p	2
1555	3091	7 32 31.5	1.739	1	137 18 28.9	7.93	1	eF; L; pmE; gmbM; 2 st inv	1
1556	VIII. 47	7 32 55.5	2.715	1	106 11 49.4	7.98	1	Cl; vL; vIC.....	1
1557	VIII. 46	7 33 4.6	2.719	1	106 1 50.0	8.00	1	Cl; vL; vIC.....	1
1558	III. 829	7 34 26.7	4.715	1	36 34 59.8	8.16	1	eF; vS; R; bM	1
1559	3092	VI. 36	7 34 41.6	2.656	1	108 45 38.6	8.12	1	Cl; pL; pC; E 0°; st L & S... ..	3
1560	462	7 35 4.7	+3.277	1	80 24 50.1	8.17	1	eF; *15, 300°.0, 90''	1
1561	3096	7 35 9.9	-0.117	5	158 58 3.7	8.09	5	pB; S; R; pmbM; 3 st 11 n... ..	5
1562	II. 616	7 35 11.2	+3.833	1	57 59 59.0	8.20	1	F; S; lbM	1
1563	461	7 35 19.3	4.653	1	37 35 56.6	8.22	1	vF; vS; R; bM	1
1564	463	M. 46	7 35 24.3	2.755	1	104 29 50.4	8.18	1	!; Cl; vB; vRi; vL; inv O... ..	4
1565	{ 464 = 3093 }	IV. 39	7 35 25.4	2.757	2	104 24 39.4	8.18	3	O; pB; pS; eE; r; 3°.75 d... ..	4†
1566	3094	7 35 26.2	2.328	1	121 19 45.1	8.17	1	Cl; B; pRi; pL; IC; st 9, 12...14.	1
1567	3095	IV. 64	7 35 41.2	+2.677	1	107 53 22.3	8.21	1	O; cB; not v well def	3†
1568	3097	7 36 43.9	-0.149	4	159 12 49.3	8.21	4	{ cL; vF; R } D neb; 40°; { pL; vF; R } * inv M	4
1569											
1570	465	7 37 36.4	+4.803	1	35 3 14.3	8.41	1	F; am 4 st	1
1571	3098	M. 93	7 38 39.2	2.542	1	113 32 43.2	8.44	1	Cl; L; pRi; IC; st 8...13	2
1572	466	Lal. 15134	7 38 41.4	2.522	1	114 21 12.2	8.44	1	Cl of 18 or 20 st 11...13	1
1573	3099	7 40 19.2	2.138	1	127 38 15.8	8.56	1	Cl; vvL; vIC; 1* 4.5 m.....	1
1574	3100	7 41 47.4	2.457	2	117 0 6.7	8.69	2	O; F; S; IE; am 60 st	2
1575	3101	7 41 56.8	2.459	1::	116 54 21.0	8.70	1::	Cl; S; pRi; pC	1
1576	3102	7 42 52.2	2.611	1	110 57 4.1	8.77	1	Cl; cL; pRi; IC; st 12	1
1577	467	7 42 56.7	4.838	1	34 9 26.9	8.83	1	vF; R; vgbM	1
1578	468	III. 479	7 44 24.4	3.280	2	80 5 29.5	8.91	2	vF; S; rr group + neb	3*
1579	469	7 45 47.6	+4.908	1	32 57 44.1	9.07	1	eF; R; p of 2	1
1580	3104	7 46 20.9	-0.423	1	161 3 33.8	8.96	1	vF; S; R; lbM	1
1581	469, a	R. 8 novæ	7 47 ±	32 57 ±	8 of 10 neb, in line with h. 469, 470.	0
1582											
1583											
1584											
1585											
1586											
1587											
1588											
1589	472	IV. 22	7 46 40.1	+2.488	1	116 2 1.1	9.07	1	pB; vL; R; er; *8 M.....	3
1590	470	III. 836	7 46 42.0	4.905	2	32 57 5.9	9.13	2	F; vS; R; *9 sf; f of 2	3
1591	471	III. 830	7 47 11.2	4.660	1	36 46 9.8	9.16	1	F; pS; E?; bM vS*? L* nf	2
1592	471, a	R. nova	7 47	2	36 46	Makes D neb with h. 871	0
1593	3103	Δ. 535	7 47 18.9	2.133	2	128 11 8.3	9.11	2	!; Cl; B; Ri; L; IC; st 12	3
1594	M. 47	7 48 20.5	2.751	W.	105 3 19.3	9.21	W.	Place from Wollaston's Cat.	0*
1595	VII. 58	7 48 39.8	2.700	1	107 21 4.6	9.22	1	Cl; pL; pRi; pC; st S.....	1
1596	473	II. 302	7 48 50.3	3.602	1	65 52 5.8	9.26	1	F; S; IE; bM; er	3
1597	473, a	R. nova	7 48 ±	65 52 ±	vF; E; * inv near N	0
1598	{ 474 = 3106 }	VII. 10	7 49 0.6	2.544	2	113 56 10.5	9.25	2	Cl; L; cRi; vIC	5
1599	3105	7 49 1.8	2.453	1	117 29 59.5	9.25	1	Cl; L; IC.....	1
1600	475	III. 837	7 50 26.3	4.881	1	33 3 54.6	9.42	1	vF; vS; R; glbM	2
1601	{ 479 = 3107 }	VII. 23	Δ. 626	7 50 36.9	2.396	3	119 41 58.1	9.37	3	Cl; pL; cRi; pC; st 11...13. ..	4
1602	477	7 50 53.9	3.682	1	62 35 59.9	9.43	1	vF; S; R; bM.....	1
1603	477, a	R. nova	7 50	62 35	F; S.....	0
1604	476	III. 750	7 50 55.0	4.067	(1)	49 47 50.2	9.44	(1)	cB; S; R; sbM	2
1605	476, a	R. nova	7 50 +	49 47 ±	Follows III. 750 (h. 476)	0
1606	III. 838	7 51 11.3	4.903	1	32 42 17.4	9.48	1	eF; vS.....	1
1607	478	III. 709	7 51 25.4	+4.530	1	38 52 4.7	+9.49	1	F; L; R; vgbM; r; am st	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1608	h. 3108	H.	h m s 7 52 3.2	s +2.778	2	103° 59' 2.7"	+ 9.49	2	cF; S; vIE 90°; glbM; am st.	2
1609	3109	7 52 12.0	1.565	1	141 55 10.1	9.47	1	pF; S; R; vgpmbM	1
1610	III. 839	7 53 15.3	4.672	1	36 10 28.9	9.63	1	eF; vS	1
1611	480	VI. 37	7 53 15.8	2.858	1	100 14 21.4	9.58	1	Cl; pL; vRi; C; st 11...20...	2*
1612	481	II. 544	7 53 39.5	3.409	1	73 54 23.9	9.63	1	pB; pL; iR; vgbM; er; *225° 5, 60".	3
1613	VIII. 1	7 54 31.8	2.675	3	108 41 12.4	9.68	3	Cl; B; pRi; iC; stS	4
1614	482	III. 605	7 54 45.3	3.586	1	66 13 22.6	9.72	1	vF; S; iR	2
1615	483, a	R. nova	7 54 32.6	3.273	...	80 7 43.3	9.71	...	γ in Lord Rosse's diagram ..	0*
1616	D'Arrest, 51	7 54 40.8	3.27	[2]	80 12 48	9.70	[2]	eF; ILL 512 f10s.5; n50''	0
1617	483	III. 512	7 54 47.2	3.273	2	80 11 58.3	9.71	2	F; S; R; psmbM; r	3
1618	484	III. 7	7 54 48.7	3.255	1	81 4 16.3	9.71	1	F; vS; vIE; 2 st p	3
5066	7 55 12.5	65 25 19.4	See No. 5066.	
1619	3111	7 56 1.4	1.004	6	150 29 10.5	9.75	6	Cl; vB; vL; pRi; st 7...13...	6
1620	3110	7 56 17.4	2.825	1	101 54 45.3	9.81	1	F; vS; R; bet 3 st 13, 14	1
1621	3112	7 56 48.9	2.457	1	117 47 40.5	9.85	1	Cl; B; pRi; pC	1
1622	485	7 57 22.1	4.938	1	31 49 50.5	9.95	1	pF; pL; R; psbM; *9, np 3'	1
1623	486	III. 877	7 59 0.5	2.845	1	101 2 6.9	10.03	1	cF; pL; R; vglbM; am st...	2
1624	488	VIII. 30	7 59 28.2	2.461	(2)	117 46 23.5	10.05	1	Cl; vL; pRi; iC; st 10...15...	3
1625	487	III. 752	7 59 54.2	3.451	1	71 46 29.0	10.10	1	eF; iE; vS*n	2
1626	489	II. 726	8 1 15.2	3.857	2	55 38 18.6	10.22	2	pB; pL; R; vglbM; r; 2 st nf	4
1627	3113	8 1 22.2	2.419	2	119 29 53.7	10.19	2	Cl; pL; Ri; C; st 9, 13...14.	2
1628	490	III. 840	8 1 50.7	4.775	1	33 55 30.7	10.29	1	pF; pL; R; psbM; *8, 164° 3'	2
1629	491	IV. 55	8 3 18.1	4.271	1	43 35 28.4	10.38	1	⊕; pB; pL; R; rrr st 20...	3
1630	3114	VII. 11	8 4 7.8	2.819	2	102 24 59.0	10.40	2	Cl; vL; Ri; iC; st 11...13	3
1631	492	III. 710	8 4 26.2	4.404	1	40 30 20.4	10.48	1	F; L; E; vgbM	2
1632	3115	B.A.C. 3073	8 4 42.2	2.817	1	102 30 54.5	10.45	1	Nebulous * 6.7	1
1633	493	II. 719	8 4 51.0	3.919	1	53 19 26.7	10.49	1	F; pL; iR; vgbM; * nr	3*
1634	494	II. 627	8 6 2.9	3.524	1	68 13 36.1	10.57	1	F; S; iE 45°; *8, np 4'	4
1635	3116	Δ. 563	8 6 26.4	2.215	3	126 58 24.1	10.57	3	Cl; B; L; iC; iE; st 9...12...	3
1636	3117	Δ. 411	8 6 31.6	1.769	2	138 51 3.1	10.57	2	Cl; B; L; iC; st 7...16	2
1637	496	VI. 22	C. H.	8 6 50.1	2.964	2	95 22 30.6	10.62	3	Cl; vL; pRi; pmC; st 9...13.	7
1638	495	8 7 37.5	4.892	1	31 46 48.9	10.73	1	pB; S; mE 0°; psmbM	1
1639	III. 711	8 8 46.0	4.438	1	39 32 33.9	10.73	1	eF; cL; iE 45°±	1
1640	497	II. 303	8 10 33.7	3.568	1	66 6 1.0	10.90	1	F; S; R; mbM; r	4
1641	498	III. 256	8 10 43.9	3.094	1	88 48 46.3	10.91	1	vF; cS; iF; 3Sst inv?	3
1642	499	III. 606	8 11 6.8	3.500	1	69 3 9.5	10.95	1	vF; S; R; sbM; stellar	3
1643	3118	8 11 21.1	2.499	1	117 2 10.9	10.93	1	F; pL; gmbM; am 60 st	1
1644	D'Arrest, 52	8 11 42	3.51	[3]	68 34 36	10.98	[3]	F; pL	0
1645	500	III. 607	8 12 13.6	3.512	1	68 26 14.6	11.02	1	vF; cS; R	2
1646	501	II. 634	8 12 26.5	3.510	1	68 30 37.2	11.04	1	cF; S; R; bM	2
1647	3119	8 12 27.9	2.633	1	111 22 33.6	11.02	1	vF; S; R; gbM; am 60± st.	1
1648	III. 288	8 12 48.8	2.549	1	115 1 53.2	11.04	1	vF; cL; er	1
1649	{ 503 = 3120 }	VII. 64	8 12 58.1	2.421	3	120 12 19.8	11.06	3	{ Cl; pL; pRi; iC; iR; } st 11...14.	4
1650	D'Arrest, 53	8 13 10	3.50	[2]	68 41 48	10.98	[2]	vF; cE; 3vSst f	0
1651	502	VI. 39	8 13 15.1	+2.444	1	119 18 33.1	11.07	1	Cl; vL; cRi; iC; st 9, ...	3
1652	3176	8 13 25.4	-140.624	1	179 41 7.5	7.74	1	F; S; R; glbM; Polaris.	1*
1653	II. 259	8 14 16.8	+3.543	1	66 57 59.4	11.18	1	F; S; iF; r	1
1654	3121	III. 902	8 14 49.0	2.818	2	102 52 35.0	11.20	2	F; vIE; gbM; r; am 50 st	3
1655	3122	8 15 35.1	2.272	4	125 46 36.2	11.24	4	* = h. 4083 in pS neb; am 70 st.	4
1656	3123	8 15 44.3	2.435	1	119 52 8.5	11.25	1	Cl; cL; pRi; pC; R; st 12...	1
1657	504	III. 753	8 17 3.0	3.488	2	69 13 21.1	11.37	3	vF; pS; R; glbM; * p 75''	4
1658	3124	8 17 39.0	2.461	1	119 2 32.8	11.46	1	Cl; pmCM; iF; st 9, 10...13.	1
1659	3125	8 17 39.2	2.369	1	122 31 10.1	11.47	1	Cl; F; S; R; gbM; st 15	1
1660	505	II. 315	8 18 41.2	3.615	1	63 35 7.7	11.49	1	pF; S; R; vsbM*	2
1661	506	III. 599	8 19 34.0	3.511	1	68 3 46.8	11.56	1	vF; pL; iF; r; * sp 2'	2
1662	507	III. 234	8 23 58.8	+3.529	1	66 58 8.1	+11.87	1	vF; S; stellar	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1663	h. 3126	H.	h m s 8 24 27.7	s +0.438	1	157° 39' 23.2"	+11.84	1	F; pS; R; gbM	1
1664	508	8 24 32.8	4.510	2	36 41 28.9	11.93	2	eF; S; R; *95°; p of 2	2
1665	509	III. 292	8 24 48.6	3.691	1	59 59 13.9	11.93	2	vF; pL; R; lbM; r; *nr ...	4
1666	510	8 25 8.2	4.500	2	36 44 52.5	12.05	2	cF; S; R; f of 2; *310°	2
1667	510, a	R. nova	8 25 8.2	4.500	::	36 40 52.5	12.05	::	Place from 510 h. by MS. ...	0*
1668	510, b	R. nova	8 25 ±	36 44 ±	No description or place	0
1669	511	8 25 27.2	3.628	1	62 32 51.4	11.98	1	eF	1
1670	512	II. 318	8 26 43.8	3.661	2	61 3 35.1	12.07	2	F; vLE; mbM; r	3
1671	3130	8 26 48.7	1.170	2	150 38 29.3	12.01	2	Cl; pS; lRi; lC	2
1672	{ 513 = 3127 }	IV. 35	8 26 57.1	2.771	2	105 39 51.8	12.06	2	F; S; att to *13; *7 nf, 10 ^s ...	3
1673	3128	8 27 10.4	2.830	1	102 41 44.4	12.08	1	B; S; E; psbM; bet 2 st.....	1
1674	3129	II. 266	8 27 14.9	2.628	3	112 29 47.4	12.08	3	cB; L; vmE 110° 3	5
1675	515	III. 257	8 28 48.7	3.093	1	88 48 50.0	12.20	1	eF; pL; iF	2
1676	514	II. 319	8 29 0.5	3.662	3	60 49 18.6	12.22	3	F; pS; R; bM; r.....	4
1677	3131	8 30 27.5	2.178	3	130 11 3.7	12.29	3	*9 inv in pB, pL, R, neb ...	3†
1678	{ 516 = 3132 }	VII. 63	8 31 28.3	2.476	2	119 28 1.1	12.37	2	Cl; cL; pRi; pC; st 11...13..	4
1679	III. 982	H. O. N.	8 31 32.3	6.536	1	16 45 45.1	12.47	1	vF; S; stellar	1
1680	III. 235	8 31 51.1	3.539	1	65 57 15.6	12.42	1	eF; S	1
1681	517	M. 44	8 32 9.0	3.462	1	69 32 36.2	12.44	1	Præsepe Cancri	3
1682	III. 983	H. O. N.	8 32 33.3	6.501	1	16 51 47.2	12.54	1	vF; S; stellar	1
1683	3133	8 32 58.2	2.357	1	124 16 9.1	12.47	1	Cl; pmC; irr Δ; st 13	1
1684	518	I. 204	8 33 37.9	4.342	4	39 17 45.8	12.56	4	cB; S; E 130° ±; psmbM*?..	5
1685	3134	8 33 41.5	1.596	2	144 37 47.0	12.50	2	pB; S; R; 3 or 4 vS st p nr ...	2
1686	519	8 34 20.5	3.005	1	93 37 53.4	12.50	1	vF; pL; gbM; r; 2 pB st s, sf	1
1687	3136	8 34 20.5	2.000	1	135 43 55.5	12.55	1	Cl; S; st L & S	1
1688	{ 521 = 3135 }	III. 49	8 34 51.4	3.345	3	75 13 2.6	12.62	3	F; S; vLE 135° ±; psbM.....	5
1689	522	II. 727	8 35 17.8	3.802	2	54 47 16.5	12.65	3	F; L; R; r	4
1690	II. 908	8 35 58.9	6.024	1	19 11 53.8	12.76	1	pB; pL; iF; er	1
1691	520	I. 288	8 37 4.6	8.188	1::	11 15 26.7	12.89	1	vB; cL; lE 90° ±; g, svmbM.	2
1692	523	8 37 28.1	4.503	1	35 36 46.6	12.82	1	eF; psbM	1
1693	4017	Δ. 609	8 37 45.6	2.423	1	122 9 4.7	12.79	1	Cl; pS; lRi; lC; viF; st 12,13	1
1694	3137	8 37 50.2	2.060	2	134 27 15.7	12.79	2	Cl; L; Ri; pmE; st 11...14...	2
1695	3138	8 38 1.3	1.977	2	136 42 16.0	12.80	2	Cl; pS; mC; iR; gbM; st 13...15.	2
1696	III. 50	8 38 43.9	3.309	1	76 54 43.4	12.88	1	eF; cL; R; lbM	1*
1697	3139	8 38 59.6	2.800	1	104 47 19.7	12.89	1	vF; vS; R; bM; *15m nr ...	1
1698	524	8 39 29.7	3.309	1	76 53 9.2	12.94	1	Cl; st 9...10	1
1699	D'Arrest, 54	8 40 20	3.43	[1]	70 27 48	12.99	[1]	eF	0
1700	525	8 40 20.3	4.183	1	42 25 39.3	13.01	1	Cl; lC	1
1701	3140	8 40 50.8	1.694	1::	143 27 29.7	12.99	1::	Cl; L; P; lC; st 10...13	1
1702	3142	8 41 3.4	1.928	1	138 16 31.0	13.00	1	Cl; pL; P; lC; st 13	1
1703	3141	Δ. 489? 490?	8 41 7.1	2.174	1	131 22 30.3	13.01	1	Cl; pRi; lCM; st 12...13 ...	1
1704	526	II. 80	8 41 23.5	3.432	3	70 24 45.8	13.06	3	B; pL; lE 10° or biN; mbM*	5
1705	526 a	R. nova	8 41 23 ±	70 24 ±	Nearly in contact with h. 526 (see description of h. 526).	?
1706	D'Arrest, 55	8 41 52	4.46	[1]	35 58 48	13.11	[1]	vF; R; *15 p 12 ^s , 270°	0
1707	527	II. 48	8 42 1.3	3.430	1	70 28 19.0	13.10	1	eeF; pL; lbM; r	2*
1708	528	VIII. 10	8 42 32.9	3.283	3	78 8 45.2	13.14	4	Cl; vIC; P	5
1709	529	III. 294	8 42 59.4	3.685	1	58 36 38.1	13.17	1	pF; vS; R; bM	2
1710	529, a	R. nova	8 43 ±	58 36 ±	Makes e close D neb with h. 529.	0
1711	530	I. 242	8 43 26.1	4.348	2	38 9 35.3	13.21	2	vB; L; vg, svmbM*10	3
1712	531	M. 67	8 43 34.3	3.291	4	77 40 36.0	13.20	5	l; Cl; vB; vL; eRi; lC; st 10...15.	8*
1713	532	I. 200	8 43 58.0	+3.746	3	56 3 38.9	+13.23	3	vB; vL; vmE 40° 9; gmbM..	5

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1714	h. 533	H. III. 712	h m s 8 45 2.1	s +4.247	2	40° 18' 38".6	+13.32	2	F; pL; R; gbM; 4 S st nr ...	3
1715	} 533, a	R. 3 novæ	8 45 ±	40 18 ±	4 (incl h. 533) nearly in a line	0
1716											
1717											
1718	II. 658	8 45 42.6	3.909	1	49 55 12.8	13.36	1	pF; vS; mbM	1
1719	534	III. 831	8 46 48.6	4.366	1	37 23 56.2	13.44	1	vF; S; R; psbM	2
1720	535	II. 823	8 46 53.6	4.333	1	38 7 1.2	13.44	1	pB; mE 0° ±; psbmM	2*
1721	536	II. 280	8 47 20.4	3.027	1	92 32 2.2	13.44	1	pF; cS; E 90° ±; bet 2 st ...	2†
1722	} 536, a	R. 3 novæ	8 47 ±	92 32 ±	13.48	...	No description	0
1723											
1724											
1725	536, b	R. nova	8 48 0.0	3.025	...	92 28 2.2	MS. No description	0
1726	538	8 48 33.4	3.025	1	92 39 34.6	13.52	1	vF; pS; R; r; *9 p.....	1
1727	D'Arrest, 56	8 48 43	3.025	[2]	92 34 48	13.53	[2]	vF; S; R; *15 p, 44" n; h. 538 nr.	0
1728	537	IV. 66	8 48 46.5	4.438	1	35 41 54.1	13.57	1	pB; fan-shaped; *11 att.....	2†
1729	III. 625	8 48 54.1	3.892	1	50 7 22.8	13.56	1	vF; vS	1
1730	II. 281	8 48 58.2	3.022	2	92 50 24.5	13.55	1	vF; pS, R	2
1731	III. 841	8 49 15.0	4.536	1	33 46 55.0	13.60	1	vF; S	1
1732	540	8 50 3.8	4.065	1	44 33 26.5	13.65	1	pB; L; E; vgbM*18	1
1733	3143	8 50 5.7	1.451	1	148 41 21.4	13.58	1	eF; S; R; pslbM.....	1
1734	3144	8 50 52.5	2.632	2	114 8 0.1	13.67	2	pF; S; R; vgpmbM	2
1735	542	8 51 19.6	3.189	??	83 7 51.0	13.70	1	F; pL; R.....	1*
1736	II. 557	8 51 19.6	3.189	1	83 9 32.0	13.70	1	F; pL; mE	1*
1737	541	III. 540	8 51 25.7	3.785	1	53 44 3.6	13.72	1	vF; S; E 110° ±; 2 vF st inv.	4
1738	539	8 51 28.6	8.388	1:	10 14 54.9	13.83	1:	pB; S; E 45° ±; *nf	1
1739	543	II. 529	8 51 59.2	2.997	2	94 21 30.2	13.74	2	cF; pL; R; vgbM	3
1740	III. 264	8 52 29.4	3.017	2	93 11 6.1	13.77	2	vF; vS; stellar	2
1741	544	8 53 6.7	3.781	1	53 42 41.9	13.83	1	eF; S; stellar	1
1742	545	II. 834	8 53 51.4	4.769	1	29 30 44.0	13.90	1	cF; pS; iR; er.....	2*
1743	546	8 54 2.0	3.023	1	92 50 15.1	13.87	1	vF; L; R; bM	1*
1744	547	8 55 0.5	3.018	1:	93 10 51.9	13.93	1	eF; R	1
1745	3145	8 55 30.8	2.097	1	135 20 55.2	13.94	1	!; eeF; vL; vvmE 19°.....	1†
1746	D'Arrest, 57	8 55 58	3.47	[2]	67 28 48	14.00	[2]	D neb; pB; S, not R; comes s 4'.	0
1747	D'Arrest, 58	8 55 59	3.46	[2]	67 32 48	14.00	[2]	vF; vS	0
1748	549	8 56 5.6	4.305	1	37 41 23.9	14.03	1	4S st in neb	1
1749	549, a	R. nova	8 56	37 41	Makes D neb with h. 549 ...	0
1750	550	I. 249	8 56 26.3	4.788	1	28 58 21.8	14.06	1	cB; cL; E 90° ±; er	3
1751	III. 608	8 56 34.0	3.527	1	64 26 53.5	14.05	1	eF; S; R; vlbM	1
1752	551	III. 60	8 56 43.6	3.399	2	70 59 30.5	14.05	2	vF; S; R; r; *nr	3
1753	548	8 57 9.1	7.170	1:	12 56 4.8	14.16	1:	eF; pL; E; vglbM	1
1754	552	III. 825	8 57 15.0	3.760	4	54 3 51.7	14.09	4	eF; S; R; vglbM; *12. 345°, 50''.	5
1755	D'Arrest, 59	8 57 28	3.39	[4]	71 7 5	14.09	[4]	pF; S; R; bMN=*15	0
1756	} 552	III. 291	8 57 34.4	3.514	3	64 0 27.3	14.11	3	vF; cL; R; bMN; 2 est p ...	3*
					D'Arrest, 60	8 57 35	3.53	[2]	64 0 24	14.10	[2]
1757	D'Arrest, 61	8 57 55	3.53	[1]	64 6 12	14.12	[1]	vF; vS	0
1758	III. 626	8 58 51.7	3.928	2	47 44 29.7	14.19	2	vF; S; iF; lbM; r	2
1759	553	II. 828	8 58 52.0	4.387	1	35 35 23.3	14.21	1	pB; pS; E; vgbM	2
1760	554	III. 647	8 59 32.7	3.811	1	51 48 43.9	14.23	1	vF; cS; R	3
1761	560	III. 275	9 0 13.6	2.820	1	104 56 21.5	14.25	1	vF; pS; bM; S* 30'' n	2
1762	557	III. 236	9 0 16.2	3.450	1	67 59 30.1	14.27	1	cF; vS; R; er; bet 2 pB st ...	3
1763	558	II. 520	9 0 16.9	3.136	1	86 2 37.8	14.26	1	vF; pL; E; gbM; er	3
1764	556	9 0 29.6	4.225	1	39 2 40.0	14.30	1	eF; sbM*15; 1st of 3	1
1765	555	I. 250	9 0 41.1	4.728	1	29 23 40.6	14.32	1	cB; cL; 1E; psbmMLBN ...	2
1766	559	9 0 51.3	4.224	2	39 0 48.6	14.32	2	pF; S; E; pslbM; 2nd of 3...	2
1767	559, a	R. nova	9 0 ±	39 ±	One of 4 (h. 556, 559, 561); one vF; one E.	0
1768	562	II. 490	9 0 56.7	+3.694	3	56 18 26.3	+14.31	3	F; L; mE 150°; r; 2 st n.....	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1769	h. 561	H.	h m s	s	2	39° 4' 21"	+14.27	2	vF; vS; lE; 3rd of 3	2
1770	3146	9 1 0.0	+4.226	1	113 5 19.6	14.32	1	eF; lE; lbM?	1
1771	564	I. 2	9 1 27.6	2.671	2	82 23 38.9	14.43	2	cB; cL; R; vg, vsmbM; r?	7
1772	563	9 2 52.2	3.195	1	44 28 15.5	14.45	1	pB; L; R; vgbM; r	1
1773	565	III. 61	9 2 55.9	4.014	1	71 44 55.2	14.44	1	vF; S; R; am 5S st; (PD 70°)	2*
1774	566	II. 564	9 3 1.4	3.377	4	54 24 3.7	14.49	4	pB; S; R; psmbM	6
1775	566, a	R. nova	9 3	54 24	eF; companion of h. 566, 567.	0
1776	567	III. 826	9 4 3.5	3.728	1	54 29 46.3	14.51	1	vF; S; R; S* 7.5 p	2
1777	569	I. 66	9 4 50.4	2.837	2	104 14 58.2	14.54	2	B; S; pmE 90° ±; psmbM...	3
1777	=										
1778	3147	I. 167	9 5 16.1	3.858	(1)	49 21 24.4	14.58	(1)	cB; R; mbMBN	2
1779	568	III. 295	9 5 21.2	3.613	1	59 23 25.4	14.58	1	vF; vS; R; 2pB st sp	1
1780	571	I. 59	9 6 5.8	2.670	3	113 36 36.3	14.61	3	B; L; mE63° 7; gmbM	5
1780	=										
1781	3148	I. 216	9 6 25.9	5.535	1	20 12 11.7	14.69	1	B; pL; lE90° ±; mbM; r; vS*sf inv.	4
1782	570	9 6 36.0	0.866	1::	157 21 50.0	14.60	1	vF; vS; mE105°	1
1783	3150	9 7 10.6	2.253	3	131 51 37.1	14.67	3	!; O; pB=*9; vS; R; am st.	4
1784	572	9 8 11.6	3.705	2	54 59 4.8	14.76	2	vF; S; R; * pl ^s , n5'	2
1785	573	III. 296	9 8 17.0	3.624	1	58 31 50.8	14.76	1	eF; S; R; lbM	2
1786	575	III. 62	9 8 21.5	3.392	(1)	70 27 55.8	14.76	1	vF; S; R; r	2
1787	575	III. 63	9 8 21.8	3.392	(1):	70 28 25.8	14.76	1	vF; S; R; r	2
1788	II. 708	9 8 31.6	3.899	1	47 27 36.4	14.78	1	pB, S; stellar	1*
1789	574	III. 832	9 8 39.8	4.271	1	36 54 12.0	14.80	1	vF; S; lE; *att; *inv	2
1790	III. 878	9 8 45.5	4.971	2	25 18 38.6	14.82	2	vF; L; R; mbM	2
1791	577	9 8 54.8	3.413	1	69 13 19.7	14.79	1	vF; S; R; np of 2	1*
1792	D'Arrest, 62	9 9 5	3.34	[3]	69 22 9	14.80	[3]	vF; vS; h. 578 f7.5; Δ.P.D. 118".	0*
1793	3152	Δ. 265	9 9 9.9	1.185	4	154 17 18.8	14.76	4	!; ⊕; vL; eRi; vgeCM; 45°d; st 13...15.	4
1794	578	9 9 12.3	3.410	1:	69 21 43.3	14.81	1	vF; S; R; sf of 2	1
1795	III. 749	9 9 35.0	5.953	1	17 34 40.4	14.88	1	F; cS; bM	2
1796	580	II. 505	9 9 37.1	2.817	2	105 43 14.6	14.82	2	pB; pS; E 45° ±; psmbM ...	3
1796	=										
1797	3151	II. 868	9 9 39.0	4.975	(1)	25 9 42.1	14.87	(1)	F; S; iF; 1st of 2	1
1798	576	II. 869	9 9 41.2	4.974	1	25 10 11.1	14.87	1	F; S; E; 2nd of 2	2
1799	3153	III. 242	9 10 3.6	2.688	1	113 2 16.5	14.85	1	F; S; lE; gbM	3
1800	579	9 10 17.0	4.684	1	28 57 59.7	14.89	1	F; pmE	1
1801	3154	Δ. 564	9 10 21.5	2.417	2	126 1 53.8	14.86	2	!; O pB; pL; R; vglbM; in L, C, Cl.	2†
1802	3155	9 10 39.7	2.628	1	116 14 46.4	14.88	1	eF; *11 att	1
1803	3156	9 10 41.4	0.713	1::	159 3 42.5	14.85	1	pF; vS; R; glbM	1
1804	581, a	R. nova	9 10 43.8	3.680	::	55 49 34.9	14.88	::	R.MS. No description	0*
1805	581, b	R. nova	9 10 43.8	3.680	::	55 19 34.9	14.88	::	R.MS. No description	0*
1806	581	9 10 50.9	3.680	1	55 40 39.3	14.91	2	vF; E; l. 113 f	2*
1807	D'Arrest, 63	9 10 52	3.68	[2]	55 47 48	14.91	[2]	vF; vS; R; h. 581, 6' n	0*
1808	581, c	R. nova	9 10 53.4	3.680	::	55 29 4.9	14.92	::	R.MS. No description	0*
1809	581, d	R. nova	9 10 58.4	3.680	::	55 47 34.9	14.93	::	R.MS. No description	0*
1810	582, a	R. nova	9 11 3.1	3.680	::	55 28 34.9	14.93	::	R.MS. No description	0*
1811	582	I. 113	9 11 12.9	3.680	2	55 39 34.9	14.93	2	cB; cL; lE; mbf; 3 st s	4*
1812	582, b	R. nova	9 11 14.5	3.680	::	55 40 20.3	14.92	::	β in Lord R.'s diag. } v nr l.	0*
1813	582, c	R. nova	9 11 16.0	3.680	::	55 27 34.9	14.92	::	α in Lord R.'s diag. } 113.	0*
1814	582, d	R. nova	9 11 17.6	3.680	::	55 27 34.9	14.92	::	R.MS. No description	0*
1815	582, e	R. nova	9 11 32.6	3.680	::	55 42 29.3	14.92	::	ε of Lord R.'s diagram	0*
1816	3157	9 11 41.7	+0.760	1	158 45 49.0	+14.90	1	F; pS; R; glbM	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1817	h. 582, f	H.	R. 2 novæ	9 11 ±	55 40 ±	2 of 15 seen.....	0*
1818	585	9 11 45.3	+2.817	1	105 53 11.2	+14.94	1	eF; R; bM; *f8 ^s .5.....	1
1819	583	III. 627	9 11 50.3	3.813	1	50 7 15.1	14.97	1	vF; vS; R	3
1820	582, g	R. nova	9 12 6.0	3.680	::	55 46 35.3	14.98	::	ζ of Lord R.'s diagram	0*
1821	586	III. 827	9 12 13.5	3.714	1	54 2 26.7	14.99	1	cF; S; R; *10 np 2'	3
1822	584	I. 205	9 12 19.6	4.187	1	38 25 33.3	15.01	1	vB; L; vmE 150° 8; vsmbM =*10.	3
1823	3158	9 12 29.8	1.357	1	152 28 51.8	14.96	1	F; vS; bet 2 st.....	1
1824	III. 64	9 12 41.2	3.385	1	70 28 52.3	15.01	1	S* and neb	1
1825	III. 628	9 12 53.2	3.832	1	49 15 51.9	15.03	1	cF; cS	1
1826	3159	9 13 2.9	2.392	1	127 25 37.3	15.01	1	vF; S; R; *12 att sf	1
1827	587, a	R. nova	9 13 30	105 55	np 587 h; close	0
1828	587	III. 488	9 13 30.4	2.819	1	105 55 14.5	15.05	1	vF; cL; E 45° ±; glbM; *11 sf 9 ^s .	2
1829	3160	9 13 52.9	2.330	1	129 57 21.8	15.06	1	eF; cL; R; vglbM rr	1
1830	588	III. 629	9 14 20.7	3.826	1	49 16 29.6	15.12	1	vF; cS; R; *10 p 2'; 1st of 2	2
1831	590	III. 630	9 14 24.2	3.827	1	49 14 39.6	15.12	1	vF; S; vgbM; 2nd of 2	2*
1832	589	III. 714	9 14 30.9	4.108	2	40 11 39.9	15.13	2	cF; cS; vIE; pglbM; 1st of 2	3
1833	589, a	R. nova	9 14 ±	40 11 ±	Seen with h. 589, 591	0
1834	592	I. 132	9 14 41.5	2.895	1	101 18 57.6	15.12	1	pB; pL; R; gmbMN	4
1835	591	III. 713	9 14 44.2	4.108	2	40 9 20.2	15.14	2	cF; cS; IE; bM; 2nd of 2 ...	3
1836	593	I. 137	9 15 45.1	3.684	3	54 53 13.7	15.19	3	vB; pL; R; smbM	4
1837	594	III. 520	9 16 48.2	2.920	1	99 50 3.2	15.24	1	cF; S; E; bet 2 st 12, 16 ...	2
1838	II. 57	9 17 8.3	3.259	1	77 46 1.8	15.26	1	F; vS, p of 2	1
1839	II. 58	9 17 11.2	3.258	1	77 46 16.1	15.27	1	B; S, f of 2	1
1840	3161	9 17 11.3	2.710	2	112 34 33.8	15.26	2	pF; S; R; gbM.....	2
1841	3162	9 17 21.9	2.016	1	140 30 21.5	15.25	1	Cl; IC	1
1842	3163	9 17 28.1	1.694	7	147 42 57.5	15.25	7	!!; O =*8; vS; R; *15, 59°-13, 13'.	8
1843	595	III. 846	9 17 37.9	4.454	1	32 1 44.8	15.26	1	cF; S; E; vglbM	2
1844	597	II. 546	9 18 8.2	3.255	5	77 57 53.6	15.32	6	pF; pS; R; bM; p of 2, 109°	7
1845	597, a	R. nova	9 18	77 58	Forms Δ with 2 E neb	0
1846	598	II. 547	9 18 13.1	3.255	6	77 58 34.9	15.33	6	vF; pL; R; bM; f of 2	7
1847	596	I. 260	9 18 33.5	4.758	2	26 54 27.1	15.37	2	B; cS; R; mbM; am st	3
1848	3164	9 19 30.4	2.502	1	123 29 36.4	15.38	1	vF; S; vglbM; rrr; st 11m...	1
1849	599	9 19 42.4	3.445	1	66 22 59.3	15.41	1	eF; vS; E 90° ±	1
1850	3165	9 20 8.5	2.740	1	111 8 37.6	15.42	1	eeF; pL	1
1851	3168	9 20 11.4	1.370	2	153 12 29.0	15.40	3	F; S; R; pmbM; B*nr	3
1852	3166	9 20 12.5	2.627	2	117 25 24.9	15.43	2	cF; S; R; gmbM	2
1853	600	II. 555	9 20 24.7	2.904	1	101 2 12.2	15.44	1	pF; pS; vIE; vglbM; r	2
1854	3167	9 20 41.0	2.688	1	114 11 43.5	15.45	1	F; S; R; bM	1
1855	602	III. 297	9 21 59.9	3.562	2	59 49 57.2	15.54	2	vF; S; R; vsbM*12	3
1856	603	III. 8	9 22 2.3	3.195	1	81 39 42.2	15.54	1	vF; E; er; 2 or 3 st inv	4
1857	601	9 22 12.0	4.421	1	31 54 24.1	15.57	1	vF; vS; R; vgbM; *7' s ...	1
1858	3169	9 22 42.2	1.839	1	145 30 15.5	15.55	1	F; pL; R; gmbM; am 80 st ..	2
1859	III. 276	9 24 5.4	2.861	1	104 6 31.2	15.64	1	vF; vS; stellar.....	1
1860	604.1	I. 56	9 24 14.6	3.409	2	67 53 46.8	15.66	3	cB; vL; E; gmbM; r; sp of 2	4+
1861	3170	9 24 15.4	2.592	1	119 46 47.5	15.65	1	F; S; IE; psbM	1
1862	604.2	I. 57	9 24 16.3	3.410	2:	67 52 50.1	15.67	2:	vF; cL; R; psbM; r; nf of 2	4+
1863	606	II. 495	9 24 37.6	3.203	2	80 57 11.4	15.68	2	F; pS; IE; gbM	3
1864	607	II. 506	9 25 2.9	2.830	1	106 7 27.0	15.70	1	pF; S; IE 90...180; mbsf ...	2
1865	III. 977	9 25 15.9	7.880	1	9 37 30.0	15.80	1	eF; vS	1
1866	605	9 25 26.7	4.972	1	23 26 15.8	15.76	1	eF; S; psbM	1
1867	3171	9 25 37.0	1.992	3	142 17 26.0	15.70	3	Cl; cL; pRi; pC; st 10...14.	3*
1868	608	II. 40	9 26 15.3	3.228	1	79 13 53.1	15.77	1	F; pL; R; gbM; p of 2	4
1869	609	III. 513	9 26 33.1	+3.227	1	79 16 23.4	15.78	1	vF; S; R; bMN, f of 2	2
1870	3174	9 26 54.2	-0.275	1	166 0 46.2	15.74	1	pF; pL; R; gbM	1
1871	610	II. 260	9 26 57.2	+3.408	2	67 40 32.3	15.81	2	F; S; vIE.....	3
1872	611	III. 298	9 27 24.1	3.589	1	57 40 57.2	15.84	1	vF; cS; R; sbMN	3
1873	3172	9 27 49.1	+2.769	1::	110 13 55.2	+15.84	1::	eF; S; R; p of 2	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1875	h. 3173	H. III. 597	9 28 2.3	+2.768	2	110 17 59.8	+15.86	2	vF; pS; lE; vglbM; f of 2...	3
1876	3175	9 28 32.8	2.840	1	105 46 48.4	15.88	1	pB; S; R.....	1
1877	3177	9 28 57.4	1.993	1	142 49 22.7	15.89	1	Cl; pRi; pC; * taken.....	1
1878	D'Arrest, 64	9 29 32	3.43	[1]	66 10 24	15.94	[1]	eF; vS; lE; vlbM; 1st of 3...	0
1879	D'Arrest, 65	9 29 34	3.43	[1]::	66 12 12	15.94	[1]	eF; S; 2nd of 3	0
1880	D'Arrest, 66	9 29 42	3.43	[1]	66 8 18	15.95	[1]	eF; vS; 3rd of 3	0
1881	3179	9 30 6.6	2.221	1::	136 18 58.8	15.96	1::	Cl; eL; vRi; st L & S	1
1882	3178	II. 556	9 30 13.5	2.768	2	110 30 23.1	15.97	2	pB; pS; vLE; gmbM	5
1883	612	III. 963	9 30 28.4	6.574	1	12 47 47.8	16.06	1	eF; S; iFig; * f 3'	2
1884	614	III. 4	9 30 39.3	3.215	2	79 51 27.0	16.00	3	vF; S; vLE; bM; Δ st nf.....	5
1885	613	9 30 42.6	3.626	2	55 21 52.3	16.01	2	F; pL; vLE ⁰ ; vglbM.....	2
1886	3180	9 31 15.5	2.756	2	111 24 53.9	16.03	2	F; S; R; glbM; 2 or 3 S st nr	2
1887	615	III. 519	9 31 34.7	3.178	(1)	82 24 1.5	16.05	1::	vF; pL; vgbM	2
1888	616	IV. 68	9 32 17.7	4.410	2	30 31 15.3	16.11	2	eF; vS; R; vgvmbMN	3
1889	3182	9 32 38.7	3.291	1::	74 32 3.3	16.11	1::	eeF; susp	1
1890	3181	9 32 44.3	3.292	1	74 26 8.3	16.11	1	vF; S; R; n of 2	1
1891	620	III. 541	9 32 45.9	3.659	1	53 29 0.6	16.12	1	eF; pS; iR; glbM; r	4
1892	617	9 33 7.8	5.746	1	16 22 12.4	16.18	1	eF; *13 nr	1
1893	618	9 33 12.5	5.149	1	20 45 24.1	16.17	1	F; pL; R; vglbM; * n	1
1894	621	9 33 24.7	3.131	1	85 46 2.7	16.19	1	vF; R; gbM	1
1895	619	III. 315	9 33 39.4	5.734	1	16 23 43.3	16.21	1	vF; vS; R; bM	2
1896	622	I. 114	9 34 34.3	3.571	4	57 31 54.6	16.22	3	B; vL; lE; vgbM; p of 2 ...	5
1897	623	III. 751	9 34 46.9	3.660	1	53 6 33.9	16.23	1	eF; vS; R; bM; r	3
1898	626	II. 275	9 34 49.3	3.084	1	89 1 56.6	16.22	2	pF; pL; R; vglbM	4
1899	624	II. 491	9 34 51.5	3.572	4	57 25 34.9	16.23	4	pB; pL; lE; vglbM; f of 2...	5
1900	628	III. 527	9 34 59.1	2.961	1	97 57 26.9	16.23	1	vF; pS; iR; vglbM.....	3
1901	627	9 35 7.4	3.573	1	57 21 1.2	16.24	1	F; nf of 3.....	1
1902	3183	Δ. 397	9 35 16.9	2.143	1	139 41 32.9	16.23	1	Cl; S; lRi; pC; st 13	1
1903	4018	9 35 22.9	2.628	1	119 24 42.2	16.24	1	eF; pS; B*8m f	1
1904	630	I. 61	9 35 27.7	3.029	1	93 4 16.5	16.25	1	B; cS; iR; bM; *9 sp 3 ^o	3
1905	625	I. 285	9 35 35.6	5.048	1	21 26 21.7	16.29	1	B; vL; mE 152 ^o .4; st inv ...	2
1906	I. 282	9 35 47.6	6.111	1	14 15 5.9	16.33	1	eB; pL; iF	1
1907	631	III. 521	9 36 17.2	2.937	1	99 44 57.7	16.29	1	pF; pS; vLE; psbM.....	2
1908	632	III. 528	9 36 18.4	2.948	1	98 58 2.7	16.29	1	vF; pS; lE 0 ^o ±; vglbM.....	3
1909	629	I. 78	9 36 41.2	5.576	1	17 4 29.8	16.36	1	vB; eL; R; psmbM; * inv f..	3
1910	3184	9 36 52.2	2.330	1	133 33 40.3	16.31	1	Cl; P; E; st 10...11	1
1911	3185	III. 289	9 37 7.5	2.791	1	109 49 54.9	16.33	1	F; pS; R; bM; r; stellar ...	4*
1912	633	III. 34	9 37 23.0	3.231	(1)	78 20 10.5	16.35	2::	eF; vS; R; bM (? P.D. 15)..	3
1913	II. 311	9 38 10.3	2.780	3	110 37 58.7	16.39	3	pB; pS; iR; mbM	3
1914	II. 624	9 38 39.5	3.156	1	83 41 19.3	16.41	1	F; pS; lE 90 ^o ±	1
1915	3186	9 38 52.0	2.825	1	107 44 26.6	16.42	1	F; R; gbM; * f	1
1916	634, a	R. nova	9 38	67 20	Makes a D neb with h. 634; which follows it.	0
1917	634	9 38 55.9	3.391	1	67 20 5.9	16.43	1	F; vS; bM; sp of 2	1
1918	635	III. 277	9 38 58.1	2.884	1	103 41 8.9	16.43	1	eF; S; R; bM; stellar; p of 2	2
1919	637	III. 278	9 39 4.6	2.884	1	103 43 23.9	16.43	1	eF; S; R; bM; stellar; f of 2	2
1920	636	9 39 18.1	3.391	1	67 16 32.5	16.45	1	F; S; R; bM; nf of 2.....	1
1921	3189	9 39 21.6	2.008	1	144 8 1.9	16.43	1	Cl; P; lC; st mm.....	1
1922	3187	9 39 28.0	2.778	1	110 56 47.5	16.45	1	vF; S; *20 f 1'	1
1923	3188	V. 50	9 39 32.5	2.619	2	120 32 50.5	16.45	2	l; vF; vL; vg, vsbMN 4'; 19 ^o .5 d.	3
1924	638	II. 717	9 40 2.0	3.823	1	45 15 27.0	16.50	1	pF; pL; iR; bM; r.....	2
1925	638, a	R. nova	9 40 4.8	3.823	...	45 16 27.0	16.48	...	RMS	0
1926	638, b	R. nova	9 40 4.8	3.823	...	45 16 57.0	16.48	...	Suspected; MS	0
1927	3191	Δ. 397	9 40 5.0	2.167	1?	139 47 14.1	16.47	1?	Cl; S; lRi; iF; st 12...15 ...	1
1928	638, c	R. nova	9 40 7.6	2.823	...	45 12 27.0	16.48	...	MS	0
1929	3190	9 40 10.4	2.634	1	119 48 2.4	16.48	1	F; S; R; *12 att 320 ^o	1
1930	638, d	R. nova	9 40 13.3	3.823	...	45 21 27.0	16.49	...	MS	0
1931	639	V. 26	9 40 15.0	3.584	1	55 56 14.0	16.50	1	l; eB; L; vimE 90 ^o	3+
1932	638, e	R. nova	9 40 30.2	3.823	...	45 13 27.0	16.52	...	MS	0
1933	640	9 41 1.2	+3.825	1	45 1 50.5	+16.55	1	pF; R; bM; r; p of 2.....	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1934	h. 638, f	H.	R. nova	h m s 9 41 4.0	s +3.823	...	45° 15' 27.0"	+16.55	...	MS	0
1935	641	9 41 7.2	3.825	1	45 0 16.5	16.55	1	F; psbM; rr; f of 2.....	1
1936	} 641, a	R. novæ	9 41 ±	45 0 ±	Several near	0
1937											
1938	D'Arrest, 67	9 41 32	3.49	[1]	54 38 7	16.56	[1]	vF; pL; R; cometary	0
1939	642	9 41 32.1	3.251	1	76 32 0.8	16.56	1	F; pL; R; glbM	1
1940	642, a	R. nova	9 41	76 32	3 "novæ," with 642 (Vide h. 646, 648).	0
1941	644	9 42 3.2	3.011	1	94 33 38.4	16.58	1	eF; L; p of 2	1
1942	646	III. 51	9 42 31.0	3.250	3	76 31 58.3	16.61	3	vF; pS; IE 0°±; r	4
1943	647	9 42 38.9	3.012	1	94 31 9.3	16.61	1	F; R; vglbM; f of 2	1
1944	645	I. 115	9 42 39.6	3.579	2	55 47 35.6	16.62	2	pB; pS; vlE; mbM; *10 sf 100°.	4
1945	648	III. 52	9 42 54.1	3.249	1	76 36 18.9	16.63	1	eF; pL; E; r	2
1946	3192	9 42 58.1	2.782	1	111 5 19.6	16.62	1	eF; vS; R; *9s	1
1947	643	V. 23	9 43 9.9	5.466	1	17 8 23.4	16.68	1	vF; vL; IE; r	3
1948	3193	9 43 17.7	2.821	1	108 31 38.2	16.64	1	F; S; R; lbM	1
1949	649	M. 81	9 43 48.9	5.066	1	20 16 10.0	16.70	1	l; eB; eL; E 156°0; g, svmbMBrN.	4
1950	{ IV. 79 = 4H. ON }	M. 82	9 43 52.3	5.142	1	19 34 16.3	16.71	1	vB; vL; vmE "a beautiful ray."	2*
1951	650	9 44 0.3	3.497	2	60 7 7.4	16.68	2	F; S; sbM *12; bet 2B st ...	2
1952	3194	B. 2686	9 44 1.8	1.975	2	145 45 42.8	16.66	2	Cl; pL; pRi; iF; st 11...12 ...	2
1953	W. H. nova?	M. 81??	9 44 38.0	5.064	1	20 12 18.9	16.73	1	vB; cL; mE; 5 or 6 st (?) inv	1*
1954	3197	9 45 5.8	1.674	1	152 2 12.3	16.71	1	Cl; cL; IC	1
1955	3195	9 45 6.2	2.705	1	116 22 6.6	16.72	1	F; pS; R; lbM	1
1956	3196	II. 98	9 45 25.6	3.300	1	72 39 56.5	16.75	1	⊕; F; L; R; vglbM; rr; 2B st sp.	4
1957	651	II. 835	9 46 19.3	4.327	2	30 2 40.3	16.81	2	cF; pS; IE; vgbM; *10 n 7'	4
1958	652	III. 254	9 46 26.5	3.097	1	87 55 58.0	16.80	1	vF; vL; vmE 111°5	3
1959	3198	9 46 38.8	2.834	1	107 59 3.0	16.80	1	vF; pS; R; lbM	1*
1960	3199	9 47 1.9	2.704	1	116 40 25.6	16.82	1	pF; R	1*
1961	3201	9 47 41.3	2.707	1	116 37 56.5	16.85	1	pF; S; R; gbM	1*
1962	3202	III. 272	9 47 50.8	2.836	1	107 58 44.8	16.86	1	F; pL; R; glbM	2*
1963	3200	III. 600	9 47 53.6	3.293	1	72 54 4.1	16.87	1	vF; S; vlE; gbM	2
1964	656	VI. 4	9 47 59.1	3.133	2	85 4 19.1	16.87	2	F; pL; vlE; vgbM; rr; *7f90°.	4
1965	3203	9 48 15.8	2.692	2	117 38 37.4	16.88	2	pB; S; R; vgbM; *11 att 203°8.	2
1966	III. 978	9 48 24.1	7.497	1	9 3 41.8	16.96	1	eF; pL; vlbM; 2 S st s	1
1967	3205	9 48 31.2	0.647	1	163 16 15.5	16.85	1	F; L; iR; glbM; S * inv ...	1
1968	3204	III. 601	9 48 39.8	3.297	1	72 30 4.3	16.91	1	vF; cS; vlE; er	2
1969	654	II. 333	9 48 42.7	5.376	1	17 9 25.2	16.94	1	pF; vS; R; bM; *11 nr.....	3
1970	653	II. 903	9 48 44.4	6.102	1	13 10 9.5	16.95	1	vF; pL; r	2
1971	655	II. 334	9 48 57.4	5.366	1	17 12 16.5	16.95	1	vF; vS; vglbM	3
1972	II. 909	H. ON 5	9 49 50.5	5.382	1	17 12 24.7	16.99	1	F; pL; R; 3rd of 3.....	1
1973	657	II. 492	9 50 7.9	3.533	3	56 57 50.4	16.98	3	pB; pL; E 90±; gbM; *9 nf	4
1974	III. 293	9 50 32.8	3.474	1	60 21 51.7	16.99	1	eeF; eS; stellar (?).....	1*
1975	659	II. 59	9 50 39.1	3.209	1	78 58 42.7	16.99	1	pB; pS; R; gmbMN; 3 st nr	2
1976	3206	III. 273	9 50 47.0	2.831	2	108 40 49.7	16.99	2	vF; pS; IE; glbM	3
1977	III. 853	9 51 10.7	4.126	1	33 42 23.9	17.03	1	vF; S; vglbM	1
1978	660	III. 542	9 51 18.7	3.585	1	53 55 47.9	17.03	1	vF; pL; iR; vglbM	4
1979	3207	9 51 19.5	3.261	1	74 53 56.9	17.03	1	vvF; *14 att; *11 f	1
1980	3208	9 51 32.2	2.848	1	107 30 25.9	17.03	1	eF; S; R	1
1981	3209	II. 268	9 52 4.6	2.731	1	116 15 22.5	17.05	1	pB; S; R; mbM	2
1982	658	I. 286	9 52 7.4	4.925	(2)	20 35 17.7	17.09	1	eB; cL; mbM; R with ray...	3
1983	V. 47	9 52 24.3	4.119	1	33 38 27.7	17.09	1	vB; L; mE 135°±	1
1984	III. 934	9 52 34.6	3.241	1	76 19 59.7	17.09	1	vF	1
1985	III. 596	9 52 37.0	2.786	2	112 7 58.4	17.08	2	vF; eS; lbM; ΔS st np	2
1986	3210	9 52 40.5	2.671	1	119 41 50.4	17.08	1	vF; S; R; * att	1
1987	3211	9 52 45.4	+2.722	1	116 28 7.7	+17.09	1	vF; S; R; *13 att sf	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1988	h. 3212	H.	9 52 50.7	+2.832	1	108° 50' 56.7"	+17.09	1	vF; S; R.....	1
1989	3213	9 53 4.5	2.607	1	123 33 46.0	17.10	1	pB; S; R; pmbM; bet 2 st...	1
1990	661	III. 24	9 53 17.7	3.367	1	66 55 52.6	17.12	1	vF; S.....	2
1991	3214	9 53 19.7	2.705	1	117 38 50.3	17.11	1	pF; pS; R; vS st inv.....	1
1992	3215	II. 293	9 53 38.9	2.831	1	108 57 52.9	17.13	1	pB; pS; iR; bM; p of 2.....	2
1993	3216	9 53 53.3	2.655	1	120 52 53.2	17.14	1	F; L; E; vgvbM.....	1
1994	3217	9 54 2.5	2.832	1	108 57 58.5	17.15	1	eF; R; lbM; f of 2.....	1
1995	663	9 54 22.3	3.398	2	64 37 15.1	17.17	2	pB; S; mE 90°±; psbMN...	2
1996	664	III. 478	9 54 25.4	3.525	1	56 37 10.4	17.18	1	eF; S.....	3
1997	3218	9 54 27.6	2.655	1	120 59 37.8	17.16	1	pB; pS; R; gpmbM.....	1
1998	662	III. 916	9 54 36.7	4.298	1	29 13 26.7	17.19	1	vF; vS; R; bM; *11, 142°-2	2
1999	665	IV. 48	9 55 19.4	3.675	1	48 35 27.3	17.21	1	eF; pL; E 45°±; vF * inv...	2
2000	3219	9 55 46.4	2.120	2	144 6 16.3	17.21	2	Cl; C; lE; st 13...16.....	2
2001	666	II. 320	9 55 58.4	3.504	1	58 8 19.2	17.24	1	F; S; R; sbM.....	2
2002	3220	9 56 17.6	2.660	2	121 0 33.2	17.24	2	F; S; R; glbM.....	2
2003	3221	9 56 40.1	2.746	2	115 29 25.8	17.26	2	cF; vL; vmE 82°-3; lbM...	2+
2004	II. 898	9 56 54.3	3.242	1	75 49 0.4	17.28	1	F; L red * n 3'	1
2005	667	9 57 14.8	3.818	3	42 3 28.0	17.30	3	pB; S; R; SmbM *12.....	3
2006	3222	9 58 3.2	2.716	1	117 46 13.6	17.32	1	eF; L; Δ 2 st 8m.....	1
2007	3224	Δ. 297	9 58 11.2	1.934	3	149 26 46.6	17.32	3	Cl; eL; iC; B; st 9...14.....	3
2008	668 = 3223	I. 163	9 58 14.3	2.988	2	97 2 32.9	17.33	2	vB; L; vmE 45°; vg, vsmbMEN.	3
2009	3225	9 59 14.8	2.627	1	123 32 36.4	17.38	1	F; pS; R; gbM.....	1
2010	Auw. N. 26	9 59 18.4	3.247	...	74 55 35.7	17.39	...	F; (Lassell, Mar. 31, 1848)...	0
2011	II. 305	9 59 26.0	3.003	1	95 51 17.9	17.43	1	F; S; lE; er.....	2
5067	9 59 51.1	89 15 7.3	See No. 5067.	...
2012	3226	10 0 2.2	2.848	1	108 33 36.3	17.41	1	F; pL; R; lbM; * s.....	1
2013	3227	10 0 14.8	2.700	2	119 15 12.6	17.42	2	cF; S; R; vgbM.....	2
2014	669	III. 65	10 0 40.4	3.299	(1)	70 53 43.5	17.45	1:	eF; cS; vLE; r.....	2*
2015	670	10 0 45.2	3.195	1	79 20 43.5	17.45	1	eF; S; psbM; 31 Leon sf 100"	1
2016	671	10 0 54.9	3.296	1	71 4 49.8	17.46	1	pB; pS; pmE; gbM.....	1
2017	3228	10 1 8.2	2.523	4	129 45 5.5	17.45	4	l; O; vB; vLE; *9M; 4°-0 d.	4+
2018	3229	10 1 46.2	1.540	1?	156 41 44.1	17.47	1?	B; R; bM.....	1
2019	672	10 2 9.8	3.759	1	43 21 12.6	17.52	1	F; S; R; gbM.....	1*
2020	3231	10 2 9.9	1.545	2	156 41 30.4	17.48	2	pB; pS; R; gbM; * 13 n...	2
2021	3230	10 2 30.4	2.719	1	118 22 22.3	17.51	1	vF; S; lE.....	1
2022	3232	10 3 9.9	2.982	1	97 47 52.5	17.55	1	F; R.....	1
2023	673	III. 518	10 3 17.7	2.936	1	101 44 15.8	17.56	1	F; pL; R; vg, slbM; f of 2...	4
2024	674	I. 79	10 4 47.5	5.285	1	15 54 35.5	17.65	1	vB; L; R; vg, vsmbM.....	2
2025	675	10 4 48.3	3.861	1	38 49 21.9	17.63	1	* 7 m in photosphere 2' or 3' d	1
2026	677	III. 53	10 5 8.4	+3.222	1	76 38 47.9	17.63	1	eF; pL; vLE; r; st inv.....	3
2027	3234	10 5 10.7	-0.511	1	169 43 54.4	17.58	1	F; S; lE; vlbM; * 15 inv...	1
2028	680	III. 255	10 5 25.3	+3.113	1	86 10 27.2	17.64	1	F; cS; R; psbM; Δ B st f...	3
2029	3233	10 5 27.7	2.700	1	120 16 4.2	17.64	1	vF; pS; E; * 8.9 sp.....	1
2030	678	II. 639	10 5 31.3	3.592	1:	50 33 28.5	17.65	1:	cB; cS; R; psbM; r.....	2
2031	678, a	R. 2 novæ	10 5 ±	50 33 ±	3 seen; one(? which)=h.678	0
2032			
2033	676	10 5 42.7	5.424	1?	14 53 50.7	17.69	1:	vF; S; R.....	1
2034	682	II. 43	10 5 43.5	3.345	2	66 34 30.8	17.66	3	pF; cL; R; vglbM; r; S * inv	5
2035	681	II. 640	10 5 48.2	3.589	1:	50 40 11.1	17.67	1:	F; S; R; gbM.....	2
2036	679	10 5 49.2	4.054	1	32 38 47.1	17.67	1	eF; S; R; vglbM.....	1
2037	684, a	R. nova	10 6 15.4	3.116	::	85 56 15.7	17.69	::	vvF; mE 0°±.....	0
2038	684	I. 3	10 6 29.6	3.116	3	85 52 53.7	17.69	4	B; pS; R; psmbM; p of 2...	8
2039	D'Arrest, 68	10 6 36	3.34	[1]	59 42 7	17.69	[1]	F; S; ?? Cl of vS st.....	0
2040	683	10 6 45.3	4.194	1	29 4 36.3	17.71	1	F; psbM; stellar; * 7.8 np 5'	1
2041	685	I. 4	10 6 58.5	3.116	3	85 50 20.3	17.71	4	B; pL; vLE; pgmbM; * 11, 78°-2, 80".	8
2042	686	10 7 32.0	3.745	1	42 42 39.2	17.74	1	F; S; R.....	1
2043	250	10 8 7.9	+87.502	1::	0 6 46.2	+19.47	1::	vF; R; gbM; *11, 2' s; P _o -larissima Borealis.	1*

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
2044	3235	10 8 9.4	+2.754	1	116 59 43.5	+17.75	1	eF; S; R; 2 B st f	1
2045	III. 964	10 8 16.1	5.415	1	14 38 38.0	17.80	1	cF; S; stellar; S * f nr	1
2046	3236	10 8 18.9	2.738	2	118 11 3.8	17.76	2	cB; L; mE 50°5; vglbM ...	2
2047	687	III. 25	10 8 51.5	3.318	3	68 10 42.7	17.79	3	cF; S; R; psBM.....	4
2048	3237	10 9 23.5	2.905	1	105 5 56.0	17.80	1	pB; pL; gpmbM.....	1
2049	688, a	R. nova	10 9 44.5	3.623	...	47 53 13.9	17.83	...	MS; no description.....	0
2050	688, b	R. nova	10 9 47.2	3.623	...	47 54 1.9	17.83	...	MS; no description.....	0
2051	I. 265	10 9 47.5	4.082	1	31 5 44.2	17.84	1	cB; cL; iR; vgbM.....	1
2052	688	I. 168	10 9 49.9	3.623	1	47 53 1.9	17.83	1	pB; vL; R; vgbM	4
2053	689	10 9 53.3	3.627	1	47 41 6.2	17.84	1	pF; vL; R; vgbM; 12°5 d; * 11 n 2'.	1
2054	692, a	D'Arrest, 69	10 9 57	3.32	[2]	67 35 48	17.83	[2]	pF; pL; gmbM (δ in Lord R.'s diagram).	0
2055	692, b	R. nova	10 10 4.7	3.324	...	67 25 19.5	17.85	...	Marked γ in Lord R.'s diagr.	0
2056	690	III. 910	10 10 17.0	4.047	1	31 53 7.8	17.86	1	vF; pL; r	2
2057	692, c	R. nova	10 10	67 28	mE, parallel to h. 692, with which it forms D neb.	0
2058	692	II. 44	10 10 23.3	3.324	3	67 28 10.5	17.85	3	B; pS; E; psbMN; sp of 2	4+
2059	691	10 10 25.8	3.726	1	42 51 47.8	17.86	1	F; S; R; bM	1
2060	III. 704	10 10 27.6	3.728	1	42 43 44.8	17.86	1	eF; vS; (?).....	1
2061	693	II. 45	10 10 41.6	3.324	3	67 24 24.8	17.86	3	B; S; vLE; psbM; r; * 9, 352°0, 75" nf of 2.	4
2062	III. 695	10 10 46.8	+5.391	1	14 29 43.0	17.90	1	vF; vS.....	1
2063	3241	10 10 51.5	-0.506	2	170 10 11.6	17.82	2	!; ○; pB; S; lE; 13°0 d; 3S st nr.	2+
2064	694	III. 348	10 10 53.6	+3.398	1:	61 38 27.1	17.87	1:	eeF; pS; lE.....	2
2065	III. 966	10 10 57.9	6.114	1	11 4 53.6	17.92	1	vF; vS.....	1
2066	695	I. 199	10 11 14.4	3.702	1	43 44 15.7	17.89	1	pB; vL; mE 45°±; vgbM...	3
2067	3239	10 11 46.1	2.128	4	147 15 39.7	17.89	4	!; vB; vL; falcate; * N.....	4+
2068	3238	Δ. 445	10 11 52.8	2.452	2	135 42 6.0	17.90	2	⊕; vS; iR; ICM; gbM; st 13...16.	2
2069	696	II. 720	10 12 5.3	3.643	1::	46 18 42.6	17.92	1:	cF; S; R; vgbM; 1st of 3...	2
2070	3240	10 12 5.5	2.777	2	116 0 17.3	17.91	2	pB; S; cE; gbM.....	2
2071	698	10 12 12.9	3.396	1	61 28 54.6	17.92	1	eF; pL; gbM	1
2072	699	II. 721	10 12 21.7	3.641	1	46 20 50.2	17.94	1	cF; S; R; vgbM; 2nd of 3...	2
2073	697	I. 266	10 12 30.9	4.011	1	32 22 5.2	17.94	1	pB; cL; E; vglbM.....	2
2074	700	II. 722	10 12 33.3	3.641	1	46 18 50.2	17.94	1	cF; S; R; stellar; 3rd of 3...	2
2075	701	10 12 47.0	3.365	1	63 47 46.5	17.95	1	F; S; R; has a *	1
2076	3242	10 13 17.6	1.946	2	151 58 35.5	17.95	2	○; * = 10 m; R; am 150 st	2
2077	III. 979	H. ON	10 13 34.0	6.559	1	9 27 5.6	18.02	1	Stellar; 1st of 3	1
2078	III. 980	H. ON	10 13 34.2	6.565	1	9 26 5.6	18.02	1	vF; S; 2nd of 3	1
2079	III. 981	H. ON	10 13 34.3	6.571	1	9 25 5.9	18.03	1	vF; S; 3rd of 3	1
2080	702	III. 330	10 13 55.0	3.342	1	65 21 53.7	17.99	1	vF; pS; R; bM	2
2081	I. 283	10 14 23.0	5.297	1	14 37 52.9	18.03	1	cB; cL; eR	1
2082	III. 911	10 14 26.4	3.998	1	32 15 55.6	18.02	1	vF; cL; iF	1
2083	D'Arrest, 70	10 14 39	3.31	[2]	67 42 12	18.01	[2]	eF; mE; a ray	0
2084	Auw. N. 27	10 14 55.0	3.288	...	69 24 42.6	18.02	...	F; lbMr (Winnecke, June 1855).	0
2085	3243	10 15 20.5	2.683	1	123 32 59.9	18.03	1	pB; vL; vLE; psbMN	1
2086	3244	10 15 27.7	2.677	1	123 59 28.2	18.04	1	vF; pS; R; vgbM	1
2087	703	II. 882	10 15 43.0	4.028	1	31 9 19.1	18.07	1	cF; pL; lE; vgbM	2
2088	II. 28	10 15 55.1	3.289	1	69 23 27.8	18.06	1	vF; cL; R } D neb; 45°, 2' {	1*
2089	II. 29	10 15 58.5	3.289	1	69 22 28.1	18.07	1	vF; cL; R }	1*
2090	3245	Δ. 386	10 16 10.6	2.352	1::	141 1 1.8	18.06	1::	Cl; 9 L & a few S st	1
5068	10 16 14.1	89 13 46.0	See No. 5068.	
2091	705	10 16 15.7	3.207	3	76 44 21.4	18.08	3	!; * or ** in neb	3
2092	704	10 16 25.6	4.456	1	22 29 6.0	18.10	1	Cl; cL; P; lC; st 10...12 ...	1
2093	D'Arrest, 71	10 16 33	3.39	[1]	61 16 42	18.09	[1]	eeF; * 11 p, ls, 150" p of 2...	0
2094	706	10 17 5.1	3.374	1	62 16 5.0	18.10	1	pB; pS; R; psbM	1*
2095	D'Arrest, 72	10 17 7	3.38	[2]	61 16 12	18.11	[2]	F; S; f of 2	0
2096	707	10 17 16.6	+4.142	1	28 1 16.9	+18.13	1	eF; vS; psbM; 2 st 11, 12, f	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
2097	h. 709	H. III. 631	h m s 10 17 27.7	s +3.556	1:	49° 39' 49.9"	+18.13	1:	vF; vS; R; pgbM	3
2098	708	III. 883	10 17 28.5	3.980	1	32° 4' 25.9"	18.13	1	F; S; R; psbM	2
2099	710 = 3246	IV. 10	10 17 31.0	3.255	3	72° 8' 18.9"	18.13	3	vF; *9 inv nr M	5
2100	3247	10 17 53.2	2.851	2	111° 4' 56.9"	18.13	2	eF; S; R; * nr	2
2101	3249	10 17 56.8	2.717	1	121° 45' 28.9"	18.13	1	F; pmE; glbM; * 11 np ...	1
2102	3248	IV. 27	Lal. 20204	10 18 2.2	2.886	4	107° 55' 50.2"	18.14	4	!; O; vB; lE, 135°; 32'' d _± ; <i>blue.</i>	7+
2103	4019	10 19 20.9	2.612	1	129° 6' 23.4"	18.18	1	vF; * 11 m 90'' n	1
2104	711	I. 86	10 19 24.9	3.385	4	60° 47' 1.7"	18.19	5	vB; pL; E; smbMEN	6
2105	712	10 19 26.0	3.116	1	85° 26' 24.7"	18.19	2	eF; S; R; 2 st Δ; * 6, 300°, 8'	3
2106	3250	10 19 37.9	2.196	1::	147° 10' 35.7"	18.19	1	st inv in neb	3
2107	713	II. 347	10 20 2.0	3.314	1	66° 26' 28.6"	18.22	3	pB; S; R; psbM	4
2108	3251	10 20 6.9	2.690	1	124° 14' 38.3"	18.21	1	eF; pL; R; vglbM	1
2109	3252	10 20 25.6	2.625	4	129° 13' 45.9"	18.23	4	pB; pL; R; vg, psbM; *13, 45°.	5
2110	D'Arrest, 73	10 20 29	3.35	[1]	63° 11' 42"	18.23	[1]	vF; pL; 3 B st sp	0
2111	III. 316	10 20 51.1	5.054	1	15° 27' 11.9"	18.27	1	eF; pS; mE; r	1*
2112	714	I. 72	10 21 27.2	3.392	2	59° 47' 27.1"	18.27	3	cB; L; E 45° _± ; psmbMN...	5
2113	3253	10 21 32.2	2.111	1	149° 57' 49.8"	18.26	1	Cl; pS; vC; st 15	1
2114	3254	10 21 53.0	2.553	1	133° 11' 7.1"	18.27	1	cB; S; R; gmbM	2
2115	3255	10 22 29.8	2.689	3	124° 56' 49.0"	18.30	2	vF; vS; R; psbM; 1st of 4...	3
2116	3256	10 22 36.9	2.689	4	124° 53' 9.0"	18.30	4	cF; S; R; psbM; 2nd of 4...	4
2117	715	II. 870	10 22 42.9	4.272	1	24° 14' 50.9"	18.33	1	F; S; R; gbM	2
2118	3257	10 22 52.7	2.689	4	124° 52' 39.0"	18.32	4	vvF; vS; R; psbM; 3rd of 4	4
2119	3258	10 23 3.0	2.546	1	133° 56' 12.6"	18.32	1	F; S; R; am st	1
2120	3260	10 23 8.2	2.554	1	133° 29' 7.6"	18.32	1	eF; S; R	1
2121	3259	10 23 11.4	2.692	2	124° 53' 48.9"	18.33	2	vF; vS; R; psbM; 4th of 4	2
2122	3261	10 23 13.8	2.556	1	133° 24' 32.9"	18.33	1	F; S; mE 280° _± ; psbM ...	1
2123	718	III. 349	10 23 15.9	3.377	1	60° 29' 34.9"	18.33	1	pF; S; R; psbM; * sf nr ...	5
2124	716	10 23 18.3	3.895	1	33° 11' 37.5"	18.35	1	eF; bet 2 S st	1
2125	717	II. 871	10 23 29.5	4.246	1	24° 32' 16.8"	18.36	1	cF; vS; R; psmbM *	2
2126	3262	10 23 33.9	2.696	1::	124° 38' 10.2"	18.34	1::	eF; vS; R; 1st of 4	1
2127	3263	10 23 37.9	2.697	2?	124° 39' 10.2"	18.34	2?	F; S; R; 2nd of 4	2
2128	3264	10 23 41.1	2.699	1	124° 30' 10.5"	18.35	2	F; S; R; bM; 3rd of 4	3
2129	719	III. 331	10 23 44.1	3.329	1	64° 24' 28.5"	18.35	1	cF; vS; E; glbM	2
2130	3265	10 23 45.9	2.697	1	124° 39' 3.5"	18.35	1	pF; S; E; pmbM; 4th of 4...	2
2131	720	II. 358	10 24 27.7	3.360	4	61° 36' 59.4"	18.38	4	F; pL; R; glbM; * f.	5
2132	3266	10 24 34.0	2.681	2	126° 1' 22.1"	18.37	2	F; L; vIE; psbM	3
2133	3267	10 24 57.7	2.635	1	129° 13' 51.7"	18.39	1	F; S; * 8 p	1
2134	721	II. 359	10 25 4.1	3.369	3	60° 46' 20.7"	18.39	3	cB; cS; R; pgmbM	4
2135	3268	10 25 26.1	2.637	3	129° 13' 52.0"	18.40	3	F; S; R; * nf	2
2136	3269	10 25 34.3	2.711	1	124° 8' 4.3"	18.41	1	eF; pL; E; glbM	1
2137	3271	10 26 51.1	2.536	1	135° 22' 3.5"	18.45	1	pF; S; R; gbM	1
2138	III. 912	10 26 58.6	3.944	1	30° 44' 23.1"	18.47	1	eF; vS	1
2139	3270	10 27 1.3	2.808	4	116° 44' 11.8"	18.46	4	pB; S; lE; gbM; 1st of 9 ...	4
2140	722	III. 917	10 27 9.3	3.945	1	30° 40' 11.4"	18.45	1	vF; pS; R; psbM	2
2141	D'Arrest, 74	10 27 11	3.28	[2]	67° 37' 18"	18.47	[2]	F; pL; * p 24 ^s , 225 ^s	0
2142	723	III. 918	10 27 12.7	3.942	1	30° 43' 31.4"	18.48	1	eF; cS; R; vglbM	2
2143	3272	10 27 47.4	2.713	1	124° 34' 31.4"	18.48	1	eF; vS; R	1
2144	3276	10 28 7.7	2.258	1::	147° 28' 14.7"	18.49	1::	Cl; B; Ri; pL	1*
2145	724	I. 164	10 28 12.8	3.474	1	51° 57' 17.3"	18.51	1	cB; L; mE 135° _± ; glbM ...	4
2146	725	III. 767	10 28 44.6	3.698	1	39° 9' 51.9"	18.53	1	vF; pS; lE	2
2147	726	III. 54	10 28 57.8	3.193	1	76° 34' 38.9"	18.53	1	eF; cL; R; vgbM; r	2
2148	727 = 3273	III. 55	10 29 10.6	3.206	3	75° 6' 29.2"	18.54	3	cF; cS; R; pmbM; r; am B st	5
2149	II. 46	10 29 13.4	3.285	1	67° 15' 27.2"	18.54	1	pF; S; r; Δ pB st n	1
2150	728	II. 46??	10 29 18.1	3.283	2	67° 23' 45.2"	18.54	2	cB; S; lE; psbM; r	3
2151	3274	10 29 20.3	+2.757	1	121° 37' 4.2"	+ 18.54	1	eF; S; R	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
2152	h. 3275	H.	10 29 22.6	+2.756	1	121 38 41.2	+18.54	1	vF; S; R.....	1
2153	730	III. 66	10 29 27.1	3.245	2	71 8 48.2	18.54	2	vF; vS; vIE; glbM; r.....	3
2154	729	III. 615	10 29 33.9	3.470	2	51 49 25.5	18.55	2	vF; cS; psbM; er.....	4
2155	3277	10 29 38.5	2.818	1	116 26 17.5	18.55	1	vF; S; R; 2nd of 9.....	1
2156	3278	10 29 41.3	2.813	1	116 52 38.5	18.55	1	eeF; 3rd of 9.....	1
2157	3279	10 29 48.4	2.816	4	116 42 50.5	18.55	4	F; S; R; 4th of 9.....	4
2158	731	IV. 60	10 29 59.7	3.770	1	35 45 49.1	18.55	1	○? cB; pL; R; vg, vsmbMN 15".	3+
2159	3280	10 30 1.1	2.815	2	116 48 5.8	18.56	2	B; L; R; p of D neb; 5th of 9	3
2160	3281	10 30 9.5	2.815	1:	116 49 17.1	18.57	1:	B; L; R; f of D neb; 6th of 9	1
2161	3282	10 30 28.7	2.815	2	116 53 42.1	18.57	2	cF; E; gbM; 7th of 9.....	2
2162	3283	10 30 39.9	2.815	1	116 56 48.4	18.58	1	8th of 9.....	1
2163	3284	10 30 59.8	2.817	3	116 52 45.7	18.59	3	F; S; R; bM; 9th of 9.....	3
2164	3285	10 31 6.1	2.636	2	130 54 24.7	18.59	2	cF; pL; pmE; lbM.....	2
2165	III. 700	10 31 7.4	3.526	1	47 36 30.3	18.61	1	cF; L; iE; mb, s of M.....	1
2166	732	II. 745	10 31 12.8	3.626	2	41 51 29.3	18.61	2	F; pS; mE 0...90°; *10 nf.	4
2167	3286	Δ. 322?	10 32 2.9	2.277	1	147 53 48.6	18.62	1	pB; vvL; iF; * inv.....	1
2168	734	II. 348	10 32 17.5	3.299	2	65 11 2.2	18.64	2	vF; S; R; gbM; vS * att....	3
2169	733	10 32 34.7	5.264	3	12 26 32.4	18.68	3	pB; S; iE; psmbM.....	3
2170	I. 272	10 32 34.7	3.158	1:	79 59 19.5	18.65	1:	B; S; iR; mbMBN.....	2
2171	3287	Δ. 355	10 33 1.6	2.409	1	143 24 4.5	18.65	1	Cl; P; st 9.....	1
2172	3288	10 33 26.3	2.729	1	125 19 17.0	18.70	1	eF; vS; mE; *15 att.....	1
2173	735	II. 641	10 33 28.5	3.451	1	51 58 16.4	18.68	1	cF; vS; R; bM.....	4
2174	3289	10 33 43.6	2.823	1	117 1 41.4	18.68	1	vF; pL; iE; glbM.....	1
2175	737	II. 77	10 34 42.7	3.195	2	75 31 44.6	18.72	2	F; cL; E; vgbM; r; *7p10 ^s	4
2176	736	III. 317	10 35 17.6	4.724	1	15 54 50.5	18.75	1	pF; S; R; gbM.....	2
2177	III. 5	10 35 25.5	3.157	1	79 49 40.2	18.74	1	eF; eS.....	1
2178	739	I. 81	10 35 49.5	3.298	1	64 20 20.5	18.75	1	cB; L; gbM; *inv; 2st f....	3
2179	740	I. 26	10 36 8.6	3.178	1	77 16 41.8	18.76	1	cB; pL; E; mbM.....	2
2180	3290	V. 7	10 36 10.9	3.203	1	74 23 11.8	18.76	1	cF; vL; R; vgvbM; er.....	2
2181	3291	10 36 23.8	2.732	4	125 37 46.8	18.76	3	pF; S; mE 0°±; vsymbM; 1st of 3.	4
2182	738	I. 80	10 36 29.0	4.648	1	16 25 22.7	18.79	1	B; S; iE; psbM; *11, 281°-8, 20°-0.	2
2183	742	10 36 33.7	3.358	1	58 32 21.1	18.77	1	eF; vS; 2st 9.10, s.....	1
2184	743	M. 95	10 36 36.7	3.175	2	77 34 22.1	18.77	4	B; L; R; pgmbMrN.....	8
2185	741	III. 842	10 36 39.2	3.782	1	33 18 30.6	18.78	1	cS; R; pgbM; *s 90"....	2
2186	3292	10 36 41.4	2.734	4	125 38 53.1	18.77	3	F; S; vIE; psbM; 2nd of 3...	4
2187	744	III. 107	10 36 53.2	3.133	2	82 30 48.4	18.78	2	vF; pS; R; bM; *9, 150"....	4
2188	3293	10 37 11.7	2.735	...	125 38 59.7	18.79	...	cF; vS; vIE; vS*att; 3rd of 3	2
2189	745	V. 52	10 37 21.0	4.018	1	26 2 25.3	18.81	1	pB; L; E 0°; glbM.....	2*
2190	746	III. 318	10 37 57.1	4.583	1	16 49 37.9	18.83	1	vF; L; R; vgbM; r; *sf....	2
2191	747	10 38 59.5	3.091	1	87 28 32.5	18.85	1	eF; L; eE; vgvbM; a ray...	1
2192	3294	10 39 4.1	2.644	1?	132 58 43.5	18.85	1	F; E; gbM; *6, 7 vnr.....	1*
2193	748	II. 78	10 39 10.5	3.189	2	75 31 1.5	18.85	2	pB; cL; iR; vglbM; r; 1st of 3	5
2194	749	M. 96	10 39 20.4	3.173	4	77 26 55.8	18.86	4	vB; vL; iE; vsymbM; r.....	8
2195	750	II. 81	10 39 34.8	3.219	2	71 59 40.1	18.87	2	cB; cL; iR; vglbM; r.....	3
2196	751	10 39 37.3	3.188	1	75 28 25.1	18.87	1	F; R; 2nd of 3.....	1
2197	3295	Δ. 309	10 39 36.8	2.313	3	148 56 44.8	18.86	3	η Argūs. The great neb....	Mon.†
2198	753	10 39 49.3	3.187	1	75 35 15.1	18.87	1	F; R; 3rd of 3.....	1
2199	752	III. 701	10 39 49.4	3.506	1	46 4 18.4	18.88	1	vF; cS; iR.....	2
2200	D'Arrest, 75	10 40 12	3.12	[1]	83 12 24	18.88	[1]	vF; S.....	0
2201	754	II. 99	10 40 16.7	3.189	2	75 16 24.7	18.89	2	vB; cL; R; vsymbMBN.....	5*
2202	3296	10 40 22.4	2.702	3	129 17 2.7	18.89	3	cF; S; R; glbM.....	4
2203	757	I. 17	Mechain.	10 40 25.7	3.177	3	76 41 10.7	18.89	3	vB; cL; R; psbM; r.....	8
2204	755	II. 360	10 40 27.4	3.323	5	60 39 47.7	18.89	5	pB; pS; R; sbM.....	6
2205	756	II. 565	10 40 33.2	3.390	2	54 33 18.7	18.89	2	F; cL; R; vglbM; 1st of 3	3
2206	3297	10 40 39.5	2.874	2	113 41 36.7	18.89	2	pF; pL; iR; glbM.....	2
2207	758	I. 18	10 40 52.2	3.177	3	76 38 27.0	18.90	3	vB; L; R; psmbM; 2nd of 3	8
2208	759	10 40 54.9	3.116	1	84 15 58.0	18.90	1	vF; R.....	1
2209	760	10 40 55.7	+3.116	2	84 20 22.0	+18.90	2	F; S; iE; bM.....	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
2210	h. 762	H.	h m s 10 41 2·5	s +3·116	1	84 18 1·3	+18·91	1	Suspected; *nr	1
2211	761	II. 41	10 41 3·8	3·176	2	76 43 38·3	18·91	2	F; L; E 90°±; vglbM; 3rd of 3.	9
2212	3298	10 41 32·5	2·807	1	120 48 44·6	18·92	1	F; S; pmE 0°	1
2213	763	III. 881	10 41 36·0	4·082	1	23 28 40·9	18·93	1	vF; S; psbM; st nr	2
2214	3299	10 41 41·8	2·870	3	114 25 38·6	18·92	3	F; S; R; psbM; 2st 10f	3
2215	764	II. 872	10 41 59·8	4·075	1	23 30 51·2	18·94	1	cB; S; lE; vgbM	2
2216	765	I. 116	10 42 2·4	3·364	2	56 16 38·2	18·94	2	cB; pS; iE; 1st of 2	4†
2217	766	I. 117	10 42 7·5	3·364	2	56 15 59·2	18·94	2	pB; pS; iE; 2nd of 2	4†
2218	I. 284	10 42 34·7	5·103	1	11 57 49·1	18·97	1	cB; vS; iF	1
2219	III. 792	10 42 57·4	3·717	1	33 49 54·1	18·97	1	vF; S; E; er	1
2220	768	II. 361	10 43 2·3	3·314	3	60 47 45·1	18·97	3	pF; S; R; bM	4
2221	767	II. 335	10 43 15·4	4·609	1	15 34 35·7	18·97	1	pF; L; iE; vgbM	2
2222	771	10 43 17·6	3·619	1	38 13 52·4	18·98	1	pB; R; pgbM	1
2223	769	III. 919	10 43 20·3	3·884	1	27 53 7·4	18·98	1	vF; vS; R; vS* nr	2
2224	770	III. 913	10 43 21·3	3·792	1	30 49 46·4	18·98	1	vF; cS; R; 2pB st s	2
2225	772	II. 718	10 43 33·3	3·496	1	45 32 51·4	18·98	1	pB; S; vLE; stellar; 3S st nr.	3
2226	772, a	R. nova	10 43 ±	45 ±	3' dist. from h. 772	0
2227	773	II. 362	10 43 33·5	3·307	4	61 17 10·4	18·98	4	B; pL; R; mbM	5
2228	776	III. 522	10 43 28·5	2·982	(3)	102 6 30·0	19·00	1:	F; S; R; lbM	4
2229	774	I. 27	10 43 29·8	3·180	2	75 50 51·4	18·98	2	B; S; lE 135°±; smbMN	5
2230	775	II. 363	10 43 41·7	3·308	2	61 9 14·7	18·99	4	cF; S; R; bM	5
2231	IV. 6	10 43 49·0	3·117	4	83 26 44·4	18·99	4	B; vL; R; bM; r	5*
2232	=	10 43 49·0	3·117	4	83 26 44·4	18·99	4	B; vL; R; bM; r	5*
2232	II. 131	10 43 52·8	3·355	1	56 29 54·7	18·99	1	F; S	1
2233	II. 493	10 43 52·8	3·355	1	56 29 54·7	18·99	1	F; S	1
2233	I. 118	10 43 54·7	3·327	1	57 16 54·7	18·99	1	cB; cL; iR; mbM (758°P.D.)	1*
2234	777	III. 88	10 43 57·2	3·121	3	83 25 0·7	18·99	3	F; vL; R; vgbM; rr	5*
2235	778	II. 494	10 44 1·8	3·356	1	56 21 52·0	19·00	1	pF; pL; lE; sp of 3	3
2236	779	I. 118?	10 44 27·8	3·355	1	56 18 10·3	19·01	1	pB; L; iE; gbM; 2nd of 3...	2*
2237	III. 108	10 44 33·7	3·141	1	80 44 57·3	19·01	1	eF; eS; R	1
2238	780	I. 172	10 44 35·5	3·395	2	52 38 29·3	19·01	2	pB; pL; vmE 42°5; *inv?	3*
2239	782	10 44 40·7	3·356	1::	56 10 30·3	19·01	1::	pB; nf of 3 in a line	1
2240	783	III. 20	10 44 40·7	3·156	1	79 6 47·3	19·01	1	vF; vL; R; vgbM	2
2241	784	III. 497	10 44 43·3	3·105	1	85 28 16·3	19·01	1	F; pS; R; vglbM	3
2242	781	II. 887	10 44 54·9	3·865	1	27 58 21·6	19·02	1	cF; pS; lE; vgbM	2
2243	786	II. 47	10 44 59·2	3·257	2	66 19 33·6	19·02	3	pB; pL; lE 120°; gbM	5
2244	785	III. 914	10 45 10·9	3·740	1	32 8 27·9	19·03	1	vF; S; lE	2
2245	787	I. 267	10 45 58·4	3·730	1	32 16 18·5	19·05	1	cB; pL; iR; vglbM; *10nf2'	2
2246	3301	10 45 58·5	2·660	2	134 24 21·2	19·04	2	Cl; pL; P; lC; iF; st 9...13	2
2247	3300	10 45 59·6	3·203	1	72 29 30·5	19·05	1	eF; vL; vglbM; B *sp	1
2248	788	I. 233	10 46 5·2	3·666	1	34 56 48·5	19·05	1	B; pL; vmE 67°0; gbM	3
2249	3302	10 46 20·2	2·809	1	122 10 58·5	19·05	1	F; S; R; *6·7 sf	1
2250	3303	10 46 25·7	2·919	1	110 6 14·5	19·05	1	vF; L; R; vglbM; r	1
2251	789	II. 364	10 46 40·2	3·291	4	62 1 7·1	19·07	4	F; pL; vLE; vlbM	5
2252	3304	10 46 52·3	2·912	1	111 2 13·1	19·07	1	F; S; R; bM	1
2253	790	10 47 1·0	3·206	1	71 54 33·4	19·08	1	pF; lE; np of 2	1
2254	791	II. 82	10 47 3·0	3·205	1	71 58 7·4	19·08	2	pF; S; E; gbM; r; sf of 2...	3
2255	792	IV. 29	10 47 11·5	2·958	1	105 16 57·4	19·08	1	eF; att to *12f	2
2256	793	10 47 21·7	3·207	1	71 38 54·7	19·09	1	2 or 3 S st & neb	1
2257	I. 268	10 47 23·7	3·721	1	32 8 1·7	19·09	1	vB; vS; R; stellar	1
2258	794	II. 16	10 48 2·0	3·132	1	81 33 42·0	19·10	1	vF; vS; lE; psbM	5
2259	3305	10 48 29·5	2·879	1	115 23 58·3	19·11	1	F; S; R; glbM	1
2260	795	10 48 57·1	4·652	2	14 3 47·2	19·14	2	eF; pL; R; vglbM; *nf	2
2261	796	10 48 57·6	3·146	1	79 30 48·9	19·13	1	vF; *9, 90°; p of 2	1
2262	798	10 49 26·2	3·145	1	79 29 49·2	19·14	1	vF; R; vsmB*12; f of 2...	1
2263	797	III. 632	10 49 33·9	3·425	2	48 18 8·2	19·14	2	F; eS; R; bM	4
2264	3306	10 49 58·9	2·976	1	103 33 21·5	19·15	1	eeF; S	1
2265	799	II. 888	10 50 5·6	3·759	1	29 44 47·8	19·16	1	vF; S; R; vgbM	2
2266	III. 972	10 50 15·1	3·819	1	27 39 6·1	19·17	1	vF; S; R; bM	1
2267	III. 67	10 50 35·3	+3·197	1	72 9 7·1	+19·17	1	vF; E; bet 2 st	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
2321	833	11 3 12.0	+3.253	1	60 36 21.1	+19.47	2	vF; pS; R; bM; 3rd of 4 ...	2
2322	3319	11 3 19.8	2.830	2	126 46 48.1	19.47	2	B; S; R; pgmbM; 1st of 3...	2
2323	III. 79	11 3 25.7	3.144	1	77 19 26.1	19.47	1	eF; pS; lE; r	1
2324	834	11 3 26.1	3.139	1	78 3 34.1	19.47	1	F; S; R; gbM.....	1
2325	830	II. 337	11 3 30.4	4.167	1	16 22 0.4	19.48	1	pF; pS; lE; gbM; *15, 22°1, 70''.	2
2326	835	11 3 36.1	3.252	3	60 32 50.1	19.47	3	vF; pL; 4th of 4	3
2327	3320	11 3 56.5	2.832	2	126 46 43.4	19.48	2	pF; S; R; bM; 2nd of 3.....	2
2328	836	III. 89	11 4 2.4	3.108	2	83 25 3.4	19.48	2	eF; R; sbM; r	3
2329	3321	11 4 9.7	2.835	1	126 42 10.4	19.48	2	vF; pL; R; *inv; 3B st nr...	2
2330	II. 819	11 4 32.1	2.972	1	107 31 27.7	19.49	1	pF; pL; iF; bM	1
2331	3323	11 4 32.5	2.534	1	149 28 52.7	19.49	1	Cl; pRi; lC	1
2332	3322	11 4 39.9	2.842	1	126 5 31.7	19.49	1	eF; S; R; glbM; 3 st 11 f ...	1
2333	3324	11 5 52.5	2.522	1	150 36 57.3	19.51	1	F; lE; 1st of 6	1+
2334	III. 723	11 5 53.4	3.424	1	40 53 27.6	19.52	1	eF; vS; p of 2	1
2335	837	11 5 54.4	2.988	1	105 11 45.6	19.52	1	::: Neb (?)	1
2336	3325	11 5 56.2	2.526	2	150 28 3.6	19.52	2	F; lE; sbM; 2nd of 6.....	2+
2337	3326	11 6 3.2	2.526	2	150 32 58.6	19.52	2	*12 with fan-shaped neb att; 3rd of 6.	2+
2338	3327	11 6 14.4	2.528	2	150 30 34.6	19.52	2	B; bM*; 4th of 6	2+
2339	II. 728	11 6 18.3	3.422	2	40 51 28.9	19.53	2	pF; pL; R; vgmbM	2
2340	3329	11 6 21.4	2.541	2	150 26 42.6	19.52	2	F; L; E0°; bM; 5th of 6 ...	2+
2341	3328	II. 269	11 6 28.1	2.923	2	115 59 51.9	19.53	2	B; pL; E; vsmbMN; 2 BstΔ	4
2342	3330	11 6 31.0	2.540	2	150 35 29.9	19.53	2	eF; S; E160°±; 6th of 6 ...	2+
2343	838	M. 97	11 6 34.8	3.514	1	34 13 38.2	19.54	1	!; O; vB; vL; R; vvg; vsbMO; 19°0 d.	4+
2344	839	III. 921	11 6 51.9	3.621	1	28 32 29.2	19.54	1	vF; L; E; vgmbM; in Δ of L st	2
2345	3332	11 6 59.1	2.546	1	150 2 19.2	19.54	1	Cl; pRi; C; E.....	1
2346	3331	III. 529	11 7 4.5	3.000	1	103 19 48.2	19.54	1	vF; S; iR; lbM	3
2347	840	I. 29	11 7 16.1	3.144	3	76 25 29.5	19.55	3	B; cL; E90°±; psmbM.....	4
2348	III. 770	11 7 26.0	3.520	1	33 29 29.5	19.55	1	vF; vS; stellar.....	1
2349	III. 706	11 7 30.1	3.404	2	41 45 30.5	19.55	2	vF; vS; vL; stellar; cB* n...	2
2350	841	II. 102	11 7 43.8	3.154	1	74 26 53.8	19.56	2	pF; L; R; glbM	2
2351	3333	11 7 49.1	2.946	1	112 57 42.8	19.56	1	vF; pS; R; bM	1
2352	843	II. 49	11 8 4.2	3.172	1	71 7 42.8	19.56	1	B; pS; R; pgmbM	2
2353	842	II. 709	11 8 6.6	3.339	1	47 38 28.8	19.56	1	pF; S; lE0°±; vgmbM	4
2354	3334	11 9 7.0	2.558	1	150 29 39.4	19.58	1	⊕ and neb; st 15...18	2
2355	II. 626	11 9 7.9	3.098	1	84 42 34.4	19.58	1	pB; S; lE; mbM.....	1
2356	844	III. 27	11 9 21.8	3.169	2	71 14 8.7	19.59	2	F; S; R; sp of 3	3
2357	3335	11 9 29.9	2.885	1	123 3 54.7	19.59	1	eF; S; R; gbM	1
2358	845	II. 50	11 9 30.6	3.169	2	71 11 38.7	19.59	2	vB; L; R; vmbM; 2nd of 3...	3
2359	846	II. 51	11 9 35.3	3.169	2	71 5 49.7	19.59	2	B; pL; R; psbM; 3rd of 3 ...	3
2360	847	I. 270	11 10 11.5	3.548	3	30 27 29.0	19.60	3	vB; pS; lE90°±; vsmbMSN.	5
2361	849	II. 521	11 10 16.1	3.098	2	84 40 50.0	19.60	2	pF; cS; iR; psmbM; *10, 330°, 3'.	5
2362	848	I. 271	11 10 20.9	3.533	1	31 14 40.3	19.61	1	vB; cL; mE305°0; smbMN	2
2363	850	II. 729	11 10 29.9	3.366	1	43 29 5.3	19.61	1	F; pL; lE90°±; glbM; r ...	3
2364	851	III. 333	11 10 39.5	3.197	1	65 49 46.3	19.61	1	cF; vS; smbM; stellar; p of 2	2
2365	851, a	R. nova	11 10 ±	65 49 ±	F; S; bM; place from MS ...	0
2366	III. 76	11 10 53.0	3.149	1	74 29 36.6	19.62	2	eF; pL.....	1
2367	3336	11 10 59.5	2.939	1	115 21 55.6	19.62	1	F; S; R; gbM.....	1
2368	III. 334	11 11 0.0	3.197	1	65 42 34.6	19.62	1	vF; S; f of 2	1
2369	852	I. 244	11 11 9.6	3.520	1	31 30 10.6	19.62	1	cB; cL; R; vgmbM	3
2370	3338	11 11 21.4	1.992	1	165 27 17.6	19.62	1	F; pS; pmE; gbM	1
2371	3337	I. 241	Δ. 617	11 11 29.6	2.899	1	122 2 45.9	19.63	1	cB; vL; E160°±; am 4 st...	2
2372	853	II. 879	11 11 32.8	3.745	1	21 59 31.9	19.63	1	pB; S; R; gbM	2
2373	854	M. 65	11 11 36.7	3.139	3	76 8 46.9	19.63	3	B; vL; mE165°±; gbMBN.	8*+
2374	855	11 11 40.5	3.111	1	81 42 52.9	19.63	1	eF	1
2375	II. 885	11 12 17.2	3.512	1	31 22 36.2	19.64	1	F; S; lE; 135°±	1
2376	856	II. 52	11 12 43.1	+3.164	1	70 52 55.5	+19.65	1	B; S; vL; sbM	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
2377	h. 857 = 875?	H.	M. 66	h m s 11 12 48.3	s +3.137	2	° ' " 76 15 17.5	" +19.65	3	B; vL; mE150°; mbM; 2stnp	9*†
2378	859	V. 8	11 12 57.4	3.140	4	75 38 28.5	19.65	4	pB; vL; vmE102°0	8†
2379	858	I. 226	11 13 1.7	3.435	1	36 4 6.5	19.65	1	pB; L; R; svmbMrN	2†
2380	860	II. 338	11 13 3.3	3.211	7	62 16 12.5	19.65	7	cF; L; R; vgvbM	9
2381	861	11 13 5.2	3.089	2	86 16 11.5	19.65	2	pB; S; R; smbMN	3
2382	II. 30	11 13 6.1	3.163	1	71 4 37.5	19.65	1	pB; *inv	1*
2383	862	II. 550	11 13 21.8	3.027	2	99 30 44.8	19.66	2	F; vS; R; lbM; *7 f; p of 2.	4
2384	863	II. 551	11 13 36.0	3.028	2	99 28 56.8	19.66	2	F; vS; R; psbM; *7p; f of 2	4
2385	856, a	R. nova	11 13 46.1	3.164	::	70 52 55.5	19.66	::	pF; S; R; vlbM; foll h. 856, 15'.	0
2386	864	II. 33	11 13 53.9	3.090	1	85 59 53.1	19.67	1	B; pL; R; psbM	3
2387	865	I. 245	11 14 10.0	3.515	4	30 9 43.1	19.67	4	pB; pL; R; vgbM	7
2388	867 = 861?	II. 32	11 14 20.6	3.088	1?	86 16 50.1	19.67	1?	pB; S; E; bM	1*
2389	866	III. 15	11 14 20.9	3.170	2	69 4 20.4	19.68	2	cF; cL; lE; gbM; sp of 2 ...	3
2390	868	11 14 51.7	3.290	1	49 22 36.7	19.69	1	pB; S; pmE; bMN=close*	1
2391	869	III. 16	11 14 52.5	3.169	2	69 1 43.7	19.69	2	vF; pS; R; gbM; nf of 2 ...	3
2392	870	III. 335	11 15 1.4	3.190	1	64 56 3.7	19.69	1	cF; vS; R; bM; nf of 2 ...	2
2393	872	III. 336	11 15 6.1	3.190	1	64 57 48.7	19.69	1	vF; vS; sf of 2	2
2394	871	II. 775	11 15 6.6	3.274	1	51 27 49.7	19.69	1	pF; cL; lE; vgbM	2
2395	II. 880	11 15 24.0	3.766	2	19 47 39.0	19.70	2	F; S; lE15°±	2
2396	873	I. 5	11 15 33.5	3.150	3	72 38 30.0	19.70	3	pB; pS; iR; bM; r	5
2397	874	II. 782	11 15 40.7	3.423	1	35 23 23.0	19.70	1	pB; S; R; vgbM; *12p	2
2398	876	III. 768	11 15 58.6	3.409	1	36 18 44.3	19.71	1	cF; vS; R; stellar	3
2399	878	II. 53	11 16 22.1	3.154	1	71 25 28.3	19.71	1	cF; S; lE; r	3
2400	877	IV. 59	11 16 23.3	3.273	1:	50 42 2.3	19.71	1:	{ H. cB; S; R; svmbMN } h. F; R; *17M.	3
2401	II. 635	11 16 27.0	3.037	1	97 52 41.3	19.71	1	F; pL; iR; vgbM	1
2402	3339	III. 530	11 16 37.0	3.014	1	103 3 31.3	19.71	1	F; S; R; stellar; p of 2	2
2403	879	IV. 4	11 16 39.9	3.070	1	90 19 53.3	19.71	1	vF; S; att to *13 m	3
2404	881	I. 219	11 17 10.2	3.271	2	50 28 40.6	19.72	2	cB; cL; iR; pgbM	3
2405	882	I. 20	11 17 11.3	3.123	3	77 53 21.6	19.72	3	F; E90°±; B* f 34s	5*
2406	3340	III. 531	11 17 11.8	3.015	2	103 4 29.6	19.72	1	pF; pL; iR; vlbM	4
2407	880	II. 845	11 17 14.0	3.569	1	25 47 2.9	19.73	1	F; pS; iR; gbM; *9np	4
2408	883	II. 829	11 17 24.4	3.462	1	31 30 53.9	19.73	1	vF; pL; pmE135°±; er	3
2409	884	III. 337	11 17 25.2	3.182	1	65 16 34.9	19.73	1	vF; vS; R	2
2410	885	III. 922	11 17 54.2	3.503	1	28 45 10.2	19.74	1	vF; vS; 2 vS st inv	2
2411	886	I. 131	11 17 58.3	3.034	1	99 1 48.9	19.73	2	pB; L; E0°±; gbM	3*
2412	3341	11 18 20.4	2.956	1	115 58 29.2	19.74	1	F; vL; gvlbM; *7 s 6'	1
2413	887	I. 194	11 18 26.4	3.301	2	45 38 32.2	19.74	2	vB; cL; vmE0°±; vsmbMN; st p.	4
2414	888	11 18 34.5	3.328	2	42 14 27.2	19.74	2	eF; S; R; vsbM*; 2st11 nf.	2
2415	II. 886	11 18 36.9	3.442	1	32 6 42.5	19.75	1	pF; iF	1
2416	889	11 18 47.7	3.199	1	61 21 44.5	19.75	1	vF; S; R; psbM; *12 nf	1
2417	III. 112	11 18 51.5	3.051	2?	95 4 52.5	19.75	2?	eF; cL; R; r (v near vB*)	3*
2418	3342	Δ. 481	11 18 56.6	2.858	2	132 27 50.5	19.75	2	Cl; cL; pRi; lC; st 10...14..	2
2419	{ 891 = 3343 }	II. 159	11 19 10.3	3.145	4	72 22 2.5	19.75	4	B; pS; R; bM	5
2420	890	I. 262	11 19 19.3	3.618	1	22 38 33.8	19.76	1	cB; S; iR; spmbMN	2
2421	892	I. 246	11 19 32.8	3.431	1	32 20 57.8	19.76	1	cB; pL; E	3
2422	893	11 19 52.5	3.144	2	72 12 19.8	19.76	3	pB; pL; E; vgbM	3
2423	894	{ II. 160 = III. 28 }	11 20 23.8	3.144	2	72 0 27.1	19.77	2	pB; L; vIE; vgbM; r	5
2424	895	II. 770	11 20 32.3	3.202	5	59 42 52.1	19.77	5	pB; pS; R; lbM; r	6
2425	896	I. 247	11 20 44.8	+3.441	1	30 40 47.4	+19.78	2	pB; pS; vIE80°±; pgbM; Sst sf nr.	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
2426	897	II. 339	11 20 45.7	+3.182	1	63° 34' 36".4	+19.78	1	pB; pL; IE; bM	2
2427	898	II. 54	11 20 49.6	3.142	2	72 18 40.4	19.78	2	F; pS; IE; r	4
2428	II. 152	11 20 59.6	3.111	1	79 41 44.4	19.78	1	F; mE; r	1
2429	3334	III. 532	11 21 9.0	3.023	1	102 24 55.4	19.78	1	cF; S; E; gbM	2
2430	899	11 21 22.8	3.231	1	53 48 40.7	19.79	1	cF; S; R; sbM*?	1
2431	900	11 21 30.0	3.158	1	68 25 51.7	19.79	1	eF; vS; E 90°	1
2432	3345	11 21 33.5	2.706	1	149 11 3.7	19.79	1	B; pL; iR; pgpmbM	1
2433	901	II. 349	11 22 5.3	3.171	1	65 7 52.0	19.80	1	pF; pL; IE	2
2434	902	II. 13	11 22 53.7	3.109	1	79 57 4.3	19.81	1	pF; pL; R; vsmbM; r	4
2435	3346	11 22 53.9	2.920	3	125 37 34.3	19.81	3	pB; cS; R; psmbM	3
2436	903	11 23 0.3	3.109	1::	79 52 19.3	19.81	1	pB; cS; E 90°	1
2437	904	II. 350	11 23 43.7	3.161	1	66 27 39.6	19.82	1	F; S; *7.8 nf 5'	2
2438	905	11 24 15.2	3.185	1	60 44 5.6	19.82	1	F; vS; R; sbmM	2
2439	906	II. 367	11 24 17.1	3.183	3	61 4 23.6	19.82	3	F; cS; R; sbMN	4
2440	907	III. 353	11 24 28.7	3.183	3	60 52 0.9	19.83	3	F; S; R; psbM	5*
2441	3347	II. 562	11 24 29.0	3.024	2	103 27 38.9	19.83	2	pF; S; R; vglbM	5
2442	3348	11 24 36.6	2.958	2	119 29 4.9	19.83	2	pB; S; mE; * 13 att	2
2443	908	I. 221	11 24 46.4	3.342	2	36 9 21.9	19.83	2	pB; vL; R; vglbM	4
2444	909	II. 836	11 25 43.2	3.442	2	27 21 32.2	19.84	2	cF; S; R; vglbM; r	3
2445	910	II. 730	11 25 44.5	3.284	2	42 10 57.2	19.84	2	pB; vL; IE 0°; vsmbM*15; *11 n.	3+
2446	912	II. 351	11 26 2.1	3.162	1	64 46 58.5	19.85	1	F; S; R; bM	2
2447	911	I. 222	11 26 2.3	3.333	1	36 5 47.5	19.85	1	pB; pL; IE 0° ±; gbM; *12nr	3
2448	III. 80	11 27 5.9	3.115	1	76 43 50.8	19.86	1	vF; vS; R	1
2449	913	II. 552	11 27 8.0	3.042	2	99 3 46.8	19.86	2	F; S; R; psbM; *14 sp 225°	3
2450	III. 771	11 27 10.4	3.340	1	34 23 50.8	19.86	1	eF; S; iR; L * in field	1
2451	3349	III. 935	11 27 37.1	3.038	1	103 19 8.1	19.87	1	eF; S; R; gbM	2
2452	914	I. 287	11 27 44.4	3.602	1	18 41 33.1	19.87	1	pB; L; mE 130°.4; mbM ...	2
2453	III. 772	11 27 44.9	3.340	1	34 15 51.1	19.87	1	vF; stellar	1
2454	II. 783	11 28 6.8	3.331	1	34 41 52.1	19.87	1	pB; pL; bM	1
2455	915	III. 847	11 28 23.8	3.387	1	29 15 14.4	19.88	1	vF; vS; R; vgbM	2
2456	916	11 28 33.7	3.255	1	43 56 23.4	19.88	1	vF; S; R; vgbM	1
2457	3350	11 28 37.1	2.938	2	127 10 35.4	19.88	2	pF; pL; vLE; glbM	2
2458	III. 969	11 28 56.6	3.748	1	14 14 50.7	19.89	1	eF; S	1
2459	3351	11 28 59.4	2.939	2	127 12 45.4	19.88	2	F; cS; IE; gvlbM	2
2460	917	II. 905	11 29 3.2	3.784	1	13 56 57.4	19.88	1	pB; pL	2
2461	918	II. 784	11 29 4.5	3.319	1	34 55 43.7	19.89	1	pF; L; IE	2*
2462	920	11 29 4.7	3.203	1	52 49 3.7	19.89	1	eF; pL; pmE; gbM	1
2463	919	III. 843	11 29 8.5	3.360	1	30 49 18.7	19.89	1	vF; R; stellar; vS* 1 d sf ...	1
2464	D'Arrest, 78	11 29 16	3.14	[1]	67 24 12	19.88	[1]	B; pS; mbMN=*13; *11 p 4 ^s , s 175".	0
2465	921	II. 837	11 29 29.6	3.398	1	27 29 1.7	19.89	1	F; vLE; gbM	2
2466	D'Arrest, 79	11 29 39	3.13	[2]	71 20 42	19.89	[2]	F; S; R	0
2467	922	11 29 41.9	3.150	1	65 7 26.7	19.89	1	vF; S; R	1
2468	3352	Δ. 289	11 29 42.0	2.764	2	150 49 29.7	19.89	2	Cl; pL; pRi; pC; st 8...13 ..	2
2469	923	III. 29	11 29 54.8	3.128	1	71 23 48.7	19.89	1	vF; eS; stellar	2
2470	924	11 29 55.8	3.125	1	72 20 37.7	19.89	1	vF; S; bM	1
2471	925	II. 731	11 30 10.0	3.262	2	41 19 6.0	19.90	2	pB; S; pmE	4
2472	926	II. 838	11 30 13.1	3.364	2	29 36 37.0	19.90	2	pF; S; R; gbM; r	3
2473	927	II. 352	11 30 27.9	3.143	2	66 32 34.0	19.90	2	vF; S; E; r	4
2474	928	III. 81	11 30 58.4	3.109	1	77 6 36.3	19.91	1	cF; cS; R; psbM	4
2475	3353	11 31 34.0	2.881	1	139 55 51.3	19.91	1	cF; S; R; am 50 Sst	1
2476	929	I. 227	11 31 41.7	3.315	2	32 57 27.3	19.91	2	pF; L; vLE; vgbM; r	4
2477	930	II. 732	11 32 1.9	3.241	1	42 45 34.6	19.92	1	F; S; att to *15; another * cont.	2
2478	3354	11 32 6.8	2.954	1	126 57 46.6	19.92	2	cB; R; sbMN*; *9 sf	2
2479	931	11 32 18.7	3.171	3	57 18 46.6	19.92	3	pB; pL; pmE; gbM; p of 2...	3
2480	932	11 32 21.7	3.171	3	57 17 20.6	19.92	3	cB; pL; pmE 0°; pgbM	3
2481	933	III. 109	11 32 29.4	3.123	2	71 31 5.6	19.92	2	cF; vS; pmE; sbM; 2 Sst f; 1st of 3.	3
2482	935	III. 609	11 32 35.2	+3.048	1	98 35 14.6	+19.92	1	vF; vS; R; gbM; *8 s 6' ...	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.		
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.										
2483	h.	H.		h m s	s								
2484	III. 773	11 32 45.1	+3.305	1	32 59 54.6	+19.92	1	cF; pS; vS*v nr	1		
2485	937	II. 839	11 32 47.4	3.327	1	30 36 24.9	19.93	1	vF; S; mE	1		
2486	{ 934 = 3355 }	11 32 47.9	3.345	2	28 56 14.9	19.93	2	F; cS; R; mbM	3		
2487	938	II. 340	11 32 54.0	3.116	2	73 53 41.9	19.93	3	cF; R; p of 2	3+		
2488	{ 936 = 3356 }	II. 103	11 32 54.5	3.144	1	64 31 35.9	19.93	1	F; cS; lE; stellar; r	3		
2489	936, a	R. nova	11 32 57.8	3.116	4	73 52 36.9	19.93	6	F; pS; E; pglbM; r; f of 2...	8+		
2490	939	II. 161	11 33 ±	73 52 ±	Smaller than h. 936	0		
2491	939, a	}	R. 3 novæ	11 33 0.4	3.122	3	71 29 46.9	19.93	3	pF; pL; R; bM; r; 2nd of 3	4		
2492	939, b			11 33 ±	71 29 ±	No description (for 939, c, see No. 5069).	0	
5069	
2493	940	III. 30	11 33 5.0	3.122	3	71 27 36.9	19.93	2	vF; pS; r; 2 vB st p; 3rd of 3	4		
2494	II. 830	11 33 23.7	3.300	1	32 58 24.9	19.93	1	pB; E	1		
2495	941	III. 375	11 33 26.1	3.129	1	68 53 9.9	19.93	1	cB; cS; R; bM; r	3		
2496	D'Arrest, 80	11 33 29	3.12	[1]	71 25 42	19.93	[1]	F; pL; *9.10, s 5'	0		
2497	III. 338	11 33 35.4	3.135	1	66 46 41.9	19.93	1	vF; vS	1		
2498	942	II. 737	11 33 45.6	3.237	2	41 30 46.9	19.93	2	F; S; vIE; glbM	4		
2499	943	I. 21	11 33 45.7	3.104	3	77 45 10.9	19.93	4	B; L; vIE	8		
2500	944	III. 320	11 33 48.3	3.142	1	64 24 10.9	19.93	1	cF; vS; R; p of 2; * 6 sf 3'	3		
2501	945	I. 94	11 33 53.4	3.183	3	52 40 24.2	19.94	3	cB; pL; pmE 90°±; bM	5*		
2502	946	III. 339	11 34 20.2	3.140	1	64 25 31.2	19.94	1	eF; S; f of 2	2		
2503	947	11 34 40.4	3.099	1	78 55 43.2	19.94	1	F; 1st of 4	1		
2504	948	III. 284	11 34 47.6	3.058	1	95 23 14.2	19.94	1	F; pS; R; psbM	3		
2505	950	11 34 51.9	3.099	1	78 54 43.2	19.94	1	vF; 2nd of 4	1		
2506	949	III. 376	11 34 52.5	3.126	2	68 54 26.5	19.95	2	vF; cS; R; bM; bet 2 st.	5		
2507	951	II. 153	11 34 58.4	3.099	1	78 57 13.5	19.95	1	pF; pS; 3rd of 4	2		
2508	3357	11 35 9.4	3.039	1	103 4 35.5	19.95	1	F; cS; lE; pslbM	1		
2509	952	III. 774	11 35 10.2	3.258	1	36 26 51.5	19.95	1	vF; cS; pmE	3		
2510	953	II. 154	11 35 11.4	3.099	1	78 58 3.5	19.95	1	pF; pS; 4th of 4	2		
2511	954	II. 341	11 35 14.4	3.143	3	62 44 1.5	19.95	3	pF; S; R; psbM; stellar	4		
2512	D'Arrest, 81	11 35 21	3.12	[1]	70 22 42	19.94	[1]	F; S; lbM	0		
2513	955	III. 775	11 35 51.3	3.253	1	36 29 51.5	19.95	1	vF; vS	2		
2514	956	11 35 56.5	3.141	1	62 43 32.5	19.95	1	eF	1		
2515	957	11 36 11.6	3.043	1	102 5 22.8	19.96	1	F; vS; R; bM	1		
2516	III. 340	11 36 15.0	3.130	1	66 24 56.8	19.96	1	vF; pL; 2 suspected neb nr	1		
2517	III. 102	11 36 15.9	3.097	1	79 6 56.8	19.96	1	eF; pS	1		
2518	D'Arrest, 82	11 36 21	3.12	[1]	70 7 42	19.95	[1]	vF; vS; slbMN* 13 m	0		
2519	958	11 36 24.4	3.306	2	29 6 47.8	19.96	2	pB; E; gbM; *8 nf 5'	1		
2520	959	II. 831	11 36 29.2	3.286	2	31 16 21.8	19.96	2	pB; cS; E; psbM*12	3		
2521	960	11 36 32.6	3.121	2	69 15 52.8	19.96	2	cF; S; R; 1st of 5	1		
2522		
2523	} 960, a	R. 4 novæ	11 36 ±	69 15 ±	8 "knots" (vide h. 960, 1, 2, 3)	0		
2524			
2525			
2526	961	11 36 39.1	3.120	2	69 20 12.8	19.96	2	cF; S; R; 2nd of 5	1		
2527	962	III. 377	11 36 44.2	3.120	3	69 17 13.8	19.96	3	F; S; R; vglbM; 3rd of 5	5		
2528	963	11 36 46.7	3.120	1	69 14 9.8	19.96	1	vF; pS; 4th of 5	1		
2529	964	11 36 53.3	3.264	1	33 34 40.8	19.96	1	F; pL; R; vgbM	1		
2530	965	11 36 53.6	3.159	1	55 42 18.8	19.96	1	F; S; R; psbM	1		
2531	III. 35	11 36 57.9	3.098	1	78 55 57.8	19.96	1	eF; vS	1		
2532	III. 776	11 37 2.1	3.265	1	33 19 56.8	19.96	1	eF; pL; lE	1		
2533	966	III. 378	11 37 6.3	3.120	1	69 15 33.8	19.96	1	eF; vS; R; 5th of 5	3		
2534	III. 36	11 37 11.9	3.098	1	78 55 57.8	19.96	1	eF; vS	1		
2535	III. 386	11 37 37.2	3.118	1	69 27 57.1	19.97	1	vF; vS; r	1		
2536	970	11 37 45.7	3.118	1	69 14 20.1	19.97	1	F; S; R; bM(?)	1		
2537	III. 385	11 37 53.2	3.117	1	69 35 57.1	19.97	1	vF; vS; r	1		
2538	971	D'Arrest	11 38 30.2	+3.096	2	78 23 56.4	+19.98	2	F; S; iR; psbM	1		

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
2539	h. 972	H. III. 833	h m s 11 38 34.3	s +3.218	3	39 1 19.4	+19.98	3	cF; cS; R; psbM	4
2540	967	11 38 34.6	3.152	1?	56 6 59.4	19.98	1?	eF; R; gbM; 1st of 4	1*
2541	973	II. 104	11 38 35.6	3.102	3	75 27 32.4	19.98	3	B; S; R; smbM*	5
2542	III. 104	11 38 36.0	3.091	1	80 40 29.4	19.98	1	vF; vS; suspected	1
2543	III. 387	11 38 37.2	3.116	1	69 27 58.4	19.98	1	vF; vS; r	1
2544	III. 103	11 38 37.6	3.093	1	80 1 58.4	19.98	1	vF; r	1
2545	I. 201	11 38 55.9	3.202	2	41 43 58.4	19.98	2	B; L; mE 25° ±	2
2546	974	11 38 56.4	3.150	1	56 1 20.4	19.98	1	vF; R; 2nd of 4	1
2547	II. 881	11 39 0.8	3.391	1	19 49 57.4	19.98	1	F; pL; mE 105° ±	1
2548	{ 968 = 975 969	11 39 1.4	3.150	2	56 3 20.4	19.98	2	vF; R; gbM; 3rd of 4	2
2549	{ = 976	11 39 14.9	3.149	2	56 7 5.4	19.98	2	vF; R; gbM; 4th of 4	2
2550	3358	11 39 16.6	2.903	2	145 36 20.4	19.98	2	vF; lE; 2 st inv	1
2551	III. 372	11 39 21.8	3.112	1	68 35 58.4	19.98	1	vF; cL	1
2552	977	III. 388	11 39 30.9	3.116	1	68 49 39.4	19.98	1	cF; S; iR; gbM; r; *7 sp 6'	4
2553	3359	III. 828	11 39 45.1	3.014	2	117 8 33.7	19.99	2	cF; vS; vIE; bM; vF*sf ...	2
2554	{ 979 = 3360	I. 120	11 39 56.8	3.040	2	106 4 52.7	19.99	2	pB; L; iR; vgpmbM	3
2555	978	II. 785	11 40 6.3	3.238	2	33 15 24.7	19.99	2	pB; S; lE; pgbM	3
2556	980	II. 723	11 40 45.5	3.136	3	58 51 30.7	19.99	3	pB; S; bM	4
2557	{ 981 = 3361	II. 553	11 40 53.3	3.053	2	100 10 27.7	19.99	2	pB; pL; R; gbM; r	4
2558	III. 940	11 40 54.2	3.462	1	14 52 59.0	20.00	1	vF; S; R; bM	1
2559	982	II. 738	11 41 12.9	3.192	1	40 30 26.0	20.00	1	B; pL; R; mbM	2+
2560	983	I. 248	11 41 18.7	3.250	2	29 48 43.0	20.00	2	B; pL; iR; pgmbM; p of 2 ..	4
2561	984	II. 832	11 41 29.7	3.248	1	29 47 51.0	20.00	1	pF; pL; vIE; gbM; f of 2 ...	3
2562	II. 739	11 41 36.1	3.189	1	40 30 59.0	20.00	1	F; vS	1
2563	986	II. 408	11 41 41.5	3.144	3	54 11 35.0	20.00	3	F; S; R; bM	5
2564	985	I. 228	11 41 45.6	3.224	4	33 8 26.0	20.00	4	B; pL; lE; svmbM	6
2565	987	11 41 51.5	3.123	1	62 46 31.0	20.00	1	pB; R; smbM	1
2566	988	I. 82	11 41 53.3	3.124	5	62 11 43.0	20.00	5	B; pL; vIE 0° ±; bMN	8
2567	III. 970	11 41 58.8	3.564	1	11 7 57.0	20.00	1	pF; pL; r	1
2568	989	III. 321	11 42 1.2	3.122	2	63 5 50.0	20.00	2	F; pS; lE; vglbM	3
2569	3362	11 42 1.4	2.998	2	126 44 21.0	20.00	2	pB; cS; vIE; lbM	2
2570	3363	II. 864	11 42 9.2	3.019	1	118 32 27.0	20.00	1	pB; S; R; mbM	2
2571	III. 715	11 42 11.9	3.185	1	40 48 0.0	20.00	1	eF; pL	1
2572	990	11 42 19.8	3.065	1	90 19 7.0	20.00	1	eF; S; psbM	1
2573	3364	11 42 34.3	2.967	1	137 29 23.0	20.00	1	Cl; vL; lC; st 9...14	2
2574	991	III. 341	11 42 42.2	3.117	(1)	64 17 48.0	20.00	1:	vF; S; p of 2	2
2575	992	II. 342	11 42 49.0	3.120	3	62 44 0.3	20.01	3	F; pL; R; pgbM	4
2576	II. 786	11 43 0.7	3.209	1	33 53 0.3	20.01	1	F; E	1
2577	III. 113	11 43 4.8	3.066	1	94 21 0.3	20.01	1	eF; eS; bet 2 st	1*
2578	993	II. 787	11 43 20.3	3.206	1	34 4 32.3	20.01	1	eF; R; gbM	2
2579	994	11 43 20.4	3.190	1	37 23 43.3	20.01	1	F; L; vmE; vgbM	1
2580	995	III. 90	11 43 20.5	3.084	2	82 38 55.3	20.01	3	F; vS; R; lbM; *13 np 80'	4
2581	3365	11 43 23.9	2.934	4	146 24 8.3	20.01	5	○; l; S; R; blue; =*7m; 1*5=d.	6
2582	996	11 43 33.5	3.115	1::	64 17 48.3	20.01	1	Neb; f of 2	1
2583	II. 824	11 43 36.9	3.193	1	36 21 30.3	20.01	1	pB; L; mE	1
2584	997	II. 788	11 43 37.1	3.203	1	34 8 32.3	20.01	1	pF; S; R; pspmbM	2
2585	III. 716	11 43 47.7	3.180	1	39 1 0.3	20.01	1	vF; vS	1
2586	3366	I. 259	11 43 57.8	3.025	2	118 3 5.3	20.01	2	B; pL; lE; gmbM; r; vS*sp inv.	3
2587	II. 825	11 43 59.4	3.189	1	39 8 0.3	20.01	1	pB; S; iF; bM	1
2588	D'Arrest, 83	11 44 6	3.11	[1]	67 19 48	20.01	[1]	vF; vS	0
2589	998	III. 379	11 44 11.2	+3.108	1	67 12 20.3	+20.01	1	eF; eS; vIE; er; st nr.....	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
2590	h. 999	H. II. 740	h m s 11 44 24.7	s +3.170	1	40° 31' 56.6	+20.02	1	pF; S; R; pspmbM	2
2591	1000	III. 616	11 44 29.7	3.139	2	51 13 20.6	20.02	2	eF; cL; iF; glbM; *6n5'; *7f	4*
2592	III. 769	11 44 39.8	3.182	1	37 16 0.6	20.02	1	eF; S	1
2593	D'Arrest, 84	11 44 49	3.17	[1]	40 35 48	20.01	[1]	vF; v diffie; H. II. 740 np	0
2594	1001	11 45 8.9	3.124	3	56 48 50.6	20.02	3	pF; S; IE; psbM	3
2595	3367	11 45 19.8	3.033	1	116 7 21.6	20.02	1	vF; cL; vmE 59° 3'	1
2596	1003	III. 389	11 45 28.9	3.103	1	68 35 6.6	20.02	1	vF; cS; R	2
2597	1002	I. 203	11 45 30.5	3.150	2	45 5 35.6	20.02	2	R; vL; R; bMpBN; er	3+*
2598	III. 971	11 45 31.1	3.378	1	14 7 0.6	20.02	1	eF; vS; R	1
2599	1004	III. 380	11 45 34.2	3.103	1	68 14 0.6	20.02	1	vF; cS; R	3
2600	1005	I. 173	11 45 38.3	3.132	3	52 13 50.6	20.02	3	vB; pL; R; smbM*9	4
2601	1007	III. 322	11 45 50.3	3.111	3	63 0 39.6	20.02	3	pF; pS; R; psbM	4
2602	1006	I. 251	11 45 50.5	3.211	2	28 33 14.6	20.02	2	B; pL; R; gmbM; r; *f.	3
2603	1008	II. 403	11 46 7.9	3.101	1	68 28 0.6	20.02	2	F; pS; iE; lbM; *p	5
2604	1009	I. 202	11 46 21.7	3.155	2	41 21 59.9	20.03	2	cB; pL; pmE; vgbM	4*
2605	1010	III. 342	11 46 22.4	3.105	1	65 49 28.6	20.02	1	vF; cS; vIE	2
2606	1011	V. 45	11 46 27.4	3.169	1	36 53 3.9	20.03	1	cB; L; E 0° ±; vsbMLrN	3+
2607	1012	III. 612	11 46 28.1	3.068	2	93 13 23.9	20.03	2	cF; cS; IE 90° ±; bM; r	4
2608	1013	III. 381	11 46 30.0	3.101	1	68 20 44.9	20.03	1	eF; R	2*
2609	II. 623	11 46 53.7	3.043	2	112 24 0.9	20.03	2	cF; S; E 170° ±; lbs	2
2610	3368	III. 290	11 46 56.8	3.047	1	109 47 29.9	20.03	1	cF; pL; pmE 56° 8'	2
2611	II. 294	11 46 58.4	3.048	1	108 47 0.9	20.03	1	F; S; E; r	1
2612	1014	II. 833	11 47 1.8	3.188	2	30 51 20.9	20.03	2	pF; pS; pmE; vgbM	3
2613	1015	IV. 67	11 47 27.9	3.184	2	30 43 51.9	20.03	2	pF; cL; R; vg; sbM	3
2614	3369	Δ. 349	11 47 29.3	2.976	1::	144 56 21.9	20.03	1	Cl; pL; pRi; gplmbM; st 13	3
2615	III. 905	11 47 31.4	3.254	1	19 54 0.9	20.03	1	eF; vS	1
2616	3370	I. 67	11 47 32.4	3.056	2	103 11 30.9	20.03	2	cB; pL; iR; gmbM; Δ 2 st.	6
2617	1016	11 47 39.1	3.108	2	60 57 8.9	20.03	2	vF; S; E; *10nf att	3
2618	II. 789	11 48 4.2	3.166	1	33 54 0.9	20.03	1	pB; E	1
2619	II. 790	11 48 4.2	3.166	1	33 54 0.9	20.03	1	F; S	1
2620	1017	IV. 62	11 48 6.5	3.165	1	34 6 0.9	20.03	1	B; pL; R; g; sbM disc	2
2621	1018	II. 162	11 48 16.5	3.086	2	77 15 16.9	20.03	2	pB; L; iR: bM; *10, 25°, 5'	4
2622	1018, a	R. nova	11 48 +	77 15 -	nf h. 1018	0
2623	1020	11 48 18.9	3.060	1	101 15 54.9	20.03	1	F; S; R; psbM; p of 2	1
2624	1019	II. 724	11 48 20.0	3.109	1	59 13 39.9	20.03	1	pF; vS; R; bM	2
2625	1021	11 48 32.4	3.060	1	101 12 44.9	20.03	1	vF; S; R; bM; f of 2	1
2626	1022	II. 132	11 48 45.9	3.079	3	82 28 4.2	20.04	3	B; pL; cE 30°; vsmbMN	4
2627	1023	II. 840	11 48 54.9	3.178	2	28 42 30.2	20.04	2	cF; S; IE; bM; *8, 90°, 6'	3
2628	III. 274	11 49 3.7	3.052	1	109 7 1.2	20.04	1	vF; pL; iF	1
2629	1024	III. 343	11 49 10.1	3.098	1	65 20 55.2	20.04	1	cF; cS; R; psbM	2
2630	1026	11 49 23.3	3.104	1	60 13 26.2	20.04	1	eF; S; R; bM	1
2631	1025	III. 707	11 49 23.7	3.136	1	40 52 57.2	20.04	1	vF; cS; another suspected	3
2632	1027	11 49 30.3	3.107	3	57 11 59.2	20.04	3	pF; S; pmE 90° ±; *11 nr	1
2633	1028	11 50 9.6	3.100	1	61 20 43.2	20.04	1	vF; S; R; bM*; p of 2	2
2634	1029	II. 791	11 50 14.3	3.147	1	33 46 31.2	20.04	1	pF; S; IE; psbM	2
2635	1030	IV. 61	11 50 15.6	3.142	1	35 50 34.2	20.04	1	cB; vL; pmE; sbMBrN	3
2636	1032	11 50 34.2	3.085	3	74 55 29.2	20.04	2	vF; pL; R; 2 st f.	3
2637	1031	I. 229	11 50 34.3	3.145	1	33 46 21.2	20.04	1	cB; pS; R; vg; smbM	2
2638
2639	1033, a	R. 3 novæ	11 50 ±	63 57 ±	one S, R; the other two E	0
2640
2641	1033	III. 323	11 50 36.1	3.095	1	63 57 34.2	20.04	1	pF; vS; E 25° bet 2 st	2
2642	1040, a	R. nova	11 50 44.7	3.121	::	41 54 13.2	20.04	::	S; R; 7' np h. 1040	0
2643	1034	III. 344	11 50 45.6	3.093	1	66 0 39.2	20.04	1	vF; vS; R; n of 2	2
2644	1035	III. 345	11 50 45.6	3.093	1	66 4 59.2	20.04	1	vF; vS; R; s of 2	2
2645	1036	III. 354?	11 50 52.9	3.097	2	61 20 22.2	20.04	2	F; vS; R; *12 near	3
2646	III. 324	11 50 53.4	3.095	1	63 53 51.2	20.04	1	eF; suspected	1
2647	1037	11 50 56.3	3.070	1	91 21 8.2	20.04	1	F; S; R; bM; *11 nf	1
2648	III. 325	11 50 57.6	3.087	1	66 6 2.2	20.04	1	eF; vS	1
2649	1038	II. 368	11 51 3.0	3.097	4	61 1 33.2	20.04	3	pB; pS; R; psbM; r	5
2650	1039	11 51 4.6	+3.097	1	61 10 ± ?	+20.04	0	vB; mE; mbM (?P.D.)	1*

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
2651	h. 1040	H.	h m s 11 51 14.4	s +3.121	1	41 59 10.2	+20.04	1	F; pL; mE; vglbM	1
2652	1041	II. 733	11 51 19.2	3.116	2	45 17 9.2	20.04	2	B; cL; mE 62°3; vsvmbM*10	3*+
2653	1042	11 51 24.9	3.086	2	73 2 32.2	20.04	2	pB; pS; R; psbM	2*
2654	1043, a	R. nova	11 51 33	3.095	:	61 39 50	20.04	:	vF	0
2655	1043	II. 369	11 51 33.0	3.095	(1)	61 44 50.2	20.04	2	F; L; E; gbF M	4
2656	1044	11 51 42.8	3.083	1	75 0 46.2	20.04	1	eF; *9 sf 5'	1
2657	1045	II. 725	11 51 43.9	3.097	1	58 47 57.5	20.05	1	pB; pL; E 19°5; biN	2
2658	II. 295	11 51 56.2	3.059	1	107 35 2.5	20.05	1	F; vS; iF; bM	1
2659	1046	III. 617	11 51 57.7	3.104	1	51 24 39.5	20.05	1	eF; pL; R	3
2660	1047	I. 223	11 52 6.8	3.122	1	38 15 44.5	20.05	1	vB; cL; mE 160°±; vsvmbMBN.	3
2661	3371	II. 296	11 52 23.1	3.059	1	108 29 25.5	20.05	1	⊕; pF; pL; R; rr; st 16	2
2662	III. 3	11 52 45.0	3.083	2	73 0 1.5	20.05	2	vF; vS; vL; r	2
2663	1048	I. 121	11 53 12.3	3.072	2	90 19 7.5	20.05	2	cB; L; vLE; psmbM; B st nr	4
2664	1049	II. 404	11 53 21.0	3.084	6	69 8 34.5	20.05	6	pF; pL; R; gbM; *12 nf...	7
2665	II. 508	11 53 21.4	3.062	1	107 3 2.5	20.05	1	pB; S; iE; bM	1
2666	III. 903	11 53 45.4	3.155	1	19 52 1.5	20.05	1	eF; S; iF; vglbM	1
2667	3372	III. 279	11 53 50.5	3.064	1	105 10 13.5	20.05	1	eF; pL; *945°±	2
2668	1050	I. 253	11 54 12.4	3.125	1	27 19 59.5	20.05	1	{ H. vb; vL; E } { h. pB; 25"; R }	2*
2669	1051	III. 77	11 54 13.1	3.079	2	75 49 13.5	20.05	2	eF; pL; R; r	3
2670	1052	IV. 28.1	11 54 43.7	3.064	1	108 5 11.5	20.05	1	pB; cL; R; vgbM	2+
2671	1053	IV. 28.2	11 54 43.7	3.064	1	108 7 11.5	20.05	1	pF; pL	2+
2672	1054	I. 252	11 54 58.6	3.117	2	27 5 10.5	20.05	2	B; cL; R; g, psvmbMrN	3
2673	1055	11 55 10.9	3.074	1	84 52 38.5	20.05	2	pF; S; R; psbM; * f 30"	2
2674	1056	III. 491	11 55 18.9	3.072	1	89 25 53.5	20.05	1	cF; cS; R; bM	3
2675	1057	II. 276	11 55 32.9	3.072	1	87 14 48.5	20.05	1	pF; L; R; gbM; *sf	4
2676	1058	II. 741	11 55 36.5	3.095	1	40 35 7.5	20.05	1	pB; pS; R	2
2677	1059	11 55 38.9	3.079	2	71 12 12.5	20.05	2	vF; vS; R; psbM	2
2678	1060	III. 390	11 55 41.8	3.079	1	70 28 21.5	20.05	3	eF; pS; R; glbM	4
2679	II. 509	11 55 47.7	3.066	1	105 36 2.5	20.05	1	F; cL; iR; lbM	1
2680	1061	IV. 56	11 55 58.3	3.089	4	44 41 3.5	20.05	4	B; vL; E; vg, vsmbM * 11..	5+
2681	3373	11 55 59.6	3.039	1	152 24 13.5	20.05	1	Cl; pRi; IC	1
2682	III. 794	11 56 8.6	3.099	1	31 18 2.5	20.05	1	eF; S	1
2683	1062	11 56 32.1	3.078	1	68 10 1.5	20.05	1?	pB. P.D. very doubtful	1*
2684	1063	11 56 36.6	3.078	1::	68 8 1.5	20.05	1?	pB. P.D. very doubtful	1*
2685	1064	11 56 40.6	3.078	1::	67 55 1.5	20.05	1?	pB. P.D. very doubtful	1*
2686	1065	11 56 51.4	3.077	3	68 59 39.5	20.05	3	vF; S; R; D neb pos 70°	3*
2687	1066	I. 174	11 56 53.2	3.080	4	57 19 32.5	20.05	4	pB; vL; mE 97°; vgbM	5
2688	D'Arrest, 85	11 56 57	3.07	[1]	70 46 42	20.06	[1]	B; E; gbM * 17 p, 82' dist.	0
2689	1067	11 56 57.5	3.077	2	68 59 58.5	20.05	2	pF; R	2*
2690	1068	11 56 59.7	3.076	4	68 51 54.5	20.05	4	pB	4*
2691	1069	III. 37	11 57 1.0	3.074	5	78 21 53.5	20.05	5	F; pS; R; gbM	6
2692	II. 781	11 57 1.6	3.087	2	36 40 2.5	20.05	2	pF; S; stellar	2
2693	1070	III. 392	11 57 1.7	3.076	1	68 53 38.5	20.05	1	vF; vS	2*
2694	1071	III. 391	11 57 2.0	3.076	2	68 49 13.5	20.05	2	F; vS	2*
2695	3374	11 57 4.9	3.046	1	156 31 32.5	20.05	1	vF; vS; R; bM*; am st	1
2696	1072	II. 277	11 57 17.2	3.072	4	87 19 26.5	20.05	4	F; pS; R; pgbM; np of 2	7
2697	1073	III. 393	11 57 20.8	3.076	1	68 54 1.5	20.05	1	eF; vS	2*
2698	1074	11 57 20.9	3.072	1	87 9 8.5	20.05	1	F; S; R	1
2699	1075	III. 394	11 57 24.7	3.076	1	69 3 31.8	20.06	1	vF; vS	2*
2700	1076	III. 258	11 57 29.5	3.072	1	87 26 4.8	20.06	1	cF; cS; vLE; bM; sf of 2	3
2701	III. 395	11 57 32.3	3.075	2?	69 7 1.8	20.06	2?	vF; vS	1*
2702	III. 396	11 57 32.3	3.075	2?	69 7 1.8	20.06	2?	vF; vS	1*
2703	1077	11 57 38.1	3.072	1	91 36 8.8	20.06	1	F; L; R; * 10 n 60"	1
2704	1078	III. 355	11 57 42.0	3.076	4	62 13 27.8	20.06	4	cF; pS; E; gbM	5
2705	D'Arrest, 86	11 58 13	3.08	[1]	38 54 7	20.06	[1]	F; iE; I. 206 nr	0
2706	3375	III. 754	11 58 24.4	3.070	1	115 44 43.8	20.06	1	pB; S; R; bM	2
2707	I. 224	11 58 40.3	3.074	2	38 56 2.8	20.06	2	B; pL; pmE; vsbM	2
2708	I. 206	11 58 42.2	3.074	3	38 42 22.8	20.06	3	B; cL; pmE 135°±; lbM	3
2709	3376	11 58 44.2	+3.072	1	103 45 28.8	+20.06	1	eF; L; pmE; vgbM; 2 st 11 nr	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
2710	h. 1079	H. III. 382	h m s 11 58 45.3	s +3.072	1	68° 38' 1.8"	+20.06	2	vF; vS.....	4
2711	1081	I. 207	11 58 51.9	3.072	1	41 44 34.8	20.06	1	pB; vL; mE 32° 0'	4
2712	1080	III. 400	11 58 53.1	3.072	2	52 21 4.8	20.06	2	eF; vS; R; stellar; *10 sp 2'	6
2713	1082	III. 383	11 58 54.0	3.072	2	68 36 5.8	20.06	3	eF; eS; R; bM	5
2714	III. 384	11 58 57.0	3.072	1	68 35 2.8	20.06	1	eF; eS	2
2715	1084	III. 717	11 58 59.8	3.072	1	39 38 47.8	20.06	1	pB; vL; vmE 166° 5'; vglbM	3
2716	1083	III. 326	11 59 0.1	3.072	2	63 39 44.8	20.06	2	eF; vS; R; vgbM	3
2717	1085	I. 225	11 59 15.4	3.070	1	36 30 34.8	20.06	1	B; pS; R; bMBrN; *12 sp, v, nr	3
2718	3377	Δ. 291	11 59 28.0	3.077	3	150 27 56.8	20.06	3	Cl; pL; pC; iR; st 10...14...	3
2719	1086	II. 370	11 59 28.8	3.070	6	61 2 37.8	20.06	6	pB; pS; lE; bM	7
2720	3378	II. 865	11 59 29.4	3.074	2	119 0 26.8	20.06	2	pF; pS; R; psbM; r; p of 2	3
2721	3379	II. 866	11 59 34.4	3.074	2	119 0 41.8	20.06	2	pF; pS; R; pgbM; f of 2	3
2722	1087	11 59 39.8	3.062	1	22 3 42.8	20.06	1	B; S; R; gbM	1
2723	1088	I. 195	11 59 52.9	3.067	2	46 9 26.8	20.06	2	vB; pS; mE 151° 0'	4
2724	3380	11 59 55.8	3.077	2	129 25 18.8	20.06	2	F; S; vIE; glbM; 3Bst nr	2
2725	1089	12 0 0.5	3.068	1	55 13 46.8	20.06	1	eF	1
2726	3381	III. 533	12 0 0.9	3.074	1	102 24 23.8	20.06	1	eF; S; iR; gbM	2
2727	1090	12 0 1.7	3.070	1	74 49 13.8	20.06	1	eF; suspected	1
2728	1092, a	R. nova	12 0 19.8	3.072	...	86 30 33.5	20.06	...	Hook-shaped; h. 1092 is nf 45°; 14' dist.	0
2729	1091	III. 708	12 0 26.9	3.064	1::	46 12 1.8	20.06	1::	vF; vS	2
2730	II. 14	12 0 42.1	3.070	1	79 41 2.5	20.05	1	lE	1*
2731	III. 904	12 0 56.3	3.040	1	19 38 2.5	20.05	1	eF; vS; E	1
2732	1093	12 0 59.2	3.064	1	56 13 7.5	20.05	1	eF; vS; R; mbM	1
2733	1092	V. 4	12 0 59.4	3.072	2	86 20 39.5	20.05	2	eF; vL; E 90° ±; bM *16...	4+
2734	1094	{ I. 33 = II. 60 }	12 1 1.2	3.070	2	78 50 36.5	20.05	3	pB; pL; mE 120°; bM; r ...	5
2735	Auw. N. 28	12 1 3.1	3.045	...	24 2 50.6	20.05	...	pB; pL; cE; mbMN (Hind, Jan. 5, 1850).	0
2736	1095	III. 68	12 1 28.2	3.067	1	73 5 11.5	20.05	1	vF; S; R; psbM; bet 2 vS st	2
2737	1096	I. 279	12 1 29.2	3.005	2	12 25 10.5	20.05	2	F; pL; vIE; glbM	4
2738	I. 263	12 1 33.2	3.032	1	20 25 2.5	20.05	1	cB; lE; bM	1
2739	{ 1097 = 3382 }	II. 548	12 1 42.3	3.074	3	98 15 19.5	20.05	3	F; pL; pmE 95° ±; vglbM	4
2740	1098	III. 356	12 1 47.9	3.063	2	59 56 7.5	20.05	2	eF; S; R; 1st of 3	3
2741	1099	III. 357	12 1 52.9	3.063	3	59 58 57.5	20.05	3	eF; S; iR; 2nd of 3	4
2742	1100	I. 278	12 1 59.4	3.003	2	14 19 20.5	20.05	2	pB; cL; R; gmbM	3
2743	1101	II. 371	12 2 1.4	3.062	2	60 3 7.5	20.05	2	pF; pL; lE; 3rd of 3	4
2744	1108	II. 321	12 2 6.7	3.061	1	59 18 5.5	20.05	1	F; vL; vgbmB	3
2745	I. 196	12 2 23.2	3.053	2	45 32 32.5	20.05	2	B; pL; lE; vgbM; * np	2
2746	1102	III. 795	12 2 29.8	3.037	1	30 22 0.5	20.05	1	vF; pS; lE; gbM; r	3
2747	1103	III. 814	12 2.31.4	3.044	1	36 6 6.5	20.05	1	vF; S; iF; vglbM; er	2*
2748	1104	IV. 54	12 2 43.5	3.051	1::	46 44 3.5	20.05	1	cB; R; vg; vsbMN	2
2749	1107	II. 747	12 2 54.1	3.047	2	42 46 19.5	20.05	2	pF; cL; vmE 109° 0'; vgbM.	3
2750	1105	I. 169	12 2 54.5	3.053	1	49 20 37.5	20.05	1	B; vL; vglbM	2
2751	1106, a	R. nova	12 2 ±	70 40 ±	S; prec h. 1106	0
2752	1106	I. 19	12 2 55.5	3.064	3	70 40 17.5	20.05	3	⊕; vB; pL; R; gbM; rrr	4
2753	III. 327	12 2 56.8	3.060	1	62 48 2.5	20.05	1	vF; pS	1
2754	1109	II. 802	12 3 22.6	3.030	1	30 56 13.5	20.05	1	F; S; E	2
2755	1110	I. 73	12 3 26.7	3.057	1	58 49 5.5	20.05	1	B; S; R; pgmbM	3
2756	1111	I. 165	12 3 27.0	3.050	2	49 49 31.5	20.05	2	vB; S; R; vsmbMBN; p of 2	4+
2757	1112	II. 83	12 3 28.7	3.064	3	73 11 28.5	20.05	3	pB; pL; R; pgmbM; r	4
2758	I. 11	12 3 36.9	3.063	1	70 52 2.5	20.05	1	B; pL; E; bM	1
2759	III. 845	12 3 42.6	3.026	1	30 53 2.5	20.05	1	vF; S; E 90° ±	1
2760	1113	II. 642	12 3 46.5	3.049	2:	49 45 45.5	20.05	2:	pF; S; E; vgbM; f of 2	4+
2761	1114	I. 208	12 4 0.9	3.036	1	38 43 54.5	20.05	1	pF; cL; vmE 60° ±	4
2762	1115	II. 405	12 4 4.2	3.061	1	69 2 58.5	20.05	1	F; pS; lE; bM; pB* nf	3
2763	1116	III. 941	12 4 4.7	2.945	1	13 5 48.5	20.05	1	eF; pS; R; Δ 2 st	2
2764	II. 803	12 4 39.4	+3.018	2	31 27 2.5	+20.05	2	F; S; R	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	+ 20.05			
2765	1117	II. 353	12 4 46.9	+3.057	1	65 5 49.5	+20.05	1	B; L; iE; bM.....	2
2766	III. 399	12 4 56.4	3.046	2	53 2 2.5	20.05	2	vF; pL; vIE; er.....	2
2767	1118	12 5 4.5	3.045	1	52 43 3.5	20.05	1	F; pL; R; vgbM; * sp 10'...	1
2768	1119	II. 105	12 5 8.6	3.063	1	76 1 5.5	20.05	1	pB; pL; iF; psbM; r; * inv	3
2769	1120	III. 358	12 5 14.5	3.051	2	60 3 17.5	20.05	2	F; S; 1st of 4.....	4
2770	1123	II. 792	12 5 16.6	3.016	1	33 2 58.5	20.05	1	F; S; iE; gbM.....	2
2771	1121	II. 372	12 5 17.2	3.051	2	60 0 1.5	20.05	1	F; S; 2nd of 4.....	4*
2772	1122	III. 359	12 5 18.7	3.051	2	60 5 31.5	20.05	2	F; S; 3rd of 4.....	4
2773	1124	III. 360	12 5 27.2	3.050	2	60 3 31.5	20.05	2	F; eS; 4th of 4.....	4*
2774	3383	III. 534	12 5 30.9	3.081	1	103 14 38.5	20.05	1	vF; pL; R; vgbM.....	2
2775	1125	12 5 35.2	2.993	1	78 22 7.5	20.05	2	vF; vL; E 45°±; *7 f.....	2
2776	1126	I. 9	12 5 42.9	3.070	1	87 55 28.5	20.05	1	pB; pS; pmE 135°±; bMN	5
2777	1127	II. 133	12 5 53.0	3.067	3	82 11 3.5	20.05	3	pF; S; iE 0°±; r.....	5
2778	III. 777	12 6 4.8	3.016	1	36 20 2.5	20.05	1	eF; S; stellar.....	1
2779	1128	III. 697	12 6 13.1	3.031	2	45 32 36.5	20.05	2	vF; cL; mE 170°±.....	5
2780	3384	12 6 14.4	3.152	1	151 56 12.2	20.04	1	Cl; mC; st eS.....	1
2781	1129	II. 373	12 6 15.1	3.048	3	60 42 57.2	20.04	3	cF; L; R; gbM.....	5
2782	II. 813	12 6 28.8	3.018	1	38 30 2.2	20.04	1	pB; S; iE.....	1
2783	1131	II. 106	12 6 38.7	3.061	1	75 48 11.2	20.04	1	F; L; iE; vglbM; r.....	3
2784	1133	II. 409	12 6 39.4	3.038	1	52 35 28.2	20.04	1	cF; pS; R; vglbM; r.....	3
2785	1130	12 6 39.6	3.066	4	82 1 3.2	20.04	4	cF; R; bM; near S*.....	4
2786	1132	M. 98	12 6 40.0	3.060	3	74 19 1.2	20.04	4	B; vL; vmE 152°1; vsymbM	7
2787	1134	II. 163	12 6 44.6	3.061	1	76 3 5.2	20.04	1	vF; pL; E; vgbM.....	2
2788	1135	II. 867	12 7 6.6	3.004	1	34 40 46.2	20.04	1	pB; vS; vsbM *12.....	2
2789	III. 796	12 7 8.8	2.988	1	29 34 2.2	20.04	1	eF.....	1
2790	1136	II. 374	12 7 26.4	3.045	4	60 48 15.2	20.04	4	pB; S; R; vsmbM *.....	5
2791	1137	II. 134	12 7 28.5	3.066	1	83 24 54.2	20.04	1	pF; pmE; vgbM.....	2
2792	1139	II. 793	12 7 29.7	2.997	1	33 12 18.2	20.04	1	pF; pS; iE; gbM.....	3
2793	III. 797	12 7 32.0	2.983	2	29 16 2.2	20.04	2	vF; S.....	2
2794	1138	II. 164	12 7 32.7	3.061	2	77 3 13.2	20.04	2	cF; { H. vmE h. R, 2 obs. } lbM ...	3
2795	II. 165	12 7 44.7	3.060	1	76 4 2.2	20.04	1	F; vmE.....	1
2796	1140	I. 175	12 8 1.5	3.037	2	56 1 26.2	20.04	2	vB; S; R; psmbM.....	3
2797	1141	III. 397	12 8 8.0	3.051	1	68 33 36.2	20.04	2	vF; cL; iR; vgbM.....	4
2798	D'Arrest, 87	12 8 9	2.96	[1]	25 25 42	20.04	[1]	pB; pS; R; *12 f; ln.....	0
2799	1142	II. 107	12 8 19.4	3.058	1	75 19 25.2	20.04	1	vF; pL; R; gbM.....	2
2800	II. 375	12 8 21.7	3.041	1	60 43 22.2	20.04	1	F; pS.....	1
2801	1143	III. 850	12 8 23.4	2.945	1	23 14 29.2	20.04	1	pF; pS; R; vgbM.....	2
2802	1144	II. 108	12 8 31.3	3.057	2	75 19 1.2	20.04	2	B; L; E 90°±; g; sbM; r...	3
2803	1145	II. 354	12 8 31.6	3.047	1	65 13 59.2	20.04	1	cF; vS; R.....	2
2804	1146	I. 95	12 8 36.0	3.030	1	52 54 3.2	20.04	1	cB; cL; iE; biN.....	3+
2805	1147	II. 135	12 8 44.8	3.065	3	82 49 9.2	20.04	3	B; pS; E; sbM *11.....	4
2806	1148	I. 35	12 8 46.7	3.058	2	76 4 2.2	20.04	3	vB; vL; vmE 17°±; sbMN	5+
2807	1149	II. 748	12 8 49.1	3.009	4	42 8 55.2	20.04	5	pF; L; mE 45°0; * n, p of 2	6+
2808	III. 718	12 8 55.0	3.006	1	41 5 2.2	20.04	1	vF; vS.....	1
2809	3385	12 9 7.3	3.126	4	132 32 44.2	20.04	4	pF; pL; pmE; vglbM.....	4
2810	1150	12 9 9.1	2.931	1	22 59 24.2	20.04	1	pB; S; R; psbM.....	1
2811	1151	I. 209	12 9 12.9	3.005	1	41 20 34.2	20.04	1	cB; pL; pmE 134°4; psbM..	3
2812	1152	II. 137	12 9 17.5	3.065	1	82 31 31.9	20.03	1	pF; pL; R; r (? R.A. 10 ^m)...	3
2813	1153	II. 136	12 9 21.7	3.063	2	81 45 38.9	20.03	2	pB; pS; iE; gb, not M; r ...	4
2814	II. 109	12 9 23.0	3.057	1	76 8 1.9	20.03	1	r.....	1*
2815	1154	12 9 25.1	3.084	1	101 31 49.9	20.03	1	F; eS; R; *170°, 60''.....	1
2816	1155	12 9 28.9	3.005	1	42 12 32.9	20.03	1	F; S; iE; f of 2.....	1
2817	1156	II. 518	12 9 31.4	3.030	2	55 42 6.9	20.03	2	F; vS; vIE; psbM; sp of 2...	4
2818	1157	12 9 32.3	3.026	1	52 53 41.9	20.03	1	vF; L; R; gbM.....	1
2819	1158	II. 519	12 9 36.9	3.030	2	55 39 37.9	20.03	2	cF; vS; iE; psbM; nf of 2...	4
2820	3386	12 9 41.0	3.160	1	144 31 20.9	20.03	1	Cl; F; pL; iF; st 13...15 ...	1
2821	1159	II. 17	12 9 58.6	3.063	2	82 1 40.9	20.03	2	pB; pL; pmE; iM; p of 2 ...	5
2822	1160	12 9 59.5	3.067	1	85 32 30.9	20.03	1	pB; L; R; gbM.....	1
2823	1161	II. 496	12 9 59.5	3.063	1	81 35 59.9	20.03	1	pF; R; vsbMSN.....	2
2824	1162	II. 11	12 10 3.8	+3.054	2	73 54 23.9	+20.03	2	pB; pL; iE; vgbM; r.....	5

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1830, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
2825	h. 1163	H. V. 51	h m s 12 10 4.1	s +2.894	1	19 48 0.9	+20.03	1	vF; eL; mE 160°±; vgbM..	3
2826	1164	III. 851	12 10 5.7	2.939	1	25 48 56.9	20.03	1	vF; pS; iR; vglbM.....	2
2827	III. 719	12 10 13.6	2.999	1	41 44 31.9	20.03	1	vF; vS; n of D neb	1
2828	III. 720	12 10 13.6	2.999	1	41 45 31.9	20.03	1	vF; vS; s of D neb.....	1
2829	1165	III. 480	12 10 17.5	3.063	2	82 32 50.9	20.03	2	vF; L; vgbM; *7 s	3
2830	1166	III. 725	12 10 20.5	3.003	2	43 35 49.9	20.03	2	vF; cL; iR; vgbM; r.....	4
2831	1167	V. 41	12 10 27.9	3.019	2	51 24 39.9	20.03	2	pB; vL; eE 43°±; vgbM ...	3
2832	1168	I. 74	12 10 32.5	3.033	3	59 36 54.9	20.03	3	cB; pL; vIE; smbM; r	5
2833	III. 91	12 10 43.7	3.063	1	82 6 1.9	20.03	1	eF	1
2834	1169	II. 742	12 10 48.8	2.995	2	41 49 7.9	20.03	2	vF; S; pmE; psbM.....	4
2835	1170	I. 264	12 10 57.2	2.864	1	18 25 17.9	20.03	1	pB; S; R; pgbM	2
2836	1171	I. 89	12 11 3.5	3.034	2	61 2 42.9	20.03	3	vB; S; E; vsvmbMN; *6.7 f 90°.	4
2837	1172	III. 702	12 11 30.9	3.029	1::	59 23 0.9	20.03	1::	vF; vS; R	2
2838	1173	M. 99	12 11 41.8	3.052	3	74 48 7.6	20.02	4	!!; {(H. h.)B; L; R; gbm; r} {(L) 3-branched spiral}	6+
2839	1174	II. 846	12 11 56.0	2.898	1	23 19 23.6	20.02	1	pB; L; cE 38°±; bMBN ...	2
2840	D'Arrest, 88	12 12 1	3.06	[1]	83 29 42	20.02	[1]	vF pS; R; *18 s 2'	0
2841	1175	V. 43	12 12 1.7	2.988	3	41 55 40.6	20.02	4	vB; vL; vmE 0°; sbMBN...	8+
2842	1176	II. 139	12 12 4.6	3.063	1	83 23 51.6	20.02	1	F; pS; R; gbm	4
2843	1177	II. 138	12 12 12.9	3.063	2	83 9 22.6	20.02	2	pB; E; psbM	5
2844	1178	12 12 13.0	3.064	1	83 53 0.6	20.02	1	neb; "1st of 5"	1
2845	1179	II. 110	12 12 22.0	3.051	1	74 20 51.6	20.02	1	B; S; R; r	3
2846	III. 535	12 12 26.0	3.090	1	101 28 31.6	20.02	1	vF; pL; iF	1*
2847	1180	II. 140	12 12 26.3	3.063	1	83 22 30.6	20.02	1	F; pS; R; gbm	4
2848	1181	II. 166	12 12 38.0	3.053	2	76 26 0.6	20.02	2	pB; vS; R; vsmbM.....	3
2849	D'Arrest, 89	12 12 41	3.06	[2]	83 12 42	20.02	[2]	pF; S; R; *9 f 18.7, n 85°...	0*
5070	12 12 41	83 46 42	See No. 5070	0
2850	1182	III. 299	12 12 45.0	3.024	1	58 52 49.6	20.02	1	cF; S; iR; gmbM	3
2851	1185	I. 75	12 12 47.4	3.025	1	59 36 15.6	20.02	1	vB; vL; E 90°±; mbMN ...	3
2852	1183	II. 568?	12 12 48.2	3.064	1	83 53 1.6	20.02	1	B; L; E; gbm	1*
2853	II. 804	12 12 48.3	2.946	1	32 29 1.6	20.02	1	pB; pL; iF	1
2854	1184	II. 376	12 12 49.1	3.029	1	61 36 0.6	20.02	2	F; S; vIE; gbm; *15 nr ...	3
2855	1186	I. 90	12 13 3.4	3.025	2	59 56 25.6	20.02	2	vB; pL; R; mbM; r; p of 2	5*
2856	II. 322	12 13 8.7	3.062	1	82 51 1.6	20.02	1	4 neb sc about. Place of the last (see note).	1*
2857	1187	II. 571??	12 13 12.5	3.063	1	83 50 34.6	20.02	1	vB; vL; R; pgbM; "3 more seen."	3*
2858	1188	II. 569?	12 13 18.6	3.024	2	59 54 20.6	20.02	2	B; S; R; bM; 2nd of 3	4
2859	II. 323	12 13 23.0	3.025	1	60 0 39.6	20.02	1	1*
2860	II. 377	12 13 25.7	2.933	1	31 6 1.6	20.02	1	cF; lE; p of 2.....	1
2861	III. 798	12 13 35.7	3.023	2	59 51 31.3	20.01	2	vF	2
2862	1189	III. 300	12 13 35.9	3.063	1::	83 54 0.6	20.02	1::	vF; S	1*
2863	1189	II. 570?	12 13 35.9	3.063	1::	83 54 0.6	20.02	1::	vF; S	1*
2863	1188, a	R. nova?	12 13 38.6	3.024	...	59 51 20.6	20.02	...	Most probably = H. III. 300	0?
2864	1191	III. 726	12 13 41.0	2.980	3	42 56 17.3	20.01	3	vF; pS; R; vgbM; r	5
2865	1190	II. 571?	12 13 46.0	3.063	1	83 52 34.3	20.01	1	vB; R; central of 4.....	1*
2866	1193	II. 805	12 13 48.1	2.930	1	31 7 37.3	20.01	1	pB; L; R; gbm	2
2867	1195	V. 5	12 14 4.2	3.042	2	70 50 17.3	20.01	2	F; vL; E; lbM; r	3
2868	1192	I. 275	12 14 4.8	2.720	3	13 51 7.3	20.01	4	pB; vS; R; lbM; 3 st f	6
2869	1194	12 14 6.1	3.063	2	83 49 57.3	20.01	2	B; pL; lE; bM; 4th of 4.....	3*
2870	1196	12 14 6.1	3.064	1	84 38 4.3	20.01	2	F; S; R; vglbM; B*340°, 60"	2+
2871	1197	II. 61	12 14 8.4	3.053	4	77 42 56.3	20.01	4	F; L; mE 135°±; bi-N; p of 2	6
2872	III. 92	12 14 23.6	3.061	2	82 34 1.3	20.01	2	vF; vS	2
2873	III. 93	12 14 23.6	3.061	2	82 34 1.3	20.01	2	eF; eS	2
2874	1198	II. 111	12 14 24.9	3.048	2	74 36 52.3	20.01	2	F; L; E 0°±; vgbM, p of 2	4
2875	1200	II. 62	12 14 32.7	3.053	2	77 43 1.3	20.01	2	F; L; lE; vgbM; f of 2	4
2876	1201	II. 572	12 14 33.1	3.063	1	83 50 23.3	20.01	1	F; lE; vgbM	2
2877	1199	II. 112	12 14 33.2	+3.047	2	74 36 53.3	+20.01	2	L; vmE 0°±; f of 2	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
2878	h. 1202	H. I. 139	M. 61	12 14 45.1	+3.064	2	84 44 55.3	+20.01	2	vB; vL; vsbM*; biN.....	5*†
2879	3387	12 14 51.8	3.132	1	122 41 54.3	20.01	1	vF; vL; R; vglbM; r	1
2880	1203	12 14 56.7	3.050	1	76 29 35.3	20.01	1	vF; R	1
2881	1204	I. 76	12 15 23.5	3.016	1	59 19 46.0	20.00	1	cB; L; E 150°±; sbM; * np	3
2882	1205	II. 378	12 15 24.5	3.017	1::	60 0 34.0	20.00	2	B; cL; lE; np of 2	3
2883	1206	12 15 24.5	3.017	1	60 1 8.0	20.00	1	F; sf of 2	1
2884	1202, a	R. nova?	12 15 30	3.06	...	84 35 40	20.00	...	F; E; 10' nf h. 1202	0†*
2885	1207	II. 63	12 15 30.2	3.051	2	77 26 25.0	20.00	2	vF; L; E 135°±; r.....	4
2886	1209	II. 628	12 15 31.6	3.044	(1)	73 40 57.0	20.00	1	pB; cL; E; gbM.....	2
2887	II. 324	12 15 34.9	3.011	1	58 11 0.0	20.00	1	F; S.....	1
2888	1210	I. 276	12 15 36.3	2.686	3	13 53 59.0	20.00	3	pB; pS; vlE; sbM	5
2889	1208	12 15 37.4	3.057	1	81 1 20.0	20.00	1	eF; *8 n 5'	1
2890	1211	M. 100	12 15 50.6	3.043	3	73 23 54.0	20.00	4	!!; { (H, h) pF; vL; R; } vg, psbMrN (L) 2-branched spiral	5
2891	1212	II. 85	12 16 1.2	3.041	1	72 30 5.0	20.00	1	pB; S; R; psbM	2
2892	D'Arrest, 90	12 16 2	3.06	[4]	83 58 42	19.99	[4]	pB; R or lE; bM	0*
2893	1213	II. 141	12 16 2.0	3.060	1	83 8 51.0	20.00	1	vF; S; R; bM; 1st of 3	2
2894	II. 84	12 16 4.6	3.043	1	73 25 0.0	20.00	1	F; S; R; r	1
2895	1216	II. 847	12 16 6.6	2.843	1	23 22 43.0	20.00	1	pF; S; vlE; vgbM	2
2896	1214	12 16 8.6	3.093	1	101 45 29.0	20.00	1	vF; vS; R; bMN.....	1
2897	1217	II. 806	12 16 8.9	2.905	2	30 47 2.0	20.00	2	pB; S; E; gbM	3
2898	1220	III. 942	12 16 11.2	2.637	1	13 3 7.0	20.00	1	eF; E 0°±	2
2899	1215	II. 142	12 16 12.5	3.060	1	83 11 1.0	20.00	1	F; pS; R; bM; 2nd of 3.....	2
2900	1218	12 16 16.0	3.058	1	81 45 7.0	20.00	1	pF; S; R; * v nr.....	1
2901	1219	II. 406	12 16 21.1	3.035	2	69 48 9.0	20.00	2	vF; pL; iR; biN?	4
2902	3388	12 16 23.3	3.231	1	147 20 30.0	20.00	1	Cl; pRi; lC; st 12...14	1
2903	1221	II. 86	12 16 23.7	3.040	1	72 31 5.0	20.00	1	cB; vS; mE; vsbM.....	2
2904	1222	II. 143	12 16 26.6	3.059	3	83 8 10.0	20.00	3	B; pL; R; bM; 3rd of 3.....	4
2905	III. 95	12 16 29.8	3.058	2	82 14 31.0	20.00	2	eF; vS; R	2
2906	III. 96	12 16 29.8	3.058	2	82 14 31.0	20.00	2	eF; vS; R	2
2907	1223	III. 94	12 16 30.1	3.060	1	82 16 59.0	20.00	1	pF; S; E; ? D	3
2908	1224	III. 31	12 16 32.7	3.038	1	71 41 1.0	20.00	1	vF; pS; R; vglbM; Δ 2 st ...	2
2909	1200, a	R. nova	12 16 34.7	3.053	::	77 43 1.3	20.00	::	vF; vmE	0
2910	1225	I. 210	12 16 34.8	2.960	5	42 14 8.0	20.00	6	vF; S; mE 100°±; vsmbMBN	8†
2911	1226	II. 625	12 16 43.1	3.077	1	92 40 18.0	20.00	1	F; pL; E 70°±; vlbM	4
2912	3389	Δ. 292	12 16 49.2	3.262	3	151 7 11.0	20.00	3	Cl; vB; vL; lC; st 12...14...	3
2913	III. 481	12 17 3.3	3.055	1	80 42 0.7	19.99	1	vF	1
2914	1230	III. 799	12 17 9.0	2.896	1	30 50 31.7	19.99	1	cF; cS; lE (? 18 th R.A.).....	2
2915	1228	I. 123	12 17 9.3	3.061	1	84 17 31.7	19.99	1	B; S; * 8.9 sf 3'	3
2916	1229	III. 648	12 17 10.3	3.006	1	57 42 34.7	19.99	1	cF; pmE 90°; vlbM	2
2917	1231	I. 65	12 17 16.6	3.107	1	108 0 3.7	19.99	1	vB; L; R; vsmbMn; r	2
2918	1233	III. 800	12 17 17.0	2.895	1	30 51 31.7	19.99	1	vF; cS; R; r	2
2919	III. 938	12 17 17.6	2.659	1	14 17 0.7	19.99	1	eF; pL; iF	1
2920	III. 801	12 17 18.5	2.894	1	30 48 59.7	19.99	1	cF; cS; R	1
2921	1232	I. 30	12 17 21.3	3.057	2	81 54 17.7	19.99	3	cB; pL; vlE; gl, smbM	5
2922	III. 97	12 17 23.3	3.057	1	81 50 0.7	19.99	...	eF	2
2923	III. 38	12 17 35.0	3.050	1	78 37 59.7	19.99	1	vF; vS	1
2924	1234	I. 166	12 17 40.3	2.981	2	49 51 4.7	19.99	2	cB; S; R; mbMN; r	4
2925	1235	I. 22	12 17 45.8	3.048	3	77 31 34.7	19.99	3	B; pS; R; gbM	6
2926	1236	II. 144	12 17 46.2	3.057	2	81 46 52.7	19.99	2	pF; pS; lE; bM	3
2927	3390	Δ. 67??	12 17 50.1	3.412	1	161 53 19.7	19.99	1	⊕; pF; L; R; st 12...16 ...	2
2928	3391	12 17 55.9	3.162	1	128 58 12.7	19.99	1	pB; S; R; pgvmbM	1
2929	1227	II. 64	12 17 56.9	3.050	1	77 59 51.7	19.99	2	cF; cS; lE	3
2930	1237	M. 84	12 17 57.6	3.045	1	76 20 8.7	19.99	1	vB; pL; R; psbM; r	2
2931	1238	II. 379	12 17 59.0	2.994	2	60 40 3.7	19.99	2	F; S; R; bM; * nf 90'	3
2932	} 1237, a	R. 9 novæ	12 18 ±	+3.045	...	76 20 ±	+19.99	...	"Twelve knots exam." (see h. 1237, 1244, 1250).	0
2933											
2934											
2935											
2936											

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.		
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.										
2937	h.	H.		h m s	s		° ' "	"					
2938	1237, a	R. 9 novæ (continued)	12 18 ±	+3.045	...	76 20 ±	+19.99	...	"Twelve knots exam." (see h. 1237, 1244, 1250).	0		
2939													
2940													
2941													
2942	II. 530	12 18 2.6	3.060	1	83 29 59.7	19.99	1	F; S.....	1		
2943	1239	I. 12	12 18 7.3	3.041	1	74 27 43.7	19.99	1	B; S; R; smbM	4		
2944	1240	II. 87??	12 18 13.0	3.039	1	73 35 30.7	19.99	1	pS; R; psbMN	2		
2945	1241	12 18 17.0	3.051	1	79 13 0.4	19.98	1	vF; pL; R; lbM	1		
2946	II. 743	12 18 19.0	2.940	1	40 23 59.4	19.98	1	F; S.....	1		
2947	1242	M. 85	12 18 19.5	3.033	2	71 2 10.4	19.98	2	vB; pL; R; bM; * np	3		
2948	1243	III. 879	12 18 22.3	2.910	2	34 43 1.4	19.98	2	cF; S; iR.....	3		
2949	1244	I. 277	12 18 36.7	2.609	2	13 42 3.4	19.98	2	B; cL; lC; psmbM	4		
2950	1245	12 18 41.2	3.045	1	76 34 33.4	19.98	1	vF; E; p of 2	1		
2951	II. 749	12 18 41.2	2.950	3	43 32 16.4	19.98	3	pB; pL; iE; vglbM	5+		
2952	II. 87	12 18 43.3	3.039	1	73 41 59.4	19.98	1	S; bM; r	1*		
2953	1248	III. 852	12 18 44.3	2.819	2	24 17 14.4	19.98	2	cF; S; R; sbM; ** sp	3		
2954	1249	III. 729	12 18 48.4	2.951	1	43 25 20.4	19.98	1	cF; S; R; vgbM	2		
2955	1246	III. 361	12 18 48.5	3.010	1:	61 39 49.4	19.98	1	vF; vL; iF; B*p	2		
2956	1250	II. 167	12 18 51.3	3.044	1::	76 29 1.4	19.98	1	Northern of 2; no description	2		
2957	1251	II. 168	12 18 51.3	3.044	1::	76 29 1.4	19.98	1	Southern of 2; E	2		
2958	1252	II. 55	12 18 52.0	3.032	2	71 0 52.4	19.98	2	pB; lE; bM	3		
2959	V. 29.1	12 18 52.0	2.993	1	55 40 35.4	19.98	1	eF; vL; np of D neb	3+		
2960	1252, a	R. 2 novæ	12 18 ±	2.993	...	55 40 ±	19.98	0		
2961				M. 86	12 19 0.2	3.044	1	76 17 9.4	19.98	1	vB; L; R; gbMN; r	5*
2962				V. 29.2	12 19 1.1	2.993	2	55 42 28.4	19.98	2	vF; vL; pvlbM; sf of D neb	3+
2963				III. 755	12 19 2.5	3.086	1	96 54 29.4	19.98	1	vF; vS; E	1
2964	III. 756	12 19 2.5	3.086	1	96 54 29.4	19.98	1	vF; vS; E	1			
2965	Auw. N. 30	12 19 2.6	3.041	A	76 6 32.4	19.98	A	F; L; mE 90° (Auwers, Mar. 5, 1862).	0		
2966	1254	II. 88	12 19 3.4	3.037	1	73 3 10.4	19.98	1	pF; S; R; vsbM; r	2		
2967	III. 39	12 19 5.0	3.050	1	78 42 59.7	19.98	1	vF; B* nr	1		
2968	1255	12 19 11.1	3.044	1	76 34 59.4	19.98	1::	f of 2 neb	1		
2969	1256	12 19 14.6	3.052	1	80 12 44.4	19.98	1	eF; vL; R; gbM	1		
2970	III. 17	12 19 17.0	3.065	1	86 43 59.4	19.98	1	vF; pS; r	1		
2971	1257	II. 34	12 19 26.2	3.063	2	85 15 53.4	19.98	2	F; pL; R; gbM; r	6		
2972	1258	I. 77	12 19 29.4	2.997	1	57 30 19.4	19.98	2	vB; L; E; g; vsmbM*	5+		
2973	III. 482	12 19 30.1	3.053	1	80 47 0.4	19.98	1	eF	1		
2974	1259	II. 169	12 19 32.4	3.044	1	76 39 31.4	19.98	1	cF; S; gbM; 2 st n, np	2		
2975	1260	12 19 37.2	3.054	1	81 18 20.1	19.97	1	vF; L; R; * sp 5'	1		
2976	1261	III. 492	12 19 44.6	3.072	1	90 6 39.1	19.97	2	{ H. vF; cL; mE } { h. F; S; R; * nr }	3*		
2977	1262	II. 113	12 19 49.3	3.038	1	74 10 51.1	19.97	1	B; pmE 135°±; sbM	3		
2978	1263	II. 23	12 19 50.7	3.065	2	86 43 44.1	19.97	2	F; pL; lE; r; (?=III. 17)...	4		
2979	II. 155	12 19 56.1	3.052	1	79 37 0.7	19.97	1	F; pL; E; lbp	1		
2980	1265	III. 114	12 19 59.7	3.083	2	95 3 10.1	19.97	2	F; vS; R; psbM; 2S st nr	5		
2981	1264	II. 89	12 19 59.8	3.037	2	73 45 10.1	19.97	2	pB; pL; pgbM; B* np	5		
2982	1266	II. 145	12 20 1.6	3.058	1	83 20 42.1	19.97	1	vF; vS; E	3		
2983	1267	II. 170	12 20 5.0	3.043	2	76 30 10.1	19.97	2	pF; S; R; bM	3		
2984	1268	II. 171	12 20 8.2	3.044	1	76 55 44.1	19.97	1	vF; vS; cE; gbM	2		
2985	1269	12 20 16.0	3.088	1	97 24 14.1	19.97	1	vF; pL	1		
2986	1270	II. 146	12 20 19.4	3.057	2	82 58 0.1	19.97	2	cF; L; R; gbM	3		
2987	1271	II. 65	12 20 20.7	3.046	2	78 7 49.1	19.97	2	B; L; cE; psbM; *10 nf	3		
2988	1272	II. 172	12 20 22.2	3.043	1	76 53 54.1	19.97	1	cF; S; gbM	2		
2989	II. 497	12 20 25.0	3.053	1	81 0 59.1	19.97	1	pF; vS	1		
2990	1273	12 20 26.1	3.089	1	97 30 49.1	19.97	1	pF; pL; lE	1		
2991	1274	I. 28, 1	12 20 33.7	3.041	2	76 9 14.1	19.97	3	vB; cL; R; p of 2	2		
2992	1274, a	R. nova	12 20 ±	3.041	...	76 9 ±	19.97	...	See note	0*		
2993	1276	II. 173	12 20 35.2	3.043	1	76 55 49.1	19.97	1	B; pS; R; bM; r	2		
2994	1275	I. 28, 2	12 20 36.7	3.041	3	76 13 17.1	19.97	3	B; cL; vIE; r; f of 2	5		
2995	1275, a	R. nova	12 20 ±	+3.041	...	76 13 ±	+19.97	...	See note	0*		

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
3045	h. 1309	H. II. 36	12 24 29.8	+3.060	3	85 17 47.9	+19.93	3	F; cL; biN or D neb	6
3046	III. 42	12 24 33.4	3.040	1	77 37 57.9	19.93	1	vF	1
3047	3397	12 24 40.2	3.195	2	129 12 18.9	19.93	2	vF; L; R; vglbM	2
3048	1311	I. 234	12 24 41.5	2.827	1	31 16 14.9	19.93	1	B; cS; E; pgbM; * 9 f 30''	3
3049	1312	M. 88	12 24 54.5	3.031	3	74 48 26.9	19.93	3	B; vL; vmE; p of D neb	3
3050	II. 118	12 24 +	3.031	...	74 48 -	19.93	...	F; S; f of D neb (not obs by h.).	8
3051	III. 69	12 24 57.5	3.024	1	72 25 57.9	19.93	1	vF; vS	1
3052	1313	II. 66	12 24 58.5	3.040	2	78 3 15.9	19.93	2	pB; S; R; gbM	3
3053	1314	II. 92	12 24 59.4	3.025	1	72 32 23.9	19.93	1	vF; S	2
3054	3398	II. 771	12 25 4.1	3.090	1	96 46 40.6	19.92	1	pB; cL; iE; gvlbM; er	3
3055	1315	III. 18	12 25 6.8	3.060	1::	85 14 53.6	19.92	1	vF; cL; r; f of 2	2
3056	1316	II. 631	12 25 6.8	3.034	1	75 48 23.6	19.92	1	cF; pmE 90°±; gbM; * 9 f 8s	2
3057	3399	12 25 9.9	3.197	2	129 8 6.6	19.92	2	pB; S; R; psmbM * 16	2
3058	1317	12 25 9.9	3.055	1	83 24 1.6	19.92	1	vS; R; sbM * 13	1
3059	1318	12 25 14.6	2.974	1	57 7 59.6	19.92	1	vF; S; R; lbM	1
3060	1196, a	R. nova	12 25 30	3.058	???	84 47	19.92	???	Query R.A.; vF; 10' s of scarlet *.	0
3061	1319	III. 834	12 25 33.9	2.833	1	32 45 54.6	19.92	1	pF; vS; iR; vgbM	2
3062	1321	12 25 45.5	2.748	1	25 29 50.6	19.92	1	pB; S; R; psbM	1
3063	1320	III. 302	12 25 50.1	2.980	2	59 30 52.6	19.92	2	eF; vS; R; bM	3
3064	1322	12 26 3.1	3.049	1	81 22 31.6	19.92	1	F; S; R; bM	1
3065	1323	III. 78	12 26 3.1	3.029	1	74 38 42.6	19.92	1	F; pS; R; r	3
3066	IV. 5	12 26 3.3	3.070	3	89 8 56.6	19.92	3	cB; vL; vmE 95°±; B * in cont.	3
3067	1324	II. 93	12 26 4.0	3.024	1	72 56 23.6	19.92	1	F; vS; bM*	2
3068	II. 158	12 26 20.2	3.046	3	80 33 36.3	19.91	3	F; pL; R; bM; r	3
3069	II. 849	12 26 23.0	2.732	1	25 12 55.5	19.91	1	pB; vS; lE; sbMSN	1
3070	II. 757	12 26 24.2	3.091	2	96 36 57.3	19.91	2	vF; S; 2 vS st inv	2
3071	1326	12 26 26.9	2.741	1	25 37 18.3	19.91	1	pB; S; pmE; pgbM; * 9 inv	1
3072	1325	12 26 32.9	3.044	1	80 2 47.3	19.91	1	eF; pL; lE; vlbM	1
3073	1327	12 26 38.0	3.104	1	101 14 26.3	19.91	1	vF; iF; bM	1
3074	1328	II. 325	12 26 53.3	2.975	1::	58 57 57.3	19.91	1::	F; pL; iR; bM	3
3075	1329	I. 31	12 26 56.3	3.048	3	81 31 57.3	19.91	3	vB; vL; mE 120°±; psmbM; L * f; * 9 p.	5*
3076	1330	II. 37	12 27 0.4	3.063	2	86 34 26.3	19.91	2	pB; L; pmE 60°±; mbM	4
3077	1331	II. 67	12 27 2.1	3.037	3	77 54 52.3	19.91	4	pF; cS; R; bM; * 9 f 30s	5
3078	III. 26	12 27 2.2	3.009	1	68 41 56.3	19.91	1	eF; L	2*
3079	1332	8 Canum	12 27 7.7	2.925	4	47 52 34.0	19.90	4	Nebulous *	4*
3080	II. 500	12 27 8.5	3.047	1	81 1 56.0	19.90	1	vL; er	1
3081	1333	II. 175	12 27 10.6	3.032	2	76 9 20.0	19.90	2	F; pL; R; vgbM	3
3082	1334	II. 147	12 27 11.7	3.051	1	82 46 13.0	19.90	1	pB; pL; pmE; vgbM; r	2
3083	1336	II. 410	12 27 13.1	2.952	2	53 42 9.0	19.90	2	cF; L; lE; vglbM; r	4
3084	1335	II. 94	12 27 14.9	3.024	2	73 40 25.0	19.90	2	F; pS; bM; r	5
3085	1337	V. 2	12 27 19.6	3.064	1	87 2 47.0	19.90	1	B; vL; mE 110°; sbM; er	5†
3086	1338	12 27 32.8	3.015	1	71 1 23.0	19.90	1	pB; pmE	1
3087	1341	12 28 6.8	2.860	1	38 25 19.7	19.89	1	eF; pL; R	1
3088	1340	12 28 12.2	3.052	1	83 6 37.7	19.89	1	pF; cS; R; bM	1
3089	1346	II. 850	12 28 13.8	2.721	1	25 42 2.7	19.89	1	F; L; iR; vgbM; S * nf	2
3090	1342	III. 493	12 28 15.1	3.071	1::	89 28 3.7	19.89	1:	eF; S; R; gbM	3
3091	1344	III. 802	12 28 15.7	2.783	2	30 19 21.7	19.89	2	vF; pS; E; vgbM; * 9 f 2'; p of 2.	4
3092	1339	I. 160	12 28 17.8	3.081	4	93 1 7.7	19.89	4	vB; cL; pmE 63°±; vsmbMN.	6
3093	1345	II. 120	12 28 22.7	3.026	3	74 43 45.7	19.89	3	B; L; lE; lbM	5
3094	1347	III. 807	12 28 23.5	2.781	2	30 17 43.7	19.89	2	eF; pS; E; f of 2	3
3095	1343	I. 36	12 28 24.2	3.033	2	77 0 46.7	19.89	3	pB; S; vLE; sp of 2	4
3096	1349	I. 37	12 28 32.3	+3.033	2	76 58 38.7	+19.89	2	pB; S; R; bM; nf of 2	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
3097	h. 1348	H.	M. 89	12 28 35.2	+3.032	1	76° 40' 32.7"	+19.89	4	pB; pS; R; gmbM	5*
3098	3400	12 28 36.9	3.211	1	128 39 49.7	19.89	1	F; vIE; glbM	1
3099	1350	II. 343	12 28 44.3	2.983	2	62 42 18.7	19.89	2	B; pS; iR; vsmbM * 12	3
3100	1351	II. 380	12 28 50.3	2.981	1	62 18 47.7	19.89	2	F; pL	3
3101	1352	I. 92	12 28 59.6	2.980	3	61 16 7.4	19.88	3	vB; vL; mE 150°; gbM; 3st f	4+
3102	1354	12 29 0.3	2.980	1	62 16 6.4	19.88	1	vF; nf of 2 or ?3	1
3103	1353	I. 119	12 29 0.4	3.046	1:	81 33 21.4	19.88	1:	cB; pL; R; gbM	2*
3104	1355	II. 407	12 29 5.7	3.008	2	69 54 20.4	19.88	2	pB; pL; vIE; lbM; r	4
3105	1356	II. 68	12 29 21.6	3.034	2	77 47 21.4	19.88	2	pB; S; iE; psbM	4
3106	1357	V. 24	12 29 22.9	2.993	4	63 14 30.4	19.88	4	B; eL; eE 136° 1'; vsbMN = * 10, 11.	5+
3107	1360	III. 880	12 29 26.1	2.821	1	35 0 1.4	19.88	1	pF; S; iR; gbM	2
3108	{ 1358 = 1363 = 1359 = 1363 }	IV. 8	12 29 26.5	3.035	4	77 59 26.4	19.88	4	vF; L; np of D neb } pos 160° ±	6*+
3109	{ 1358 = 1363 = 1359 = 1363 }	IV. 9	12 29 28.0	3.035	2	78 0 26.4	19.88	2	vF; L; sf of D neb } pos 160° ±	6*+
3110	1361	I. 32	12 29 45.6	3.047	5	81 59 3.1	19.87	5	cB; pS; mE 0° ±; sbMrN ...	9
3111	M. 90	12 29 52.8	3.028	2	76 4 18.1	19.87	2	pL; bMN	2*
3112	1364	III. 939	12 29 55.6	2.392	1	14 59 52.8	19.86	1	eF; S	2
3113	1362	III. 602	12 29 59.1	3.024	1	74 58 8.1	19.87	1	vF; L; E; vgbM; cB * att... ..	2+
3114	3401	12 30 6.6	3.241	1	132 51 14.1	19.87	1	vF; S; * 10 n 30"	1
3115	3402	12 30 17.6	3.199	1	124 44 3.1	19.87	1	vF; L; iE; vglbM	1
3116	3403	12 30 19.4	3.224	1	129 45 59.1	19.87	1	F; S; pmE; 2 st p	1
3117	III. 13	12 30 23.9	3.051	1	83 9 55.8	19.86	1??	vF; vS	1
3118	1365	II. 15	12 30 26.2	3.039	1	79 40 5.8	19.86	1	pF; pS; R; sbMN; * np ...	3
3119	1366	12 30 26.7	3.039	1	79 45 21.8	19.86	1	F; R; bM (?=II.15+5/P.D.)	1
3120	1367	M. 91??	12 30 30.8	3.025	1?	75 26 55.8	19.86	1?	np this place is a F neb; not M. 91, whose existence? ..	1
3121	1368	M. 58	12 30 36.6	3.031	3	77 24 52.8	19.86	3	B; L; iR; vmbM; r	6
3122	1369	I. 124	12 30 40.5	3.053	1	83 51 55.8	19.86	1	pB; L; vgbM	3
5071	12 31 0.5	89 2 46.5	See No. 5071.	
3123	1370	III. 495	12 31 14.1	2.945	2	55 46 15.8	19.86	2	cF; S; iE; bM	3
3124	D'Arrest, 91	12 31 17	3.02	[3]	76 7 12	19.85	[3]	vF; S; R	0
3125	1371	I. 125	12 31 18.5	3.056	2	84 54 40.5	19.85	2	pB; L; E; psbM	4
3126	III. 98	12 31 45.7	3.047	1:	82 25 53.5	19.85	1:	vF; eS	1
3127	1374	I. 273	12 31 53.3	2.369	3	15 2 24.5	19.85	4	cB; L; iE; pgmbM	7*
3128	3404	M. 68	12 32 5.1	3.166	1	115 58 45.2	19.84	1	⊕; L; eRi; vC; iR; rrr; st 12, red.	4
3129	1372	III. 504	12 32 6.7	3.050	1	83 12 29.2	19.84	1	vF; cS	3
3130	1373	II. 31	12 32 8.0	3.072	1	89 16 17.2	19.84	1	F; L; E 90° ±; vgbM	4
3131	1375	II. 183	12 32 26.6	3.088	2	94 34 23.2	19.84	2	pB; cL; E; sbMN=*	4
3132	1376	I. 43	12 32 44.2	3.110	1	100 50 14.2	19.84	1	!; vB; vL; eE 92°; vsmbMN.	3+
3133	1377	II. 632	12 32 49.6	3.016	3	73 56 8.2	19.84	4	pF; pL; R; gbM	5
3134	1378	I. 24	12 32 50.8	3.034	5	79 2 56.2	19.84	5	B; pS; R; gmbM; r; 3 st f... ..	7
3135	II. 636	12 32 53.7	3.089	1	95 2 22.9	19.83	1	F; vL; bM	1
3136	III. 105	12 33 11.7	3.040	2	80 51 21.9	19.83	2	eF; L; R; vlbM	2
3137	III. 509	12 33 13.9	3.065	1	88 0 51.9	19.83	1	vF; vS	1
3138	1379	II. 577	12 33 14.1	3.059	2	86 6 33.9	19.83	2	F; S; R; 2 st 8 f	3*
3139	3405	12 33 23.5	3.242	1	130 8 57.9	19.83	1	eF; L; R; psbM; p of 2	1
3140	1380	II. 184	12 33 26.5	3.088	1	94 21 50.9	19.83	1	F; L; E; vglbM	3
3141	3406	12 33 31.0	3.242	1	130 12 2.6	19.82	1	F; L; R; vgbM; r	1
3142	1381	I. 254	12 33 42.8	2.689	1	27 36 49.6	19.82	1	B; L; vmE 118° 6'; glbM ...	2
3143	1382	III. 43	12 33 53.3	3.027	2	77 20 9.6	19.82	3	vF; pS; E; 2 or 3 vS st inv ..	5
3144	1383	II. 69	12 34 10.3	3.033	2	79 3 58.6	19.82	2	pB; pL; R; psbM; r; * 12 np 1'.	4
3145	3407	Δ. 272	12 34 13.2	3.463	1	152 12 8.6	19.82	1	Cl; pL; pC; cE; st 10... ..	1
3146	I. 7	12 34 16.3	3.042	1:	81 24 51.6	19.82	1:	vB; vL (no doubt a comet) ..	1
3147	II. 19	12 34 20.3	3.042	1	81 30 51.6	19.82	1	F; vL	2
3148	1384	II. 148	12 34 23.9	+3.043	3	81 55 10.3	+19.81	3	pB; S; R; psmbM	6*

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
3149	h. 3408	H.	h m s 12 34 44.6	s +3.247	1	129° 54' 10.3"	+19.81	1	eF; vS; R; * att nf; p of 2...	1
3150	II. 744	12 34 46.0	2.815	2	38 48 50.3	19.81	2	pF; S; iR; er	2
3151	} 1385 {	I. 178 I. 179	}	12 34 48.0	2.886	4	48 4 47.3	19.81	4	{ B; L; R; mbM } D neb; { pb } pos 160°	5†
3152				12 34 56.3	3.021	1	76 17 9.3	19.81	1
3153	1387	12 34 56.5	2.922	3	54 9 54.3	19.81	3	F; pS; R; lbM; * 8.9 f	5
3154	1388	II. 411	12 34 56.9	3.026	3	77 34 1.3	19.81	3	B; pL; lE; vsymbM; 2 st p ..	5
3155	1386	M. 59	12 35 3.9	3.249	1	129 58 30.3	19.81	1	pF; S; R; psbM; f of 2.....	1
3156	3409	12 35 3.9	3.041	1	81 33 55.3	19.81	1	cF; pL; E; psbM; r	5
3157	1389	II. 149	12 35 7.3	3.059	1	86 10 20.3	19.81	1	B; E	1
3158	1390	12 35 8.7	2.935	1	56 39 40.3	19.81	1	F; S; R; np of 2.....	2
3159	1391	II. 659	12 35 9.8	2.883	2	47 56 36.3	19.81	2	pF; S; R	3
3160	1392	II. 660	12 35 10.4	3.095	2	96 16 10.3	19.81	2	vF; cS; lE; gbm	4
3161	{ 1393 = 3410 1394 = 3411 }	II. 772								
3162	{ 3411 = 3410 1394 = 3411 }	II. 773	12 35 11.3	3.095	2	96 11 25.3	19.81	2	cF; S; E; gbm	4
3163	D'Arrest, 92	12 35 21	3.07	[1]	91 2 12	19.80	[1]	pB; pL; E; lbM; ? biN	0
3164	1395	II. 532	12 35 21.7	3.055	1	85 16 24.0	19.80	1	cF; S; R; lbM	3
3165	1397	V. 42	12 35 22.0	2.934	2	56 41 21.0	19.80	2	!; vB; vL; eE70°±; bMN; B*nr.	3†
3166	1396	I. 14	12 35 23.8	3.070	1	89 18 56.0	19.80	1	pB; L; E45°±	4
3167	1398	III. 603	12 35 38.1	3.015	2	74 55 47.0	19.80	2	vF; L; mE135°±; vgbM ...	3
3168	1400	12 35 39.8	2.992	2	69 17 27.0	19.80	2	vF; L; vglbM.....	2
3169	1399	II. 38	12 35 41.1	3.060	2	86 32 30.0	19.80	2	B; L; iR; vgvmbM; r	4
3170	1401	12 35 41.9	3.055	1	85 32 15.0	19.80	1	vB; cS; R; smbM	1*
3171	} 1402 {	II. 70 = II. 176	}	12 35 43.8	3.026	1::	77 49 6.0	19.80	1	F; R; gbm	3
3172				1402, a	R. nova	12 35 43±	77 49 ±
3173	1403	II. 125	12 35 49.4	3.019	1	75 58 44.0	19.80	1	pB; S; E; r; * 12 sf 1'	2
3174	II. 20	12 35 56.9	3.043	1	81 53 10.3	19.79	1	vS	1*
3175	III. 494	12 36 3.7	3.072	2	89 53 20.7	19.79	2	vF; cS; E	2
3176	1402	I. 10	12 36 12.1	3.062	1	87 14 46.7	19.79	1	eB; pS; lE; mbM	5
3177	} 1406 {	II. 794 No. 1	}	12 36 15.9	2.754	1	34 4 23.7	19.79	1	vF; S; R; gbm	1*
3178				3412	12 36 21.6	3.262	1	130 58 46.7	19.79
3179	1407	II. 794 No. 2	12 36 23.0	2.756	1	34 22 23.7	19.79	1	F; S; 4 vS st sp	2*
3180	1405	III. 44	12 36 25.5	3.025	4	77 39 25.7	19.79	3	vF; pL; lE115°±; np of D neb	6*†
3181	1410	I. 274	12 36 33.0	2.258	4	14 48 37.7	19.79	5	pB; cS; R; gbm; * p	8
3182	1408	M. 60	12 36 34.8	3.025	5	77 40 38.4	19.78	6	vB; pL; R; f of D neb	8†
3183	3413	12 36 41.2	3.256	1	129 57 35.4	19.78	1	vF; R; bM; r	1
3184	1409	II. 12	12 36 42.3	3.005	3	72 50 28.4	19.78	4	eB; L; E90°; gbm; r	6
3185	1413	12 36 48.8	2.698	1	30 16 19.4	19.78	1	pF; pL; gbm; 2 B st f	1
3186	III. 662	12 36 50.2	3.072	1	89 47 49.4	19.78	1	vF; pL	1
3187	1411	II. 126	12 36 54.0	3.018	2	76 7 8.4	19.78	2	F; vL; pmE; ?D; 3 st nr ...	4
3188	1412	II. 661	12 36 54.5	2.876	2	48 12 25.4	19.78	3	vF; vS; stellar; * 15, 16 f ...	3
3189	1414	I. 176	12 37 7.8	2.929	4	57 3 48.4	19.78	5	!; pB; L; vmE34°.3; sp of 2	6*†
3190	1415	I. 177	12 37 16.6	2.929	3	57 1 24.1	19.77	3	!; pF; L; E90°±; nf of 2 ...	6*†
3191	3414	II. 558	12 37 23.1	3.109	1	99 18 53.1	19.77	1	vF; L; E; * 16 att; * 9 p ...	2
3192	1416	II. 127	12 37 27.8	3.016	1	75 43 35.1	19.77	1	F; cS; R; bM; r	3
3193	1417	II. 71	12 37 29.4	3.025	1	78 2 34.1	19.77	1	vB; S; vsymbMN	2
3194	3415	12 37 37.9	3.264	1::	130 19 53.1	19.77	1::	F; pL; R; gbm	1
3195	1418	II. 643	12 37 42.5	2.898	1	52 5 58.1	19.77	1	pF; pL; R; gbm; r	2
3196	II. 39	12 37 46.2	3.057	1	86 0 49.1	19.77	1	pB; 2 S st in M; S * p	1
3197	1419	I. 142	12 37 58.3	3.057	2	86 11 14.8	19.76	2	B; pL; iR; mbM; * 10 sp ...	3
3198	1420	I. 15	12 37 59.0	3.072	1	89 41 29.8	19.76	1	B; vL; mE45°±; psbM ...	4
3199	1421	12 38 12.3	+3.023	1	77 47 31.8	+19.76	1	B; S; R; psbM	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. O.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. O.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos	Other Authorities.								
3200	h.	H. III. 663	h m s 12 38 15.2	s +3.072	1	89 46 47.8	+19.76	1	vF; S; iF.....	1
3201	1422	III. 328	12 38 23.9	2.952	3	62 6 37.8	19.76	4	pF; cS; R; bM; r; p of 2 ...	6
3202	1423	II. 774	12 38 32.4	3.098	1	96 18 3.8	19.76	1	pF; S; R; psmbM	3
3203	3416	12 38 36.0	3.273	1	130 56 55.5	19.75	1	eF; S; R; vgbM.....	1
3204	1424	III. 329	12 38 43.7	2.951	1	62 10 24.5	19.75	2	F; vS; R; sbM*10; f of 2 ...	4
3205	3417	12 38 47.3	3.104	1	97 52 44.5	19.75	1	vF; cS; R; glbM	1
3206	III. 778	12 39 11.1	2.733	2	34 28 43.5	19.75	2	cF; S; lE.....	2*
3207	1425	II. 326	12 39 17.7	2.928	1	58 30 6.2	19.74	1	vF; pmE; ?biN	3
3208	3418	12 39 18.0	3.274	1	130 49 55.2	19.74	1	eF; lE; vgbM.....	1
3209	3419	12 39 36.0	3.263	1	128 48 5.2	19.74	1	eeF; pL; R	1
3210	3420	12 39 36.4	3.118	1	100 52 34.2	19.74	1	eF; S; 1 or 2 st inv	1
3211	3421	12 39 45.5	3.291	1	132 34 50.2	19.74	1	pF; S; R; gbM	1
3212	3423	III. 523	12 39 58.2	3.111	1	99 17 29.9	19.73	1	cF; L; E 45°±; gvlbM	2
3213	3422	12 40 2.9	3.279	1	130 47 24.9	19.73	1	eF; pS; R; vgbM ; S * sp ...	1
3214	1426	II. 181	12 40 6.3	3.081	1	91 58 8.9	19.73	1	B; pL; pmE 25°	5*
3215	1427	III. 398	12 40 13.3	2.984	3	69 46 52.9	19.73	3	F; S; R; sbM*; rr	4
3216	1428	II. 795	12 40 16.6	2.728	1	34 41 11.9	19.73	1	vF; vS; vmE; vsmbM	3*
3217	1430	12 40 39.3	2.897	1	53 52 53.6	19.72	1	pF; vS; R; psbM	1
3218	1429	III. 543	12 40 43.4	3.051	1	84 53 57.6	19.72	2	eF; pL; *9.10 p 10 ^s	3
3219	1431	II. 128	12 40 44.5	3.010	3	75 28 34.6	19.72	3	pB; vL; E; vglbM; r	5
3220	III. 664	12 40 48.3	3.076	1	90 55 44.6	19.72	1	vF; S	1
3221	1432	II. 182	12 40 58.2	3.081	1	92 33 48.6	19.72	1	pB; pL; E 90°±; mbM.....	5
3222	1433	II. 381	12 41 2.8	2.943	2	62 0 41.6	19.72	3	F; cS; R; bM	4
3223	III. 906	12 41 5.5	2.332	1	18 3 44.6	19.72	1	vF; pL; E	1
3224	1435	II. 796	12 41 9.5	2.731	1	34 51 21.6	19.72	1	cF; pS; vlE; mbMN	3*
3225	1434	II. 72	12 41 11.2	3.021	6	78 14 47.6	19.72	6	pF; S; vlE	7
3226	3424	Δ. 510 ?	12 41 12.2	3.282	1	130 32 29.3	19.71	1	pB; L; R; gbM; r	1
3227	1436	I. 39	12 41 21.8	3.094	2	95 2 2.3	19.71	2	vB; L; lE 45°±; smbMrN ..	5
3228	I. 8 =	}	12 41 28.1	3.032	6	80 44 53.3	19.71	6	cB; pL; iR; bM; r.....	6*
		III. 6									
3229	{ 1437 = } { 3425 = } { 1438 = } { 3426	I. 129	12 41 46.4	3.107	2	97 54 3.3	19.71	2	vB; R; vmbMrN; r	3
3230	{ 1437 = } { 3425 = } { 1438 = } { 3426	III. 524	12 41 49.6	3.119	2	100 38 16.3	19.71	2	F; L; mE40°; vlbM; B*p ...	4
3231	II. 578	12 42 2.1	3.054	1	85 50 43.0	19.70	1	F; S.....	1
3232	III. 514	12 42 5.8	3.109	2	98 21 44.0	19.70	2	eF; cS; pmE	2
3233	1439	II. 662	12 42 8.9	2.842	2	47 18 39.0	19.70	2	cF; S; R; gbM	3
3234	3427	12 42 15.5	3.288	1	130 31 49.0	19.70	1	vF; vS; R; psbM	1
3235	III. 815	12 42 19.7	2.753	1	38 2 43.0	19.70	1	S; stellar	1
3236	III. 722	12 42 22.8	3.118	1	100 20 43.7	19.69	2	eF; S	1
3237	3428	Δ. 511	12 42 26.6	3.289	1	130 36 13.7	19.69	1	pB; cS; R; gbM.....	1
3238	1440	III. 610	12 42 29.3	3.092	1	94 26 6.7	19.69	1	eF; pL; lE	2
3239	1451, a	R. nova	12 42 35.9	2.95	::	63 45 13.4	19.69	::	E	0
3240	1441	II. 95	12 42 39.1	3.000	2	74 4 17.7	19.69	2	cB; pL; vmE 28°5; sbMN ..	4+
3241	1442	12 42 39.4	2.948	1	63 45 43.7	19.69	1	vF; pL.....	1
3242	1443	II. 412	12 42 40.1	2.888	1	53 54 17.7	19.69	1	F; S; E; glbM; er	4
3243	1444	I. 140	12 42 51.5	3.045	4	83 55 22.7	19.69	4	pB; L; vlE; glbM	6
3244	1445	III. 536	12 43 1.6	3.130	2	102 33 51.7	19.69	2	F; pS; R; gbM	3
3245	1446	12 43 19.9	3.093	1	94 30 56.4	19.68	1	eF; vS; bet 2 st	1
3246	1448	III. 424	12 43 22.9	2.900	1	56 4 35.4	19.68	1	vF; stellar	2
3247	1475, a	R. nova	12 43 26	2.92	::	60 23 ±	19.68	::	E; bMN	0
3248	1447	III. 611	12 43 26.4	3.088	1	93 23 9.4	19.68	1	eF; S; bM	2
3249	1451	I. 84	12 43 32.0	2.945	1	63 44 13.4	19.68	1	vB; vL; E; vg; vsymbMeBN ..	2+
3250	1449	III. 280	12 43 34.1	3.135	1	103 34 27.1	19.67	1	F; vS; R; stellar; np of 2 ...	2
3251	1450	II. 298	12 43 37.5	3.135	1	103 34 57.1	19.67	1	F; pL; R; lbM; sf of 2	2
3252	3430	12 43 38.8	3.293	1::	130 19 50.1	19.67	1::	neb; 1st of 3	1
3253	3431	12 43 38.8	3.293	1::	130 19 50.1	19.67	1::	2nd of 3	1
3254	1452	I. 41	12 43 45.4	+3.098	1	95 38 6.1	+19.67	1	vF; pL; E	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
3255	h.	H.	h m s 12 43 53.8	s +2.720	1	36 20 41.1	+19.67	1	F; S; vsmbM	1
3256	1453	II. 73	12 44 3.4	3.019	4	78 19 45.1	19.67	5	cF; pL; { ^R mE90°} r; *12p.	6*
3257	1454	12 44 5.1	3.047	1	84 23 15.1	19.67	1	vF; vS; R	1
3258	1456	M. 94	12 44 17.2	2.838	5	48 6 31.8	19.66	5	vB; L; iR; vsmbMBN; r...	10+
3259	1457	III. 496	12 44 18.7	2.890	1::	55 5 14.8	19.66	1	eF; vS; pmE	2
3260	1455	III. 515	12 44 20.7	3.108	1	97 39 9.8	19.66	1	F; pL; iE; pglbM	2
3261	1458	III. 721	12 44 30.1	2.776	1	41 33 55.8	19.66	1	vF; S; R; psbM	2
3262	3432	I. 133	12 44 30.5	3.118	1	99 41 33.8	19.66	1	cB; vS; vbMN=*9; *10 sf	2
3263	3429	12 44 31.3	3.300	1	130 37 52.8	19.66	1	F; R; gbM	1
3264	3433	12 44 37.2	3.297	1	130 18 42.8	19.66	1	F; L; E; gbM; 3rd of 3	1
3265	1460	12 44 52.3	3.012	1	77 9 54.5	19.65	1	pB; mE; r	1
3266	II. 344	12 44 54.5	2.940	1	63 28 40.5	19.65	1	F; pL; iE	1
3267	1459	III. 537	12 44 55.1	3.132	1	102 38 47.5	19.65	1	F; vS; iR; gbM	2
3268	III. 907	12 44 55.5	2.244	1	17 36 39.8	19.66	1	vF; cL; E 135°±	1
3269	1451, b	R. nova	12 45 0.0	2.95	::	63 23 0.4	19.65	::	E 0°	0
3270	1463	IV. 78	12 45 0.2	2.175	1	16 21 38.5	19.65	1	pB; L; R; vg, vsbM	2
3271	3434	12 45 6.7	3.313	1	131 54 27.5	19.65	1	B; pS; R; vg, vsmbM	1
3272	III. 82	12 45 10.2	3.005	1	75 44 40.5	19.65	1	vF; S; E; r	1
3273	1461	I. 16	12 45 11.6	3.075	1	90 26 26.5	19.65	1	cB; L; vIE; vgibM.....	5
3274	1462	I. 25 =	12 45 15.0	3.015	5	77 55 36.5	19.65	5	B; pL; R; psbM; p of 2.....	7
3275	3435	II. 74	12 45 22.2	3.533	2	149 35 20.2	19.64	3	Cl; vL; st vB (κ Crucis).....	3+
3276	1464	Δ. 301	12 45 35.3	3.143	1::	104 38 49.2	19.64	1::	vF; pS; r.....	2
3277	1465	III. 70	12 45 46.1	2.992	3	73 23 18.2	19.64	3	vF; pL; E?	4
3278	1466	II. 75	12 45 53.4	3.015	4	78 0 38.2	19.64	4	pB; vmE 34°0; 3B st s; f of 2	6+
3279	III. 489	12 45 55.7	3.152	1	106 13 39.9	19.63	1	vF; S; lbM	1
3280	1467	III. 544	12 46 6.4	3.048	1	84 46 58.9	19.63	1	F; cS; R; gbM	2
3281	3436	12 46 11.3	3.295	2	128 58 3.9	19.63	2	B; pS; iE; mbM.....	3
3282	III. 525	12 46 12.7	3.115	1	98 45 39.9	19.63	1	vF; vS	1
3283	1468	II. 535	12 46 13.2	3.063	1	87 58 25.9	19.63	1	F; pL; mE; *9 p 90°	2
3284	III. 516	12 46 22.4	3.111	1	97 54 38.9	19.63	1	vF; S	1
3285	1469	II. 24	12 46 22.4	3.058	2	87 3 50.9	19.63	2	pF; pS; R; mbM	5
3286	1471	III. 618	12 46 26.5	2.862	1	52 25 5.9	19.63	1	eF; cS; R; bM	2
3287	1470	II. 186	12 46 29.7	3.101	1	95 51 34.9	19.63	1	F; cL; R; vglbM; r	3
3288	3437	II. 559	12 46 46.5	3.114	1	98 26 38.6	19.62	1	F; S; R; vlbM; p of D neb...	2
3289	III. 517	12 46 47.5	3.112	1	98 1 38.6	19.62	1	vF; S	1
3290	3438	12 46 47.5	3.114	1	98 26 17.6	19.62	1	vF; S; R; vlbM; f of D neb	1
3291	1472	III. 106	12 46 48.1	3.020	2	79 31 37.0	19.60	2	vF; pL; R; r	3
3292	I. 134	12 47 4.9	3.120	1	99 46 38.3	19.61	1	cB; vL; mE	1
3293	I. 135	12 47 15.1	3.131	2	101 49 8.3	19.61	2	pF; pS; R; mbM; p of D neb	2
3294	I. 136	12 47 16.1	3.131	2	101 50 8.3	19.61	2	pF; pS; R; mbM; f of D neb	2
3295	III. 526	12 47 16.9	3.121	1	99 51 38.3	19.61	1	eF; eS	1
3296	3439	12 47 17.2	3.384	1	137 58 54.3	19.61	1	vF; S; R; glbM	1
3297	II. 187	12 47 21.4	3.102	2	96 6 8.3	19.61	2	pB; pS; mbM; r.....	2
3298	1473	II. 345	12 47 29.9	2.925	3	62 9 58.3	19.61	2	F; R; *9 att 1/n	4
3299	II. 560	12 47 31.8	3.120	1	99 28 38.3	19.61	1	pF; pS; iR	1
3300	1475	I. 93	12 47 54.0	2.911	2	60 17 53.0	19.60	2	pB; pS; iE; *8nf 1'	4
3301	II. 538	12 47 55.2	3.132	2	101 52 8.0	19.60	2	vF; S; 2 or 3 st near	2
3302	1474	II. 21	12 47 59.1	3.029	3	81 10 55.0	19.60	3	pF; pL; R; bM; r	7
3303	1477	II. 382	12 48 6.7	2.921	2	61 49 20.0	19.60	2	pF; pS; gbM	3
3304	1476	III. 548	12 48 7.8	3.054	1	86 20 18.0	19.60	1	cF; S; vS* att	2
3305	1478	I. 211	12 48 9.1	2.765	1	42 43 7.0	19.60	1	pB; cS; R; psbM; *14 p ...	5
3306	1479	III. 816	12 48 26.6	2.683	1	36 8 27.7	19.59	1	eF; S; iE.....	2
3307	IV. 40	12 48 33.9	3.136	1	102 17 37.7	19.59	1	S; att to pB*	1
3308	3440	12 48 39.2	3.230	1	118 45 0.7	19.59	1	F; cS; R; gvlbM	1
3309	1509, a	R. 2 novæ	12 48 45	3.05	::	86 42 ±	19.58	::	{ F; D neb; E at right angles to each other.	0
3310											
3311	1480	I. 141	12 48 53.4	+3.047	1::	84 56 47.4	19.58	1::	{ H. vB } { h. pF } ; cL; E135°± ...	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
3312	h. 3441	H.	12 49 6.0	+3.326	1	131° 2' 20.4"	+19.58	1	eF; cS; R; gbM; p of 2.....	1
3313	3442	12 49 6.1	3.326	1	131 3 50.4	19.58	1	eF; S; R; gbM; f of 2	3
3314	1481	II. 383	12 49 9.7	2.915	1	61 29 48.4	19.58	1	vF; pL.....	2
3315	1483	I. 243	12 49 10.9	2.590	2	30 54 17.4	19.58	2	B; pS; vIE; vgbM	2
3316	1482	II. 777	12 49 20.0	3.104	1	96 3 28.1	19.57	1	F; S; R; bM	2
3317	3443	12 49 23.7	3.680	3	154 11 48.1	19.57	3	Cl; pL; pRi; iF; st 10 ...18	3
3318	{ 1484 = 3445 }	II. 549	12 49 33.0	3.112	2	97 46 5.1	19.57	2	pB; L; pmE 0°; gbM	3
3319	1485	II. 384	12 49 48.9	2.917	1	62 3 50.8	19.56	1	F; cL	2*
3320	II. 563	12 49 51.1	3.140	1	102 54 35.8	19.56	1	pB; iF; bM	1
3321	1486	M. 64	12 49 51.8	2.951	3	67 33 15.8	19.56	3	!; vB; vL; vmE 120°±; bMSBN=* ?.	10+
3322	3447	12 50 5.6	3.220	1	116 32 13.8	19.56	1	F; S; R; gbM	1
3323	3446	12 50 6.4	3.321	1	129 59 36.8	19.56	1	pF; vS; R; sbM*17';*10,70°3	1
3324	1487	II. 346	12 50 6.5	2.917	1	62 14 32.8	19.56	1	vF; pL; iF	3
3325	3444	Δ. 164	12 50 7.9	3.899	2	160 6 53.5	19.55	2	⊕; B; L; R; g; vsbM; st 12	2
3326	1488	III. 817	12 50 11.1	2.680	1	36 56 44.8	19.56	1	vF; S; iR; bM	2
3327	3448	12 50 15.0	3.375	2	135 30 1.5	19.55	3	F; pL; mE; vgbM	3
3328	1489	12 50 23.4	2.725	1	40 26 12.5	19.55	1	Neb; ?	1
3329	1490	12 50 42.3	3.138	1	102 17 56.2	19.54	1	vF; 3St sp	1
3330	1491	II. 536	12 50 42.5	3.060	1	87 40 23.5	19.55	1	pF; pL; pmE; vgbM; *nf30°	2
3331	1493	II. 387	12 50 45.5	2.906	2	60 46 1.2	19.54	2	pF; pL; R; vS* att	3
3332	1492	III. 613	12 50 47.5	3.087	1	92 51 26.2	19.54	1	cF; E; er; *sf 30''	2
3333	1494	II. 386	12 50 55.4	2.912	1::	61 49 52.2	19.54	1	F; pL; R.....	2
3334	1495	12 51 9.4	2.837	1	51 52 3.2	19.54	1	eF	1
3335	3449	Δ. 311	12 51 45.5	3.571	2	148 50 34.2	19.54	2	Cl; L; pRi; iR; st 10... ..	2
3336	1496	II. 385	12 51 48.6	2.908	2	61 38 24.6	19.52	2	F; S; R; psbM	3
3337	1497	I. 68	12 51 59.8	3.151	1	104 17 25.6	19.52	2	B; R; psmB; *13 np	3*
3338	II. 299	12 52 1.7	3.152	1	104 31 34.6	19.52	1	pB; pL; mbM.....	1*
3339	III. 908	12 52 1.7	2.172	1	19 1 32.6	19.52	1	eF; vS; iR; vlbM	1
3340	1499	IV. 30	12 52 22.9	2.852	2	54 23 0.3	19.51	2	vF; pL; vmE 30°±; bet 2 st	5+
3341	II. 644	12 52 24.2	2.832	1	51 56 32.3	19.51	1	pB; S; R; mbM.....	1
3342	1498	I. 162	12 52 28.6	2.990	6	75 4 23.3	19.51	6	B; pL; mE 90°; sbMN; S* inv.	7
3343	1500	12 52 41.1	2.903	1::	61 16 45.3	19.51	1::	1st of 5; *7 n	1
3344	1501	II. 388	12 52 49.6	2.903	2	61 17 56.0	19.50	3	cF; S; R; *7 n; 2nd of 5 ...	4
3345	III. 758	12 52 58.9	3.102	1	95 20 33.0	19.50	1	vF; vS; p of 2.....	1
3346	III. 759	12 52 58.9	3.102	1	95 20 33.0	19.50	1	vF; vS; F of 2	1
3347	1502	II. 389	12 53 0.1	2.902	1::	61 16 45.0	19.50	1::	3rd of 5	2
3348	1503	III. 83	12 53 9.2	2.999	2	76 46 17.0	19.50	2	cF; pL; R; vglbM; r.....	3
3349	1504	12 53 17.4	3.106	1	96 5 50.7	19.49	1	vF; S; E	1
3350	1505	II. 778	12 53 20.0	3.102	1	95 19 0.7	19.49	1	pF; cS; E; psbM; *np.....	2
3351	1507	II. 391	12 53 24.0	2.901	2	61 15 23.7	19.49	2	pB; pmE; bM; *7 n; 4th of 5	3
3352	1506	III. 614	12 53 24.1	3.094	1	93 50 3.7	19.49	1	cF; S; iR; bM	2
3353	1508	II. 390	12 53 28.3	2.908	1	62 21 3.7	19.49	1	vF	2
3354	1510	III. 363	12 53 29.7	2.900	1	61 17 14.7	19.49	2	pF; S; R; *7 n; 5th of 5 ...	3
3355	II. 300	12 53 33.9	3.147	2	103 11 32.7	19.49	2	2
3356	1509	I. 143	12 53 33.9	3.055	2	86 45 0.7	19.49	3	cB; cE; *10 att 135°±	4*+
3357	1512	12 53 35.2	2.724	1	42 1 50.7	19.49	1	pF; S; R; gbM	1
3358	1511	I. 69	12 53 37.9	3.150	1	103 45 56.7	19.49	1	pB; pL; iR; st nr	2*
3359	3450	12 53 44.3	3.257	1	120 12 2.4	19.48	1	vF; cS; R; * att; p of 2.....	1
3360	3451	12 53 51.3	3.257	1	120 10 2.4	19.48	1	vF; vS; R; slbM; f of 2.....	1
3361	II. 517	12 53 51.5	3.069	2	89 18 31.4	19.48	2	pB; pS; R; bM	2
3362	3452	12 54 9.0	3.361	1	132 0 55.1	19.47	1	eF; 3 or 4 st 11, 12, f	1
3363	V. 3	12 54 9.5	3.060	1	87 35 32.4	19.48	1	eF; vL; rr	1*
3364	II. 392	12 54 10.0	2.899	1::	61 23 44.4	19.48	1::	1st of 3	1
3365	1514	II. 645	12 54 10.5	2.823	2	51 54 16.4	19.48	2	pB; cS; R; smbM; *17 np..	3
3366	1513	IV. 47	12 54 12.7	3.094	1	93 47 23.4	19.48	1	pB; S; R; bM; stellar?.....	3
3367	1515	12 54 26.8	2.719	1	42 1 52.1	19.47	1	eF; S; E; bM	1
3368	1516	II. 393	12 54 39.4	+2.897	1	61 21 47.1	+19.47	2	F; pL; 2nd of 3	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
3369	h. 1518	H. II. 394	h m s 12 54 42.9	s +2.897	1	61° 24' 48".1	+19.47	1	vF; 3rd of 3.....	2
3370	1517	12 54 50.7	3.154	2	104 13 10.8	19.46	2	cF; L; vIE 45°±	4
3371	1519	II. 779	12 54 51.3	3.112	1	96 57 43.8	19.46	1	cF; S	2
3372	III. 364	12 55 1.9	2.895	1	61 12 29.8	19.46	1	vF	1
3373	3453	II. 190	12 55 43.0	3.115	1	97 19 31.2	19.44	1	F; pS; vIE; glbM	4
3374	III. 760	12 55 51.7	3.115	1	97 22 30.2	19.44	1	cF; vS; R	1
3375	3454	12 56 14.8	3.358	1	130 39 45.9	19.43	1	vF; R; Δ2st 8, 9, f	1
3376	III. 818	12 56 20.2	2.663	1	38 47 27.9	19.43	1	cF; S; R; vglbM	1
3377	II. 191	12 56 27.9	3.136	1	100 44 59.9	19.43	1	pB; pL; iR	2
3378	3456	12 56 38.5	3.263	2	119 46 26.6	19.42	2	pB; S; R; bM; *f 6 ^s	1
3379	3455	12 56 42.6	3.425	1	136 28 2.6	19.42	1	eeF; S; R; p of 2	1
3380	1521	12 56 48.3	2.646	1	37 55 20.6	19.42	1	eF; R; psbM	1
3381	3458	II. 561	12 56 56.5	3.129	1	99 35 40.6	19.42	1	pB; L; R; gmbM	2
3382	3457	12 56 58.2	3.426	1	136 29 22.3	19.41	1	F; S; R; f of 2	1
3383	1520	I. 40	12 56 59.1	3.101	2	94 48 23.6	19.42	2	pF; L; E; gbMBN; r	3
3384	III. 761	12 56 59.5	3.113	1	96 55 28.6	19.42	1	vF; S	1
3385	1522	II. 395	12 57 6.0	2.887	1	61 3 42.6	19.42	1	F; S; R; bM; *9 nf 1'	2
3386	3459	Δ. 411	12 57 14.1	3.455	1	138 32 6.3	19.41	1	B; vL; vmE 38°.7	1
3387	3460	12 57 32.1	3.388	2	132 50 51.0	19.40	2	B; pS; R; gpmbM; p of 2...	2
3388	3461	12 57 34.1	3.306	1	124 35 6.0	19.40	1	F; pL; R; vglbM	1
3389	3462	12 57 40.2	3.387	2	132 45 46.0	19.40	2	eF; S; R; psbM; f of 2.....	2
3390	1523	II. 188	12 57 43.2	3.107	(2)	95 45 18.0	19.40	2	F; pL; iE; r	3
3391	1524	II. 396	12 58 15.4	2.877	4	60 7 28.7	19.39	5	F; S; R; psbM*11	6
3392	3463	12 58 18.4	3.329	1	126 48 31.4	19.38	2	vF; pS; am 3S st.....	2
3393	1527	12 58 23.9	1.656	2	13 50 36.4	19.38	2	vF; S; R; vgbM.....	2*
3394	3464	12 58 26.1	3.263	1	119 0 14.4	19.38	1	F; cS; R; gbM	1
3395	1525	II. 413	12 58 26.5	2.825	2	54 4 20.7	19.39	2	pB; cS; R; smbM	4
3396	1526	II. 397	12 58 27.1	2.888	1	61 40 48.4	19.38	1	F; S; R	2
3397	3465	I. 130	12 58 31.6	3.116	1	97 15 49.4	19.38	1	vB; pS; E 0°±; bMBN.....	3
3399	1528	12 59 2.4	2.840	1	56 4 6.1	19.37	2	eF; S; R	2
3400	1529	II. 398	12 59 4.5	2.885	1	61 30 58.1	19.37	1	F; S; iF	2
3401	III. 303	12 59 9.8	2.874	1	60 10 26.1	19.37	1	eF; vS	1
3402	1530	II. 663	12 59 27.3	2.754	2	47 31 10.8	19.36	2	F; vS; R; stellar; vS*s	3
3403	1532	III. 779	12 59 28.4	2.537	1	32 55 57.8	19.36	1	eF; S; iE.....	3
3404	3466	12 59 33.6	3.255	2	117 28 37.8	19.36	2	vF; vL; cE; vgbM	2
3405	1531	III. 304	12 59 35.3	2.873	2	60 12 10.8	19.36	3	vF; vS; vIE; vglbM; *sp...	4
3406	1533	III. 783	12 59 35.4	2.588	1	35 40 40.8	19.36	1	vF; S; E; *att	2
3407	3467	12 59 37.3	3.221	1	112 55 45.8	19.36	1	F; pL; R; glbM	1
3408	III. 765	12 59 57.7	3.224	1	113 15 24.5	19.35	1	vF; pL; iF	1
3409	III. 937	13 0 17.3	1.670	1	13 57 22.5	19.35	1	vF; S; iR; bM	1
3410	III. 781	13 0 18.4	2.582	1	35 38 40.2	19.34	1	vF; S	1
3411	III. 782	13 0 32.3	2.580	1	35 36 40.2	19.34	1	vF; S	1
3412	1534	13 0 36.8	3.099	1	94 16 0.9	19.33	1	vF; vS; R; psbM	1
3413	3468	13 0 40.4	3.480	1	138 45 23.9	19.33	1	B; pL; R; gmbM	1
3414	III. 780	13 0 41.9	2.540	1	33 34 22.2	19.34	1	cF; S	1
3415	1535	13 0 57.8	2.948	2	70 50 8.9	19.33	2	F; vS; R; sbM; stellar	2*
3416	III. 346	13 1 8.2	2.902	1	64 29 21.6	19.32	1	eF; pL; iE	1
3417	3469	13 1 29.8	3.264	1	117 53 57.3	19.31	1	eF; cS; R	1
3418	1537	II. 189	13 1 32.2	3.111	1	96 1 35.3	19.31	1	B; pL; R; *9 sf	4
3419	III. 365	13 1 32.5	2.873	1	60 56 21.3	19.31	1	vF	1
3420	1536	II. 301	13 1 33.1	3.168	2	104 45 42.3	19.31	2	B; pL; R; psmbM	3*
3421	II. 185	13 1 40.9	3.102	1	94 36 21.3	19.31	1	F; S; iF; pB*nr.....	1*
3422	1539	III. 654	13 1 50.4	2.743	1	47 34 57.3	19.31	1	vF; vS; R; lbM	2
3423	1538	III. 401	13 1 51.4	2.811	2	54 3 8.3	19.31	2	vF; S; R; stellar	3
3424	1542	II. 815	13 1 54.5	2.599	1	37 18 52.3	19.31	1	vF; vS; stellar	2
3425	3470	13 1 54.7	3.404	1	132 21 15.0	19.30	1	vF; S; E; r	1
3426	Auw. N. 31	13 2 1.0	3.099	...	94 38 51.0	19.30	...	A neb (Markree Obs. Apr. 9, 1852).	0*
3427	1541	13 2 6.3	2.993	1	77 37 1.0	19.30	1	vF; S; iE; 2Sst s	1
3428	III. 766	13 2 11.9	3.225	1	112 38 21.7	19.29	1	vF; vS	1
3429	3471	13 2 15.9	+3.219	1	111 48 8.7	+19.29	1	pF; cS; R; slbM; am st.....	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h. 1540 =	H.		h m s	s		° ' "	"			
3430	3472	I. 42	13 2 23.2	+3.118	2	97 5 2.7	+19.29	2	pB; pL; R; vgpmbM; * 8 np	6
3431	III. 819	13 2 29.4	2.616	1	38 34 19.7	19.29	1	vF	1
3432	1543	II. 537	13 2 32.9	3.057	1	87 35 35.7	19.29	1	cF; pL; R; lbM; er	2
3433	1544	III. 366	13 3 5.6	2.863	1	60 20 39.4	19.28	1	cF; pS; lE	2
3434	1545	13 3 37.6	2.558	1	35 44 28.1	19.27	1	pF; S; iR; gbM	1
3435	III. 655	13 4 15.2	2.704	1	47 27 18.5	19.25	1	vF; pS; lbM	1
3436	1546	III. 305	13 4 20.5	2.852	1	59 36 40.5	19.25	2	vF; vS; vLE	3
3437	1547	I. 96	13 4 28.5	2.781	2	52 11 9.2	19.24	2	vB; vL; vmE 25°; vsbMN...	4
3438	III. 848	13 4 34.7	2.340	1	27 7 16.5	19.25	1	vF; vS	1
3439	D'Arrest, 93	13 4 37	3.05	[1]	63 51 42	19.24	[1]	pF; pL; R	0
3440	1550	III. 820	13 4 45.5	2.610	1	39 9 53.2	19.24	1	vF; R; bet 2 vS st	2
3441	1549	I. 85	13 4 48.8	2.907	2	66 20 21.9	19.23	2	pF; cL; E 17°0; biN; * 9 f	3
3442	1548	13 4 50.5	3.175	1	105 3 9.9	19.23	1	vF; R; bM; * 10 np 5'	1
3443	3473	13 4 51.7	3.419	3	132 21 4.9	19.23	3	pB; cS; R; am 4 st	3
3444	1551	II. 414	13 5 2.2	2.787	1	52 58 43.9	19.23	1	pF; S; E; psbM	2
3445	1552	II. 637	13 5 7.5	3.097	(1)	93 36 2.6	19.22	1	F; cL; iR; lbM	2
3446	II. 356	13 5 7.9	2.897	1	65 12 15.6	19.22	1	pB; S	1
3447	1553	III. 669	13 5 28.3	3.183	1	106 0 52.6	19.22	1	vF; R; bM	2
3448	1554	II. 746	13 5 30.7	3.203	1	108 46 4.3	19.21	1	cB; S; R; mbMpBN	2
3449	1555	III. 545	13 5 36.5	3.036	1	84 31 7.3	19.21	1	cF; vS; R; er	2
3450	1556	II. 129	13 5 42.5	2.982	3	76 39 28.3	19.21	3	cF; cL; vLE; lbM	4
3451	1557	13 5 46.2	2.663	1	43 4 1.3	19.21	1	pF; cS; R; *12 nf 90"	1
3452	1559	II. 664	13 5 55.8	2.692	2	45 12 45.3	19.21	2	pF; L; mE 20°; vlbM	4
3453	1558	M. 53	13 6 2.0	2.941	5	71 5 3.0	19.20	5	!; ⊕; vB; vC; ir; vmbM; st 12...	12
3454	1560	III. 649	13 6 10.1	2.826	3	57 26 40.0	19.20	2	vF; S; lE; * 13 n	4
3455	3474	13 6 13.8	3.425	3	132 12 57.0	19.20	3	pB; pL; R; gbM; * 7 nf ...	3
3456	1561	13 6 16.8	3.027	2	83 11 34.0	19.20	2	vF; S; R; pgbM	2
3457	1562	13 6 27.1	2.646	1	42 10 39.0	19.20	1	F; vS; R; gbM	1
3458	1563	III. 367	13 6 45.9	2.861	2	61 27 30.7	19.19	2	vF; pL; iR	3
3459	1564	I. 97	13 6 59.7	2.775	2	52 39 40.4	19.18	2	vB; pl; E 166°8; smbMvBN; * np.	3
3460	III. 909	13 7 19.4	1.918	1	18 36 12.4	19.18	1	vF; vS; R	1
3461	1565	II. 510	13 7 33.0	3.185	2	105 51 1.8	19.16	2	cF; pS; vLE; bM	3
3462	II. 816	13 7 49.8	2.568	1	37 58 10.8	19.16	1	F; S; iR; vgbmM	1
3463	3477	13 7 55.8	3.243	1	113 14 29.5	19.15	1	F; L; R; vgvlbM; * 9 p ...	1
3464	3476	13 7 56.4	3.744	1	149 19 9.5	19.15	1	Cl; P; E; sc st 11...	1
3465	1566	II. 511	13 7 56.9	3.184	1	105 38 49.5	19.15	1	pB; pL; R; bM	3
3466	3475	13 7 59.4	3.845	1	152 40 25.2	19.14	1	Cl; vL; vRi; st 11...	1
3467	3478	13 8 26.1	3.283	1	117 40 52.2	19.14	1	pF; R; sp of 2	1
3468	III. 670	13 8 26.7	3.185	1	105 43 13.2	19.14	1	vF	1
3469	II. 512	13 8 31.7	3.185	2	105 37 43.2	19.14	2	cF; S	2
3470	3479	13 8 46.5	3.283	1::	117 35 33.9	19.13	1::	Neb; nf of 2	1
3471	1567	13 9 0.0	2.837	1	59 33 42.9	19.13	1	vF	1
3472	1569	VI. 7	13 9 30.5	2.938	1	71 35 10.3	19.11	2	Cl; vF; pL; iR; vgbM; st 15...	3
3473	1568	II. 513	13 9 31.1	3.188	1	105 53 46.3	19.11	1	F; pS; iR	3
3474	1570	M. 63	13 9 31.9	2.699	1	47 13 45.3	19.11	1	vB; L; pmE 120°±; vsmbMBN.	3
3475	1571	III. 306	13 9 38.0	2.823	1	58 18 9.3	19.11	1	cF; cS; R; p of 2	2
3476	1572	III. 307	13 9 52.3	2.821	1	58 13 16.0	19.10	1	cF; cS; R; f of 2	2
3477	3480	I. 138	13 10 27.4	3.274	1	116 6 47.7	19.09	1	vB; S; R; vsmbM; * 10 f...	4
3478	3482	13 10 33.7	3.358	1::	124 41 30.4	19.08	1	eF; vS; E; r	1
3479	3481	13 10 33.9	3.357	1	124 35 15.4	19.08	1	eF; vS; R; * nr	1
3480	3483	13 10 46.3	3.519	1	137 10 22.1	19.07	1	B; S; R; psbM	1
3481	1573	III. 308	13 10 57.3	2.817	1	58 10 2.4	19.08	1	vF; cS	3
3482	III. 312	13 11 14.2	3.227	1	110 22 7.8	19.06	1	F; L; iR; bM	1
3483	1574	III. 282	13 11 54.7	3.178	1	104 6 53.5	19.05	1	vF; pL; pmE 135°±	2*
3484	1575	III. 309	13 11 55.9	+2.810	1	57 47 16.5	+19.05	1	eF; vS	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
3485	{ 1576 = 3489 }	III. 117	13 12 6.9	+3.162	2	102 0 4.2	+19.04	2	vF; cS; R; 1st of 3	5
3486	{ 1577 = 3490 }	II. 193	13 12 9.0	3.161	2	101 55 11.2	19.04	2	pB; S; vIE; sbM; 2nd of 3...	5
3487	3484	II. 566	13 12 9.6	3.285	1	116 50 8.2	19.04	1	pB; pS; cE; psbM; *7.8 f...	3
3488	{ 1578 = 3491 }	III. 118	13 12 14.9	3.162	2	101 57 46.0	19.04	2	cF; pS; vIE; 3rd of 3.....	5
3489	1578, a	R. nova	13 12 ±	3.162	...	101 57 ±	19.04	...	No description, one of a group of four.	0
3490	3485	13 12 26.3	3.285	1	116 40 17.9	19.03	1	vF; S; R; 1st of 4	1
3491	1579	II. 313	13 12 42.4	3.236	1	111 4 37.6	19.02	1	cB; cS; vIE 90° ±; bf	2
3492	II. 780	13 12 44.0	3.258	1	113 40 5.6	19.02	1	F; L; R; vglbM.....	1
3493	3486	13 12 55.4	3.469	1	132 59 25.6	19.02	1	eF; vS; R; 2nd of 4	1
3494	III. 724	13 12 55.9	3.226	1	109 52 4.6	19.02	1	cF; vS; iF	1
3495	1580	II. 327	13 13 3.1	2.818	2	59 1 33.6	19.02	2	pF; pL; gbM	3
3496	3487	13 13 4.2	3.470	2	132 58 39.3	19.01	2	pB; pL; R; 3rd of 4	2
3497	3488	13 13 8.5	3.470	2	132 59 51.3	19.01	2	cF; S; vIE; 4th of 4	2
3498	1583	III. 633	13 13 20.1	2.702	1	48 51 23.3	19.01	1	vF; S; R; lbM	2
3499	1581	III. 539	13 13 22.1	3.174	1	103 20 52.0	19.00	1	cF; vS; R; gbM.....	2
3500	1582	13 13 24.5	3.085	1	91 34 7.0	19.00	1	vF; iR; * 11 sp	1
3501	1584	III. 650	13 13 39.1	2.787	1	56 11 1.0	19.00	1	vF; cS; R; bM; sp of 2	3
3502	1585	13 13 46.8	2.786	1	56 7 15.0	19.00	2	vF; S; bet 2 st; nf of 2	2
3503	3493	II. 567	13 14 3.7	3.289	2	116 41 36.7	18.99	2	cB; pS; iE; psbM *	4
3504	3492	13 14 4.5	3.385	1	125 54 9.7	18.99	1	vB; pS; R; svmbM	1
3505	II. 665	13 14 9.9	2.660	1	46 10 1.7	18.99	1	pB; cS; E	1
3506	II. 22	13 14 34.0	3.002	1	80 46 29.4	18.98	??	vF; vS	1*
3507	1586	III. 619	13 15 6.1	2.717	1	50 42 22.8	18.96	1	vF; S; cE 0° ±	2
3508	3494	13 15 12.3	3.342	1::	121 36 35.5	18.95	1	eeF; p of 2	1
3509	1588	III. 808	13 15 14.4	2.367	1	31 37 0.8	18.96	1	cF; S; cE	2
3510	1587	III. 119	13 15 32.9	3.168	2	102 14 3.2	18.94	2	cF; cS; iR; glbM	2
3511	1589	II. 646	13 15 38.3	2.712	1	50 31 29.5	18.95	1	F; L; iR; vglbM	2+
3512	II. 826	13 16 2.4	2.359	1	31 34 28.2	18.94	1	F; S; E	1*
3513	3495	13 16 12.7	3.345	1	121 36 34.6	18.92	1	F; iE; psbM; f of 2	1
3514	1590	III. 368	13 16 19.0	2.841	2	62 17 27.6	18.92	2	pF; pS; pmE; glbM; r	3
3515	1592	13 16 19.0	2.827	1	60 56 38.6	18.92	1	vF; L; Δ 2 st 11 np.....	1
3516	1591	III. 925	13 16 24.5	3.018	1	82 52 33.6	18.92	1	vF; S; R; gbM	3
3517	3497	13 16 35.7	3.164	1	101 33 30.3	18.91	1	pB; S; iE	1
3518	3496	13 16 37.7	3.939	1	152 40 50.0	18.90	1	Cl; eR; mC; st 12...16	1
3519	3498	13 16 46.4	3.409	2	126 57 18.3	18.91	2	cB; P; R; psmbM; r.....	2
3520	1594	II. 666	13 16 59.8	2.646	1	46 10 43.3	18.91	1	pF; S; R; gmbM	2
3521	3499	13 17 1.9	3.327	2	119 34 56.0	18.90	2	vF; S; vIE	2
3522	1593	13 17 2.8	2.990	1	79 33 7.0	18.90	1	pF; S; R; gbM	1
3523	3500	13 17 5.0	3.333	1	119 36 57.0	18.90	1	vF; vS	1
3524	1596	II. 328	13 17 15.0	2.790	2	57 42 15.0	18.90	3	pB; pL; R; gmbM; * p.....	5
3525	3501	Δ. 482	13 17 15.1	3.481	4	132 17 10.7	18.89	4	!!; vB; vL; vmE 122° 5; bifid	4+
3526	1595	II. 653	13 17 15.9	2.955	3	75 17 16.7	18.89	3	pB; vS; R; gmbM; * f.....	4
3527	1597	II. 314	13 17 43.5	3.240	1	110 23 21.4	18.88	1	F; pS; iE; vgbM	2*
3528	3502	13 17 56.3	3.325	1	119 6 14.4	18.88	1	pB; S; E.....	1
3529	1598	III. 84	13 17 59.6	2.953	1	75 31 20.4	18.88	1	eF; vS; R; psbM	3
3530	3503	Δ. 312?	13 18 19.5	3.811	2	148 16 44.8	18.86	2	Cl; Ri; iC; st 11...	2
3531	3504	Δ. 440	13 18 24.0	3.554	2	136 34 49.8	18.86	2	!!; ⊕; ω Centauri	2+
3532	3505	13 18 27.3	3.370	2	123 9 1.8	18.86	2	vF; S; R; glbM.....	2
3533	1599	III. 402	13 18 29.8	2.730	2	52 53 5.8	18.86	2	cF; cS; R; vsmbM*; *12sp; sp of 2.	3
3534	1600	III. 403	13 18 39.8	2.729	2	52 50 44.8	18.86	2	F; cS; R; vsmbM*; n of 2	3
3535	1600, a	R. nova	13 18 39.8	2.729	::	52 49 ±	18.86	:::	vF	0
3536	IV. 70	13 18 48.6	1.743	2	18 45 22.8	18.86	2	○?; cB; S; R; g; slbM.....	2
3537	1602	II. 667	13 19 4.9	+2.632	1	46 0 37.5	+18.85	1	pB; vS; vIE; glbM	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
3538	III. 115	13 19 8.3	+ 3.167	2	101 35 25.2	+ 18.84	2	vF; vS; stellar	2
3539	1601	II. 25	13 19 11.5	3.050	2	87 9 53.2	18.84	2	pB; pL; vIE; vsmbM * 12...	6
3540	1604	III. 404	13 19 47.8	2.730	2	53 19 49.9	18.83	2	cF; pS; E; bM; sp of 2.....	3
3541	1603	13 19 49.1	2.926	2	72 23 55.6	18.82	2	vF; S; R; * nf	1
3542	3507	13 19 51.7	3.328	2	118 50 27.6	18.82	2	cF; S; R; psbM; * f 2'.....	2
3543	3508	13 20 2.9	3.329	1::	118 54 24.3	18.81	1::	vF; S; R; p of D neb.....	1
3544	3509	13 20 4.1	3.329	1	118 53 52.3	18.81	1	pF; S; f of D neb	1
3545	1605	III. 405	13 20 7.4	2.728	2	53 16 40.6	18.82	2	vF; pL; R; nf of 2.....	3
3546	3506	13 20 8.6	3.977	1	152 41 27.0	18.80	1	Cl; vRi	1
3547	3510	13 20 17.7	3.594	2	138 10 46.0	18.80	2	pB; cS; iE; glbM; r	2
3548	1606	III. 651	13 20 49.5	2.773	3	57 17 26.7	18.79	3	F; pS; vIE; bM	4
3549	1607	13 20 59.9	2.916	1	71 29 31.4	18.78	1	vF; R	1
3550	D'Arrest, 94	13 21 13	3.02	[2]	83 16 42	18.78	[2]	pF. See note.....	0*
3551	3511	13 21 19.2	3.373	1	122 26 25.1	18.77	1	pF; L; vmE; pgbM; rr	1
3552	III. 821	13 21 34.6	2.438	1	36 28 48.1	18.77	1	cF; stellar	1
3553	1609	III. 784	13 21 38.4	2.371	1	33 47 3.1	18.77	1	cF; S; iR	2
3554	1608	13 21 47.4	2.770	3	57 14 35.8	18.76	3	pF; pL; iE; lbM; f of 2.....	3
3555	3512	13 22 3.3	3.898	2	150 12 3.1	18.77	2	Cl; vF; S; vRi; st 15.....	2
3556	1611	13 22 8.6	2.560	1	42 38 21.5	18.75	1	vF; pS; R	1
3557	1610	V. 22	13 22 17.6	3.221	2	107 14 29.2	18.74	2	cF; L; mE 128° 8; pgbM ...	4
3558	1613	13 22 28.5	2.920	2	72 13 18.2	18.74	2	F; pL; R; gbM	2
3559	1614	III. 672	13 22 28.5	2.558	2	42 41 26.2	18.74	2	F; vS; R; stellar.....	5
3560	1612	III. { 45 46 }	13 22 29.1	2.974	2	78 16 4.2	18.74	2	{ vF; pL vF; pL } D neb	4
3561											
3562	1615	III. 71	13 22 37.1	2.922	1	72 26 57.9	18.73	2	vF; S; R; am 3 st; * 7 nf...	3
3563	1616	13 22 49.1	2.953	2	75 58 15.9	18.73	2	vF; S; R	2
3564	3513	13 22 51.6	3.321	2	117 25 21.6	18.72	2	vF; pL; vIE; * 7 nf 10'.....	2
3565	1617	II. 679	13 22 54.8	3.081	1	90 59 58.6	18.72	1	F; cS; iE; gbM; p of 2.....	3
3566	1618	II. 680	13 22 59.8	3.081	1	90 56 3.6	18.72	1	pF; pL; iR; bM; f of 2.....	3
3567	1619	III. 642	13 23 9.3	2.952	1	75 54 40.6	18.72	1	vF; S; iR	2
3568	1620	III. 652	13 23 21.2	2.774	1	58 8 37.3	18.71	2	vF; vS; R; glbM	3
3569	3515	13 23 31.3	3.400	2	124 3 45.0	18.70	2	F; pL; vIE; vglbM	2
3570	3514	Δ. 252?	13 23 41.0	4.129	1	155 15 11.7	18.69	1	l; B; pL; cE; bM curved axis; 4 st inv.	1+
3571	1621	13 23 50.0	2.908	2	71 8 20.0	18.70	2	cF; S; R; bM; * f	2
3572	1622	M. 51	13 23 55.4	2.539	5	42 5 4.0	18.70	4	!!!; nucl & ring (h); spiral (R)	10+
3573	3516	13 23 57.1	3.382	1	122 30 27.7	18.69	1	pB; S; R; g; psbM	1
3574	1623	I. 186	13 24 4.4	2.536	4	42 0 50.7	18.69	3	B; pS; R; gbM; f of 2.....	6
3575	1625	IV. 63	13 24 12.2	2.264	1	30 51 32.7	18.69	1	pB; cL; iR; gmbM; r	2
3576	II. 689	13 24 14.0	2.546	3	42 35 13.7	18.69	2	pB; pL; R; mbM	3
3577	1624	III. 406	13 24 21.5	2.726	2	54 26 31.4	18.68	2	vF; vS; iE	3
5072	13 24 32.0	89 18 29.7	See No. 5072.	
3578	II. 797	13 24 39.9	2.409	2	36 12 12.4	18.68	2	pF; cS; R; vglbM	1
3579	3517	III. 507	13 24 52.2	3.142	1	98 3 15.8	18.66	1	vF; cS; R; gbM; r	2
3580	3518	13 25 18.7	3.610	1	137 24 41.2	18.64	1	F; pL; R; vgbM	1
3581	1626	III. 643	13 25 21.9	2.945	1	75 22 53.5	18.65	1	F; S; cE; * 11 att np.....	2
3582	1627	III. 9	13 25 26.6	3.003	3	81 57 44.2	18.64	4	F; vS; R; psbM; p of 2.....	6
3583	1628	III. 10	13 25 42.8	3.003	2	81 57 21.2	18.64	2	F; vS; R; stellar; f of 2.....	4
3584	1629	III. 99	13 25 48.2	3.004	1	82 6 15.9	18.63	1	F; S; R; psbMN	4
3585	1630	13 25 54.1	3.076	1	90 18 50.9	18.63	1	pB; S; R; psmbM	1
3586	1631	13 26 21.9	3.003	1	81 59 5.6	18.62	1	eF	1
3587	1632	III. 656	13 26 44.5	2.615	1	47 24 36.3	18.61	1	vF; S; R; lbM	2
3588	1633	III. 926	13 27 7.3	3.009	1	82 47 49.7	18.59	1	vF; S; * 9 nf inv? (? 28 ^m R.A.)	3*
3589	3519	13 27 10.2	3.397	1	122 45 39.4	18.58	1	eF; eS; * S and * p	1
3590	1635	II. 841	13 27 14.9	2.075	1	26 33 53.7	18.59	1	pB; S; vIE	3
3591	1634	13 27 18.2	2.904	1	71 25 25.4	18.58	1	vF; pL; R; bM.....	1
3592	1636	II. 842	13 27 21.3	2.072	1	26 30 43.7	18.59	1	pB; pS; R; gbM	3
3593	3520	13 27 47.5	3.583	1?	135 11 16.8	18.56	1::	vF; S; R; * n, nr	1
3594	3521	13 27 59.8	3.399	1	122 44 13.8	18.56	1	vF; S; R; * 10 f.....	1
3595	1637	III. 86	13 28 4.5	2.942	2	75 27 49.8	18.56	2	vF; S; vIE; 1st of 3	4
3596	1638	III. 85	13 28 4.9	+ 2.943	3	75 32 20.8	+ 18.56	3	cF; S; R; bM; 2nd of 3.....	5

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
3597	1639	III. 87	13 28 5.2	+2.943	1	75 37 21.8	+18.56	2	cF; pL; R; glbM; 3rd of 3..	4
3598	1640	III. 407	13 28 6.2	2.713	3	54 34 50.8	18.56	3	F; cS; R; * 10 p; p of 2 ...	5
3599	III. 822	13 28 9.6	2.422	1	37 46 37.8	18.56	1	cF; pS; iR; lbM.....	1
3600	1641	III. 928	13 28 15.0	3.054	(1)	87 53 16.5	18.55	1:	vF; S; R.....	2
3601	1642	III. 408	13 28 16.7	2.711	2	54 29 21.8	18.56	2	vF; vS; R; f of 2	4
3602	1643	13 28 31.7	2.941	1	75 35 24.2	18.54	1	F; L; E; vgbM	1
3603	1645	III. 425	13 28 51.6	2.711	1	54 36 47.9	18.53	1	F; S; R; vS * nr	2
3604	3522	13 28 52.3	3.666	1	139 6 59.6	18.52	1	eeF; S; lE	1
3605	1644	III. 100	13 28 58.3	3.008	3	82 41 27.9	18.53	3	vF; pS; vLE; * 9 sp	4
3606	3523	{ M. 83 = Δ. 628 }	13 29 9.0	3.360	4	119 9 31.6	18.52	4	!!; { (H, h) vB; vL; E 55°-1; esbMN (L) 3-branched spiral. }	6+
3607	3524	13 29 20.8	3.539	4	132 8 2.3	18.51	4	F; pL; cE; vglbM.....	4
3608	1646	III. 101	13 29 27.9	3.000	1	81 54 44.3	18.51	1	vF; pL; R; er.....	4
3609	III. 823	13 29 26.9	2.410	1	37 39 33.6	18.52	1	cF; pL; R; vlbM	1
3610	III. 409	13 29 29.9	2.696	1	53 43 3.3	18.51	1	vF; pL; R; lbM.....	1
3611	1647	13 30 0.9	3.041	1	86 30 48.0	18.50	1	eF; eL.....	1
3612	1648	III. 620	13 30 4.6	2.654	2	50 55 41.0	18.50	2	cF; pL; E 65°; biN?	3
3613	3525	13 30 17.1	3.591	2	135 9 0.4	18.48	2	vF; S; R; vglbM; * 13 att...	2
3614	1649	II. 297	13 30 30.2	3.234	2	107 10 3.1	18.47	2	!!; { (H, h) cF; vL; vg, psmbMLN (L) 2-branched spiral. }	4+
3615	1650	I. 34	13 30 35.1	2.985	1	80 23 32.1	18.47	1	B; L; E 150°; psbMrN	3
3616	1651	III. 72	13 30 48.6	2.916	3	73 18 27.1	18.47	3	vF; S; R; bM.....	4
3617	II. 817	13 30 49.8	2.409	1	38 1 30.1	18.47	1	pB; S; R; vgbM	1
3618	1652	III. 369	13 30 53.0	2.795	1	61 51 45.1	18.47	1	vF; S; vLE	2
3619	1653	III. 505	13 31 11.6	3.025	(1)	84 46 14.8	18.46	1	vF; S; R; bM.....	4
3620	3526	II. 638	Δ. 623	13 31 59.9	3.390	1	120 55 34.6	18.42	1	B; pL; E 45°±; psmbM ...	2
3621	3527	13 32 12.4	3.174	1	100 46 55.6	18.42	1	pB; L; pmE; glbM	1
3622	III. 803	13 32 21.9	2.236	2	32 10 56.6	18.42	2	vF; vS	2
3623	1656	III. 673	13 32 32.9	2.465	1	40 59 35.6	18.42	1	eF; vS; R; gbM.....	2
3624	1654	II. 895	13 32 45.5	3.059	1	88 26 55.0	18.40	1	vF; S; R; bM; p of D neb...	2
3625	1655	II. 896	13 32 49.9	3.058	1	88 27 15.0	18.40	1	F; S; iR; f of D neb	2
3626	1657	13 33 13.6	3.019	2	84 13 5.7	18.39	2	vF; R; am pB st.....	2
3627	1660	13 33 14.4	0.995	1	14 14 14.0	18.40	1	eF; S	1
3628	1658	III. 370	13 33 26.7	2.775	1	60 53 12.4	18.38	1	cF; S; mE 0°±; * 9 sp.....	2
3629	3528	13 33 43.5	3.374	1	119 12 28.8	18.36	1	vF; pL; R; vlbM	1
3630	1659	III. 410	13 33 58.5	2.660	2	52 25 32.8	18.36	2	F; cS; vLE; er.....	3
3631	3529	13 34 27.7	3.665	2	137 27 41.9	18.33	2	B; pL; vLE; vglbM; 3 st nr..	2
3632	1661	13 34 32.8	2.628	4	50 30 0.2	18.34	4	F; S; R; gbM; S * np	4
3633	Auw. N. 32	13 34 43.9	3.196	...	103 9 3.6	18.31	...	A nebula (Markree Obs. 1855)	0
3634	3530	13 35 3.6	4.112	2	152 11 47.3	18.31	2	Cl; P; L; iF; st 12	2
3635	1662	13 35 5.5	3.026	1	85 1 44.6	18.32	1	eF; S; bet 2 st.....	1
3636	1663	M. 3	13 35 40.8	2.769	5	60 55 6.0	18.30	5	!!; ⊕; eB; vL; vsmbM; st 11..	14
3637	1664	I. 98	13 35 56.1	2.670	3	53 37 45.7	18.29	3	cB; pL; R; g, psmbM	4
3638	1664, a	R. nova	13 36 8.6	2.670	::	53 37 45.7	18.29	::	F; S.....	0
3639	1665	II. 798	13 36 24.8	2.248	2	33 37 25.4	18.28	2	F; E 73°-0; D or biN; B * nf	3
3640	3531	Δ. 273	13 36 54.7	4.131	2	152 11 45.2	18.24	2	Cl; B; S; pC; iR; st 10...12	2
3641	3532	13 37 15.3	3.985	1	148 29 24.2	18.24	1	Cl; L; vRi; st 7...16	1
3642	3533	Δ. 388	13 37 37.5	3.757	2	140 40 6.6	18.22	2	⊕; vB; pL; R; rrr; st 15...	2
3643	3534	13 38 48.2	4.237	2	153 59 16.1	18.17	2	Cl; S; C; iR; st 14.....	2
3644	1666	II. 668	13 39 9.3	2.561	1	47 47 33.4	18.18	1	vF; vS; lE 90°±; sbM	2
3645	I. 170	13 39 17.9	2.556	2	47 33 53.1	18.17	2	cB; pL; E 90°±; bMN.....	2
3646	3535	13 39 25.8	3.397	1	119 41 12.8	18.16	1	vF; R; vlbM; * p	1
3647	3536	13 39 41.2	3.405	1	120 13 12.2	18.14	1	pF; S; R; 2 st nr	1
3648	V. 6	13 39 55.7	2.899	1	72 59 14.2	18.14	1	eF; vL; r.....	1
3649	1667	III. 785	13 39 57.1	+2.232	1	34 0 24.5	18.15	1	eF; 2 st att or inv	2
3650	III. 946	13 40 20.5	-0.151	1	9 52 6.7	18.09	1	vF; vS; R	1*
3651	1668, a	R. nova	13 40 ±	+2.511	::	45 28 ±	18.13	::	R; bM; is sp h. 1668	0
3652	1668	I. 180	13 40 31.6	2.511	1	45 27 31.9	18.13	1	cB; L; pmE 142°; gbM.....	2
3653	3538	13 40 39.5	+3.401	1	119 44 21.3	+18.11	1	F; S; R; gbM.....	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	" "			
3654	3537	13 40 51.7	+4.064	1	149 14 42.0	+18.10	1	Cl; vL; vRi.....	1
3655	1669	II. 533	13 41 9.9	3.026	2	85 21 35.0	18.10	2	vF; vL; lE; vgbM	4
3656	1670	II. 688	13 41 13.7	2.456	1::	43 9 25.1	18.07	1	cF; L; vmE	3
3657	3539	13 41 22.7	3.405	1	119 47 20.4	18.08	1	F; S; R; gbM.....	1
3658	1672	III. 681	13 41 40.0	2.605	1	51 0 17.4	18.08	1	pF; cS; lE; F* inv	2
3659	1673	III. 621	13 41 51.2	2.612	2	51 28 9.4	18.08	1	eF; S; R	3
3660	{ 1671 = 3540 }	II. 306	13 41 51.7	3.139	2	96 31 46.8	18.06	2	vF; vS; R; r	3
3661	3541	13 42 11.8	3.782	1	140 30 34.2	18.04	1	○, or vF; eS; D neb	1+
3662	1674	I. 255	13 42 19.9	1.999	1	28 18 58.8	18.06	1	B; pL; mE 57°4; psbMBEN	2*
5073	13 42 38.3	89 14 2.4	See No. 5073.	
3663	1675	II. 710	13 42 57.5	2.571	1	49 19 5.9	18.03	1	cF; cS; R; sbM; p of 2	3
3664	1676	III. 422	13 43 37.6	2.671	2	55 41 13.3	18.01	2	vF; R; stellar; 1st of 3	3*
3665	1677	II. 711	13 43 45.0	2.567	2	49 19 42.0	18.00	2	pB; pS; vIE; glbM; f of 2 ...	3
3666	3542	Δ. 282	13 44 7.0	4.155	1	151 9 39.1	17.97	1	Cl; pL; pC; st 11... ..	1
3667	1678	13 44 8.2	3.014	1	84 18 29.4	17.98	1	vF; vL; R; vgbM	1
3668	1679	III. 423	13 44 22.4	2.667	1	55 36 31.1	17.97	2	F; S; R; psbM; 2nd of 3 ...	3*
3669	1682	II. 669	13 44 27.1	2.539	1	47 56 13.1	17.97	1	cF; pL; R; gbM	2
3670	1680	13 44 28.9	2.668	1	55 39 38.1	17.97	1	eF; pL; R; svmbM*	2
3671	1684	I. 256	13 44 33.4	2.012	1	29 6 16.1	17.97	1	vB; pL; iR; psmbM	2
3672	1689	II. 899	13 44 38.6	0.426	1	12 28 21.7	17.99	1	vF; pS; lE 0°+	2
3673	1681	II. 307	13 44 43.6	3.129	1	95 21 14.5	17.95	1	cF; L; iR; bM	3
3674	1685	II. 712	13 44 51.4	2.570	2	49 44 8.8	17.96	3	cF; S; vIE; sbM	4
3675	1683	II. 685	13 44 52.1	3.089	1	91 30 32.5	17.95	1	F; pS; R; 2 st p	3
3676	3543	III. 923	13 44 56.4	3.388	2	117 46 54.5	17.95	2	pB; S; R; slbM	3
3677	1686	III. 549	13 45 5.2	3.042	1	86 58 26.2	17.94	1	F; vS; R; psbM	2
3678	1687	III. 929	13 45 8.1	3.044	1	87 12 15.2	17.94	1	vF; S; E 0°; rr	3
3679	3544	13 45 39.7	3.737	1	137 48 33.6	17.92	1	vF; vS; R; *8 f	1
3680	III. 665	13 45 44.8	3.078	2	90 25 29.6	17.92	2	cF; vL; R; lbM; r.....	2
3681	1688	13 45 49.7	3.036	1	86 28 58.6	17.92	1	F; iR	1
3682	1690	II. 670	13 46 20.2	2.494	1	46 3 51.0	17.90	1	cF; pL; R; psbM	2
3683	1691	III. 698	13 46 26.0	2.561	1	49 37 37.0	17.90	1	vF; S; iR; B* p.....	3
3684	1694	III. 849	13 46 43.8	2.006	1	29 25 55.7	17.89	1	eF; vS	2
3685	1692	II. 308	13 46 50.5	3.147	1	96 54 11.4	17.88	1	vF; S; R; lbM	3
3686	1693	II. 686	13 47 2.3	3.081	1	90 44 38.1	17.87	1	F; S; R; bM	3
3687	1695	II. 424	13 47 6.9	2.661	1	55 49 32.1	17.87	1	pF; cL; R; lbM	2
3688	1696	II. 713	13 47 25.5	2.544	2	48 56 20.8	17.86	2	cF; pL; bM; B* p; 1st of 4.	5
3689	1697	II. 697	13 47 26.3	2.588	2	51 23 43.8	17.86	2	cF; L; lE 90°; vgbM	3
3690	1697, a	R. 2 novæ	13 47 ±	51 23 ±	{ 2 neb in a line with }	0
3691			
3692	1700	II. 415	13 47 32.2	2.617	2	53 10 5.5	17.85	2	cF; S; R; lbM; *nf 90'' ...	3
3693	1698	II. 714	13 47 32.9	2.545	3:	49 2 24.5	17.85	1	pB; S; R; 2nd of 4	5
3694	1699	II. 715	13 47 33.2	2.545	3:	49 1 24.8	17.86	1	pF; S; R; 3rd of 4	5
3695	1702	III. 699	13 47 45.1	2.543	2	48 57 33.2	17.84	3	vF; pS; 4th of 4	3
3696	1701	III. 506	13 47 56.1	3.008	5	83 58 28.9	17.83	5	F; pL; vmE 15°; r	7
3697	3546	13 47 58.1	3.422	1	119 38 52.6	17.82	1	pF; S; R; glbM; bet 2 st 10.	1
3698	1703, a	R. nova	13 48 4.7	3.009	::	84 2 32.4	17.81	::	F; S; E (nisi=III. 506).....	0
3699	3545	13 48 30.8	4.752	2	159 43 18.7	17.79	2	Cl; vL; lRi; lC; st 11... ..	2
3700	III. 682	13 48 38.9	2.574	1	50 52 52.3	17.81	1	eF; cS; E	1
3701	II. 671	13 48 58.8	2.520	1	48 3 22.0	17.80	1	pB; pL; E	1
3702	1703	I. 6	13 49 5.3	3.009	4	84 3 32.4	17.78	4	B; pL; R; psbM; *8 nf.....	8
3703	1703, b	R. nova	13 49 5.3	3.009	:	84 17 32.4	17.78	:	vF; L	0
3704	1705	II. 534	13 49 12.0	3.011	(1)	84 18 20.4	17.78	2	cF; L; R; gbM	4
3705	3547	13 49 15.5	3.657	2	133 14 39.1	17.77	2	pB; cS; R; pgbM; am st ...	2
5074	13 49 18.6	89 31 9.0	See 5074.	
3706	3548	13 49 20.9	3.581	3	129 17 44.1	17.77	4	l; vB; vL; vl; vsmbM*	6+
3707	1706	III. 786	13 49 24.0	2.192	2	34 58 35.4	17.78	2	F; cS; R; stellar; *16 nf ...	3
3708	1704	III. 285	13 49 33.3	3.125	1	94 48 28.8	17.76	1	vF; vS; R	2
3709	1708	II. 843	13 49 38.6	1.943	1	28 37 31.4	17.78	1	F; S	2
3710	1707	II. 716	13 49 46.8	+2.531	1	48 49 39.8	+17.76	1	pB; L; R; bMFN	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
3711	h. 1709	H. III. 809	h m s 13 49 48.6	s +2.031	1	30 39 35.1	+17.77	1	cF; S; E; ?* inv	2
3712	1710	II. 889	13 50 29.0	2.999	4	83 12 55.9	17.73	4	cF; pL; R; vgbM; *11 np...	5
3713	1711	13 50 36.8	2.715	3	60 8 59.9	17.73	3	pB; pL; R; lbM	3
3714	II. 844	13 50 38.7	1.987	1	29 47 45.9	17.73	1	pB; cL	1
3715	I. 238	13 50 40.4	1.986	3	29 48 15.9	17.73	3	cB; pL; vIE; vgbM.....	3
3716	1712	I. 187	13 50 41.1	2.382	4	42 4 55.9	17.73	4	B; L; mE40°4; smbMN ...	5
3717	1713	13 50 47.4	2.576	1	51 31 33.6	17.72	1	pB; IE; vglbM	1+
3718	1714	II. 698	13 50 56.6	2.579	3	51 42 17.3	17.71	3	F; cS; R; smbM.....	4
3719	3549	13 50 57.8	4.119	2	148 54 29.7	17.69	2	Cl; Ri; vC; pL; st 11...12...	2
3720	I. 239	13 51 6.5	1.973	1	29 35 14.3	17.71	1	vF; pS; E; mbM	3
3721	1715	III. 546	13 51 15.6	2.996	1	83 3 41.7	17.69	2	pB; vS; R; stellar	3
3722	1715, a	R. nova	13 51 +	83 0 +	F; S; R	0
3723	1717	I. 181	13 51 17.9	2.496	1	47 28 19.0	17.70	1	cB; cL; R; gbM	2
3724	1721	13 51 19.2	0.414	1	13 7 58.9	17.73	1	Cl; P; S	1
3725	1716	III. 547	13 51 22.6	2.996	1	82 58 36.7	17.69	2	vF; vS; biN; r; stellar	3
3726	1719	I. 240	13 51 30.0	1.968	1	29 33 27.0	17.70	1	pB; pL; E; mbMN	3
3727	1718	13 51 31.1	2.524	1	48 52 36.7	17.69	1	F; L; vgbM; *9 nf.....	1
3728	1720	III. 666	13 52 10.9	3.101	1	92 31 21.8	17.66	1	vF; cS; R; gbM.....	2*
3729	3550	13 52 33.0	3.414	1	118 11 17.2	17.64	1	vF; S; R; glbM	1
3730	1722	I. 191	13 52 34.9	2.575	2	51 52 3.5	17.65	2	cF; S; np of 2	3
3731	1723	I. 190	13 52 38.2	2.575	3	51 53 52.5	17.65	3	cF; cL; E15°0; lbM; sf of 2	4
3732	III. 125	13 52 39.4	2.709	1	60 11 32.5	17.65	1	vF; S; iR; sbM*	1
3733	3551	13 52 55.7	3.492	1	123 16 22.6	17.62	1	vF; S; R; gbM	1
3734	3552	13 53 10.9	3.479	1	122 23 20.3	17.61	1	pB; pL; R; vgbM	1
3735	1724	III. 411	13 53 24.5	2.618	2	54 32 37.3	17.61	2	eF; vS; pmE90°	3
3736	III. 667	13 53 31.9	3.097	2	92 10 41.0	17.60	2	vF; cS	2
3737	1725	III. 412	13 53 39.8	2.592	1	53 4 6.0	17.60	1	cF; cS; E	2
3738	1727	III. 810	13 53 41.9	1.944	1	29 28 38.3	17.61	1	vF; vS; R	2
3739	1726	III. 683	13 53 51.9	2.557	1	51 8 1.0	17.60	2	vF; pL; iF	4
5075	13 53 58.0	89 13 53	See No. 5075.	
3740	1728	II. 699	13 54 24.3	2.541	2	50 24 2.1	17.57	2	F; pS; R; lbM	5
3741	1732	III. 684	13 54 50.2	2.535	(1)	50 9 17.8	17.56	1	vF; vS; R; bM; in Cl	2
3742	3553	13 54 50.7	3.632	1	130 44 6.2	17.54	1	eF; E bet 2 vS st.....	1
3743	1729	II. 672	13 55 0.0	2.498	1	48 19 36.5	17.55	1	pF; pS; bM.....	2
3744	III. 56	13 55 0.2	2.958	1	79 54 37.2	17.54	1	eF; vS; E; r	1
3745	1733	13 55 10.5	1.647	1	24 24 32.5	17.55	1	pF; pS; R; psbM	1
3746	1730	III. 11	13 55 17.4	2.974	4	81 17 11.9	17.53	4	cF; S; R; psbM; *p	5
3747	1731	13 55 20.0	2.978	1	81 38 22.9	17.53	1	vF; R; bM	1
3748	3554	13 55 24.3	3.501	3	123 17 44.6	17.52	3	pB; pL; R; gpmbM	3
3749	1736	I. 230	13 55 44.4	2.117	2	34 9 20.6	17.52	2	pB; S; pmE45°±; vsvmbMN.	4
3750	1734	II. 309	13 55 59.7	3.134	1	95 18 37.0	17.50	1	pF; cL; R; gmbM; np of 2 ..	3+*
3751	1735	II. 310	13 56 3.7	3.134	1	95 21 25.7	17.49	1	pF; cL; R; sf of 2	3*
3752	1738	II. 827	13 56 14.1	1.945	1	29 58 35.3	17.51	1	pB; S; iE; mbM.....	2
3753	1737	III. 653	13 56 26.8	2.646	2	56 49 5.4	17.48	2	vF; cS; iE0°; bM	3
3754	1739	II. 416	13 56 57.7	2.606	1	54 33 31.8	17.46	1	pF; cS; iE; bM; *11 sp.....	3
3755	1740	13 57 18.2	2.607	1::	54 40 40.5	17.45	1	vF; S	3
3756	1741	II. 417	13 57 19.7	2.598	3	54 11 30.5	17.45	3	pB; pL; ivIE; vsmbM	4
3757	1742	III. 413	13 57 25.7	2.599	1	54 18 33.2	17.44	1	F; *13 p	2
3758	II. 799	H. O. N.	13 57 26.0	2.080	1	33 30 33.5	17.45	1	pF; L; E.....	2
3759	III. 57	13 57 30.0	2.953	1	79 42 30.9	17.43	1	eF; cS	1
3760	1744, a	III. 787?	R. nova?	13 57 30.5	2.137	...	35 5 18.3	17.44	...	B; S; R; gmbM; conn with M. 101.	0
3761	1743	II. 691	13 57 30.6	2.293	1	40 9 16.2	17.44	1	pB; pL; vmE90°±; smbMN	2
3762	1744, b	R. nova	13 57 32.8	2.134	...	35 0 28.3	17.44	...	vF; cL; R; gvlbM } all conn	0*
3763	1744, c	R. nova	13 57 33.0	2.137	...	35 5 51.3	17.44	...	F; pS; iR; glbM } with	
3764	1744, d	R. nova	13 57 41.7	+2.131	...	34 57 41.3	17.44	...	vF; pL; iR; vlbM } M. 101.	
3765	1747	III. 947	13 57 43.7	-0.245	1::	11 5 51.1	17.47	1	vF; pL; iR; vglbM	2
3766	III. 787	13 57 50.1	+2.133	1	35 3 26.9	17.43	1	vF; vS	1+*
3767	1744, e	R. nova	13 57 59.7	2.131	...	35 2 3.3	17.43	...	F; pL; iE; vlbM } conn with	0*
3768	1744, f	R. nova	13 58 2.8	+2.134	...	35 8 16.3	+17.43	...	pB; pS; R; psbM } M. 101.	

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
3769	h.	H.	D'Arrest, 95	h m s	s	[2]	77° 27' 4"	+17'40"	[2]	F; pS	0
3770	1744	M. 101	13 58 12.9	2.127	1	34 57 22.3	17.41	1	pB; vL; iR; g, vsmbMBSN..	5*†
3771	1744, g	R. nova	13 58 14.1	2.131	...	35 2 42.3	17.41	...	vF; pL; R; vlbM; conn with M. 101.	0*
3772	3555	Δ. 431	13 58 39.9	3.807	3	137 38 52.1	17.37	3	Cl; vL; vLC; st 8...	3
3773	1744, h	III. 788?	R. nova?	13 58 49.8	2.125	...	34 59 40.3	17.39	...	B; pS; R; psbM } conn with	{ 0*
3774	1744, i	III. 789?	R. nova?	13 58 53.9	2.122	...	34 56 57.3	17.38	...	pB; pL; iR; gbM } M. 101.	{ 0*
3775	3556	13 59 1.0	3.449	1	119 20 28.8	17.36	1	pF; S; R; psbM	1
3776	1746	VI. 9	13 59 8.7	2.700	2	60 48 11.1	17.37	2	Cl; L; vRi; vmC; st 11...	4
3777	1745	III. 286	13 59 14.1	3.129	1	94 46 56.5	17.35	1	F; L; R; vgbM	3
3778	III. 788	13 59 19.3	2.120	1	34 59 23.1	17.37	1	vF; vS; 2nd of 3 } inv in {	1†
3779	III. 789	13 59 26.2	2.118	1	34 57 22.8	17.36	1	vF; vS; 3rd of 3 } M. 101. {	1†
3780	III. 58	13 59 30.1	2.955	1	79 57 25.5	17.35	1	eF; S; 1E.....	1
3781	1745, a	R. nova	13 59 30.2	3.129	::	94 46 56.5	17.36	::	pB; S; E.....	0
3782	1748	I. 231	13 59 50.7	2.096	1	34 25 50.5	17.35	1	pB; S; R; gbM	3
3783	I. 214	13 59 59.3	2.149	2	35 40 25.5	17.35	2	pB; L; bM	2
3784	1750	II. 800	14 0 20.7	2.061	1	33 35 24.6	17.32	1	pB; S; pmE; bM	2
3785	1749	14 0 34.6	2.994	1	83 17 26.0	17.30	1	F; mE; vglbM	1
3786	1751	III. 287	14 0 49.1	3.138	(2)	95 25 38.7	17.29	1::	F; pS; iR	3
3787	III. 790	14 0 54.3	2.104	1	34 52 19.0	17.30	1	vF; pL	1
3788	III. 762	14 0 55.8	3.085	1	91 1 21.4	17.28	1	vF; vS	1
3789	II. 692	14 1 24.6	2.225	1	38 37 18.1	17.27	1	F; pS; vgbM; np of 2	1
3790	II. 693	14 1 42.5	2.223	1	38 37 47.8	17.26	1	F; vS; smbM; stellar; sf of 2	1
3791	III. 59	14 1 43.2	2.958	1	80 24 17.5	17.25	1	eF; S	1
3792	3557	14 1 46.3	3.704	1	132 39 20.2	17.24	1	pF; vL; R; vgbM	1
3793	III. 791	14 2 26.5	2.073	1	34 19 14.9	17.23	1	vF; S; R; p of 2	1
3794	I. 232	14 2 26.5	2.073	1	34 19 14.9	17.23	1	cB; R; vgbM; f of 2	1
3795	II. 801	14 2 34.8	2.071	2	34 16 45.6	17.22	2	F; pL	2
3796	3558	14 2 45.6	3.520	1::	122 58 35.7	17.19	1::	F; R; *8 s nr	1
3797	3559	14 3 10.0	3.775	2	135 25 33.1	17.17	2	vF; S; R; bM.....	2
3798	1752	III. 32	14 3 17.4	2.846	1	71 47 18.4	17.18	2	cF; cS; R; sbMF*	4
3799	1753	II. 890	14 3 59.6	2.988	3	82 58 10.5	17.15	3	pB; pS; R; gbM; r	5
3800	1754	II. 876	14 4 1.8	2.817	1	69 43 45.5	17.15	1	pB; vS; E	3
3801	1755	IV. 46	14 4 12.2	3.126	1	94 22 49.9	17.13	1	pB; vS; R; psmbM*; *18 inv.	2
3802	3560	14 4 16.2	3.475	1	119 59 46.6	17.12	1	pB; L; R; gbM; rr	1
3803	3561	14 4 23.5	3.419	1	116 26 51.6	17.12	1	vF; S; R; bM; *sf	1
3804	III. 674	14 5 5.6	2.265	2	40 46 9.3	17.11	2	cF; cS; iR	1
3805	1756	14 5 13.3	3.050	1	88 5 4.7	17.09	1	vF; S; rr	1
3806	1757	II. 687	14 5 59.1	3.105	1	92 32 43.5	17.05	1	pB; L; E 20° ±; lbM	3
3807	1757, a	R. nova	14 6 ±	3.105	...	92 33 ±	17.05	...	3' dist from h. 1757	0
3808	1758	IV. 49	14 6 5.9	3.104	1	92 29 28.5	17.05	1	cF; S; R; stellar	3
3809	1759	II. 877	14 6 36.4	2.801	1	68 54 40.9	17.03	1	pB; pL; iR	2
3810	1760	III. 685	14 6 51.3	2.482	3	50 1 53.6	17.02	3	vF; S; vLE	5
3811	3562	14 6 54.8	3.848	2	137 27 28.0	17.00	2	pF; S; R; psbM; S *nf	2
3812	III. 676	14 7 49.9	2.196	2	38 59 31.4	16.98	2	F; S; 1E; stellar	2
3813	1761	14 8 20.7	3.010	1	84 56 4.5	16.95	1	F; S; R; bM	1
3814	III. 644	14 8 23.6	2.871	1	74 13 59.5	16.95	1	vF; vS; E; a D neb	1
3815
3816	1762, a	R. 3 novæ	14 8 ±	2.725	...	64 0 ±	16.94	...	3 "knots" near h. 1762	0
3817
3818	1762	III. 134	14 8 31.0	2.725	3	64 0 55.2	16.94	3	F; pL; pmE 90°; *10 np	5
3819	1764, a	R. nova	14 8 47.0	2.53	::	52 56 57.7	16.93	::	eeF	0
3820	1763	III. 804 = III. 835	14 9 17.5	1.899	1	31 34 24.3	16.91	1	vF; S; E; r	3*
3821	1764, b	R. nova	14 9 18.9	2.53	::	53 6 33.7	16.89	::	vF	0
3822	1764	III. 414	14 9 40.1	2.533	1	53 7 33.7	16.89	1	cF; pL; vmE 110° 3; vgvmbM.	2
3823	3563	14 9 41.8	3.742	1	132 43 31.1	16.87	1	l; vF; pmE; esvmbM *12	1
3824	D'Arrest, 96	14 9 53	2.92	[2]	78 28 0	16.87	[2]	F; S; R; III. 47 f 10° 4	0
3825	1765	III. 47	14 10 2.1	+2.926	1	78 32 18.1	+16.87	1	vF; vS; R; gbM; r	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
3826	h. 1766	H. II. 418	h m s 14 10 10.3	s +2.549	3.	53 59 51.1	+16.87	3	pB; R; vsmbM; 2 or 3 st inv	4
3827	1768	III. 731	14 10 40.3	2.462	1	49 51 34.2	16.84	1	cF; vS; R; p of 2	2
3828	1767	14 10 43.5	2.960	1	81 10 12.2	16.84	1	F; pL; iF; gbM	1
3829	III. 805	14 10 44.8	1.789	3	29 20 6.5	16.85	3	eF; vS; R; stellar	3
3830	1770, a	R. nova	14 10 45.3	2.968	::	81 39 52.2	16.85	::	vF	0
3831	1769	III. 732	14 10 48.4	2.459	3	49 45 41.2	16.84	4	cF; S; R; gbM	5
3832	1770, b	R. nova	14 10 53.9	2.968	:	81 46 46.3	16.83	:	vF	0
3833	1771	II. 419	14 11 8.5	2.521	2	52 46 41.6	16.82	3	F; pS; E80°; D or biN	4
3834	1771, a	R. nova	14 11 ±	2.521	...	52 46 ±	16.82	...	Makes D or biN neb with h. 1771.	0
3835	1770	14 11 13.9	2.968	2	81 46 56.3	16.81	2	pB; cS; gbM	2
3836	III. 551	14 11 29.6	+2.970	2	81 59 ±	16.80	2	2*
3837	III. 948	14 11 36.1	-0.747	1	10 44 46.7	16.79	1	eF; vS; E0°±	1
3838	1773	II. 194	14 11 38.1	+2.720	2	64 12 39.0	16.80	2	cF; pS; R; vsmbM*	4
3839	1772	III. 552	14 11 43.6	2.970	1	81 58 40.7	16.79	1	vF; vS; R	2
3840	1774	14 11 43.8	2.897	1	76 27 52.7	16.79	1	vF; cS; pmE	1
3841	1775	14 12 12.9	2.701	1	63 4 49.8	16.76	1	vF; S; lE	1
3842	3564	14 12 24.5	3.475	1	118 36 20.5	16.75	1	eF; L; S* inv	1
3843	1776	I. 99	14 12 32.5	2.517	2	52 51 11.8	16.76	2	cB; S; R; vsbM*	4
3844	1777	III. 347	14 12 50.8	2.723	1	64 32 41.9	16.73	1	vF; S; vLE; bM	2*
3845	1778	II. 579	14 13 1.0	3.013	2	85 21 16.6	16.72	2	pF; cL; E; gbM	3
3846	1779	I. 144	14 13 16.5	3.014	3	85 25 7.3	16.71	3	B; pL; R; psbM; r; *12 nf.	4*
3847	1780	14 13 20.8	2.541	1	54 13 56.6	16.72	1	pF; R	1
3848	1779, a	R. nova	14 13 28.5	3.014	...	85 24 7.3	16.72	...	Place from MS.	0
3849	1781	III. 12	14 13 31.7	2.967	(1)	81 50 21.0	16.70	2	F; S; iR	4
3850	1782	I. 145	14 13 52.1	3.023	1	86 6 53.4	16.68	1	pF; pS; lE; p of 2	2
3851	1783	I. 146	14 14 1.1	3.022	1	86 5 8.4	16.68	1	B; S; R; vsmbM; f of 2	2
3852	1784	III. 415	14 14 31.5	2.536	2	54 9 46.8	16.66	2	vF; cL; p of 2	3
3853	1785	14 14 45.4	2.535	1	54 8 52.2	16.64	1	pB; S; f of 2	1
3854	1786	II. 754	14 15 1.5	2.440	3	49 39 28.2	16.64	3	pB; pS; R; bMFN; *sp	5
3855	1783, a	R. nova	14 15 5.4	3.02	...	85 53 50	16.64	...	L; F; vmE	0
3856	1790	I. 235	14 15 18.1	1.909	1	32 38 5.9	16.63	1	pF; L; iR; vgmbM; r	3
3857	1787	III. 110	14 15 30.3	2.879	1	75 26 38.0	16.60	1	F; cS; vLE; *8 sf	3
3858	1789	14 15 30.7	2.534	1	54 13 57.3	16.61	1	vF; R; gbM; *8 sf	1*
3859	1788	III. 416	14 15 30.8	2.531	2	54 5 13.3	16.61	2	vF; S; R; np of 2	3*
3860	1791	III. 417	14 15 43.3	2.532	3	54 9 17.0	16.60	3	cF; S; R; bM*; sf of 2	4*
3861	3565	III. 924	14 15 47.9	3.474	1	118 1 55.4	16.58	1	F; S; E; gvlbM; r	2
3862	3566	Δ. 357	14 16 13.4	4.121	1	144 9 51.1	16.55	1	Cl; vlRi; vlC; st 10	1
3863	III. 135	14 16 25.3	2.689	1	62 58 37.8	16.56	1	eF; vS; stellar	1*
3864	1792	III. 121	14 16 32.5	3.291	1	106 5 6.5	16.55	2	F; pL; R; vgbM; p of 2	4
3865	1795	III. 418	14 16 41.3	2.489	1	52 14 23.5	16.55	1	eF; S; R; stellar	2
3866	1793	III. 122	14 16 46.6	3.292	1	106 7 34.2	16.54	2	vF; L; vLE; vglbM; f of 2	4
3867	1796	III. 733	14 16 49.7	2.419	1	49 2 30.5	16.55	1	F; vS; R; bM	3
3868	1794	III. 927	14 16 53.9	2.978	4	82 46 50.2	16.54	4	F; S; lE	5
3869	1797	II. 177	14 17 7.8	2.866	1	75 43 22.6	16.52	1	pB; pS; gbM	3
3870	II. 694	14 17 17.0	2.134	1	38 49 32.9	16.53	1	pF; pS; lE; mbM	1
3871	1800	III. 734	14 17 24.4	2.414	2	48 58 44.6	16.52	2	cF; pS; R; gbM	3
3872	1799	III. 668	14 17 28.1	3.107	1	92 34 12.0	16.50	1	F; pS; R; vgbM*; r	2
3873	{ 1798 = 3569 }	III. 120	14 17 32.1	3.243	2	102 32 27.0	16.50	2	vF; pL; R; vgbM	3
3874	3568	Δ. 313	14 17 36.5	4.345	1	148 59 58.4	16.48	1	Cl; S; pC; st L & S	1
3875	II. 331	14 17 43.9	0.726	2	17 47 31.9	16.53	2	pF; cS; iR; bM; er	2
3876	1801	II. 673	14 17 48.1	2.380	1	47 35 13.0	16.50	1	F; pL; lE; vglbM	2
3877	1802	III. 136	14 18 3.6	2.714	1	64 45 34.4	16.48	1	vF; S; pmE0°±; *9 f	4
3878	1803	14 18 6.3	2.566	1	56 18 59.4	16.48	1	F; S; R; bM	1
3879	3567	14 18 9.9	6.643	1	167 46 0.3	16.41	1	vF; E; gbM; r	1
3880	1804	II. 420	14 18 12.4	2.530	2	54 29 56.4	16.48	2	pB; S; R; smbM	3
3881
3882	1804, a	R. 3 novæ	14 18 ±	+2.530	...	54 30 ±	+16.48	...	h. 1804 is D; 2 others near ..	0
3883

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
3884	h. 1805	H. III. 419	14 18 31.3	+2.496	1	52 54 5.8	+16.46	1	vF; S; cF; vgbM; er.....	2
3885	3570	Δ. 302	14 19 23.2	4.417	3	150 5 16.7	16.39	3	Cl; L; pRi; pCM; st 8... ..	3
3886	III. 763	14 20 1.3	3.095	1	91 37 28.4	16.38	1	eF; S	1
3887	1806	14 20 15.9	3.000	1	84 33 53.8	16.36	1	vF; S; R; vgbM	1
3888	III. 319	14 20 17.6	0.680	1	17 44 24.0	16.40	1	eF; vS	1*
3889	1807	III. 14	14 20 54.8	2.952	1	81 7 26.2	16.34	1	eeF; L; r.....	2
3890	1809	III. 677	14 21 3.9	2.177	1	40 48 47.2	16.34	1	vF; pS; vLE; vglbM	2
3891	1808	II. 329	14 21 12.5	2.552	1	56 7 10.9	16.33	1	vF; S; R; vsmbM; r	4
3892	1810	14 21 25.6	2.408	1	49 24 57.6	16.32	1	vF; S; R; gbm	1
3893	3571	14 21 35.0	3.508	1	119 7 31.7	16.29	1	eF; S; R	1
3894	1811	14 21 47.6	2.907	1	77 59 10.7	16.29	1	vF; vS; R; *9sp	1
3895	1812	14 21 59.6	2.685	2	63 31 30.7	16.29	2	pF; S; R; gbm	2
3896	1814	II. 674	14 22 4.2	2.374	4	48 6 50.7	16.29	4	F; S; E 90°±; gbm	5
3897	1820	I. 236	14 22 10.2	1.865	2	32 47 47.7	16.29	2	B; S; R; psbMrN	5
3898	Auw. N. 33	14 22 12.0	3.068	...	89 49 7.8	16.26	...	Neb *11f 150 ^a (Bond, May, 1853).	0
3899	1818	I. 185	14 22 14.2	2.243	2	43 13 13.4	16.28	2	cB; pS; R; pglbM	4
3900	1813	I. 70	14 22 14.9	3.146	1	95 20 26.8	16.26	1	⊕; vB; cL; R; gbm; rrr; st 19; *17 sf.	3
3901	1815	III. 132	14 22 20.8	2.657	1	61 57 59.1	16.27	1	F; S; E; sbM	2
3902	1816	II. 580	14 22 34.6	3.020	1	86 5 58.2	16.24	1	eF; cL; R; np of 2	2
3903	1819	II. 357	14 22 35.7	2.728	2	66 11 4.5	16.25	2	vF; S; R; vgbM	3
3904	1817	II. 581	14 22 36.3	3.020	2	86 8 14.2	16.24	2	cB; pL; R; sf of 2	3
3905	1817, a	R. nova	14 23 +	3.020	...	86 8 ±	16.24	...	Makes a BD neb with h. 1817	0
3906	1821	14 22 41.4	+2.602	1	58 58 19.5	16.25	1	vF; R; *7 p; *11s	1
3907	III. 949	14 23 2.9	-1.635	1	9 18 14.0	16.30	1	eF; S; IE	1
3908	1822	III. 126	14 23 11.4	+2.608	2	59 21 35.9	16.23	2	cF; S; * inv; *12 nf	3
3909	3572	Δ. 469	14 23 40.3	3.826	2	133 34 30.1	16.17	2	pB; L; R; vglbM; st inv ...	2
3910	1823	II. 150	14 23 43.8	2.964	2	82 5 56.7	16.19	2	cF; pL; iR; gbm	5
3911	1824	III. 645	14 23 54.1	2.867	1	75 22 18.4	16.18	1	eF; vS; np of 2	2
3912	Auw. N. 34	14 24 3.0	3.066	...	89 42 5.1	16.17	...	Neb R (Bond, May, 1853)...	0
3913	1825	II. 891	14 24 4.0	2.982	2	83 23 45.1	16.17	2	pB; pL; vLE; bM	3
3914	1826	II. 330	14 24 7.8	2.583	1	58 9 43.1	16.17	1	pF; pS; R; bM	2
3915	1828	III. 420	14 24 14.9	2.478	1	53 0 46.8	16.16	1	F; S; E?; * inv?	2
3916	1827	14 24 17.5	2.868	1	75 27 56.8	16.16	1	eeF; sf of 2	1
3917	1829	II. 421	14 24 34.5	2.499	3	54 2 59.5	16.15	4	pF; pL; R; mbM; r	5
3918	Auw. N. 35	14 24 48.0	3.067	...	89 45 4.8	16.16	...	Neb; F; E (Bond, May, 1853)	0
3919	1831	14 24 52.7	2.686	1	63 59 8.9	16.13	1	eF;	1
3920	1832	II. 695	14 24 53.3	2.122	1	39 45 49.2	16.14	1	pB; L; iR; vgbM	2*
3921	1830	II. 892	14 24 59.9	2.977	2	83 7 34.6	16.12	2	vF; pS; iE	3
3922	3573	Δ. 342	14 25 7.9	4.256	2	145 56 31.0	16.10	2	Cl; L; pRi; iC; st 9... ..	2*
3923	1833	II. 27	14 25 31.6	2.951	2	81 18 15.0	16.10	2	pB; pL; R; gbm; r	5
3924	1834	14 26 19.7	2.916	1	78 51 33.5	16.05	1	vF; vS; R; stellar	1
3925	II. 807	14 26 22.6	1.686	1	29 54 6.1	16.07	1	pB; pS; E 0°	1
3926	1835	II. 574	14 26 22.9	3.002	2	84 55 54.5	16.05	2	F; pS; vLE; *14 inv	3
3927	II. 79	14 26 28.2	2.924	1	79 28 9.2	16.04	1	F; L; R; lbM; r	1
3928	3574	14 26 28.4	3.887	2	135 20 29.9	16.03	2	vF; S; cE; bet 2 st	2
3929	III. 882	14 26 35.4	0.855	1	19 44 5.4	16.08	1	vF; pL; R; bM	1
3930	1836	III. 310	14 26 39.7	2.563	1	57 43 21.2	16.04	1	vF; vL; iR; lbM; * p	3
3931	1838	II. 696	14 26 42.1	2.100	1	39 26 14.5	16.05	1	F; S; cE; *15 np	2
3932	1837	II. 893	14 26 53.5	2.988	4	83 55 1.9	16.03	4	cF; pS; R; gbm	5
3933	1839	II. 422	14 26 53.6	2.470	1	53 4 35.9	16.03	1	F; pS; E; bM	2
3934	1843	I. 237	14 27 51.2	1.758	1	31 27 43.7	15.99	1	B; L; IE 0°; vgbmB	3
3935	1842	I. 189	14 27 51.3	2.110	1	39 55 14.7	15.99	1	B; L; E 45°±; pgbM; r ...	2
3936	1840	III. 283	14 27 55.2	2.678	1	63 54 56.1	15.97	1	vF; vS; R; r; 3 st 9, 10 np...	2
3937	1841	II. 894	14 28 8.6	2.988	2	84 1 3.8	15.96	4	vF; S; R; *12 att	6
3938	1844	III. 421	14 30 6.5	2.454	4	52 50 57.8	15.86	4	F; cS; R; bM; p of 2	5
3939	1845	14 30 18.9	2.454	1	52 53 2.5	15.85	1	vF; S; R; f of 2	2
3940	1849	II. 808	14 30 26.8	1.902	1	34 53 55.5	15.85	1	pF; S; iF; r; *10 f	2
3941	3575	14 30 28.4	3.880	1	134 25 46.3	15.81	1	F; S; vgbM; am st	1
3942	1848	I. 188	14 30 29.8	+2.120	1	40 38 43.5	+15.85	1	cB; S; E 90°±; psmbM ...	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
3943	1848, a	R. 3 novæ	14 30 ±	+2.120	...	40 38 ±	+15.85	...	3 novæ, one mottled	0
3944				14 30 39.7	3.032	1	87 6 6.6	15.82	3	vF; mE or biN 140° ±; *6.7 p	4
3945				14 30 42.5	3.070	1	89 46 46.6	15.82	1	pB; pS; IE; gbM	3
3946	1846	II. 582	14 30 39.7	3.032	1	87 6 6.6	15.82	3	vF; mE or biN 140° ±; *6.7 p	4
3947	1847	II. 681	14 30 42.5	3.070	1	89 46 46.6	15.82	1	pB; pS; IE; gbM	3
3948	1851	II. 423	14 31 38.1	2.448	2	52 49 33.4	15.78	3	pB; cS; R; bM; r	4
3949	1851, a	R. 2 novæ	14 32 ±	2.488	...	52 50 ±	15.78	...	No description	0
3950				14 32 2.2	2.488	...	52 50 ±	15.78	...	No description	0
3951				14 31 38.2	2.321	(2)	47 36 0.4	15.78	1	cF; cS; R; lbM; r	3
3952	1853	II. 675	14 31 38.8	2.325	1::	47 45 53.4	15.78	1::	F; vS; R; bM; 4B st p	2
3953	1852	II. 700	14 31 38.9	2.404	2	50 55 57.4	15.78	3	cF; cS; IE; in Δ of st	4
3954	3576	II. 196	14 32 4.2	3.477	1	115 55 37.2	15.74	1	cB; cS; R; psbM; r; * nr ...	3
3955	III. 127	14 32 12.5	2.594	1	59 53 49.2	15.74	1	eF; vS	1
3956	1854	II. 575	14 32 12.8	2.986	4	84 1 23.2	15.74	4	cB; pS; R; mbM; *15 p ...	5
3957	III. 894	14 32 20.6	2.754	2	68 53 19.9	15.73	2	vF; vS	1
3958	III. 128	14 32 26.5	2.593	1	59 53 48.9	15.73	1	vF; vS; iR	1
3959	1855	II. 649	14 32 26.6	2.351	1	48 52 33.2	15.74	1	F; cS; IE 0° ±	4
3960	1859	14 32 46.2	2.351	2	48 56 49.6	15.72	2	F; pL; E 0° ±; gbM	2
3961	1856	III. 895	14 32 50.2	2.760	1	69 21 2.0	15.70	1	vF; S; vgbM; * f; p of 2 ...	2
3962	1858	14 32 55.7	+2.761	1	69 24 52.0	15.70	1	eF; vS; * att; f of 2	1
3963	III. 950	14 32 57.9	-1.318	1	10 33 44.4	15.78	1	vF; S; R; S Cl p	1
3964	1857	I. 182	14 32 59.9	+3.068	1	89 40 43.0	15.70	1	cB; pL; R; psmbM; r	3
3965	1861	III. 675	14 33 2.8	2.175	1	42 44 30.0	15.70	1	vF; pS; iE; * n; 1st of 3 ...	2
3966	3577	Δ. 333	14 33 15.1	4.353	1	146 56 49.8	15.66	1	Cl; L; pRi; CM; st 11...13...	1
3967	VI. 8	14 33 16.5	3.198	1??	98 30 48.4	15.68	1??	Cl; pL; eRi; vmC	1*
3968	1860	III. 671	14 33 19.0	3.326	2	106 52 1.1	15.67	2	vF; pL; R	3
3969	1864	14 33 29.2	2.172	2	42 42 29.7	15.69	1	vF; S; R; * nr; 2nd of 3 ...	2
3970	1862	III. 550	14 33 39.6	3.014	1	85 56 1.8	15.66	2	vF; S; R; vglbM; *8.9 nf ...	3
3971	1863	II. 682	14 33 44.9	3.069	1	89 42 24.5	15.65	1	pF; S; IE; bM	3
3972	1865, a	R. 3 novæ	14 34 ±	2.170	...	42 41 ±	15.67	...	h. 1865 is quadruple; ? F neb connecting.	0
3973				14 33 49.6	2.170	1	42 40 59.1	15.67	1	vF; S; R; psbM; 3rd of 3 ...	1
3974				14 33 53	3.03	[1]	87 12 42	15.64	[1]	○?; vF; S; disc; *15 n 95'' ...	0
3975	1865	14 33 49.6	2.170	1	42 40 59.1	15.67	1	vF; S; R; psbM; 3rd of 3 ...	1
3976	D'Arrest, 97	14 33 53	3.03	[1]	87 12 42	15.64	[1]	○?; vF; S; disc; *15 n 95'' ...	0
3977	1866	I. 184	14 34 38.1	3.325	2	106 38 34.0	15.60	2	pF; pL; pmE 45° ±; mbM; *10s.	3*
3978	3578	III. 508	14 34 40.0	3.197	2	98 24 7.0	15.60	2	F; pL; E; r	3
3979	1867	III. 657	14 34 41.8	2.281	1	46 36 11.3	15.61	1	vF; cS; E 90° ±	2
3980	1868	III. 658	14 34 51.9	2.281	1	46 37 41.0	15.60	1	vF; eS; IE	2
3981	1869	III. 686	14 35 2.8	2.387	2	50 45 12.7	15.59	2	vF; S; R; lbM	3
3982	1870	III. 133	14 36 29.1	2.598	1	60 40 46.3	15.51	1	vF; L; iR; lbM	2
3983	1871	III. 896	14 36 42.8	2.773	1	70 31 5.7	15.49	1	vF; cS; R; vglbM	2
3984	1873	I. 171	14 37 12.6	2.297	1	47 34 26.1	15.47	1	pB; S; R; smbM; r; * nr ...	3
3985	1872	II. 538	14 37 17.6	3.039	1	87 43 15.8	15.46	1	pB; L; iR; gbM; r	2
3986	3579	14 37 23.1	3.276	1	103 20 50.5	15.45	1	vF; S; E; psbM	1
3987	1874	I. 126	14 37 48.7	3.035	1	87 26 59.9	15.43	1	B; L; vmE; bMBN	2
3988	III. 48	14 38 25.5	2.880	1	77 18 32.0	15.40	1	eF; S	1
3989	3580	Δ. 356	14 38 57.4	4.258	1	143 56 9.2	15.34	1	Cl; pL; pRi; IC; st 10...11...	1
3990	1875	I. 183	14 39 9.0	3.068	1::	89 38 40.5	15.35	1::	pF; pS; vLE; r	3
3991	1877	II. 809	14 39 24.3	1.892	1	35 59 5.8	15.36	1	F; S; vLE; Δ 2 st 10.11 ...	2
3992	1878	III. 687	14 39 47.5	2.368	3	50 40 19.9	15.33	3	cF; cS; R; bM	4
3993	3581	14 39 54.0	3.293	1	104 15 56.3	15.31	1	pB; pL; pmE; gpmbM	1
3994	1876	III. 690	14 39 54.2	3.362	1	108 29 45.3	15.31	1	vF; S; iR; lbM	2
3995	1879	III. 885	14 41 13.2	2.773	1	70 54 33.2	15.24	1	vF; vS; cE 90°; vglbM	2
3996	3582	14 43 44.7	4.208	1	142 5 25.1	15.07	1	Cl; vF; vS; vC	1
3997	1880	14 43 52.3	2.985	1	84 17 59.7	15.09	1	D neb; both eF	1
3998	III. 373	14 44 58.6	3.104	3	91 57 9.6	15.02	3	F; R; bMFN; S* s	3*
3999	1881	II. 576	14 46 13.3	3.003	1	85 27 35.5	14.95	1	eF; S; vLE; bM; ? biN	4*
4000	1882	III. 129	14 46 16.8	2.551	1	59 35 3.5	14.95	1	vF; S; R; pgbM	2
4001	1883	14 46 17.5	2.295	3	48 49 8.8	14.96	3	pB; pL; IE; psbM; *8 np ...	3
4002	1884	III. 130	14 46 34.3	+2.551	1	59 37 52.9	+14.93	1	vF; S; R; pgbM	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
4003	h. 1885, <i>a</i>	H.	R. nova	h m s 14 46 50.9	s +3.009	::	85 53 7.0	+14.91	:	eF; 2' p h. 1885	0
4004	1885	III. 554	14 46 58.9	3.009	1	85 53 7.0	14.90	1	F; pS; vME 148° 4; gvlbM...	4
4005	D'Arrest, 98	14 47 33	3.02	[1]	86 27 8	14.87	[1]	vF; pL; vlbM; *8.9p; 225'' s	0
4006	III. 806	14 47 33.0	1.555	1	30 26 56.7	14.89	1	vF; vS; lE	1
4007	1886	14 48 52.2	3.344	1	106 40 4.7	14.79	1	F; S; R; bM; *16 sp.....	1
4008	1887	II. 676	14 49 5.8	2.229	2	46 52 39.0	14.80	2	pB; S; R; smbM; stellar ...	3
4009	3583	14 49 57.8	3.877	1	131 27 21.3	14.71	1:	F; mE; L* sf	1
4010	1888	II. 677	14 50 2.4	2.227	1	46 56 4.2	14.74	1	F; cS; R; psbM	2
4011	1890	III. 976	14 50 41.1	2.532	1	59 12 55.0	14.70	1	eF; pS; iF	2
4012	1889	III. 691	14 50 51.7	3.382	1	108 42 27.1	14.67	1	pF; S; R; stellar.....	2
4013	II. 683	14 51 12.5	3.081	1	90 31 47.8	14.66	1	pB; pL; R; mbM; cB*npatt	1
4014	1891	14 51 21.5	1.979	1	39 44 42.1	14.67	1	pF; S; vsbM*13; 1st of 3 ...	1
4015	1893	III. 678	14 51 46.4	1.977	1	39 44 55.2	14.64	1	F; S; vsbM*13; 2nd of 3 ...	2
4016	1892	III. 131	14 51 48.3	2.534	1	59 28 16.9	14.63	2	F; S; R; vgbM; *nf (2 var)	4*
4017	3584	14 51 49.4	5.894	1	161 52 27.1	14.57	1	eF; S; R; bM.....	1
4018	3585	14 52 5.8	4.224	1	141 21 29.4	14.58	1	Cl; pL; pRi; lC	1
4019	1895	III. 679	14 52 31.1	1.973	1	39 45 19.0	14.60	1	vF; vS; vsmbM*13; *6 nr; 3rd of 3.	2
4020	1895, <i>a</i>	R. nova	14 52 ±	1.973	...	39 45 ±	14.60	...	S	0
4021	1894	II. 539	14 52 53.9	+3.034	1	87 32 48.8	14.56	1	cB; cL; E 165° ±; sbMN ...	2
4022	III. 311	14 53 0.6	-0.023	1	16 24 35.3	14.61	1	vF; S; iR; bet 2 st	1
4023	3586	14 53 11.2	+3.296	1	103 36 47.9	14.53	1	vF; S; E; glbM	1
4024	3587	I. 71	14 53 28.2	3.185	2	96 53 29.6	14.52	2	cB; S; R; svmbM	4
4025	II. 756	14 53 51.4	1.781	2	35 37 32.1	14.48	2	pF; pS; iE; r	2*
4026	1896	I. 127	14 54 5.4	3.037	1	87 44 16.7	14.49	1	B; pS; R; psmbM	2
4027	1897	14 54 15.4	3.038	1	87 49 46.1	14.47	1	vF; vS; R	1
4028	III. 811	14 54 22.9	1.784	1	35 35 32.7	14.49	1	vF; S; E	1
4029	1898	II. 756?	14 54 29.9	1.781	1	35 32 4.1	14.47	1	B; R; sbM; splendid * f ...	1*
4030	1898, <i>a</i>	R. nova	14 54 50.6	1.781	::	35 32 4.1	14.47	::	vF	0
4031	3588	14 55 2.0	4.347	1	143 47 24.0	14.40	1	Cl; vL; Ri; lC; st 9...12 ...	1
4032	3589	14 55 20.2	4.409	1	145 2 17.4	14.38	1	Cl; cL; Ri; lCM; st 13...14	1
4033	1899	II. 540	14 57 2.4	3.045	(1)	88 14 41.0	14.30	1	pB; S; mbM	2
4034	II. 332	14 57 32.3	0.165	2	17 45 19.6	14.32	1	pB; cL; iR; bp; r	2
4035	3590	14 57 32.5	6.045	3	162 19 4.3	14.21	3	F; cS; lE; glbM; am st	3
4036	1900	14 57 46.9	+3.675	1	122 34 35.2	14.24	1	eeF(?)	1
4037	III. 312	14 57 54.8	-0.262	1	15 36 18.3	14.31	1	eF; vS; lE; 2 st inv	1
4038	II. 542	14 58 21.5	+3.030	1	87 21 21.6	14.22	1	pB	1
4039	II. 541	14 58 21.9	3.037	1	87 49 21.6	14.22	1	F	1
4040	3592	14 58 41.7	3.755	2	125 47 8.4	14.18	2	vF; S; lE; vlbM; r	2
4041	3591	14 58 51.0	5.033	2	154 8 9.5	14.15	2	pB; pL; R; vglbM	2
4042	III. 511	14 59 1.9	3.036	1	87 46 20.4	14.18	1	vF; R; p of 2	1
4043	1901, <i>a</i>	R. 2 novæ?	14 59 ±	3.038	...	87 51 ±	14.16	...	2 of 6	*
4044											
4045	1901	I. 128	14 59 23.5	3.038	1	87 50 41.8	14.16	1	vB; pL; R; psbMN; f of 2...	2
4046	D'Arrest, 99	14 59 29	3.03	[1]	87 26 48	14.15	[1]	eF; S; v close * f 7'	0
4047	1902	II. 543	15 0 2.6	3.039	1	87 54 47.6	14.12	1	cF; S; lE; psbM	2
4048	III. 886	15 0 16.5	2.848	1	76 36 4.3	14.11	1	eF; vS; np of 2	1*
4049	III. 887	15 0 16.5	2.848	1	76 36 4.3	14.11	1	eF; vS; sf of 2	1*
4050	1903	II. 544	15 0 44.0	3.021	3	86 53 20.4	14.08	3	pB; S; v lE; lbM; am st	5
4051	1905	II. 751	15 1 1.8	2.725	(3)	69 54 1.8	14.06	1:	cF; cS; E; p of D neb	4*+
4052	1905	II. 752	15 1 9.2	2.725	(3)	69 56 1.8	14.06	1:	pF; pS; E; f of D neb	4+
4053	1904	IV. 71	15 1 14.4	2.746	1:	71 1 0.5	14.05	1	*6 in vL neb	2
4054	1906	15 1 29.3	2.179	2	46 49 50.5	14.05	2	F; S; R; psbM	2
4055	II. 192	15 1 37.4	3.255	2	100 46 50.6	14.02	2	F; L; E; r	2
4056	1907	II. 585	15 2 30.8	3.013	3	86 24 19.8	13.96	3	pF; cS; i lE; gbM; *14 f ...	4
4057	II. 684	15 2 36.1	3.057	1	89 0 5.5	13.95	1	pB; S; iE	1
4058	1909	I. 215	15 2 36.3	1.639	1	33 41 39.4	13.98	1	vB; cL; pmE 146° 0; gbM...	2+
4059	1909, <i>a</i>	R. nova	15 2 ±	1.639	...	33 42 ±	13.98	...	vS	0
4060	D'Arrest, 100	15 2 41	3.05	[1]	88 55 48	13.95	[1]	eF; H. II. 545, s 3' 15''	0
4061	1908	II. 545	15 2 41.0	3.057	1	88 59 25.5	13.95	1	pF; S; E; psbM	5
4062	II. 755	15 5 10.7	+1.792	1	36 55 21.6	+13.82	1	pB; pL; lE	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.			Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.	h	m	s							
4063	h. 3593	H. III. 736	15	6	3.3	+3.312	1	103° 44' 18.9"	+13.73	1	pB; pL; pmE0°; psmbM; * inv	2
4064	1910	II. 757	15	6	9.4	1.548	1	32 27 30.8	13.76	1	cB; S; E; mbMRN; r	4
4065	II. 818	15	6	27.0	1.128	1	26 30 47.5	13.75	1	pF; cS; R; vgbM	1*
4066	3594	15	7	19.4	4.059	2	135 7 30.2	13.64	2	○; vS; R; quite sharp	2†
4067	3595	III. 116	15	7	30.1	3.238	1	99 33 7.2	13.64	1	F; cL; R; vgbM	2
4068	1911	15	7	38.4	2.199	1	48 14 58.2	13.64	1	F; vS; R; bM	1
4069	1912	III. 659	15	7	58.6	2.198	1	48 12 57.2	13.64	1	cF; vS; R; bM; r	2
4070	1912, a	R. nova	15	8	18.6	2.198	...	48 8 27.2	13.62	...	vF; place from MS.	0
4071	1913	II. 678	15	8	29.8	2.175	1	47 31 15.0	13.60	1	F; S; R; r; 3 st nr	2
4072	} 1913, a	R. 2 nova	15	8	±	2.175	...	47 31 ±	13.60	...	2 nova apparently connected.	0
4073	
4074	II. 763	15	8	37.8	1.353	2	29 40 9.0	13.60	2	pF; pS; E 0° ±	2
4075	3596	VI. 19	15	9	22.4	3.443	1	110 29 44.6	13.52	1	⊕; pF; L; viR; vgbM; rrr	2
4076	3597	III. 138	15	9	53.3	3.504	1	113 31 44.7	13.49	1	F; S; R; gbM	4
4077	1914	II. 650	15	10	0.6	2.167	2	47 26 25.3	13.51	2	cB; pL; pmE; smbMN	5
4078	1915	III. 660	15	10	2.9	2.162	2	47 16 49.0	13.50	2	vF; S; vE; gbM	3
4079	1915, a	R. nova	15	10	2.9	2.162	...	47 17 ±	13.50	...	Close to 1915 pos 0°	0
4080	III. 737	15	10	15.1	1.863	1	39 12 3.0	13.50	1	vF; vS; stellar	1
4081	3598	III. 139	15	10	24.3	3.505	2	113 31 3.5	13.45	2	cF; S; R; gpmbM	5
4082	II. 758	15	10	56.2	1.606	1::	34 0 29.5	13.45	1::	pF; pS; iR	1
4083	1916	M. 5	15	11	27.2	3.028	1	87 23 8.7	13.39	1	!; ⊕; vB; L; eCM; st 11...15	11†
4084	R. nova	15	11	39	1.568	1::	33 51 55	13.39	1::	One of 2, 15' apart np & sf...	0
4085	II. 760	15	11	57.2	1.606	1::	34 6 25.7	13.39	1::	pF; pS; R	1
4086	1917, a	R. nova	15	12	±	15.52	...	32 10 ±	13.38	...	A ray, vmE, par. to h. 1917	0
4087	1917	II. 759	15	12	11.7	1.552	2	33 10 5.4	13.38	2	cB; vL; vmE 155° 0; vg; psbMN.	3†
4088	R. nova	15	12	53	+1.568	2::	34 2 31	13.39	2::	One of 2, 15' apart np & sf...	0
4089	III. 943	15	13	10.9	-0.870	1	14 7 6.1	13.37	1	vF; vS	1
4090	II. 400	15	13	15.7	+2.681	2	68 36 53.4	13.28	2	vF; S; er	2
4091	III. 944	15	13	35.7	-0.876	1	14 7 4.9	13.33	1	vF; vS	1
4092	} 1915 = 3599	III. 374	15	13	38.5	+3.109	2	92 4 2.5	13.25	2	vF; pL; vE; r	3
4093		3600	15	13	51.6	3.297	1	102 34 48.6	13.22	1	B; S; R; glbM; p of 2
4094	3601	15	13	56.2	3.299	1	102 39 13.6	13.22	1	F; S; IE; glbM; f of 2	1
4095	3602	15	14	3.2	3.194	1	96 51 8.3	13.21	1	eF; vS; psbM	1
4096	1920	15	14	36.6	2.018	2	43 36 42.0	13.20	2	cF; L; pmE; glbM; * s	2
4097	1919	I. 148	15	14	59.9	2.975	4	84 25 34.8	13.16	4	cB; cL; iR; vsbM * 12; amst	5
4098	1922	III. 661	15	16	12.3	2.158	(2)	47 50 32.7	13.09	1::	eF; S	2
4099	1921	15	16	13.2	2.156	4	47 46 29.7	13.09	4	vF; pL; vE; vgbM	4
4100	3603	Δ. 357	15	17	15.2	4.480	3	144 1 39.4	12.98	3	Cl; vL; vRi; IC; st 11...14.	3
4101	3604	Δ. 389	15	17	55.5	4.300	2	140 10 47.2	12.94	2	⊕; cB; L; R; vgbM; rrr; st 15	2
4102	1923	II. 874	15	19	40.8	2.729	1	71 25 32.5	12.85	1	pB; cS; R; psbM; * 7 n	2
4103	1924	15	21	5.7	2.141	1	47 50 27.1	12.77	1	vF; vS; sp of D neb	1
4104	1925	II. 651	15	21	8.7	2.141	4	47 50 15.1	12.77	4	pF; pS; R; nf of D neb	6
4105	II. 130	15	23	5.1	2.825	2	76 33 15.6	12.62	2	F; pL; iR; vgbM; r	2
4106	} 1926 = 3606	II. 401	15	23	29.0	3.116	2	92 20 47.7	12.59	2	pB; pS; R; vgbM; 3 st f	3
4107		3605	15	23	30.4	5.466	1	156 22 34.9	12.53	1	F; S; am st
4108	3607	15	25	20.2	4.334	2	140 11 14.9	12.43	2	⊕; cB; pL; R; vglbM; rrr; st 16.	2
4109	II. 906	15	26	3.2	0.808	1	24 45 29.8	12.46	1	F; S; IE 45° ±; vglbM	1
4110	II. 654	15	27	17.7	2.782	1	74 32 25.2	12.34	1	F; pS; E 150° ±	1
4111	1927	II. 178	15	28	2.2	2.777	1	74 20 50.4	12.28	1	pB; cS; p of D neb	4
4112	1927	II. 179	15	28	2.2	2.777	1	74 20 50.4	12.28	1	pB; cS; f of D neb	4
4113	II. 399	15	28	57.7	2.483	1	60 51 21.6	12.22	1	pF; pL; iR; bM; r	1
4114	II. 761	15	29	26.4	1.445	1	33 1 16.3	12.21	1	pF; pS; iF	1
4115	1931	II. 762	15	29	57.4	1.431	(1)::	32 50 58.1	12.17	1	cF; cL; IE	2
4116	1928	II. 96	15	30	6.8	2.746	1	72 55 36.2	12.14	1	pF; pL; iE; gbM	2
4117	1931, a	R. nova	15	30	31.4	+1.431	2::	32 46 22.1	+12.14	2::	No description	0

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
4118	h. 1929	H.	15 30 38.0	+2.953	1	83° 33' 9.7"	+12.09	1	⊕; vF; vL; R; vgbM; rrr...	1+
4119	1930	III. 634	15 30 48.4	2.172	2	49 46 12.3	12.11	2	vF; S; R; gbM; 2 st 8 f.....	3
4120	3608	15 31 12.7	7.159	1	165 13 11.1	11.97	1	F; pL; R; vgbM	1
4121	3609	15 31 20.1	3.694	1	120 5 47.2	12.04	1	vF; L; R; gbM; r	1
4122	II. 76	15 32 10.2	2.834	1	77 24 9.7	11.99	1	pF; pL; R; rr.....	2
4123	1932	15 33 25.5	2.396	1	57 46 40.6	11.92	1	vF; vS; R; bM	1
4124	1934, a	R. nova?	15 34 29.2	1.214	...	30 6 57.2	11.76	...	R; psbM (by diagram)	0
4125	3610	15 34 53.7	5.014	2	150 46 24.5	11.75	3	!; O; pF; vS; R; r? am 150 st	3+
4126	1933	II. 655	15 35 0.3	2.759	1	73 45 35.7	11.79	1	F; pS; E 0°	2
4127	1934, b	R. nova?	15 35 33.7	1.214	...	30 8 46.2	11.77	...	F; mE	0
4128	1934	II. 764	15 36 18.6	1.214	1	30 11 55.2	11.74	1	cB; S; R; psbM; r.....	2
4129	II. 656	15 36 21.3	2.790	1	75 20 47.7	11.69	1	pB; S; E 135°±; bM	1
4130	II. 765	15 36 24.1	1.306	1	31 29 46.9	11.73	1	pF; cS	1
4131	II. 766	15 36 32.5	1.215	1	30 13 46.6	11.72	1	pB; cL; iE; r	1*
4132	3611	Δ. 552	15 36 53.4	3.904	2	127 19 24.9	11.63	2	!; ⊕; vB; L; R; vgbM; st 13...15.	2
4133	1934, c	R. nova	15 37 18.3	1.214	...	30 14 28.0	11.62	...	(? if not =II. 766).....	0
4134	III. 378	15 38 47.1	1.169	1	29 47 35.8	11.56	1	vF; vS	1
4135	1935	II. 425	15 39 16.4	3.019	2	87 8 54.4	11.48	2	vF; vS; R; gbM	5
4136	1936	III. 635	15 39 29.9	2.103	2	48 26 39.7	11.49	2	vF; vS; R; bM; sp of 2.....	3
4137	1937	III. 636	15 39 33.0	2.102	1	48 26 5.4	11.48	1	cF; vS; R; bM; nf of 2.....	2
4138	3613	15 40 38.9	3.334	1	103 19 13.4	11.38	1	eF; S; R; vS * p	1
4139	1938	II. 97	15 40 39.7	2.709	2	71 40 24.0	11.40	2	pF; cS; R; r; bet 2D st.....	4
4140	VII. 29	15 40 56.1	3.664	1	118 10 31.5	11.35	1	Cl; pL; pRi; st vS	1
4141	3612	Δ. 343	15 41 7.4	4.716	1	146 2 39.3	11.31	1	Cl; L; pRi; st 12...14	1
4142	3614	15 41 11.3	3.684	1	118 57 17.9	11.33	1	vF; S; R; sbM	1
4143	III. 371	15 42 3.8	2.459	1	60 54 51.7	11.29	1	vF; S; R	1
4144	3615	Δ. 334	15 44 34.9	4.795	3	147 0 57.8	11.06	3	Cl; pS; pRi; mC; st 16	3
4145	1939	II. 583	15 47 11.0	+3.054	1	89 1 54.3	10.91	1	pF; S; E 90°±; gbM; r ...	3
4146	III. 313	15 47 35.7	-0.483	2	17 24 55.8	10.96	2	vF; S; E 90°±; vS * f	2
4147	II. 657	15 47 49.1	+2.772	1	74 59 59.1	10.87	1	F; bet 2 B st	1
4148	1940	15 49 3.7	2.948	1	83 39 30.4	10.78	1	pB; pL; E	1
4149	III. 739	15 49 24.0	0.892	1	27 15 50.0	10.80	1	vF; pL; R; vgbM	1
4150	1941	15 50 19.6	2.946	2	83 35 40.4	10.68	2	!; vF; vS; R; g; smbM	2
4151	1942	III. 646	15 51 5.5	2.742	1	73 42 40.9	10.63	1	vF; S; iE; p of 2	2
4152	1943	III. 73	15 51 8.4	2.741	1	73 37 40.9	10.63	1	eF; vS; iE; f of 2	3
4153	3516	Δ. 304	15 51 50.9	5.056	4	150 6 0.6	10.52	4	Cl; B; vL; pRi; iC; st 7... 4	4
4154	3617	15 52 18.5	3.846	1	124 8 38.0	10.50	1	F; S; R; gpmbM; Δ of st np	1
4155	3618	Δ. 359	15 56 43.3	4.641	1	143 38 28.8	10.16	1	Cl; S; mC; st 11...14	1
4156	1944	III. 622	15 57 35.3	2.182	2	52 15 33.5	10.15	2	vF; S; R; * 10 sf	4
4157	III. 33	15 58 43.2	+2.658	1	70 16 11.9	10.14	1	eF; (?)	1
4158	II. 873	15 58 43.5	-0.284	1	18 55 6.9	10.13	1	F; R; bM	1
4159	1945	15 58 49.9	+2.901	1	81 31 23.2	10.04	1	*7 in photosphere	1
4160	1946	III. 637	15 59 36.9	2.065	2	48 55 52.0	10.00	2	pF; vS; R; stellar	3+
4161	III. 140	16 0 51.0	2.629	1	69 4 1.0	9.90	1	vF; vS; r; pB * sf	1
4162	3619	Δ. 360	16 2 19.2	+4.675	2	143 50 42.2	9.74	3	Cl; vB; vL; vRi; iC; st 10... 3	3
4163	III. 973	16 2 40.3	-3.081	1	10 38 44.0	9.90	1	vF; vS; iE 0°; r	1
4164	1947	III. 553	16 2 48.9	+3.051	1	88 55 4.2	9.74	1	F; L; pmE; vgbM; r.....	2
4165	III. 883	16 3 32.5	-0.266	1	19 13 44.8	9.76	1	eF; vS	1
4166	3620	16 3 44.7	+3.922	1	125 52 42.5	9.65	1	pF; R; vgvbM; r	1
4167	1948	III. 74	16 4 5.4	+2.713	1::	72 55 51.5	9.65	1	vF; S; r	2*
4168	III. 884	16 5 50.0	-0.147	1	19 59 34.4	9.58	1	vF; vS	1
4169	3621	16 6 33.5	+3.866	1	123 53 2.9	9.43	1	eF; S; E; lbM	1
4170	3622	Δ. 326	16 7 16.0	4.936	2	147 32 42.5	9.35	2	Cl; B; L; iC; st 7...10	2
4171	III. 812	16 7 18.1	1.199	1	32 8 30.2	9.44	1	vF; vS; iE	1
4172	1949	III. 889	16 7 18.4	2.293	1	56 35 35.9	9.33	1	vF; S; R; bM	2
417	3624	M. 80	16 8 41.9	+3.567	2	112 37 34.1	9.27	2	!!!; ⊕; vB; L; vmbM (var *); rrr; st 14.	2*
4174	III. 314	16 8 49.3	-0.740	2	17 10 21.4	9.38	2	vF; vS; iE	2
4175	3623	Δ. 68	16 9 54.3	+6.720	3	161 51 51.0	+ 9.10	3	⊕; pF; L; iR; vgbM; rrr; st 14...	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ′ ″	′ ″			
4176	1950	III. 888	16 10 18.4	+2.222	2	57 41 3.1	+9.17	2	vF; S; R; vglbM	3
4177	1951	III. 688	16 11 21.9	2.206	2	53 56 19.0	9.10	2	vF; S; iR	3
4178	1952	II. 151	16 11 58.8	2.911	1	82 15 5.9	9.03	1	F; pL; iE; vgbM; r	2
4179	3625	16 13 42.5	4.589	1	141 36 46.8	8.86	1	Cl; eL; eRi.....	1
4180	1953	II. 402	16 14 31.8	3.114	1	91 56 56.9	8.83	1	vF; cL; cE 45°±; r	2
4181	1954	16 14 43.9	2.132	1	51 52 57.9	8.83	1	vF; eS; R	1
4182	1955	III. 623	16 14 50.5	2.133	2	51 54 7.9	8.83	2	vF; vS; R; * nf.....	4
4183	{ M. 4= B.A.C.5455 }	16 15 3.6	3.665	...	116 11 11.1	8.77	...	{ Cl; 8 or 10 L st in line, with 5 st; rrr. }	4
4184	3626	Δ. 514	16 16 1.3	4.097	1	130 20 6.7	8.69	1	Cl; B; L; pRi; ICM; st 9...11	2
4185	II. 810	16 16 4.4	1.154	1	32 2 48.5	8.75	1	pF; pS; iE	1
4186	III. 891	16 16 46.1	2.121	1	51 40 47.1	8.67	1	eF; vS; R; lbM	1
4187	3627	Δ. 412	16 17 20.5	4.456	2	138 49 29.1	8.57	2	Cl; cL; pRi; ICM; st 13...15	2
4188	1956	III. 624	16 18 3.7	2.121	2	51 44 50.1	8.57	2	F; S; iR; bM	4
4189	3628	Δ. 536	16 18 14.8	4.037	3	128 30 59.3	8.51	4	B; pL; R; psbM; rr	4
4190	III. 740	16 18 19.3	0.401	1	24 17 36.7	8.59	1	eF; pL; iR	1
4191	III. 892	16 18 23.5	2.144	1	52 26 49.2	8.54	1	eF; S; bM	1
4192	II. 811	16 18 42.4	1.320	1	34 34 35.2	8.54	1	pB; iR; vglvBM	1
4193	3629	VI. 10	16 18 43.0	3.657	1	115 43 5.4	8.48	1	Cl; cL; mC; gbM; rrr	2
4194	1957	16 20 20.8	2.009	1	48 44 41.0	8.40	1	F; R; bM	1
4195	1958	III. 638	16 20 27.2	2.010	2	48 47 10.4	8.38	2	cF; vS; R; bM	3
4196	1958, a	R. 2 novæ	16 20 ±	2.010	...	48 47 ±	8.38	...	2 novæ, one eF; one S	0
4197											
4198	1959	III. 639	16 21 10.4	2.025	1	49 13 2.9	8.33	1	vF; vS; R	2
4199	3630	16 21 41.1	7.067	1	162 57 2.8	8.16	1	vF; vS; *9 nr	1
4200	3631	16 21 49.9	4.654	1	142 18 59.3	8.21	1	Cl; L; iC; st L	1
4201	III. 680	16 22 6.9	1.623	2	39 48 49.8	8.26	2	vF; S; R; lbM; er	2
4202	II. 690	16 22 34.5	1.687	2	41 17 48.6	8.22	2	F; pS; iF; gbM	2
4203	3632	16 22 36.1	5.219	2	150 18 1.9	8.13	2	pF; pL; vIE; gbM	2
4204	1960	II. 652	16 23 0.8	2.004	2	48 45 15.4	8.18	2	eF; pL; R; gbM; r	3
4205	II. 647	16 23 2.7	2.057	2	50 13 17.4	8.18	2	F; S; iF	2
4206	3633	16 23 22.7	4.423	1::	137 48 2.7	8.09	1::	eF; (?); * f nr	1
4207	3634	16 23 39.8	4.426	1	137 51 20.8	8.06	1	F; cS; iE; vglbM; * p	1
4208	1961	II. 875	16 23 50.4	2.053	1	50 8 9.6	8.12	1	pF; S; vIE; vgbmM	2
4209	3635	Δ. 400	16 23 50.9	4.506	1	139 27 48.5	8.05	1	Cl; L; iC; iF	1
4210	3636	16 24 7.3	4.246	1	133 44 18.2	8.04	1	Cl; μ Normæ inv	1
4211	3637	VI. 40	Mechain	16 24 43.0	3.350	1	102 44 17.3	8.01	1	⊕; L; vRi; vmC; R; rrr ...	2
4212	1962	III. 640	16 24 58.0	2.004	1	48 52 16.6	8.02	2	eF; vS; R; bM	3
4213	1962, a	R. nova	16 25 ±	2.004	...	48 52 ±	8.02	...	No description; near h.1962	0
4214	1963	III. 641	16 25 18.7	2.010	1	49 3 57.0	8.00	1	vF; vS; R	3
4215	1964	III. 890	16 25 34.0	2.203	1	54 38 1.1	7.97	2	vF; pL; iE; rr; * nr	3
4216	3638	16 25 36.3	4.316	2	135 19 19.6	7.92	2	Cl; B; S; st pL	2
4217	1964, a	R. nova	16 25 47.9	2.203	...	54 35 11.1	7.91	...	No description; 4' nf 1964...	0
4218	II. 753	16 26 37.0	2.624	2	69 55 19.4	7.88	3	pB; pL; vIE; pgmbM	2
4219	III. 813	16 26 50.0	1.264	1	34 9 56.7	7.89	1	vF; vS; iR	1
4220	3639	16 26 50.4	6.305	1	159 4 52.8	7.76	1	eF; eS; R; gbM.....	1
4221	1965	16 28 15.0	2.190	1	54 21 40.8	7.76	1	F; S; R; gbM; * 11 np.....	1
4222	III. 730	16 28 18.6	2.579	1	68 9 52.2	7.74	1	eF; vS; E	1
4223	3640	16 29 14.5	4.483	1::	138 43 48.6	7.62	1::	!; F; vL; viE; B * inv	1
4224	3641	Δ. 483	16 30 29.3	4.233	2	133 5 18.9	7.53	2	Cl; cL; pRi; iR; st 11...14..	2+
4225	3642	Δ. 413	16 30 52.1	4.475	1	138 29 3.7	7.49	1	Cl; vL; iRi; iC; rrr; F neb inv.	1
4226	1966	III. 893	16 31 43.6	2.058	2	50 41 28.4	7.48	2	vF; S; R; gbM; bet 2 st ...	4
4227	1967	16 32 19.0	2.155	1	53 31 20.9	7.43	1	vF; vS; sbM * 12	1
4228	3643	16 33 50.3	4.420	1	137 11 50.2	7.24	1	Cl; (in M. Way).....	1
4229	3644	Δ. 442	16 36 6.6	4.404	2	136 45 36.8	7.06	2	Cl; pRi; eiCM; st 11...12...	2+
4230	1968	M. 13, Halley	16 36 41.2	2.140	3	53 16 19.4	7.08	3	!!; ⊕; eB; vRi; vgeCM; st 11...20.	14+
4231	1969	II. 701	16 38 12.4	2.125	(1)	52 54 7.5	6.95	1	pB; pL; E 45°±; vgbmM...	2
4232	3646	Δ. 364	16 38 18.4	4.776	3	143 33 29.1	6.87	3	Cl; L; Ri; ICM; st 9...12 ...	2
4233	3645	16 38 23.8	+7.032	1	162 20 32.0	+6.80	1	vF; pL; vglvBM	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
4234	h. 1970	H.	Σ.5 = Lal. 30510	16 38 36.0	+2.513	1	65 56 10.0	+6.90	1	{ O; vB; vS; R; disc and border.	1*
4235	3647		16 39 3.5	+5.147	1	148 44 47.7	6.79	1	pF; R; vglbM; *5 p 79 ^s ...	1
4236	I. 280	16 39 16.1	-3.036	3	11 33 21.3	7.01	3	B; cL; lE; slbM	3
4237	3648	Δ. 454	16 39 19.8	+4.308	3	134 28 15.7	6.79	3	Cl; pS; pRi; pC; st 12...15..	3
4238	1971	M. 12	16 39 58.1	3.110	1	91 41 47.4	6.78	2	!!; ⊕; vB; vL; iR; gmbM; rrr; st 10...	7
4239	3649	16 40 39.6	5.171	2	148 58 2.1	6.67	2	⊕; pB; cL; R; glbM; rr ...	3
4240	3650	Δ. 456?	16 40 41.4	4.311	1	134 28 33.4	6.68	1	Cl; vL; vRi; lbM; st 12...13	1
4241	D'Arrest, 101	16 41 24	0.72	[2]	28 9 18	6.72	[2]	F; S; R; mbM	0
4242	D'Arrest, 102	16 41 47	0.68	...	27 46 0	6.69	...	F; S; makes Δ with 2 S st 12 and 14 m.	0
4243	3651	16 41 51.9	4.170	1	130 58 41.7	6.59	1	Cl; eL; eRi (in M. Way) ...	1
4244	IV. 50	16 43 6.4	1.678	1	42 8 38.8	6.56	1	vB; L; R; Disc + F, r, border	1
4245	3652	Δ. 499	16 44 14.5	4.197	1	131 33 37.7	6.39	1	Cl; B; cL; pRi; st 10...13 ..	1
4246	3653	II. 584	16 45 1.0	3.584	1	111 55 47.5	6.35	1	pB; cL; iR; rrr; st 14...16 ..	2
4247	III. 727	16 45 54.8	1.932	1	48 1 25.6	6.32	1	eF; S; E90°	1*
4248	III. 735	16 46 0.4	1.775	1	44 20 25.3	6.31	1	aF; pS	1
4249	3654	Δ. 520	16 46 3.5	4.112	2	129 16 12.2	6.24	2	Cl; B; L; Ri; st 8...11	3
4250	D'Arrest, 103	16 47 1	0.56	...	26 47 5	6.26	...	F; pS; irr	0
4251	3655	16 47 33.9	4.327	2	134 33 17.6	6.12	2	Cl; pRi; vLC; iF; st L & S...	2
4252	3656	16 47 40.8	+4.378	2	135 42 21.0	6.10	2	Cl; L; lRi; lC; st 8...12	2
4253	III. 974	16 47 55.2	-6.988	1	7 9 0.3	6.41	1	cF; S; bM; p of 2	1
4254	III. 975	16 47 58.3	-7.058	1	7 5 59.7	6.39	1	vF; vS; f of 2	1
4255	3657	Δ. 374?	16 48 3.0	+4.734	1	142 28 51.8	6.06	1	Cl; S; Δ ar; st 13	1
4256	{ 1972 = 3659 }	M. 10	16 49 47.6	3.159	3	93 52 6.8	5.96	3	!; ⊕; B; vL; R; gvmbM; rrr; st 10...15.	7
4257	1973	III. 689	16 49 47.6	2.121	1	53 16 46.7	5.99	1	eF; cL; E90°	2
4258	3658	16 50 8.6	4.032	2	126 53 32.3	5.91	2	⊕; vF; vL; iR; vgbM; rrr; st 20.	2
4259	1974	16 50 11.5	2.008	1	50 9 37.8	5.96	1	vF(?)	1*
4260	3660	Δ. 456	16 50 37.1	4.326	3	134 26 53.8	5.86	3	!; Cl; B; vL; vRi; st 11...	2
4261	3661	{ M. 62 = Δ. 627 }	16 52 18.7	3.810	5	119 53 42.9	5.73	5	!; ⊕; vB; L; gmbM; rrr; st 14...16.	5+
4262	III. 123	16 52 22.2	2.523	1	66 46 53.8	5.76	1	vF; pL; R; lbM	1
4263	3662	Δ. 521	16 52 26.0	4.130	2	129 30 46.3	5.71	2	Cl; B; pL; cRi; st 10	2
4264	{ 1975 = 3663 }	M. 19	16 53 59.2	3.701	3	116 3 13.0	5.60	2	⊕; vB; L; R; vCM; rrr; st 16... red.	7
4265	3664	Δ. 556	16 55 16.3	4.067	1	127 40 56.4	5.48	1	Cl; L; pRi; lC; st 9...11 ...	1
4266	III. 124	16 55 22.1	2.521	1	66 46 39.3	5.51	1	vF; stellar	1
4267	III. 728	16 55 52.8	1.534	1	39 51 36.0	5.50	1	vF; cS; iR	1
4268	{ 1976 = 3665 }	VI. 11	16 55 55.3	3.661	2	114 33 37.9	5.43	2	⊕; B; L; R; gCM; rrr; st 16.	3
4269	3666	II. 195	16 56 43.5	3.606	1	112 30 9.8	5.36	1	⊕; cB; L; R; gpmCM; rrr; st 16.	3
4270	{ 1977 = 3667 }	VI. 12	17 1 28.2	3.716	2	116 23 13.1	4.97	2	⊕; vB; L; R; psbM; rrr; st 16; F neb f.	4
4271	1978	17 1 34.2	3.715	1	116 22 33.5	4.95	1	F; S; vgbM; ⊕ p	1
4272	D'Arrest, 104	17 3 40	0.57	[1]	27 21 54	4.86	[1]	vF; vS; R	0
4273	3668	17 3 51.8	5.588	2	152 39 3.3	4.71	2	F; vL; vIE; am st; 2 st inv...	2
4274	IV. 57	17 4 10.8	1.883	2	47 29 24.4	4.78	2	F; stellar	2
4275	3670	I. 147	17 5 40.2	3.803	2	119 17 18.0	4.60	2	⊕; vB; cL; R; s, vglbM; rrr; st 16...17.	3
4276	3669	17 5 46.4	5.242	1	149 0 21.5	4.55	1	vF; vS; R; glbM	1
4277	D'Arrest, 105	17 6 3	0.73	[1]	29 3 54	4.59	[1]	pF; vS; R; *13 nr	0
4278	D'Arrest, 106	17 6 15	+0.71	[1]	28 49 54	+4.58	[1]	F; pL; lE	0

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
4279	h. 3671	H. I. 45	h m s 17 7 49.0	s +3.765	2	117° 58' 27".9	+4".43	2	⊕; cB; pS; R; gvmbM; rrr; st 16...17.	4
4280	3672	Δ. 522	17 8 4.2	4.140	1::	129 17 11.4	4.38	1::	Cl; pL; Ri; R; gbM; st 12...14.	1
4281	3673	17 8 45.0	+4.279	2	132 43 29.9	4.33	2	Cl; vL; pRi; IC (* nf taken)	2
4282	III. 945	17 9 14.5	-2.000	1	14 23 21.1	4.47	1	vF; S; E; S * s	1
4283	3676	17 9 30.9	+3.643	1	113 35 51.4	4.28	1	pF; L; R; rr	1
4284	3675	17 9 41.0	4.722	2	141 35 30.9	4.23	2	!!!; O; pB; vS; R	3+
4285	3674	17 9 57.7	+5.861	2	154 51 25.4	4.18	2	vF; vS; vLE; glbM	2
4286	III. 951	17 10 45.4	-3.497	...	11 12 41.4	4.38	1	eF; S	1
4287	{ 1979 = 3677 }	M. 9	17 10 57.6	+3.507	2	108 22 59.8	4.16	2	⊕; B; L; R; eCM; rrr; st 14	6
4288	3678	17 11 31.9	4.021	2	125 54 52.0	4.10	2	cF; vL; icE; vglbf; * 8 inv.	2
4289	3679	17 11 35.8	3.828	1::	120 0 3.0	4.10	1::	Diffused neb in patches	1
4290	3680	17 12 40.1	4.109	3	128 20 0.0	4.00	3	!!!; ⊙; eF; S; am St ₁	3+
4291	II. 812	17 12 40.7	+1.011	1	32 24 7.4	4.08	1	F; S; R; vglbM	1
4292	1980	II. 767	17 12 44.1	-1.062	1	17 32 6.2	4.14	1	cF; pL; R; vgmbM	2
4293	I. 149	17 12 47.8	+3.535	1	109 25 12.0	4.00	1	cB; pS; lE; er	1
4294	M. 92	17 12 56.9	1.840	...	46 43 31.2	4.04	...	⊕; vB; vL; eCM; rrr; st S.	8*
4295	3681	I. 46	17 15 16.2	3.719	1	116 12 43.4	3.78	1	cF; L; R; gbM; rrr	2
4296	3683	I. 48	17 15 23.2	3.491	1	107 40 23.4	3.78	1	⊕; vB; cL; vgvmbM; rrr; st 20.	3
4297	3682	17 15 28.0	3.960	1	124 3 6.8	3.76	1	F; L; E; vglbM; * inv	1
4298	D'Arrest, 107	17 16 23	0.61	[1]	28 4 54	3.77	[1]	pB; S; R; bMN=*12	0
4299	3685	17 16 28.0	3.827	1	119 51 37.1	3.67	1	Neb in patches (M. Way)	1
4300	3684	Δ. 225	17 17 23.3	6.165	2	156 55 47.9	3.53	2	⊕; cB; L; vgmbM; rrr; st 14...17.	2
4301	Auw. N. 36	17 20 19.2	3.184	...	94 57 2.0	3.37	...	F; L; vlbM (Winnecke, April 12, 1860).	0
4302	{ 1981 = 3686 }	IV. 11	17 20 50.4	3.650	2	113 37 47.3	3.21	3	!!; ⊙; pB; S; R	5*+
4303	III. 137	17 22 2.6	2.412	1	63 25 22.2	3.24	1	vF; pL; iF	1
4304	3687	17 22 58.1	3.913	1	122 28 48.3	3.11	1	Cl; S; P; B * inv	1
4305	3688	17 24 47.4	4.142	1	128 58 1.5	2.95	1	eF; pS; lE; * 9 att	2+
4306	3689	17 25 34.1	3.915	3	122 28 38.7	2.89	3	Cl; st 6-7, 13	2
4307	3690	Δ. 457	17 26 3.2	4.380	2	134 38 32.2	2.84	2	⊕; vB; L; R; pg, psvmbM; rrr; st 17...	2
4308	II. 901	17 26 3.6	2.680	1	73 29 2.4	2.88	1	F; S; iF; er	1
4309	3691	17 27 57.9	6.657	2	159 41 19.0	2.60	2	cF; S; R; glbM; * 13 sp	2
4310	3693	17 28 51.0	3.998	1	124 54 49.0	2.60	1	Cl; pL; lRi; IC	1
4311	3692	Δ. 366	17 29 17.3	4.871	2	143 35 5.2	2.54	2	⊕; B; vL; Ri; st 13	3
4312	3694	17 29 48.6	5.527	1	151 36 5.1	2.47	1	eF; S; R; p of 2	1
4313	3696	Δ. 568	17 30 1.9	4.067	1	126 51 8.0	2.50	1	Cl; pL; pRi; iR; st 9...10...	1
4314	{ 1982 = 3697 }	I. 44	17 30 6.9	3.658	4	113 48 55.0	2.50	4	pB; pL; R; * 12 f inv	6
4315	{ 1983 = 3698 }	M. 14	17 30 16.0	3.146	2	93 9 25.0	2.50	2	!; ⊕; B; vL; R; eRi; vgmbM; rrr; st 15...16.	7
4316	3695	17 30 18.8	5.528	1::	151 36 3.9	2.43	1	eeF; f of 2	1
4317	4020	h. o. n.	17 30 26.3	3.939	1	123 9 16.8	2.46	1	Cl; F; L; pRi; IC; st 13...15	1
4318	3699	M. 6	17 30 55.8	3.905	1	122 7 4.9	2.43	1	Cl; L; iR; IC; st 7, 10...	4
4319	3700	17 32 15.0	5.436	1	150 39 39.1	2.27	1	eF; S; R; 3 st nr	2
4320	D'Arrest, 108	17 33 56	+0.69	[2]	29 6 12	2.26	[2]	vS; gbM	0
4321	VI. 41	17 34 7.6	-2.177	1	14 11 11.9	2.33	1	⊕; cL; R; vgbM; rr	1
4322	3701'	17 35 0.1	+4.002	1::	124 56 32.1	2.07	1	Nebulous portion of M. Way	1
4323	3702'	Δ. 612	17 35 9.3	3.912	2	122 17 0.5	2.05	2	Cl; vL; Ri; IC	2
4324	3702'	17 36 30.1	3.957	1	123 37 33.2	1.94	1	Cl; vL; pRi; st 8...12	1
4325	II. 587	17 37 53.6	+2.997	1	86 44 25.5	+1.85	1	F; cL; iF	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by H. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
4326	3703	17 37 56.4	+ 3.887	1	121° 27' 58.6"	+ 1.82	1	Cl; pS; lRi; IC; st 10...12 ..	1
4327	1984	17 38 46.0	+ 3.689	1	114 49 51.5	1.75	1	Cl; st vS	1
4328	1987	III. 741	17 39 11.8	- 1.069	1	17 49 22.7	1.69	1	vF; vS; R; stellar; * 8 s.....	2
4329	3704	17 39 39.5	+ 4.019	1;	125 22 49.8	1.66	1;	Cl; F; eL; vS st + neb	1
4330	3701	17 39 43.4	19.744	1	175 24 24.0	1.20	1	pB; R; vgbM.....	1
4331	1985	I. 150	17 40 32.5	3.565	1	110 18 21.0	1.60	1	pB; pL; R; bM	3
4332	3705	Δ. 557	17 40 41.6	4.077	2	127 0 0.1	1.57	2	⊕; vB; pL; R; vgbmB; rrr; st 18...20.	2
4333	1986	II. 586	17 40 54.7	3.557	2	109 58 3.8	1.56	2	pB; pS; R; gbM; r; * 15 np	3
4334	3706	Δ. 597?	17 40 55.0	3.999	1;	124 48 46.5	1.55	1;	Cl; vL; vRi; st 12...13	1
4335	3707	VI. 13	17 41 40.5	3.847	2	120 10 12.7	1.49	2	Cl; pL; pRi; bifid; st 12...	3+
4336	3708	17 41 59.7	3.991	1;	124 34 43.8	1.46	1;	eL; iR; pmbM; r	1
4337	3709	17 42 37.5	4.018	1;	125 20 42.0	1.40	1;	Cl; rr; st eS + neb	1
4338	1988	17 44 16.8	3.705	1	115 21 32.1	1.27	1	eF; S; (?)	1
4339	3711	17 44 29.7	3.620	1	112 18 23.5	1.25	1	Cl; pRi (in M. Way).....	1
4340	3710	M. 7 (Lacaille)	17 44 39.6	3.999	1	124 46 37.9	1.23	1	Cl; vB; pRi; IC; st 7...12 ...	3
4341	3712	17 44 54.5	3.815	1;	119 5 36.3	1.21	1;	Neb or nebulous patch of M. Way.	1
4342	3713'	17 45 36.6	3.855	1	120 23 56.5	1.15	1	Neb or nebulous patch of M. Way.	1+
4343	1989	17 45 57.6	2.502	1	66 53 15.5	1.15	1	!; vF; S; R; vsymbMvSRN.	1+
4344	3713	17 46 1.3	5.767	1	153 38 2.5	1.05	1	F; S; E; bM; bet 2 st 10 ...	1
4345	3714	17 48 31.2	6.132	1	156 24 42.9	0.83	1	pF; S; pmE 90°; * 12 f, att...	1
4346	1990	M. 23	17 48 40.9	3.532	3	108 59 43.7	0.89	3	Cl; B; vL; pRi; IC; st 9.10, 11...13.	6
4347	3715	Δ. 460?	17 48 51.4	4.372	2	134 13 52.2	0.84	2	Neb + Cl; pL; mE; gvlbM...	2
4348	III. 957	17 49 25.1	2.631	1	71 42 59.5	0.85	1	vF; vS; p of 2	1
4349	III. 958	17 49 31.1	2.629	1	71 38 58.5	0.85	1	vF; vS; f of 2	1
4350	3716	17 50 20.0	+ 5.991	2	155 23 58.1	0.67	2	vF; vS; f * of * inv	2
4351	Auw. N. 37	17 50 54.1	- 0.638	...	19 49 12.3	0.81	...	pF; L; mE (Auwers, July 22, 1854).	0
4352	3717	17 51 10.6	+ 3.685	1	114 38 24.1	0.67	1	Cl; Ri; eL; vLc	1
4353	VIII. 53	17 51 30.0	3.491	1	107 23 18.2	0.64	1	Cl; pS; lRi; IC	1
4354	D'Arrest, 109	17 53 43	0.49	[2]	27 20 24	0.53	[2]	vF; R; 1st of 3	0
4355	{ 1991 = 3718 }	{ IV. 41 V. 10, 11, 12 }	M. 20	17 53 51.8	3.640	3	113 1 39.9	0.43	3	!!!; vB; vL; trifid; * inv ...	8 Mon.†
4356	D'Arrest, 110	17 54 9	0.48	[2]	27 18 30	0.50	[2]	vF; vS; 2nd of 3	0
4357	3719	II. 199	17 54 13.9	3.282	2	98 56 37.3	0.41	2	pB; pL; R; rr.....	3
4358	3721	VII. 7	17 54 36.4	3.779	1	117 53 32.8	0.36	1	Cl; pS; Ri; IC; st 9...13 ...	3
4359	3720	I. 49	17 54 37.2	3.845	2	120 1 29.8	0.36	2	⊕; B; pL; R; gymbM; rrr; st 16...17.	3
4360	D'Arrest, 111	17 54 38	0.49	[2]	27 22 0	0.46	2	F; pL; 3rd of 3	0
4361	3722	M. 8	17 55 17.9	3.677	3	114 21 15.3	0.31	3	!!!; vB; eL; eiF; with LCl...	8 Mon.†
4362	1992	17 55 27.9	2.811	1	78 57 6.6	0.32	1	1
4363	V. 9	17 55 39.8	3.652	1	113 27 28.4	0.28	1	F; L; cE.....	1
4364	3723	II. 200	17 55 51.7	3.846	2	120 3 24.5	0.25	2	⊕; pF; cS; R; gbM; rrr; st 16...17.	3*
4365	3724	Δ. 569	17 56 1.6	4.054	1;	126 18 6.9	0.23	1;	Cl (in M. Way)	1
4366	3725	17 56 6.0	3.677	2	114 19 58.9	0.23	2	Cl; B; L; pRi; vL neb p ...	2
4367	1993	M. 21	17 56 13.8	3.626	1	112 30 8.6	0.22	1	Cl; pRi; IC; st 9...12	2
4368	V. 13	17 56 32.3	3.692	1	114 53 23.7	0.19	1	eL; eiF; st f	1*
4369	Auw. N. 38	17 56 38.5	3.076	...	90 17 44.9	0.20	...	pF; vS; vS neb * p (Hind, Ap. 1852).	0
4370	Auw. N. 39	17 57 15.0	3.247	...	97 34 58.4	0.15	...	(Brorsen, 1856.) No description.	0
4371	II. 198	17 57 27.3	3.778	1	117 49 20.6	0.12	1	pF; S; iE; er or Cl	1
4372	3726	Δ. 473	17 57 50.9	+ 4.350	2	133 43 25.8	0.06	2	⊕; B; R; eC; gbM; rrr; st 15...16.	2*
4373	IV. 37	17 58 20.0	- 0.023	1	23 22 9.5	+ 0.15	1	○; vB; pS; sbMvSN.....	1
4374	1994	II. 197	17 58 42.4	+ 3.696	1	115 0 52.0	0.00	1	eF; pL; iR; r	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	N of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
4375	3727	17 58 45.8	+5.788	1	153 47 28.2	-0.06	1	eeF; eeS; R	1+
4376	3729	17 58 47.5	3.646	1::	113 14 0.0	0.00	1::	Cl; vL; vRi	1
4377	III. 555	17 59 39.1	2.624	1	71 27 7.8	-0.04	1	cF; S; IE; r	1
4378	3730	IV. 12	18 0 44.2	3.722	1	115 56 17.6	0.18	1	⊕; F; L; IE; vglbM; rr; st 20.	3
4379	1995	18 0 50.8	3.518	1	108 26 46.6	0.18	1	Cl; pRi; vIC; st L & S	1
4380	3732	18 1 9.8	3.769	1	117 32 31.7	0.21	1	F; vL; eE; lbM; rr	1
4381	3728	18 1 11.4	8.690	1	166 36 39.2	0.36	1	vF; vS; R; glbM	1
4382	3731	18 1 12.2	3.900	3	121 46 50.4	0.22	3	⊕; pB; pL; R; glbM; rrr; st 16	3
4383	II. 902	18 1 18.7	2.649	1	72 25 54.3	0.19	1	F; L; R; vglbM	1
4384	{ 1996 = 3733 }	18 1 23.9	3.671	2	114 7 30.1	0.23	2	vF; vL; IE; * inv	2
4385	1997	VIII. 54	18 2 24.7	3.476	1	106 48 57.7	0.31	1	Cl; L; IC; st cL	3
4386	3734	18 2 47.5	3.970	1	123 53 29.2	0.36	2	○; F; L; eE; hazy border...	2
4387	D'Arrest, 112	18 3 51	1.34	[1]	37 44 0	0.38	[1]	eF; vS; R; * 16 nr.	0
4388	{ 1998 = 3735 }	VII. 30	18 4 20.6	3.602	2	111 37 33.6	0.48	2	Cl; vL; IC	3
4389	3736	II. 201	Δ. 619	18 4 33.3	3.902	3	121 51 0.7	0.51	3	⊕; cB; L; R; rrr; st 15	5
4390	2000	Σ. 6	18 5 17.8	2.912	1	83 10 53.5	0.55	2	○; vB; vS; R; l hazy	3*
4391	1999	18 5 23.5	3.617	1:	112 10 47.6	0.58	1	Cl; st vS	1
4392	{ 2001 = 3739 }	VII. 31	18 7 24.1	3.616	2	112 10 21.5	0.75	2	Cl; pRi; pC; eE; st 13	3
4393	3737	Δ. 376	18 7 24.6	4.797	2	142 15 13.3	0.79	2	⊕; cB; cL; R; gmbM; rrr; st 15	2
4394	3738	18 7 56.9	5.793	1	153 51 18.2	0.86	1	eF; S; *6, sp	1
4395	2002	18 8 47.2	3.529	1	109 55 8.6	0.88	3	vF; pS; R; * inv	3+
4396	2003	VIII. 55	18 9 28.2	3.473	1	106 41 22.1	0.93	1	Cl; IC	2
4397	2004	M. 24	18 10 13.7	3.518	2	108 28 7.3	0.99	2	!; Cl; vRi; vmC; R; st 15 (M. Way).	4*
4398	3740	VIII. 15	18 10 14.2	3.363	1	102 17 10.3	0.99	1	Cl; lRi; IC	2
4399	2005	18 10 20.9	3.429	1	104 59 37.7	1.01	1	Cl; lRi; IC; st 10...12	1
4400	2006	M. 16	18 10 57.0	3.401	1::	103 50 2.2	1.06	1	Cl; at least 100 st L & S	3
4401	2007	M. 18	18 11 44.6	3.485	1::	107 11 8.1	1.13	1	Cl; P; vL	4
4402	3741	18 11 45.3	5.724	2	153 17 47.0	1.20	2	vF; S; R; gvlbM; *9 p	2
4403	2008	M. 17	18 12 33.1	3.460	2	106 13 36.0	1.20	2	!!!; b; eL; eiF; 2-hooked	9+
4404	3742	I. 50	18 14 41.9	3.855	3	120 25 26.0	1.40	3	⊕; vB; pL; R; rrr; st 16	4
4405	2009	18 15 20.7	3.358	1::	102 5 58.8	1.44	1	IC; lRi; IC; st 11...12	1
4406	{ 2010 = 3743 }	M. 28	18 15 55.4	3.692	2	114 56 30.0	1.50	2	!; ⊕; vB; L; R; geCM; rrr; st 14...16.	8
4407	3744	II. 204	18 17 13.4	3.645	1	113 16 30.4	1.62	1	○ or ⊕; pB; eeS; R	2
4408	3745	18 19 12.0	5.726	1	153 22 22.5	1.85	1	pF; S; R; gbM	1
4409	3746	18 19 23.4	3.358	1	102 6 42.3	1.79	1	Cl; pL; pRi; st 12...15	1
4410	VIII. 72	C. H.	18 20 43.1	2.921	1	83 31 15.3	1.89	1	Cl; IC; st L	3
5076	18 20 44.6	123 30 27.3	See No. 5076	0
4411	M. 69=Δ. 613	18 22 13.2	3.917	3	122 26 33.5	2.05	3	⊕; B; L; R; rrr; st 14...16	4*
4412	3748	I. 51	18 22 16.6	3.708	1	115 34 55.5	2.05	1	⊕; B; S; R; rr	2
4413	2011	18 23 4.6	3.385	1	103 15 12.7	2.11	1	Cl (in M. Way)	1
4414	{ 2012 = 3749 }	II. 205	18 23 24.2	+3.651	3	113 33 57.5	2.15	3	⊕; pB; pL; iR; gpmbM; rrr; st 16.	4
4415	Auw. N. 40	18 23 35.4	-1.719	...	15 29 47.7	2.01	...	!!; pB; pL; E 50°; 2 st p Var. (Tuttle.)	0*
4416	2013	VI. 23	18 24 31.5	+3.477	1	106 59 3.8	2.24	1	Cl; pL; vRi; pC; st 11...15	2
4417	II. 907	18 24 55.8	1.967	1	50 14 52.8	2.24	1	F; S; iF	1
4418	2014	VIII. 14	18 25 4.6	3.488	1::	107 26 2.3	2.29	1	Cl; L; Ri; IC; st vS	2
4419	Auw. N. 41	18 25 4.9	0.230	...	25 5 44.2	2.20	...	S; pmE; * inv (Σ. neb No. 7)	0
4420	3751	18 25 44.1	3.317	1	100 29 32.8	2.34	1	Cl; P; IC; pS; st 9...10, 12...13	1
4421	3752	Δ. 607	18 26 32.0	+3.936	4	123 5 5.1	-2.43	4	B; S; IE; rrr; st 15	5

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
4422	h. 3750	H.	h m s 18 27 8.1	s +7.511	1	163° 22' 49.3"	-2.59	1	vF; S; lE; glbM.....	1
4423	Auw. N. 42	18 27 19.3	3.210	...	96 4 46.7	2.45	...	pF; vS; E (Winnecke, June 1855).	0
4424	{ 2015 = 3753 }	M. 22	18 27 52.1	3.662	4	114 0 25.8	2.54	4	!!; ⊕; vB; vL; R; vRi; vmC; st 11...15.	10
4425	2016	18 28 5.5	2.496	1	66 32 22.1	2.53	1	Cl; P; IC.....	1
4426	3754	VIII. 12	18 29 5.1	3.266	1	98 20 0.1	2.63	1	Cl; L; pRi; vLc.....	2
4427	3755	18 32 2.0	5.603	2	152 25 36.5	2.95	2	pF; S; R; psbM; r.....	2
4428	3756	M. 70=Δ. 614	18 34 4.9	3.910	3	122 25 12.6	3.08	3	⊕; B; pL; R; gbM; st 14...17	4*
4429	2017	18 34 9.9	3.185	1	94 53 29.9	3.07	1	Cl; L; Ri; st 10...18	1
4430	2018	18 34 41.1	3.219	1	96 21 12.4	3.12	1	Cl; vRi; vLc (in M. Way)...	1
4431	3757	18 35 9.5	5.941	2	155 19 18.8	3.24	2	vB; pL; R; vg, psbM; *7p	2
4432	3758	M. 26	18 37 32.6	3.293	1	99 31 57.9	3.37	1	Cl; cL; pRi; pC; st 12...15.	6
4433	VI. 15	18 39 37.8	3.715	1	116 3 40.2	3.56	1	Cl (?); vF; cL	1
4434	3759	18 40 6.8	5.131	1	147 27 46.8	3.64	1	pF; pS; lE 90°; psbM	1
4435	Auw. N. 43	18 43 18.8	3.192	...	95 22 6.4	3.86	...	Cl; B; 60 st 13 (Winnecke, 1854).	0
4436	3760	18 43 36.6	5.679	1	153 19 39.2	3.96	1	Neb. No description.....	1
4437	2019	M. 11	18 43 37.2	3.219	2	96 26 7.6	3.88	2	!; Cl; vB; L; iR; Ri; st 1 L, 11.	8+
4438	3761	18 44 2.1	4.873	2	143 58 46.9	3.97	2	F; S; vLE; gbM	3
4439	4021	h. o. n.	18 44 18.1	4.866	2	143 53 19.3	3.99	2	pF; S; R; glMbM; last of gr	2
4440	2020	18 44 52.5	2.836	1	79 49 0.6	3.98	1	Cl; pRi; lC; iF	1
4441	3762	I. 47	18 45 29.2	3.276	1	98 52 8.5	4.05	1	⊕; pB; vL; ir; vglbM; rrr	2
4442	3763	M. 54=Δ. 624	18 46 6.2	3.846	3	120 38 29.7	4.11	3	⊕; vB; L; R; g, smbM; rrr; st 15.	6
4443	2021	18 46 17.5	3.549	1	110 4 8.1	4.13	1	Cl; pRi; st 9...13	1
4444	{ 2022 = 3766 }	III. 143	18 46 39.8	3.623	2	112 52 17.2	4.16	2	F; S; rr Cl + neb	3
4445	3764	18 47 25.9	6.042	2	156 17 14.0	4.30	2	vF; S; R; glbM; *9 sp	2
4446	3765	18 48 2.6	6.430	1	158 47 1.2	4.36	2	vF; pL; R; vglvbm	2
4447	2023	{ M. 57 D'Arquier }	18 48 20.1	2.228	2	57 8 57.2	4.26	3	{ !!!; ⊙; B; pL; cE (in Lyra). }	14+
4448	3767	18 48 51.4	5.153	1	147 56 49.3	4.39	1	pF; cS; R; vmbM	1
4449	3768	18 49 59.0	5.874	1	155 5 21.7	4.51	1	pF; S; E; glbM; 2 st 8 p	1
4450	3770	Δ. 573	18 50 5.5	4.047	1	126 48 45.9	4.47	1	⊕; vL; lE; vglbM; rrr; st 14...16.	1
4451	2024	18 50 25.1	2.838	1:	79 49 16.2	4.46	1	Cluster.....	1
4452	3769	18 50 26.1	4.871	1:	144 7 32.7	4.51	1	cF; pL; R	1
4453	VIII. 13	18 52 20.9	3.281	1	99 7 33.8	4.64	1	Cl; vL; P	1
4454	3771	18 52 32.0	6.472	1	159 6 45.8	4.74	1	vF; S; R; pmbM; *7, 8 nf	1
4455	3772	18 53 15.0	5.552	1	152 23 37.6	4.78	1	eeF; vglbM; v difficult	1
4456	3773	18 53 29.2	5.938	1	155 39 40.7	4.81	1	vF; S; R; glbM; p of 2	1
4457	2025	18 53 33.1	3.086	1	90 39 5.1	4.73	1	Cl; vL; P; st 12...	1
4458	3774	18 53 44.6	5.934	1	155 38 10.4	4.82	1	eF; S; R; glbM; f of 2	1
4459	2026	18 53 59.2	5.512	1	108 44 8.6	4.78	1	Cl; pL; pRi; R; st 12...15...	1
4460	2027	18 54 50.0	2.810	1	78 35 18.1	4.83	1	Cl; P; lC.....	1
4461	3775	18 55 7.2	5.463	2	151 34 23.8	4.94	2	cF; vS; cE; psbM; 3 st p	2
4462	III. 742	D'Arrest, 113	18 55 32.0	1.618	1	41 45 14.2	4.86	1	vF; stellar	1*
4463	2028	18 55 58.2	2.351	1	60 55 35.4	4.92	1	Cl; pL; P; st 11...12	1
4464	3776	Δ. 262	18 56 26.8	5.730	2	154 3 58.5	5.05	2	cB; cL; R; vg, svmbM; r	2
4465	3777	18 57 31.7	5.518	1	152 10 36.1	5.13	1	eF; cS; R; glbM	1
4466	2029	18 57 56.6	3.035	1	88 25 11.0	5.10	1	Cl; L; lC; st L & S	1
4467	3778	Δ. 295	18 58 28.0	5.324	5	150 11 49.7	5.21	5	⊕; B; vL; iR; rrr; st 11...16	5
4468	3779	18 59 37.5	5.085	1	147 15 30.0	5.30	1	pB; pL; R; gbM	1
4469	3780	19 0 33.6	4.653	2	140 51. 32.9	5.37	2	pF; pL; mE 63°; vglbM	2
4470	2030	VII. 19	19 0 50.5	2.981	3	85 59 17.5	5.35	3	Cl; vL; vRi; pC; st 12...14.	5
4471	2031	VII. 62	19 1 46.5	2.971	2	85 32 43.1	5.43	1	Cl; S; Ri; lC; st 11...12	5
4472	3781	19 2 15.9	5.014	1	146 32 24.1	5.53	1	pB; S; R	1
4473	Auw. N. 44	19 4 4.8	+3.051	Au	89 11 51	-5.62	Au	!!; pB; pL; gbM; Var.? (Hind)	0*

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ′ ″	″			
4474	3782	19 4 14.3	+4.647	2	140 53 11.6	-5.68	2	vF; pS; iR	2
4475	3786	19 6 50.4	4.160	1	130 25 44.6	5.88	1	vF; S; R; psbM	1
4476	3783	19 6 53.8	5.349	1	150 44 16.4	5.92	1	vF; S; R; lbM; 1st of 3.....	1
4477	3784	19 7 6.3	5.349	1	150 44 46.1	5.93	1	eF; vS; 2nd of 3.....	1
4478	3785	19 7 7.3	5.351	1	150 46 25.8	5.94	1	eF; S; 3rd of 3	1
4479	2032	IV. 14	19 7 14.8	3.137	2	92 57 5.3	5.89	2	vF; L; R; vvlbM; r	3
4480	2033	19 8 3.7	2.968	1	85 22 9.5	5.95	1	Cl; P; lC	1
4481	2034	19 8 38.7	3.449	1	106 30 34.4	6.02	1	Cl; vL; lC	2
4482	2035	19 9 37.9	3.097	1	91 9 48.3	6.09	1	Cl; P; lC; st 10...11	1
4483	3787	19 10 22.9	5.690	1	154 8 23.1	6.23	1	pB; S; R; pgbM	1
4484	Lac. I. 13	19 10 45.5	6.917	...	161 45 35.7	6.21	...	Neb without st (Lac. Auw. 40)	0
4485	2036	M. 56	19 11 7.2	2.339	4	60 3 41.6	6.18	6	⊕; B; L; iR; gvmCM; rrr; st 11...14.	14
4486	3788	19 11 17.6	4.954	1	146 1 45.6	6.28	1	vF; L; R; vglbM	1
4487	2037	III. 743	19 11 37.3	2.931	1	83 42 46.5	6.25	1	○; F; L; R; vsbM disc; S* nf	3*†
4488	3789	19 11 39.2	5.283	3	150 10 43.4	6.32	3	cF; cS; R; lbM; * 9 s	3
4489	3790	19 12 48.3	5.896	2	155 52 59.1	6.43	2	eeeF; pS; am S st	3
4490	2038	19 13 23.1	3.102	1:	91 21 22.0	6.40	1	eS; stellar	1
4491	3791	19 14 30.8	4.890	2	145 13 36.8	6.54	2	pB; S; mE; psbM	2
4492	Auw. N. 45	19 15 55.9	2.098	...	52 28 32.1	6.53	...	vF (Winnecke, Dec. 1853)	0
4493	2039	VIII. 81	19 17 13.9	2.565	3	68 6 5.0	6.70	3	Cl; P; lC	4
4494	3792	19 18 23.8	4.094	1	129 9 44.8	6.84	1	eF; pS; R; vglbM	1
4495	2040	19 19 6.7	3.000	1	86 44 46.9	6.87	1	Cl; Ri; bet 2 st 9	1
4496	3793	19 20 47.4	4.941	2	146 11 39.2	7.06	2	eF; vS; R; lbM; 3 vS st nr	2
4497	2041	VIII. 21	19 21 22.5	2.491	1	65 8 23.8	7.04	1	Cl; vL; pRi; vLC; st 10...	3
4498	2042	VI. 14	19 24 29.7	2.628	3	70 1 4.0	7.30	3	Cl; L; vC; E 0°; st 14...18	5
4499	2043	VI. 38	19 24 53.0	2.875	4	81 3 37.8	7.34	4	eB; S; iR; rrr	5
4500	3796	19 27 18.0	4.068	1	128 51 32.9	7.57	1	eF; R; vgbM	1
4501	3795	19 27 18.9	4.209	1	132 36 18.9	7.57	1	eF; vS; * 14 att	1
4502	3794	19 28 30.3	6.638	2	160 57 27.8	7.74	2	pB; E; biN; * 8 f	2
4503	3798	M. 55 = Δ. 620	19 31 7.9	3.817	2	121 15 44.2	7.86	2	{ ⊕; pB; L; R; vRi; } vgbM; st 12...15.	5
4504	3797	19 31 43.0	5.109	2	148 58 44.5	7.95	2	pS; R; vgbM	2
4505	2044	19 33 58.6	1.790	2	43 44 48.8	8.04	2	Cl; L; pRi; lC; st 11...14 ...	2
4506	3799	19 34 7.2	4.858	1	145 40 17.8	8.14	1	pB; pS; pmE; glbM	1
4507	2045	III. 744	19 35 0.0	3.300	...	100 38 29.9	8.17	...	pF; pL; R; bM; r	2
4508	2046	19 35 6.9	2.462	1:	63 30 55.5	8.15	1:	Cl; vL; pRi; lC; st 10...15	1
4509	3800	19 35 16.0	3.744	1	118 52 41.0	8.20	1	eF; pS; R; vlbM; * np	1
4510	2047	IV. 51	19 36 3.0	3.386	3	104 28 52.5	8.25	3	○; B; vS; R	8†
4511	2048	Harding	19 36 24.3	2.053	1	50 7 59.5	8.25	1	Cl; vL; vRi; st 11...15 (Harding, 1827).	1
4512	2049	VII. 18	19 37 13.6	2.555	1	67 1 48.4	8.32	1	Cl; cRi; E; st 11...12	2
4513	II. 878	19 40 47.3	1.298	1	34 18 29.2	8.56	1	pB; iF; bM	1
4514	2050	IV. 73	19 41 7.5	1.622	1	39 49 41.7	8.61	1	⊙; B; pL; R; * 11 M	2†
4515	2051	VIII. 73	19 43 30.0	2.912	1	82 26 25.1	8.83	2	Cl; P; lC	3
4516	2052	VII. 9	19 45 4.3	2.569	1	67 15 45.5	8.95	1	Cl; L; pRi; pC; st 11...12 ..	2
4517	2053	19 45 43.7	1.072	1	30 55 55.2	8.96	1	Cl; vL; lC; st 7,	1
4518	2054	VIII. 16	19 46 33.8	2.408	1	60 57 6.5	9.05	1	Cl; P; lC; st 11...12	2
4519	2055	VIII. 18	19 46 53.0	2.832	1:	78 40 27.3	9.09	1:	Cl; S; P	2
4520	2056	M. 71	19 47 28.8	2.674	3	71 34 55.1	9.13	3	Cl; vL; vRi; pmC; st 11...16	13
4521	2057	VI. 16?	19 48 13.9	2.696	1:	72 28 24.0	9.20	1:	Cl; vS; vC	3
4522	2058	VIII. 19	19 48 41.7	2.824	1	78 15 35.1	9.23	1	Cl; P; lC	3
4523	3802	19 48 57.9	3.818	1	122 11 30.6	9.28	1	vF; S; R; psbM	1
4524	2059	19 49 31.7	2.823	1	78 12 33.0	9.30	1	Cl; S; P	1
4525	3801	19 49 42.9	5.672	1	155 36 45.6	9.38	1	eF; vS; R; psbM; * 11 np...	1
4526	3803	19 50 49.6	4.355	1	137 27 28.8	9.44	1	vF; S; vIE; glbM	2
4527	II. 202	19 51 22.9	2.417	1	61 1 41.1	9.43	1	Neb; r	1
4528	3804	19 51 33.2	4.845	2	146 28 18.7	9.51	2	eF; eL; R; vglbM; 2 st f ...	2
4529	3805	19 51 47.6	4.079	1	130 35 26.7	9.51	1	pB; S; R; vS * np	1
4530	3806	19 52 27.6	4.760	1	145 13 47.9	9.57	1	vF; S; R; bM	1
4531	3807	19 53 19.4	+4.402	1	138 39 10.1	-9.63	1	pF; S; vIE; psbM	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos	Other Authorities.								
4532	h. 2060	H.	M. 27	19 53 29.3	+2.588	4	67° 39' 43.0"	- 9.60	3	!!!; vB; vL; BiN; iE (Dumb-bell N).	13+
4533	3808	19 54 40.9	4.723	2	144 45 44.5	9.75	2	F; S; vLE; glbM.....	2
4534	3809	19 55 31.7	4.849	1	146 47 1.7	9.81	1	pF; S; R.....	1
4535	2063	19 56 8.8	1.354	1	34 15 9.9	9.77	1	Cl; pS; pmC; iR; st 12...16	1
4536	2062	III. 144	19 56 23.1	2.309	3	56 50 48.7	9.81	3	F; am M. Way st.....	4*
4537	2061	19 56 23.6	2.847	1	79 7 24.4	9.82	1	Cl; cL; E; pRi; st 13... ..	1
4538	Auw. N. 46	19 56 39.0	3.068	...	89 56 49.0	9.85	...	F; *10 p 1 ^s , s 1' 29" (Bond, Nov. 1852).	0
4539	3810	19 56 41.0	5.207	1	151 29 30.7	9.91	1	F; pS; gbM.....	1
4540	3811	Δ. 425	19 57 5.0	4.395	2	138 46 5.7	9.91	2	B; S; cE; gpmbM.....	2
4541	3812	19 57 39.9	4.849	1	146 47 18.2	9.96	1	F; S; iE; glbM.....	1
4542	2065	19 57 46.5	3.151	1	93 57 0.8	9.94	1	Cl; S; vmC; st 19.....	1
4543	2064	M. 75	19 57 49.1	3.547	3	112 18 47.5	9.95	3	⊕; B; pL; R; vmbMBN; rr	10
4544	2066	VII. 59	19 59 11.2	1.967	1	46 23 49.7	10.01	1	Cl; L; vRi; cC.....	2
4545	3813	19 59 31.7	4.733	1	145 11 4.4	10.12	1	eeF; L; pmE.....	1
4546	3814	19 59 40.6	4.387	2	138 46 26.4	10.12	2	vB; S; R; pgvmbM.....	2
4547	3815	19 59 57.7	4.382	2	138 41 20.8	10.14	2	cF; cS; E 90°; gbM.....	2
4548	2067	Σ. 2630	20 0 38.3	2.258	1	54 37 13.1	10.13	1	Cl; st L & S; * inv.....	1
4549	3816	20 2 2.9	6.426	2	161 11 51.8	10.34	2	F; pS; iE; glbM; *9 p 10 ^s .5;	2
4550	2068	Σ. 2631	20 2 4.8	2.636	1	69 17 54.5	10.25	1	Cl; iC; st 10...13; * inv.....	1
4551	2069	VIII. 86	20 2 44.4	2.179	1	52 9 34.3	10.29	1	Cl; P; iC.....	2
4552	3819	20 3 15.0	4.278	1	136 34 21.6	10.38	1	F; vS; R; vgmbM; *7 nf ...	1
4553	3817	20 3 27.6	6.429	2	161 17 6.5	10.45	2	pB; S; R; eS*sf; 2nd of 4...	2
4554	3818	20 3 44.5	6.428	2	161 17 10.6	10.48	2	vF; vS; R; 3rd of 4.....	2
4555	3821	20 4 2.8	4.209	1	134 56 31.8	10.44	1	vF; L; R; glbM.....	1
4556	3820	20 4 40.6	6.419	2	161 17 3.8	10.54	2	F; S; R; r; vS*att; 4th of 4	2
4557	VIII. 22	20 5 58.2	2.509	1	63 42 34.8	10.54	1	Cl; P; iC.....	1
4558	2070	20 5 59.3	2.261	1	54 34 19.8	10.54	1	Cl; pRi; * inv.....	1
4559	2071	VIII. 20	20 6 7.1	2.514	1	63 55 43.5	10.55	1	Cl; vB; vL; Ri; iC; st 6...11	2
4560	3822	20 6 37.5	4.584	1	143 12 32.8	10.64	1	pF; cL; pmE; glbM.....	1
4561	IV. 72	20 7 22.0	2.185	1	52 1 28.8	10.64	1	F; vL; vmE; * att.....	1
4562	3823	20 8 5.5	4.644	1	144 23 0.5	10.75	1	vF; L; iE.....	1
4563	3824	20 8 29.7	4.206	2	135 13 51.9	10.77	2	pF; S; R; vglbM.....	2
4564	3825	20 10 43.4	4.344	2	138 40 53.1	10.93	2	pF; S; R; svbM*12.....	2
4565	2072	IV. 13	20 10 45.8	2.419	2	59 51 33.3	10.89	2	!!; ⊙; F; S; vviE.....	7+
4566	VIII. 83	20 12 23.7	1.746	1	40 12 6.0	11.00	1	Cl; pRi; iC.....	1
4567	D'Arrest, 114	20 12 26	2.42	[2]	59 47 42	11.01	[2]	Cl+neb; S; st vS.....	0
4568	3826	20 14 4.0	4.433	1	140 52 0.6	11.18	1	F; S; R; glbM; am st.....	1
4569	3827	20 14 52.7	4.143	1	134 5 51.8	11.24	1	F; cS; R; bM.....	1
4570	2073	20 15 40.2	3.468	3	109 45 13.6	11.28	3	cL; E; bM*17; *10 att.....	3*
4571	2074	20 15 46.1	2.547	1	64 41 32.2	11.26	1	Cl; S; vLC; st 10...11.....	1
4572	2075	IV. 16	20 16 7.9	2.676	3	70 20 19.3	11.29	3	!!; ⊙; B; pS; R; 4Sst nr ...	6+
4573	2076	III. 141	20 16 44.2	3.590	1	115 14 51.2	11.36	1	cF; cL; vLE; vglbM; r; 3st p	3
4574	3828	20 17 43.6	4.267	1	137 28 57.8	11.44	1	pB; pL; gbM; 2st 10 nr.....	1
4575	2077	VIII. 56	20 18 5.5	2.137	2	49 40 14.4	11.42	2	Cl; pB; pS; P; pC; st 10...12	5
4576	2078	M. 29	20 18 51.9	2.212	1	51 56 3.6	11.48	1	Cl; P; iC; st L & S.....	6
4577	3830	20 20 50.3	4.274	1	137 56 31.9	11.67	1	vF; *12 att sp.....	1
4578	3831	20 22 0.0	4.141	1	134 40 55.8	11.74	1	eF; pS; R; vglbM.....	1
4579	3829	20 22 36.0	9.441	1	170 28 50.4	11.92	1	pB; cS; R; psmbM.....	1
4580	3832	20 23 2.0	3.727	4	121 17 38.7	11.81	4	pF; cS; R; gbM; bet 2 st ...	4
4581	3834	20 25 35.6	3.752	3	122 26 56.3	11.99	3	cB; L; mE 6°0; pslbM.....	3
4582	2079	III. 142	20 25 52.1	3.118	2	92 29 44.3	11.99	2	vF; pL; E 0° or biN; p of 2	5
4583	2080	20 26 8.3	3.118	2	92 30 28.7	12.01	2	vF; vS; sf of 2.....	2
4584	3833	20 26 29.1	6.798	1	164 6 50.1	12.13	1	F; S; R; gbM; 5 st p.....	1
4585	I. 103	20 26 43.0	2.935	1	82 44 5.5	12.05	1	vB; L; gmbM; er.....	1
4586	2081	20 27 19.8	2.941	3	83 3 41.3	12.09	3	⊕; B; L; R; rrr; st 16...; *9 p	3*
4587	3835	20 27 59.4	4.457	2	142 35 10.9	12.17	2	pB; cL; R; glbM; r.....	2
4588	3836	20 28 25.0	4.458	2	142 38 11.7	12.21	2	vF; cS; R; slbM; f of 2.....	2
4589	2082	VIII. 17	20 28 35.0	+2.646	1	68 13 54.9	-12.17	1:	Cl; vL; P; vLC.....	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.			Annual Precession in Right Ascension for 1880.			No. of Obs. used.	North Polar Distance for 1860, Jan. 0.		Annual Precession in N.P.D. for 1880.		No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.	h	m	s	s	°	'		''	''	''				
4590	2083	VI. 42	20	28	36.4	+1.211	3	29	49	56.8	-12.14	3	Cl; pL; eRi; pCM; st 11...16	5		
4591	VII. 8	20	28	42.5	2.510	5	62	9	51.9	12.17	5	Cl; vB; vL; vRi; cC; st pL..	5		
4592	3837	20	30	3.7	4.563	1	144	47	44.4	12.32	1	pB; pL; R; psbM	1		
4593	3838	20	31	3.8	5.831	1	159	14	17.4	12.42	1	pF; L; mE; vglbMvS*	1		
4594	2084	IV. 76	20	31	57.3	1.269	3	30	20	17.9	12.37	3	vF; vL; vg; vsbM; rr.....	4†		
4595	3839	20	32	30.6	3.750	1	122	58	55.9	12.47	1	vF; L; R; gbM	1		
4596	3840	20	33	4.6	4.497	2	143	51	23.1	12.53	2	vF; pS; cE; lbM	2		
4597	2085	VIII. 23	20	34	40.7	2.769	1	73	59	36.0	12.60	1	Cl; P; vLC	2		
4598	2086	III. 219	20	37	14.2	2.852	1	77	59	34.9	12.77	1	vF; S; stellar; * att	2		
4599	3841	20	39	39.2	3.882	1	128	30	26.2	12.96	1	B; cS; R; pgmbM; 4 st p ...	1		
4600	2088	V. 15	20	39	53.0	2.478	3	59	47	14.8	12.94	2	!!; pB; cL; eiF; h Cygni inv	6†		
4601	2087	II. 426	20	40	7.8	3.074	2	90	11	12.9	12.97	2	eF; S; R; bM.....	4		
4602	2087, a	R. 3 novæ	20	40	±	3.074	...	90	11	±	12.97	...	Group of 5 with many st ...	0		
4603																	
4604																	
4605	2089	II. 427	20	40	13.6	3.074	2	90	12	34.6	12.98	2	F; vS; R; bM.....	4		
4606	3842	20	42	15.1	4.243	1	139	17	41.8	13.14	1	pB; S; lE; gbM	1		
4607	II. 206	20	45	16.1	2.454	1	58	23	48.0	13.30	1	F; S; iE; r	1		
4608	2090	M. 72	20	45	44.9	3.302	3	103	3	35.8	13.34	3	⊕; pB; pL; R; gmCM; rrr..	10		
4609	3843	20	47	8.1	4.356	2	142	24	21.2	13.46	2	vF; S; E; p of 2	2		
4610	3844	20	47	18.9	4.043	1	134	30	30.2	13.46	1	eF; cS; R	1		
4611	3845	20	47	48.0	4.353	3	142	24	12.7	13.51	3	F; pL; vLE; vgbM; f of 2 ...	3		
4612	3846	20	48	20.9	4.211	2	139	10	13.8	13.54	2	pF; S; vLE; pgmbM; vB * p	2		
4613	VIII. 82	20	49	13.4	2.092	1	45	15	32.5	13.55	1	Cl; cL; st pS	1		
4614	3847	20	49	27.4	4.529	2	146	6	24.7	13.61	2	eeF; vS; vmE0°; *13 att, n	2		
4615	2091	VIII. 76	20	49	51.7	2.024	1	43	15	22.6	13.58	1	Cl; L; P; vLC	3		
4616	2092	V. 14	20	50	35.2	2.478	2	58	50	11.8	13.64	1	!!; eF; eL; eE; eiF; bifurcate	4†		
4617	M. 73	20	51	16.0	3.299	...	103	10	31.0	13.70	...	Cl??; eP; vLC; no neb	3		
4618	2093	20	51	19.8	2.490	1	59	19	15.3	13.69	1	F; eL; neb & st	1†*		
4619	2094	20	51	30.9	2.095	1	45	3	59.3	13.69	1	Cl; P; lC	2		
4620	VIII. 58	20	51	33.5	2.123	3	45	53	24.0	13.70	3	Cl; P; lC; st L	3		
4621	2096	V. 37?	20	53	48.2	2.142	(1)	46	13	4.8	13.84	1:	F; eeL; diff nebulosity	3		
4622	2095	20	53	54.5	3.083	1	90	44	34.9	13.87	1	eF; S; E 0°	1		
4623	3848	20	53	57.1	4.201	2	139	35	0.3	13.89	2	eF; cS; R; bM	2		
4624	3849	20	54	13.9	4.203	1	139	40	27.7	13.91	1	eF; R; lbM; *11 f	1		
4625	2097	I. 52	20	54	43.8	2.801	(3)	74	21	51.7	13.91	1	B; pL; R; gbM	4		
4626	3850	20	55	23.1	4.345	3	143	6	3.6	13.98	3	pB; S; R; psbM; am st	3		
4627	2099	I. 192	20	56	17.5	1.749	1	35	59	39.6	13.98	2	eB; L; E 45°±; r; * att ...	5†		
4628	2098	IV. 1	Lal. 40765	20	56	31.2	3.273	4	101	55	4.8	14.04	4	!!!; O; vB; S; elliptic	23*†		
4629	2100	20	57	5.8	3.291	1	103	3	8.9	14.07	1	eF; pL; R; r	1		
4630	2101	20	57	8.1	2.054	1	43	13	48.5	14.05	1	Cluster; no description	1		
4631	3851	20	57	21.1	4.033	2	135	22	12.0	14.10	2	F; pL; E; vglbM; * p.....	2		
4632	2102	II. 203	20	57	38.9	2.535	2	60	39	30.3	14.09	2	pB; cS; R; psbM; pB*np...	4		
4633	3852	20	58	18.4	4.111	1	137	44	44.2	14.16	1	pF; S; R; bM; 2 st 12 n ...	1		
4634	IV. 74	20	59	22.9	0.771	1	22	26	51.2	14.16	1	eF; * 7 m in neb (?)	1		
4635	3853	20	59	38.5	5.136	1	154	35	45.2	14.26	1	pB; cS; lE; pgbM	1		
4636	3854	20	59	39.2	5.000	1	154	5	34.2	14.26	1	pF; cS; R; psbM; *7.8 p ...	1		
4637	3855	20	59	45.4	4.185	1	139	52	10.5	14.25	1	eeF; S; R; B *** sf	1		
4638	2103	VIII. 57	21	0	46.2	2.255	2	49	3	58.9	14.27	2	Cl; P; lC; st 10...	3		
4639	3856	21	2	7.3	4.173	1	139	51	26.0	14.40	1	B; cS; R; pgmbM	1		
4640	2105	VIII. 74	21	2	45.7	1.948	(1)	39	43	13.3	14.39	1	Cl of triple st; lC	2		
4641	3857	21	2	47.1	5.457	1	158	51	57.9	14.47	1	vF; cS; R; glbM	1		
4642	2104	21	3	36.5	2.825	1	75	7	5.2	14.46	1	Cl; lC	1		
4643	2106	21	5	3.4	2.471	1	56	50	48.0	14.60	1	Cl; pRi; iF; st 11...15	1		
4644	3858	21	5	39.3	4.079	2	137	47	45.7	14.61	2	pB; pL; lE; gbM	2		
4645	2107	21	6	14.3	2.150	1	44	53	32.7	14.61	1	Cl; vL; pRi; E; st 10... ..	1		
4646	III. 209	21	6	52.3	2.860	1	76	58	16.2	14.66	1	vF; S; R	1		
4647	3859	21	6	55.9	4.114	2	138	56	41.3	14.69	2	B; cS; cE; psmbM; *10 f...	2		
4648	2110	VI. 24	21	7	45.4	2.253	2	48	4	44.0	14.70	2	Cl; vF; pL; vRi; vC; st 15...18	4		
4649	2108	21	7	48.6	3.008	1	86	3	33.4	14.72	1	eF	1		
4650	2109	III. 858	21	7	52.9	+3.035	2	87	44	17.4	-14.72	2	eF; pL; R; lbM.....	3		

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
4651	3860	Δ. 406	21 9 26.5	+4.109	2	139 8 52.8	-14.84	2	vB; pS; E; mbM	2
4652	2111	21 9 33.0	2.424	1	54 23 16.7	14.81	1	Cl; no description	1
4653	2112	III. 145	21 12 36.6	2.640	1	64 9 17.3	14.99	1	F; S; vLE; r	3
4654	2113	21 14 20.1	3.216	1	99 22 29.7	15.11	1	vF; R; gbM; * nr	1*
4655	2114	21 15 26.3	1.717	1	32 59 50.8	15.14	1	Cl; F; pS; P	1
4656	3861	21 15 58.1	3.884	1	133 3 41.7	15.21	1	eF; vS; R; p of 2	1
4657	2115	21 16 16.6	2.021	1	39 47 4.3	15.19	1	Cl; P; IC	1
4658	3862	21 16 41.6	4.607	1	150 37 16.9	15.27	1	B; pL; IC; gpmbM	1
4659	3863	21 16 53.5	3.879	1	133 0 26.2	15.26	1	vF; pS; R; f of 2	1
4660	3864	21 17 58.9	4.085	1	139 40 10.1	15.33	1	eeF; vS; R	1
4661	2116	VII. 51	21 18 12.1	2.181	3	44 13 3.7	15.31	3	Cl; pS; pRi; pC; st 13... ..	4
4662	2117	21 18 44.9	2.447	1	54 5 44.1	15.33	1	Cl; P; st 10... ..	1
4663	3865	21 19 9.1	4.222	1	143 23 17.3	15.39	1	eF; pL; vmE 90°8; *s	1
4664	2118	VII. 50	21 19 37.1	2.135	1	42 34 45.6	15.38	1	Cl; P; ? neb	2
4665	3866	21 21 26.5	3.879	2	133 41 50.4	15.52	2	F; cL; lE; gvlbM; p of 2 ...	2
4666	2119	21 21 31.0	2.148	1	42 40 21.6	15.48	1	Cl; S; C; cE	1
4667	3867	21 21 37.5	3.880	2	133 45 57.1	15.53	2	F; S; R; vglbM; f of 2	2
4668	3868	21 22 49.4	3.757	1	129 14 1.3	15.59	2	cF; cS; R; pgbM	2
4669	III. 936	21 22 51.7	1.459	1	27 43 29.5	15.55	1	vB; er	1
4670	2120	M. 15= Lal. 40815	21 23 9.9	2.895	1	78 27 22.3	15.59	2	{ !; ⊕; vB; vL; iR; } vsmbM; rrr; st vs. }	16
4671	3869	21 23 31.9	3.898	1	134 40 55.1	15.63	1	B; R; cS; psbM	1
4672	2121	III. 859	21 24 18.3	3.043	1	88 7 23.2	15.66	1	F; S; R; mbM; *14s	2
4673	2122	VII. 52	21 24 22.5	2.188	1	43 31 7.5	15.65	1	Cl; L; cRi; IC; st 10...13 ...	2
4674	3870	Δ. 263?	21 24 38.7	4.809	2	154 31 11.7	15.71	2	pF; cL; vlE; vgpmbM; r ...	2
4675	2123	21 25 11.3	2.819	1	73 11 28.0	15.70	1	Cl; IC	1
4676	2124	VI. 32	21 25 42.8	2.044	2	39 1 55.7	15.71	2	Cl; cL; vRi; pC; st 11...16	3
4677	3871	21 25 44.2	3.800	2	131 26 32.5	15.75	2	cF; S; R; gbM	2
4678	2125	M. 2= Lal. 41928	21 26 12.5	3.091	2	91 26 37.2	15.76	3	{ !!; ⊕; B; vL; g; pmbM; } rrr; st eS. }	19+
4679	3872	21 26 36.3	4.251	1	145 10 46.0	15.80	1	pB; pL; vmE 127°1; g; pslbM	1
4680	3873	21 26 51.0	3.699	1:	127 24 5.7	15.81	1:	eF; pL; vgbM; *6 f 40s.....	1
4681	2126	M. 39	21 27 12.6	2.160	1	42 10 58.0	15.80	1	Cl; vL; vP; vIC; st 7...10...	3
4682	2127	21 29 5.1	2.244	1	44 37 6.0	15.90	1	Cl; P; IC.....	1
4683	3875	21 30 3.9	3.827	2	133 10 21.6	15.98	1	F; pL; R; vglbM; *13 inv...	2
4684	3874	21 30 20.1	4.755	1	154 32 2.4	16.02	1	vF; S; R; vS * nf	1
4685	3877	21 31 24.1	3.821	2	133 10 19.5	16.05	2	B; S; vlE; mbM.....	2
4686	3876	21 31 59.5	6.179	1	165 44 25.1	16.13	1	pF; R; g; psmbM; am st ...	1
4687	{ 2128 = 3878	M. 30	21 32 26.0	3.422	2	113 47 59.0	16.10	2	!; ⊕; B; L; lE; gpmbM; st 12...16.	3+
4688	3879	21 32 59.7	4.134	1	143 20 29.8	16.14	1	eF; cS; lE; vglbM.....	1
4689	3880	21 33 22.2	3.873	1	135 25 37.2	16.16	1	vF; cL; R; vglbM	1
4690	3881	21 33 31.6	3.626	1	125 4 56.2	16.16	1	eF; vS; am st	1
4691	3882	21 33 48.4	3.620	1	124 48 14.9	16.17	2	F; S; R; bM	2
4692	3883	21 36 32.5	3.965	2	139 3 59.4	16.32	2	F; S; R; glbM; p of 2	2
4693	3884	21 36 57.1	3.961	2	138 59 48.8	16.34	2	F; S; R; glbM; f of 2	2
4694	3885	21 37 10.8	3.904	1	137 9 38.5	16.35	1	F; S; R; gbM	1
4695	Auw. N. 47	21 38 22.8	3.296	...	99 28 19.2	16.36	...	Nebulous *10-11 or vSCL (Cooper).	0
4696	3886	21 38 39.8	5.278	1	160 58 43.5	16.45	1	pB; S; R; vgbM; *9 f	1
4697	3888	21 38 45.1	4.022	3	141 12 49.8	16.44	3	pB; L; pmE; vgbM	3
4698	3887	21 38 56.2	4.466	1	151 21 18.5	16.45	1	eF; pL; R; p of 2	1
4699	3889	21 38 58.8	4.460	1	151 15 13.5	16.45	1	pB; pS; lE; gbM; f of 2 ...	1
4700	2129	21 39 8.8	2.013	1	36 1 22.4	16.42	1	Cl; S; P; IC	1
4701	2130	VII. 40	21 39 21.8	2.049	1	36 56 49.1	16.43	1	Cl; S; pRi; has a ruby * ...	2
4702	2131	IV. 75	21 39 45.0	1.387	2	24 32 15.8	16.44	2	!; cF; pL; gbM **	4
4703	3890	21 39 51.6	3.605	1	125 5 43.6	16.48	1	pB; S; R; glbM	1
4704	3891	21 41 22.0	3.609	2	125 31 43.2	16.56	2	pB; pL; R; vgbM; *14 att p	2
4705	2132	II. 261	21 41 44.2	2.773	2	68 29 7.2	16.56	2	F; pS; R; vglbM; r	3
4706	III. 696	21 42 15.3	+1.576	4	26 49 29.9	-16.57	4	vF; cS; R; r	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
4707	h. 3892	H.		h m s 21 42 28.6	s +4.229	1	147° 12' 6.1"	-16.63	1	pF; cS; R; bM	1
4708	3893			21 42 30.4	4.187	1	146 13 34.1	16.63	1	F; L; R; g, psmbM	1
4709	2134	VII. 66		21 42 34.2	1.441	1	24 50 39.6	16.58	1	Cl; cL; cRi; pC; st 11...14	3
4710	2133			21 42 41.2	2.650	1	60 41 28.0	16.60	1	vF; ? a * inv in neb	1*
4711	3894			21 43 34.1	3.921	2	138 54 28.9	16.67	2	vB; pS; R; mbMN	2
4712	3895			21 44 13.2	3.907	1	138 32 26.0	16.70	1	B; S; R; in Δ of st 13.	1
5077				21 45 12.5		...	40 53 46.4		...	See No. 5077.	
4713	3896			21 45 46.0	3.984	3	141 19 11.6	16.78	3	vF; pL; lE; vgbM; r	3
4714	3897			21 45 50.0	3.493	1	119 56 36.9	16.77	1	eeF; vS	1*
4715	3898			21 46 30.0	3.486	1	119 41 30.0	16.80	1	eF; S; E towards eF*	1
4716	3900			21 46 58.7	3.588	2	125 28 21.1	16.83	2	B; pL; iR; glbM; r	2
4717	3899			21 46 59.2	3.940	2	140 10 49.8	16.84	2	pB; S; lE; mbM	2
4718	2135	III. 452		21 47 29.0	3.042	3	87 42 49.8	16.84	3	F; pL; R; bM; r	4
4719	2136	VIII. 67		21 49 48.4	1.722	1	28 3 3.1	16.93	1	Cl; P; vC	2
4720			D'Arrest, 115	21 49 53	3.04	[2]	87 42 42	16.95	[2]	Cl; vS; st 19 m; bet 2 st 16...	0
4721	3901			21 50 58.1	3.752	1	133 58 28.4	17.02	1	cF; cL; cE; glbM	2
4722	3902			21 51 10.5	3.521	3	122 33 8.4	17.02	3	F; pL; vLE; vglbM	3
4723	2137	III. 930		21 51 49.3	3.289	1	107 10 39.5	17.05	1	eB	2*
4724	3903			21 51 53.6	3.750	1	134 3 32.2	17.06	1	cF; S; vLE; svbMN	2
4725	3905			21 52 37.2	3.401	1	115 18 18.3	17.09	1	F; pS; R; vglbM; *10 f	1
4726	3904			21 52 54.3	3.975	3	142 25 3.4	17.12	3	pB; S; R; psbM	3
4727	3906			21 53 6.8	3.852	1	138 21 46.4	17.12	1	eF; S; R; *8 np	1
4728	2138	III. 692		21 53 29.4	3.243	1	103 57 3.1	17.13	3	vF; cL; E 135°±; vgbM	4
4729	3908			21 53 52.3	3.513	4	122 32 38.5	17.15	4	pB; pL; lE; gbM; 1st of 4...	4+
4730	3909			21 53 53.1	3.514	4	122 38 8.5	17.15	4	cB; cS; R; sbM*; 2nd of 4	4+
4731	3910			21 53 55.3	3.515	2	122 39 58.5	17.15	2	cF; S; R; p of D neb; 3rd of 4	2+
4732	2141			21 53 55.9	2.106	1	35 50 51.1	17.13	1	Cl; vL; pRi; lC	1
4733	3911			21 53 59.0	3.514	6	122 39 31.5	17.15	6	B; pL; R; f of D neb; 4th of 4	6+
4734	2139	II. 247		21 54 0.1	2.858	1	72 56 7.5	17.15	3	pB; pS; R; bMN; r; *sp...	4+
4735	3912			21 54 6.3	3.582	1	126 28 45.2	17.16	1	eF; S; R; *8 s 2'	1
4736	3907			21 54 21.4	4.525	2	154 43 13.6	17.18	2	cF; pS; vgbM	2
4737	2140	III. 693		21 54 26.8	3.339	1	111 13 26.9	17.17	1	vF; S; R; lbM; p of 2	2
4738	2142	II. 595		21 54 38.9	3.316	1	109 35 22.6	17.18	2	vF; pS; vLE 90°; lbM	4
4739	2143	II. 1		21 54 51.2	3.341	1	111 28 50.3	17.19	1	pB; pL; mE 64° 3 bet 3 st; er	3
4740	2144			21 55 8.7	3.337	1	111 8 59.7	17.21	1	vF; pL; iR; vglbM; f of 2...	1
4741		III. 165		21 55 13.6	2.599	1	55 33 55.0	17.20	1	vF; am 5 or 6 st	1
4742	3913			21 56 20.4	4.543	1	155 19 8.9	17.27	1	vF; S; lE; vgbM	1
4743	3914			21 56 24.7	4.521	2	154 59 23.6	17.28	2	pB; S; R; pmbM	2
4744	2145			21 56 42.8	2.949	1	79 51 21.9	17.27	1	Cl; lRi; lC; st 9...10	1
4745	3915			21 56 50.6	3.900	3	140 47 57.3	17.29	4	cB; S; R; am st	4
4746	2146	II. 599		21 57 7.6	2.493	1	49 37 29.6	17.28	1	F; cS; cE; vglbM; er	2
4747	3916			21 58 4.3	4.529	2	155 22 59.5	17.35	2	vF; S; R; psbM; *11 p 3'	2
4748	3917			21 58 7.9	3.839	1	140 40 49.5	17.35	1	pF; S; R; smbM	1
4749	3918			21 58 27.0	3.488	3	121 55 30.5	17.35	3	F; R; gbM; 1st of 4	3
4750	3920			21 58 39.2	3.486	1	121 51 49.2	17.36	1	eF; S; stellar; 2nd of 4	1
4751	3921			21 58 40.2	3.488	3	121 49 42.2	17.36	3	cF; R; stellar; 3rd of 4	3
4752	3922			21 58 47.8	3.484	1	121 43 31.9	17.37	1	pB; L; lE; gbM; 4th of 4	1
4753	3919			21 58 53.7	4.140	1	148 6 29.6	17.38	1	pB; L; cE; gpslbM	2
4754	3923			21 59 24.6	3.451	1	119 43 57.3	17.39	1	vF; vS; R; almost a ○	1
4755	2147	VII. 53		21 59 42.2	2.384	2	44 11 36.3	17.39	2	Cl; L; cRi; pC; st 9...12	4
4756	2148			21 59 57.7	2.741	1	63 34 17.7	17.41	1	eF; R; bM; vF * np	1*
4757	3924			22 0 29.0	3.798	1	137 50 36.5	17.45	1	vB; pS; R; gbM	1
4758	3926			22 1 8.3	3.427	1	118 29 42.9	17.47	1	⊕; pL; iR; rr	1
4759	3925			22 1 30.8	4.794	1	159 20 46.7	17.51	1	pF; S; R; gbM	1
4760	2149	II. 207		22 1 37.4	2.682	1	59 20 35.6	17.48	1	B; pL; gbM; er	2
4761	2150	II. 897		22 2 34.9	3.273	1	107 19 55.1	17.53	1	pB; lE; r	2
4762	3927			22 2 47.3	4.487	2	155 32 6.2	17.56	2	pB; S; R; 2 st nr.	2
4763	3928			22 3 13.8	3.462	4	121 14 22.2	17.56	4	F; S; R; gbM; r; 2vS st nr.	4
4764	2151	III. 862		22 4 16.2	2.525	1	49 40 48.3	17.59	1	eF; pS; lE; r; am 3 st.	2
4765	3929			22 5 15.4	3.392	1	116 50 37.8	17.64	1	pF; S; lE; bM	1
4766	3930			22 6 4.9	+3.436	4	120 4 3.9	-17.67	4	F; pL; R; vglbM	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
4767	h. 2152	H. III. 931	h m s 22 6 37.7	s +3.272	1	107 45 40.3	-17.69	1	vF; S; R; bM.....	2
4768	2153	II. 606	22 6 43.4	2.447	1	45 20 52.3	17.69	1	eF; S; er.....	4
4769	3931	22 6 57.4	3.731	1	136 32 31.7	17.71	1	pB; S; pmE; psbM; p of 2...	1
4770	3932	22 7 7.9	3.730	1	136 32 16.4	17.72	1	F; vS; R; *8 f; f of 2.....	1
4771	VIII. 63	22 7 8.2	2.127	1	33 42 56.0	17.70	1	Cl; S; P; IC.....	1
4772	2154	22 7 34.6	2.119	2	33 25 12.4	17.72	2	Cl; pC; has a ruby *10.....	2
4773	2155	VIII. 75	22 9 41.1	2.358	1	40 48 55.7	17.81	1	Cl; L; P; IC; st vL.....	3
4774	2157	VI. 29	22 10 2.5	2.236	1	36 22 2.4	17.82	1	Cl; C; st eS.....	2
4775	2156	III. 932	22 10 8.7	3.249	1	106 16 47.8	17.84	1	vF; S; vLE; vglbM; *13n ...	2*
4776	III. 863	22 11 0.4	2.563	1	50 10 13.2	17.86	1	vF; vS; mbM.....	1
4777	3933	22 11 26.0	3.957	1	145 48 53.0	17.90	1	eeF; R; (?).....	1
4778	III. 864	22 12 28.8	2.569	1	50 8 9.4	17.92	1	vF; S; mE 165°+.....	1
4779	2158	III. 933	22 12 53.8	3.247	2	106 28 21.5	17.95	2	F; pS; R; gpmbM.....	4
4780	3934	III. 458	22 12 57.6	3.353	1	115 22 44.5	17.95	1	F; S; R; er.....	2
4781	3935	22 15 6.5	3.395	1	119 3 25.8	18.04	1	vF; S; E; glbM; ?biN.....	1
4782	3936	22 15 10.0	3.403	1	119 39 16.8	18.04	1	eF; pL; R; vlbM.....	1
4783	2159	22 15 23.6	2.153	1	32 36 51.1	18.03	1	Cl; L; pRi; IC.....	1
4784	3937	22 15 26.0	3.450	1	123 3 31.5	18.05	1	eF; S; R; lbM.....	1
4785	3938	22 16 21.1	3.466	2	124 24 1.6	18.08	1	cB; pS; vLE; glbM; B ** sp	2
4786	3939	22 16 44.2	3.429	3	121 53 45.0	18.10	3	F; cS; vLE; p of 2.....	4
4787	3940	22 18 15.5	3.424	4	121 51 23.5	18.15	3	F; cS; vLE; f of 2.....	4
4788	3941	22 18 57.4	4.079	1	150 52 54.3	18.19	1	eeF; lE; vglvbm; 3 st sf ...	1
4789	3942	22 19 7.4	3.478	2	125 51 19.3	18.19	2	vF; pS; R; vglvbm.....	2
4790	2160	II. 248	22 19 38.1	2.916	2	74 33 55.0	18.20	2	F; cS; R; gbMS*; 3 st nr...	4
4791	2161	22 19 39.4	2.198	1	32 52 26.3	18.19	1	Cl; L; pRi; IC; st 10...16...	1
4792	3943	II. 469	22 20 50.7	3.336	2	115 34 20.5	18.25	2	cF; cS; lE; r; * inv.....	3
5078	22 20 50.±	115 34 ±	(See No. 5078).....	0
4793	2162	22 20 52.2	2.770	1	61 36 48.8	18.24	1	vF; S; R; am st.....	1
4794	3944	22 21 12.0	3.474	1	126 10 21.2	18.26	1	vF; S; R; gbM.....	1
4795	Auw. N. 48	22 22 5.6	3.285	...	111 32 59.8	18.29	...	!; pF; vL; EorbiN (Harding)	0
4796	2163	22 22 21.2	2.365	1	37 53 26.3	18.29	1	Cl; P; lC; st 12...13.....	1
4797	VII. 41	22 22 37.4	2.380	2	38 24 45.0	18.30	2	Cl; iR; lC; st vS.....	2
4798	3945	22 22 59.4	3.503	1	128 33 0.1	18.33	1	eF; S; R; p of 2.....	2
4799	2165, a	R. nova	22 23 13.6	3.213	::	104 40 30.3	18.34	::	vF; E np to sf (nisi=h.2164)	0
4800	3946	22 23 23.8	3.501	1	128 32 9.8	18.34	1	eF; S; R; f of 2.....	2
4801	2164	22 23 28.3	3.213	2	104 44 11.8	18.34	2	vF; cS; vglbM.....	2
4802	2165	IV. 31	22 24 53.0	3.213	4	104 50 30.3	18.39	4	F; pS; R; vsbMFSRN.....	5
4803	D'Arrest, 116	22 24 57	2.76	[1]	59 45 18	18.38	[1]	vF; pL; vlbM; h.2166 dist 2'	0
4804	2166	22 24 58.6	2.760	2	59 45 25.3	18.39	2	vF; S; R; vglvbm.....	2
4805	3948	22 25 28.5	3.352	1	117 58 4.4	18.42	1	vF; S; lE; *11 p.....	1
4806	3947	22 25 32.6	3.541	1	131 39 53.4	18.42	1	F; pL; pmE.....	1
4807	2167	II. 476	22 26 55.7	3.173	5	101 4 53.2	18.46	5	vF; pL; R; glbM; r.....	6
4808	2168	II. 428	22 27 3.9	3.027	2	85 9 17.9	18.47	2	pF; S; R; psbM; r.....	4
4809	2169	III. 180	22 27 40.9	2.886	1::	70 23 45.6	18.48	1::	eF; vS; R; *9 s.....	2
4810	3949	22 28 2.3	3.331	2	116 45 55.0	18.50	3	cB; L; mE 0°; vlbM.....	3
4811	2170	III. 237	22 29 46.2	2.877	1	69 5 50.5	18.55	1	F; S; iR; vglvbm.....	2
4812	3950	22 29 46.2	3.466	2	127 57 13.5	18.55	2	vF; S; vLE; gbM.....	2
4813	2171	22 30 32.5	2.984	3	80 11 4.6	18.58	4	eF; pS; lE 90°; vglbM.....	4
4814	3951	22 30 37.2	4.257	1	157 12 16.0	18.60	1	pB; pS; mE 90°.....	1
4815	2172	I. 53	22 30 39.5	2.736	1	56 20 5.6	18.58	3	B; pL; pmE 160°; smbM ...	4†
4816	} 2172, a	R. 5 novæ	22 30 ±	2.736	...	56 20 ±	18.58	...	5 near; positions measured; no distances.	0*
4817											
4818											
4819											
4820	} 2173	II. 233	22 30 44.0	2.857	2	66 55 34.6	18.58	2	cB; S; mE 163° 0; vsmbM *11; p of 2.	5
4821											
4822	3950	(No. 2?)	22 30 50.5	3.466	1	127 55 48.4	18.62	1	eF; ??.....	1?
4823	2174	III. 166	22 30 54.0	2.736	1	56 17 51.3	18.59	1	eF; vS; E.....	2
4824	2175	II. 234	22 31 5.7	+2.858	2	66 55 58.3	-18.59	2	F; pS; mE 90°±; vglbM; f of 2.	5

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
4825	h. 2176	H.	h m s 22 32 57.9	s +2.982	1	79° 42' 40.2"	-18.66	1	eF; pL; E	2
4826	2177	22 34 15.4	2.333	1	33 20 2.3	18.69	1	Cl; vL; pRi; vLc	1
4827	2178	II. 705	22 35 6.6	2.218	1	29 27 5.4	18.72	1	B; S; R; pglbM; er	2
4828	3952	22 35 26.3	3.451	1	128 36 27.8	18.74	1	eeF; S; R; * f 90°, 40s	1
4829	3953	Δ. 255?	22 36 9.6	4.117	1	155 51 19.9	18.77	1	F; S; R; bM	1
4830	3954	22 36 30.8	3.351	1	120 47 3.9	18.77	1	F; pL; vmE0°; vglbM	1
4831	2179	II. 442	22 37 14.0	3.079	2	90 53 42.0	18.80	2	F; S; R; psbM	5
4832	3955	22 37 30.5	3.475	1	130 4 35.7	18.81	1	F; cS; lE; glbM	1
4833	2180	II. 477	22 38 40.4	3.166	4	101 44 29.8	18.84	5	vF; pL; R; lbM	6
4834	2181	II. 598	22 40 12.8	3.262	3	113 2 10.3	18.89	3	pB; S; vLE; vgmbM; S * nr	4
4835	} 2181, a	R. 7 novæ	22 40 ±	3.262	...	113 2 ±	18.89	...	No descriptions	0
4836											
4837											
4838											
4839											
4840	2182	VIII. 77	C. H.	22 41 23.2	2.374	2	32 39 8.7	18.91	2	Cl; pL; pRi; lC; st 9...13 ..	3
4843	3956	22 42 23.0	3.408	1	127 34 56.5	18.95	1	eF; vS; R; * 12 att np	1
4844	{ D'Arrest, 117 = R. nova }	22 42 30	2.99	[2]::	79 10 30	18.94	[2]	vF; vS; R; III. 216 nf	0
4845	2183	III. 216	22 42 56.0	2.988	1	79 7 46.2	18.96	2	cF; S; R; glbM; *11 np ...	5
4846	2184	III. 217	22 43 2.5	2.987	1	79 2 36.2	18.96	2	cF; S; R; pgbM; f of 2	5
4847	D'Arrest, 118	22 43 20	2.99	[2]	79 6 0	18.97	[2]	eF; vS; R; 2 st 11, s	0
4848	2184, a	R. nova	22 43 22.5	2.99	...	79 8 2.2	18.96	...	One of 5. See h. 2183, 2184, D'Arrest, 117, 118.	0*
4849	2185	II. 443	22 43 25.0	3.088	2	92 16 54.6	18.98	2	cF; cS; R; sbM *13; * np ..	4
4850	2186	II. 702	22 44 15.4	3.238	3	111 21 1.0	19.00	3	pB; pS; lE 120° ±; mbM ...	4
4851	2187	II. 453	22 44 21.1	3.118	2	96 18 6.0	19.00	2	vF; pL; lE; vgbM; r	3
4852	2188	22 44 36.2	2.534	1	38 33 54.0	19.00	1	Cl; vP	1
4853	2189	22 45 12.8	3.069	2	89 38 56.4	19.02	2	pF; pS; R; gbM	2
4854	} 2189, a	R. 3 novæ	22 45 ±	3.069	...	89 39 ±	19.02	...	"A group of 4," incl h. 2189	0
4855											
4856											
4857											
4858											
4859	3957	22 45 42.2	3.506	1	136 5 36.8	19.04	1	pF; lE; glbM; vS * inv	1
4858	3958	22 46 23.9	3.420	1	130 3 30.2	19.06	1	vF; S; R	1
4859	3959	22 46 49.9	3.930	1	154 26 29.6	19.08	2	pB; pS; R; vglbM	1
4860	3960	Δ. 518	22 47 2.8	3.421	2	130 24 35.6	19.08	2	cB; L; vmE 43° 3; mbM ...	2
4861	3961	22 47 45.7	3.456	1	133 23 49.0	19.10	1	eF; vL; * 7 nf	1
4862	3962	22 48 42.2	3.940	1	155 46 26.1	19.13	1	pB; cS; R; gpmbM	2
4863	3963	22 48 42.4	3.383	3	127 46 23.4	19.12	3	cB; vL; vLE; vglbM	3
4864	2190	VII. 43	22 48 46.6	2.356	2	29 55 6.3	19.09	2	Cl; pRi; cC	3
4865	3964	22 49 1.9	3.385	3	128 5 31.1	19.13	4	cB; L; vLE; gpmbM; rr	4
4866	2191	III. 745	22 49 18.9	2.466	1	33 38 14.1	19.13	1	vF; pL; iF; er	2
4867	3965	22 49 21.4	3.428	2	131 49 4.8	19.14	2	F; cL; vLE; vgmbM	2
4868	2192	III. 576	22 49 28.5	2.784	1	54 22 38.8	19.14	1	vF; cS; R; stellar; * p	3
4869	2193	22 50 18.4	2.404	1	30 45 23.2	19.16	1	Cl; P; pC; st 9...11	1
4870	2194	III. 465	22 51 3.9	2.986	1	77 37 14.6	19.18	1	eF; S; R	4
4871	2195	III. 243	22 51 12.3	2.886	1	64 37 4.6	19.18	2	F; pS; E 90°; gbM; er	3
4872	} 2195, a	R. 2 novæ	22 51 ±	2.886	...	64 37 ±	19.18	...	{ 2 of a group of 3; inv in F neb. }	0
4873											
4874	2196	22 51 36.5	2.543	1	36 23 55.3	19.19	1	Cl; vL; E	1
4875	D'Arrest, 119	22 52 32	2.97	[2]	75 12 12	19.21	[2]	pF; R; bet 2 st 16; *13 nr...	0
4876	2197	II. 450	22 52 46.6	3.163	2	103 33 10.4	19.22	2	F; vS; vLE; smbM; er; p of 2	4+
4877	2198	II. 451	22 52 46.9	3.163	2	103 34 46.4	19.22	3	F; vS; vLE; smbM; er; f of 2	5+
4878	Auw. N. 49	22 53 6.3	3.144	...	101 16 41.1	19.23	...	*11.12 in neb (Markree Obs. Oct. 8, 1855).	0
4879	2199	II. 251	22 53 8.0	2.968	2	74 46 10.1	19.23	2	pB; L; E 175°; vgbM	3
4880	II. 249	22 54 3.1	2.967	2	74 21 17.2	19.26	2	F; cS; lE; lbM; pB * p	2
4881	3966	22 54 13.6	+3.387	1	130 19 19.2	-19.26	1	F; L; mE 33° 8; vglbM	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
4934	D'Arrest, 121	h m s	s	[2]	82° 42' 24"	-19°64	[2]	F; S; R; Δ with 2 st 19, n ...	0
4935	2229	23 12 51.0	3.036	1	82 21 19.5	19.65	1	eF; eS	1
4936	2230	II. 439	23 13 9.5	3.037	2	82 33 59.5	19.65	2	cB; pS; R; psbM	3
4937	3983	23 13 21.1	3.584	1	152 53 32.2	19.66	1	eF; eS; am 5 st; (?)	1
4938	2231	III. 435	23 13 25.2	3.036	1	82 21 49.2	19.66	1	F; vS; R; psbM	2
4939	2232	II. 250	23 13 27.2	2.994	2	73 32 34.2	19.66	2	pB; cS; R; smbM	4
4940	2233	II. 440	23 13 38.0	3.037	2	82 32 58.2	19.66	1	cB; pS; R; psbM	3
4941	D'Arrest	23 14 5.8	2.998	[4]	74 12 23.9	19.67	[4]	cB; pS; bM* (D'Arrest, Resultate).	0*
4942	2231, a	R. nova	23 14 13.7	3.036	::	82 21 49	19.67	::	E p and f	0
4943	D'Arrest, 122	23 14 22	3.04	[2]	82 33 6	19.67	[2]	vF; vS	0
4944	3985	23 14 21.5	3.314	2	133 14 52.6	19.68	2	F; S; R; lbM	2
4945	3986	23 14 30.5	3.717	2	158 25 49.6	19.68	2	F; vS; E90°; psbM	2
4946	2234	II. 441	23 14 36.4	3.035	1	81 52 38.6	19.68	1	F; S; F* att	2
4947	2235	IV. 52	23 14 38.8	2.616	1	29 35 10.6	19.68	1	vF; *9 inv a l excentric	3
4948	3987	23 15 3.6	3.219	1	120 2 41.3	19.69	1	eF; S; R; sbM	1
4949	3984	23 15 6.0	4.986	1	172 40 19.0	19.70	1	vF; pL; R; vlbM; *nr	1
4950	2236	II. 600	23 15 23.6	2.858	2	49 54 46.3	19.69	2	eF; L; mE0°±; vlbM; r ...	4+
4951	3988	23 16 17.5	3.215	1	120 8 51.7	19.71	1	vF; S; R; glbM; *nr	1
4952	III. 473	23 16 47.9	3.002	1	74 0 18.4	19.72	1	eF; cL; sc st f; (?)	1
4953	III. 218	23 16 48.7	3.032	1	80 50 18.4	19.72	1	eF; pS; lE	1
4954	3989	23 17 17.8	3.464	2	148 33 48.1	19.73	2	pF; pS; R; glbM; p of 2 ...	2
4955	3990	23 17 37.9	3.462	1	148 39 59.1	19.73	1	eF; S; R; f of 2	1
4956	2237	23 17 51.8	3.010	1	75 29 12.1	19.73	1	vF; pS; R; gbM	2
4957	2238	M. 52	23 18 3.2	2.643	1	29 10 20.1	19.73	1	Cl; L; Ri; mCM; R; st9...13	8
4958	3991	23 18 14.6	3.674	1	158 47 39.8	19.74	1	eF; vS; R; psbM; *10 p 22°	2
4959	3992	23 18 40.6	3.452	1	148 34 57.5	19.75	1	eF; R	1
4960	3994	23 18 48.0	3.266	1	129 59 55.5	19.75	2	{ eF; S; R } D neb; 4 st p..	2
4961	2239	III. 212	23 18 52.9	3.016	1	76 33 33.5	19.75	1	vF; vS; R; psbM	2
4962	2240	23 18 52.9	2.956	1	63 44 7.5	19.75	1	F; vS; psmbM; *10 p	1
4963	3993	23 18 56.4	3.590	1	156 2 49.5	19.75	1	eF; cL; R; vgvlbM	1
4964	2241	IV. 18	23 19 9.9	2.864	5	48 13 57.5	19.75	5	!!!; O; vB; pS; R; blue.....	16*+
4965	III. 438	23 20 0.9	3.112	1	100 11 16.2	19.76	1	eF; S; stellar	1
4966	2242	III. 226	23 20 15.0	3.025	4	78 18 33.9	19.77	4	{ (H.) vF } S; R; vsmbM; { (h.) pB } *9 p (? var).	6*
4967	2242, a	R. nova	23 20 15.0	3.025	::	78 24 33.9	19.77	::	vF; S; 6' s of h. 2242	0
4968	2243	23 20 49.8	3.040	1	81 59 10.6	19.78	1	F; cS; gbM; p of 2	1
4969	2244	23 20 59.8	3.041	1	81 59 55.6	19.78	1	vF; S; R; gbM; f of 2	1
4970	3995	23 21 2.6	3.457	1	150 29 2.6	19.78	1	B; S; lE; vsmbM*11	1
4971	2245	II. 226	23 21 28.0	2.985	2	68 20 38.3	17.79	2	vF; pL; vLE; lbM; am 4 st...	3+
4972	2246	III. 860	23 21 36.7	2.938	(1)	58 25 6.3	19.79	1:	vF; S; R; lbM; r	2
4973	2247	II. 242	23 21 52.1	3.008	2	73 28 4.3	19.79	2	vF; S; iR; r; * f	4
4974	D'Arrest, 123	23 21 55	3.06	[1]	87 19 4	19.79	[2]	eF; *14 p 13°7, 1n.....	0
4975	2248	III. 426	23 23 21.9	3.061	1	86 52 7.7	19.81	1	eF; cL; R; gbM; *nr	3
4976	2249	VIII. 69	23 23 29.0	2.839	2	41 38 50.7	19.81	2	Cl; P; lC; st 7...11	4
4977	D'Arrest, 124	23 23 47	3.06	[2]	87 12 42	19.81	[2]	vF; vS; *11 f 1° s, n 85''	0
4978	3996	Δ. 347?	23 24 48.4	3.352	1	144 52 22.1	19.83	1	pF; L; R; vgbM	1
4979	3997	23 25 20.0	3.325	2	142 28 11.8	19.84	2	cB; S; lE; psbM; *8 f	2
4980	2250	III. 213	23 25 20.6	3.019	1	74 55 35.8	19.84	1	eF; pL; Δ with 2 st 10	2*
5079	23 25 33.1	96 22 12.0	See No. 5079.	
4981	III. 187	23 25 56.9	3.084	2	93 31 10.5	19.85	2	eF; pL; stellar	1
4982	D'Arrest, 125	23 26 4	3.09	[1]	93 39 12	19.85	[1]	eF; pL; 3 st 11 & 12 f.....	0
4983	3998	23 26 48.5	3.494	1	156 19 38.2	19.86	1	eeF; pL (certain)	1
4984	III. 188	23 27 3.0	3.084	1	93 41 10.2	19.86	1	eF; stellar	1
4985	3999	23 27 43.9	3.347	2	146 47 2.9	19.87	2	B; cS; E; g, sbM; *8.9 p ...	2
4986	2251	23 27 44.2	3.022	1	74 42 26.9	19.87	1	vF; vS; gbM; *nf 1'	1
4987	2252	23 27 53.0	3.059	2	85 52 26.9	19.87	2	eF; *12 p; sp of 2	2
4988	2253	23 28 2.0	3.058	1	85 49 1.9	19.87	1	vF; nf of 2	1
4989	2254	III. 579	23 28 2.6	2.900	2	46 27 42.9	19.87	2	eF; S; R; *9-10 p, v nr.....	3
4990	2255	VIII. 62	23 28 15.5	+2.510	2	17 51 23.9	-19.87	2	Cl; L; P; lC; st 8, 10...15...	5

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
4991	h. 2256	H. II. 244	h m s 23 28 35.2	s +3.026	1	75° 27' 48.6"	-19.88	2	F; S; R; psbM; stellar	4
4992	4000	23 28 56.7	3.209	1	128 13 2.6	19.88	1	pB; L; E; vgbM.....	1
4993	2257	23 29 2.6	3.067	1	88 36 51.3	19.89	1	pB; S; R; psbM; *12 sp ...	1
4994	2257, a	R. nova	23 29 10.6	3.067	::	88 36 51.3	19.89	::	No description	0
4995	2258	23 29 20.0	3.073	1	90 28 43.3	19.89	1	F; pL; lE; gbM; *10 s	1
4996	2259	III. 146	23 31 28.2	2.992	1	63 45 34.7	19.91	1	F; S; lE; bM; am st	2
4997	2260	II. 432	23 31 35.5	3.092	3	97 17 32.7	19.91	3	pF; cL; E 12° ±; vgbM.....	7
4998	2261	I. 110	23 31 41.7	3.111	2	103 43 56.7	19.91	2	{ (H.) cB } ; cL; E; gmbM; { (h.) eF } r (? var.)	4*
4999	III. 189	23 32 21.2	3.087	1	95 24 5.4	19.92	1	eeF	1
5000	2262	I. 111	23 32 38.8	3.107	2	103 4 12.4	19.92	2	pB; pL; iR; mbM	4
5001	4001	23 34 41.4	3.397	1	156 45 1.5	19.95	1	eF; S; R; p of 2	1
5002	4002	23 34 49.4	3.396	1	156 44 21.5	19.95	1	cF; cS; R; f of 2	1
5003	2263	II. 208?	23 35 17.3	3.006	1	64 32 13.5	19.95	1	vF; *14 att 255°	1*
5004	II. 208	23 36 45.2	3.010	1	64 38 3.2	19.96	1	cL; R; *10-11 np	1*
5005	2264	II. 255	23 37 8.5	3.049	3	80 0 26.2	19.96	5	cB; cS; gmbM; r; B * f.....	7
5006	2265	II. 256	23 37 13.4	3.051	1	80 50 20.2	19.96	3	pB; S; R; *15 sf	4
5007	4003	23 37 35.5	3.190	3	133 41 27.9	19.97	3	cF; S; vLE; svbmM *14	3
5008	2266	23 38 26.1	2.757	1	21 1 30.9	19.97	1	vL; surrounds *8	1
5009	4004	23 38 29.5	3.141	2	120 17 4.6	19.98	2	vF; S; R; gmbM; *12 f ...	2
5010	2267	III. 427	23 39 27.3	3.066	2	86 58 39.6	19.98	2	cF; pL; vLE 0°; lbM; 2 Bstr	4
5011	2268	II. 213	23 40 2.3	3.011	1	61 17 42.3	19.99	1	cF; cL; vLE; vglbM; r	2
5012	4005	23 40 35.8	3.137	2	121 17 45.3	19.99	2	B; cL; R; psbmM	2
5013	2269	III. 437	23 40 50.1	3.064	1	83 54 46.3	19.99	2	F; S; R; gbM; er	3
5014	2270	23 41 35.4	3.066	1	86 36 8.0	20.00	2	vF; cL; R; vglbM; *13 n ...	2
5015	2271	III. 854	23 42 7.7	3.015	5	59 47 45.0	20.00	5	cB; vS; R; psbM; rr	7*
5016	2272	VII. 55	23 43 4.8	2.849	1	22 45 49.7	20.01	1	Cl; pRi; pC; st 11...15	4
5017	4006	23 43 34.1	3.151	1	131 31 2.7	20.01	1	B; pL; R; gbM	1
5018	2273, a	R. nova	23 43 ±	3.028	...	63 37 ±	20.01	...	vvF; a little np h. 2273	0
5019	2273	23 43 52.2	3.028	1	63 37 55.7	20.01	1	vF; S; iF; vF * inv	1
5020	2274	II. 230	23 43 55.7	3.041	2	70 37 26.7	20.01	2	pF; pS; R; mbM; sp of 2 ...	4*
5021	2274, a	R. nova	23 44 13.9	3.04	...	70 40 43.7	20.01	...	Seen and meas. with h. 2274, 2275.	0
5022	2275	II. 231	23 44 17.1	3.042	2	70 39 16.7	20.01	2	pB; pL; E 90°; bM; nf of 2	4
5023	2276	23 44 38.0	3.049	1	74 31 22.4	20.02	1	Cl of sc st 10 m	1
5024	2277	II. 851	23 45 4.8	3.024	2	59 31 6.4	20.02	3	vF; cS; R; * nf	5
5025	2278	III. 231	23 46 11.5	3.063	2	82 54 5.4	20.02	2	cF; S; R; psbM; stellar; 1st of 4.	3
5026	2279	III. 232	23 46 19.0	3.063	2	82 53 50.4	20.02	2	pF; S; R; psbM; stellar; 2nd of 4.	3
5027	2280	23 46 38.2	3.063	1	82 54 18.1	20.03	1	F; S; R; 3rd of 4	1
5028	2281	III. 233	23 46 45.2	3.063	2	82 48 10.1	20.03	2	pF; pL; lE; glbM; 4th of 4	4
5029	2282	II. 468	23 48 10.1	3.066	4	84 51 58.1	20.03	4	pB; pS; iR; psbM; r; *7 p 30°	5
5030	2283	23 49 42.8	2.975	2	29 23 16.8	20.04	2	Cl; S; pRi; vC; st 10, 13...	2
5031	2284	VI. 30	C. H.	23 49 58.5	2.994	1	34 3 46.8	20.04	1	Cl; vL; vRi; vmC; st 11...18	3
5032	2285	VII. 56	23 50 0.7	2.979	1	29 33 50.8	20.04	1	Cl; pRi; pC.....	2
5033	2286	23 50 47.4	3.063	1	80 0 18.8	20.04	1	vF; vS; (?)	1
5034	2287	23 51 34.3	2.999	1	30 45 41.8	20.04	1	Cl; vL; P; lC; st 7, 10...	1
5035	4009	23 51 45.2	3.136	1	146 14 16.8	20.04	1	pB; cS; R; gmbM	1
5036	2288	III. 466	23 51 47.0	3.065	1:	80 2 53.5	20.05	1	vF; pS; iR	2
5037	2289	III. 867	23 51 50.3	3.070	1	87 8 41.5	20.05	1	eF; pS; iR; lbM.....	2
5038	2290	II. 232	23 52 17.4	3.058	1	70 0 13.5	20.05	1	pF; S; R; sbM; * 10 sp.....	3
5039	2291	II. 10	33 52 28.1	3.063	1	75 58 24.5	20.05	1	F; pS; iE 15° ±	3
5040	2292	23 53 17.8	3.033	1	40 3 38.5	20.05	1	Cl; pRi; pC; st 9... ..	1
5041	2293	23 53 49.6	3.069	1	84 32 7.5	20.05	1	vF; S; R; psbM	1
5042	2294	III. 855	23 54 14.4	3.056	2	59 20 47.5	20.05	3	eF; S; R; sbM; stellar sp of 2	4
5043	2295	III. 856	23 54 17.0	3.056	3	59 20 0.5	20.05	3	eF; S; R; stellar; nf of 2 ...	4
5044	2296	III. 984	23 55 10.4	3.067	2	77 48 0.5	20.05	3	eF; stellar; Δ with 2 st	4*
5045	4010	23 55 43.1	3.085	1	125 1 38.5	20.05	1	vF; S; R; am st	1
5046	2297	II. 240	23 56 4.5	3.067	1	74 37 31.5	20.05	1	cB; cL; ir; vgbM	2+
5047	2298	III. 436	23 56 32.1	+3.070	1::	83 25 16.5	-20.05	1	vF; pL; R; lbM.....	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
5048	2299	23 56 39.7	+3.070	1	83 17 43.5	-20.05	1	vF; pL; R; gbM.....	1
5049	2300	II. 227	23 56 48.7	3.067	1	70 0 3.5	20.05	1	pF; cL; mE 45°±; lbM.....	4
5050	2301	23 57 19.7	3.072	1	85 34 38.5	20.05	1	pB; vS; mE; vsmbM.....	2
5051	2302	23 57 32.6	3.051	1	22 6 18.2	20.06	1	!; eeF; eeL.....	1*
5052	4011	23 57 35.2	3.087	2	152 50 42.2	20.06	2	F; S; R; gbM.....	2
5053	2303	23 57 55.7	3.072	1	83 51 42.2	20.06	1	pF; S; R; * 9 np.....	1
5054	2304	23 57 56.2	3.072	1	85 34 12.2	20.06	1	vF; S; R; gbM.....	1
5055	2305	VIII. 29	23 58 1.9	3.070	1	111 29 37.2	20.06	1	Cl; vP; vIc.....	2
5056	2306	23 58 17.7	3.072	1	85 33 47.2	20.06	1	vF; S; R; * nf.....	1
5057	4013	III. 190	23 59 17.4	+3.072	1	94 29 48.2	-20.06	1	vF; vS; R; vg, psmbM; 2 st 9 sf.	2

SUPPLEMENTARY LIST OF NEBULÆ AND CLUSTERS.

5058	G. P. Bond	0 35 1.0	+3.073	...	89 50 6.6	-19.80	...	F; S; R; * 11 sp 1'; disc Sept. 16, 1862.	0
5059	79, b	R. nova	0 50 7.9	3.241	...	60 23 27	19.56	...	No description; γ in Lord R.'s diagram.	0
5060	S. Coolidge	3 6 55.2	3.089	...	89 4 27.5	13.67	...	F; disc Jan 25, 1860.....	0
5061	S. Coolidge	3 16 29.0	+3.086	...	89 19 3.0	13.07	...	F; disc Dec. 16, 1859.....	0
5062	(123)	4 57 18.4	-0.385	1	159 36 32.1	- 5.43	1	No description.....	1
5063	(374)	5 20 52.9	-0.469	1::	159 36 35.4	- 3.42	1::	No description.....	1
5064	J. H. Safford	6 5 7 :	+3.093	...	88 50 39 :	+ 0.43	...	2 Cls; near 2 st 9.10 & 10.11; disc Mar. 19, 1863.	0
5065	J. H. Safford	6 6 40.9	3.093	...	88 58 11.2	0.44	...	Cl; bet 2 st 9.10 & 10.11; disc Mar. 19, 1863.	0
5066	G. P. Bond	7 55 12.5	3.509	...	65 25 19.4	9.73	...	vF; cometic; disc Sept. 1, 1852.	0
5067	S. Coolidge	9 59 51.1	3.079	...	89 15 7.3	17.38	...	Neb; no description; disc Mar. 31, 1859.	0
5068	S. Coolidge	10 16 14.1	3.079	...	89 13 46.0	18.05	...	F; disc Mar. 31, 1859.....	0
5069	939, c	R. nova	11 33 ±	3.122	...	71 29 ±	19.93	...	No description.....	0
5070	2849, a	D'Arrest	12 12 41	3.06	...	83 46 42	20.02	...	A nebula; no description ...	0*
5071	S. Coolidge	12 31 0.5	3.069	...	89 2 46.5	19.87	...	*12, in F neb; disc May 3, 1859.	0
5072	S. Coolidge	13 24 32.0	3.067	...	89 18 29.7	18.67	...	*12, in F neb; disc Apr. 30, 1859.	0
5073	S. Coolidge	13 42 38.3	3.065	...	89 14 2.4	18.03	...	*12, in F neb; disc Apr. 30, 1859.	0
5074	G. P. Bond	13 49 18.6	3.068	...	89 31 9.0	17.77	...	S; R; *9'; disc June 8, 1855.	0
5075	S. Coolidge	13 53 58.0	3.065	...	89 13 53.3	+17.60	...	*12, in neb; disc Apr. 29, 1859.	0
5076	Lac. I. 11	18 20 44.6	3.952	...	123 30 27.3	- 1.89	...	Neb. without stars.....	0
5077	G. P. Bond	21 45 12.5	2.217	...	40 53 46.4	16.67	...	Neb; no description; disc Feb. 10, 1848.	0
5078	3943, a	Lassell	22 20 50 :	3.336	...	115 34 ±	18.25	...	Neb* * 1' dist from h. 3943 ..	0
5079	G. P. Bond	23 25 33.1	+3.093	...	96 22 12.0	-19.83	...	Neb; * 9'10 sf; disc Oct. 23, 1848.	0

Of this supplementary list, the objects Nos. 5058, 5060, 5061, 5064, 5065, 5066, 5067, 5068, 5071, 5072, 5073, 5074, 5075, 5077, and 5079 were communicated to me by Professor BOND, Director of the Observatory of Harvard College, U.S., too late for insertion in the body of the Catalogue.

ERRATA.

In page 7, lines 13, 14, for 5063 read 5079, and for six read 22.