

# Quasiparticles in the Kondo lattice model at partial fillings of the conduction band using the density matrix renormalization group

Sebastian Smerat\* and Ulrich Schollwöck

Institut für theoretische Physik C and JARA-Fundamentals of Future Information Technology, RWTH Aachen University, D-52056 Aachen, Germany

Ian P. McCulloch

School of Physical Sciences, University of Queensland, Brisbane, Queensland 4072, Australia

Herbert Schoeller

Institut für theoretische Physik A and JARA-Fundamentals of Future Information Technology, RWTH Aachen University, D-52056 Aachen, Germany

(Received 18 November 2008; revised manuscript received 10 February 2009; published 4 June 2009; publisher error corrected 5 June 2009)

We study the spectral properties of the one-dimensional Kondo lattice model as a function of the exchange coupling, the band filling, and the quasimomentum in the ferromagnetic and paramagnetic phases. Using the density-matrix renormalization group method, we compute the dispersion relation of the quasiparticles, their lifetimes, and the  $Z$  factor. Sigrist *et al.* [Phys. Rev. Lett. **67**, 2211 (1991)] provided the exact ground state and the quasiparticle-dispersion relation of the Kondo lattice model with one conduction electron. The quasiparticle could be identified as the spin polaron. Our calculations of the dispersion relation for partial band fillings give a result similar to the one-electron case, which suggests that the quasiparticle in both cases is the spin polaron. We find that the quasiparticle lifetime differs by orders of magnitude between the ferromagnetic and paramagnetic phases and depends strongly on the quasimomentum.

DOI: [10.1103/PhysRevB.79.235107](https://doi.org/10.1103/PhysRevB.79.235107)

PACS number(s): 71.10.Li, 71.27.+a, 73.21.Hb

## I. INTRODUCTION

The Kondo lattice model (KLM) has been a matter of constant interest for more than last three decades. In two and three dimensions it is one of the common models to describe heavy-fermion<sup>1</sup> physics and is also a possible candidate for high- $T_c$  superconductivity.<sup>2</sup> Our motivation to study the one-dimensional (1D) KLM (Ref. 2) is threefold. First it has been shown<sup>3</sup> that the spin polaron, which is a quasiparticle of the KLM, plays an important role in nonequilibrium transport in a quantum wire coupled to a ferromagnetic spin chain; our method provides the possibility to investigate the quasiparticles of the model. The spin polaron might also play an important role in the electron-spin decay process<sup>4</sup> in quantum dots induced by the hyperfine interaction due to nuclear spins. Second it might be helpful to understand the one-dimensional model in greater detail to assist investigations in higher dimensions. And lastly the model has become interesting for the description of mesoscopic systems, such as carbon nanotubes filled with fullerenes or endohedral fullerenes, the so-called peapods.<sup>5</sup> The aim of this work is to expand on the understanding of the spectral properties of the 1D KLM. We show, by means of the density-matrix renormalization group (DMRG),<sup>6,7</sup> that persistent quasiparticle states exist, which are likely to be the spin-polaron states, and extrapolate their lifetimes and their spectral weights. Furthermore we calculate the quasiparticle-dispersion relation. For the case of half-filling we show that our results qualitatively agree with the results of a strong-coupling expansion in Ref. 8. We compare dispersion relations and confirm the existence of a critical coupling constant at which the

effective quasiparticle mass diverges for large momenta.

The KLM (Fig. 1) consists of a lattice with one localized  $f$  electron on each of the  $L$  lattice sites, which do not interact with each other and a band of itinerant conduction electrons of finite filling  $n$  coupled to the localized electrons by an on-site Heisenberg exchange interaction. The Hamilton operator of the 1D KLM is given by

$$H = -t \sum_{i=1}^L \sum_{\sigma=\uparrow,\downarrow} (c_{i\sigma}^\dagger c_{i+1\sigma} + \text{H.c.}) + J \sum_{i=1}^L \mathbf{S}_i \cdot \mathbf{s}_i, \quad (1)$$

where  $t$  is the hopping parameter,  $c_{i\sigma}^\dagger$  generates an electron at site  $i$  with spin  $\sigma$  and  $\mathbf{S}_i(\mathbf{s}_i)$  are the spin operators of the localized (conduction) electrons at site  $i$ , respectively.  $J$  is the Kondo coupling constant; we will consider only  $J > 0$  here, i.e., the antiferromagnetic coupling case. With  $k$  we denote the quasimomentum in the following.




FIG. 1. (Color online) The Kondo lattice model. The conduction electrons are depicted in the upper row (green) and the localized electrons are depicted as bolt arrows in the lower row (red).

In principle, the 1D KLM supports three phases, depending on the filling  $n$  and on the coupling  $J$ : a ferromagnetic, a paramagnetic, and [at half-filling ( $n=1$ ) only] a spin liquid phase (see Fig. 9). At half-filling of the conduction-electron band the model is best understood and early works using large- $N$  methods<sup>9,10</sup> and the Gutzwiller approximation<sup>11,12</sup> revealed that the magnetic properties are due to the competition of the Ruderman-Kittel-Kasuya-Yosida (RKKY) interaction and the formation of Kondo singlets, where such a singlet is a conduction electron forming a spin singlet with a localized electron. Due to half-filling, the electrons induce an effective RKKY interaction between the localized spins, which forces antiferromagnetic power-law correlations in the ground state. The occurrence of RKKY oscillations or  $2k_F$  oscillations could be confirmed in Ref. 13 using DMRG. By means of exact diagonalization<sup>14</sup> and quantum Monte Carlo,<sup>15</sup> it was shown that the ground state is spin and charge gapped and that it is a singlet of total spin. Therefore the ground state can be associated for all  $J$  with the universality class of spin liquids. There has been a controversial discussion about the size of the Fermi volume (which is a single line in one dimension), whether it is small, and therefore the Fermi wave vector is  $k_F = \frac{\pi}{2}n$ , or whether it is large, and therefore  $k_F = \frac{\pi}{2}(n+1)$ . While a small Fermi volume would correspond to only conduction electrons contributing to the Fermi volume, a large Fermi volume would mean that the localized electrons also contribute to the Fermi volume. The idea of a large Fermi volume is borrowed from the periodic Anderson model.<sup>2</sup> There the  $f$  electrons can move back to the conduction band and therefore contribute to the Fermi volume. The KLM can be derived from the periodic Anderson model<sup>16</sup> in the case of large Coulomb interaction, where only one localized electron per site is allowed and other occupations are fully suppressed. This gives rise to the question whether the Fermi volume is also large in the KLM. Lately the authors of Ref. 8 could argue within a strong-coupling expansion and from the evaluation of the conduction-electron density that the Fermi volume in the case of half-filling is small. In the same work, Ref. 8, the quasiparticle dynamics of the half-filled KLM has been examined as well. It has been possible to calculate the quasiparticle-dispersion relation to good accuracy, where the quasiparticle mass has been found to diverge around  $k \approx \pi$  for  $t/J > t/J_c \approx 0.50 \pm 0.02$ . Therefore the quasiparticles behave like nearly localized  $f$  electrons due to the strong correlation of the conduction and localized electrons. This is consistent with an early large- $N$  approach,<sup>2,10</sup> where it could be shown that the effective electron mass is by magnitudes larger than the bare electron mass. Although the large- $N$  approach is valid for arbitrary filling, its application is best at half-filling due to an intrinsic small energy scale, which can be brought into relation with a spin gap.<sup>2</sup>

In the limiting case of vanishing conduction-electron density it could be rigorously shown<sup>17</sup> by both applying the Perron-Frobenius theorem and later exact diagonalization<sup>18</sup> that the KLM is ferromagnetic for all  $J$ . Importantly, Sigrist *et al.*<sup>17</sup> could show that the quasiparticle of the Kondo lattice model is the spin polaron, which corresponds to an excited state separated from a continuum of scattering states. Representatives of the constituent elements of the spin polaron are




FIG. 2. (Color online) Representatives of the constituent elements of the spin polaron: (a) electron spin down; (b) electron spin up and one of the lattice spins down.

shown in Fig. 2. In Fig. 2(a) the localized spin lattice is completely ferromagnetic and the electron spin is oriented in the opposite direction. Due to the antiferromagnetic exchange interaction the electron energy for quasimomentum  $k$  is reduced. Via spin-flip processes, this state is coupled to the states shown in Fig. 2(b), where the electron spin and one of the localized spins are flipped and the momentum  $q$  has been transferred to the spin lattice. These are states of higher energy since the antiferromagnetic interaction can only reduce the energy if the two flipped spins are at the same site. The coupling leads to a level repulsion between the states of Figs. 2(a) and 2(b), with the energetically lower one corresponding to the spin-polaron state and the higher ones forming the scattering states band. As proposed in Ref. 3, the spin-polaron state is expected to have a very long lifetime if its energy lies outside the band of scattering states, so that it is protected against magnon absorption and emission processes.

At partial band fillings  $n$ , ferromagnetism also survives in the strong-coupling limit,<sup>19</sup> where the KLM can be mapped to an effective Heisenberg model with a ferromagnetic exchange coupling. In this limit the formation of Kondo singlets, which move through the lattice, is sufficient to explain the occurrence of ferromagnetism, but this does not exclude RKKY interaction, which might still play an important role. From exact diagonalization studies,<sup>20</sup> it follows that the KLM is ferromagnetic for  $J \geq 0.5$  at  $n \approx 0.25$ . The critical  $J$  increases roughly linear to  $J \approx 3$  at  $n \approx 0.75$ . This raises the question of which mechanism drives ferromagnetism at couplings  $J$  of  $\mathcal{O}(1)$  and one proposal<sup>21</sup> is that double exchange might be the crucial mechanism, where one conduction electron is responsible for screening several localized electrons. The screening lowers the total energy in the antiferromagnetic KLM as long as  $J$  surmounts a critical value and forces the localized spins to align in the same direction.

At a certain  $J$  the KLM approaches a second-order transition<sup>22</sup> by lowering  $J$  to a paramagnetic phase, where the spin polaron<sup>23</sup> might play an important role. The transition line has been calculated using exact diagonalization<sup>20</sup> and has been refined later by means of bosonization.<sup>21</sup> The destruction of the ferromagnetic phase is described by a quantum random transverse-field Ising Hamiltonian.<sup>21</sup> Approaching the transition line from high  $J$  it has been proposed<sup>21,24</sup> that the large ferromagnetic cluster splits up in several small clusters each corresponding to one spin polaron. Just below the transition line the small clusters' direction of magnetization is not the same anymore for all clusters and leads to zero

net magnetization. By means of DMRG the spin-structure factor of the localized electrons could be calculated<sup>24</sup> and it has been found that the size of the Fermi volume is small for very low  $J$  and becomes large approaching the transition line from lower  $J$ . From this one can conclude that near the transition line, the localized electrons are incorporated in the Fermi volume and therefore spin polarons are formed. Lowering  $J$  the spin polarons are destroyed. In another proposal<sup>25</sup> using DMRG the size of the Fermi surface for small  $J$  has been calculated from the spin-correlation function and found to be small. The corresponding authors find for  $J \gtrsim 1$  that strong boundary charge perturbations mash the true bulk behavior and therefore a small Fermi surface is not distinguishable from a large Fermi surface. The Fermi surface size is left as an open question. The paramagnetic KLM has also been argued<sup>26</sup> to belong to the class of Tomonaga-Luttinger liquids.<sup>27</sup> This is motivated by the gapless spin and charge excitation,<sup>2</sup> which also makes the model difficult to handle with numerical methods using finite system sizes in this regime. From an analysis of Friedel oscillations, which are  $2k_F$  or  $4k_F$  oscillations, the Luttinger parameters could be determined for  $J > 1.8$  and the Fermi volume has been found to be large. For very low  $J$ , RKKY or  $2k_F$  oscillations dominate the correlation functions of the KLM. This could be attributed<sup>21</sup> to the backscattering of the conduction electrons at the localized electrons. In a recent work<sup>28</sup> the Luttinger parameter has been calculated for many values of  $J$  and  $n$  in the paramagnetic phase. Using a logarithmic correction the spin-correlation function can be fitted perfectly to DMRG data.

In this paper we consider the spectral properties of the Kondo lattice model at partial band fillings. We will calculate the dispersion relation in the ferromagnetic phase for different Kondo couplings  $J$  and various fillings  $n$  and find a well-defined quasiparticle band. Comparing the one conduction-electron case of Ref. 17 with the partial band filling case here, the latter seems to be a direct continuation of the former, meaning that the quasiparticle-dispersion relation is found to be similar in both cases. Therefore it is likely that the spin-polaron picture used in Ref. 17 suits here as well. We are able to confirm the results of Ref. 8 at half-filling. In a second step we will show from the width of the spectral densities that the lifetime of the spin polaron is very long and therefore the quasiparticle is persistent. We also examine the spectral densities in the paramagnetic phase and find unexpectedly that a quasiparticle excitation visible in the spectral density exists and can be fitted reasonably good by a Lorentzian function. It has been argued<sup>21</sup> that this quasiparticle also might be of the spin-polaron type. Its lifetime is smaller by several orders of magnitude than in the ferromagnetic phase but the ratio depends very sensitively on the values of  $J$ ,  $n$ , and the quasimomentum  $k$ . An interesting effect is found that the lifetime is maximal in the ferromagnetic phase if the quasimomentum is close to the Fermi points.

The paper is outlined as follows. In Sec. II we will discuss the method, particularly how we calculate spectral densities, how we extract the lifetimes, and how we extrapolate them. In Sec. III we will present our results. We will end up in a brief summary in Sec. IV.

## II. METHODS

In this section we describe the methods used in our calculations. First of all we briefly discuss our DMRG algorithm. Second we describe the correction-vector method, which we use to calculate the Green's functions. At last we show how to calculate the lifetime of quasiparticles using the spectral functions we obtained from the Green's functions.

### A. DMRG

For the calculation of ground states, we use a *DMRG* algorithm with Abelian and non-Abelian symmetries, whose implementation is based on a matrix-product formulation. We use open-boundary conditions for all calculations. We kept up to 1800 DMRG ansatz states in our calculations setting the discarded weight typically smaller than  $10^{-5}$ .

### B. Correction-vector method

Applying the *correction-vector*<sup>29-32</sup> method we obtain the spectral functions  $A(\omega)$ , where  $\omega$  is the energy. To calculate  $A(\omega)$ , we need the retarded Green's function  $G_A(\omega+i\eta) = G_A^+(\omega+i\eta) + G_A^-(\omega+i\eta)$ , where

$$G_A^+(\omega+i\eta) = \langle 0 | A^\dagger \frac{1}{\omega + E_0 + i\eta - H} A | 0 \rangle, \quad (2)$$

$$G_A^-(\omega+i\eta) = \langle 0 | A \frac{1}{\omega - E_0 + i\eta + H} A^\dagger | 0 \rangle \quad (3)$$

are the two branches of the retarded Green's function and  $A$  is an arbitrary operator,  $|0\rangle$  is the ground state with energy  $E_0$ , and  $\eta > 0$  is an artificial broadening factor, which is needed to lower the lifetime of the excitation to avoid boundary effects due to the finite system size. The basic rule is to choose  $\eta > \frac{c}{L}$ , where  $c$  is the velocity of the excitation, but the minimal  $\eta$  is strongly depending on the model.

In principle one would need to compute both branches of the Green's function to obtain the complete spectral properties. For the determination of lifetimes the spectral weight of the quasiparticle is nearly completely concentrated in one of the branches. Therefore we can neglect the other branch in this case. From now on, we will base all our arguments concerning the Green's function on the + branch. Concerning the spectral density the calculations for the - branch can be done similarly except for a minus sign.

The correction vector is defined as

$$|c(\omega+i\eta)\rangle = \frac{1}{\omega + E_0 + i\eta - H} A | 0 \rangle \quad (4)$$

and hence

$$(\omega + E_0 + i\eta - H) |c(\omega+i\eta)\rangle = A | 0 \rangle, \quad (5)$$

where the ground state  $|0\rangle$  is obtained from the preceding DMRG calculation. This leads to a non-Hermitian system of linear equations for real and imaginary parts, which can be solved using the generalized minimal residual method (GMRES) method.<sup>33</sup> The outcome is  $|c(\omega+i\eta)\rangle$ , which allows us to calculate the Green's function as

$$G_A(\omega + i\eta) = \langle 0|A|c(\omega + i\eta)\rangle. \quad (6)$$

The spectral density can then be obtained by applying the standard formula

$$A(\omega + i\eta) = -\frac{1}{\pi} \text{Im} G_A(\omega + i\eta), \quad (7)$$

where  $\omega$  is assumed to be real.

### C. Quasiparticle lifetime

For the calculation of *quasiparticle lifetimes* we will limit ourselves to electronic systems. It is useful to transform the Hamiltonian into the Fourier space according to

$$c_{i\sigma} = \frac{1}{\sqrt{N}} \sum_k c_{k\sigma} e^{ikr_i}. \quad (8)$$

Hence we obtain

$$H = \sum_k \sum_{\sigma=\uparrow,\downarrow} [\epsilon_0(k) c_{k\sigma}^\dagger c_{k\sigma}] + J \sum_k \mathbf{S}_k \cdot \mathbf{s}_{-k} \quad (9)$$

with  $\epsilon_0(k) = -2t \cos ka$ , where  $a$  is the lattice spacing. The one-electron Green's function is then defined as

$$G_{k\sigma}(\omega + i\eta) = \langle 0|c_{k\sigma} \frac{1}{\omega + E_0 + i\eta - H} c_{k\sigma}^\dagger |0\rangle. \quad (10)$$

The self-energy  $\Sigma_\sigma(k, \omega)$  is implicitly defined for the interacting system  $H$  as

$$G_{k\sigma}(\omega + i\eta) = \frac{1}{\omega + i\eta - [\epsilon_0(k) - \mu + \Sigma_\sigma(k, \omega + i\eta)]},$$

with  $\mu$  as the chemical potential. Note that  $\eta$  appears also in the self-energy. This is necessary, because  $\lim_{\eta \rightarrow 0}$  will not be carried out in the numerical calculations. In general, the self-energy is a complex function  $\Sigma_\sigma(k, \omega) = R_\sigma(k, \omega) + iI_\sigma(k, \omega)$ . The separation of real and imaginary parts leads to

$$G_{k\sigma}(\omega + i\eta) = \frac{1}{\omega - [\epsilon_0(k) - \mu + R_\sigma(k, \omega + i\eta)] + i[\eta - I_\sigma(k, \omega + i\eta)]}. \quad (11)$$

We now assume that the self-energy is continuous and only weakly depends on  $\omega$  in the vicinity of a resonance  $\omega_{i\sigma} = \epsilon_0(k) - \mu + R_\sigma(k, \omega)|_{\omega=\omega_{i\sigma}}$ , where  $\omega_{i\sigma}$  is one out of several resonances, which are well separated to provide the correct determination of the lifetime of the quasiparticles (see the end of this section for the explicit extrapolation scheme). In addition we assume  $|I_\sigma(k, \omega)| \ll |\epsilon_0(k) - \mu + R_\sigma(k, \omega)|$  near the resonance we are interested in, i.e., we assume long lifetimes, because we are interested in these. This leads to

$$I_\sigma(k, \omega + i\eta) \approx I_\sigma^{(i)}(k) \quad (12)$$

in the vicinity of the  $i$ th resonance. For the real part of the self-energy we apply a Taylor expansion at the resonance  $\omega_{i\sigma}$ . We find

$$\begin{aligned} & \omega - [\epsilon_0(k) - \mu + R_\sigma(k, \omega + i\eta)] \\ & \approx (\omega - \omega_{i\sigma}) \left[ 1 - \frac{dR_\sigma(k, \omega + i\eta)}{d\omega} \Big|_{\omega+i\eta=\omega_{i\sigma}} \right] \\ & - i\eta \frac{dR_\sigma(k, \omega + i\eta)}{d\omega} \Big|_{\omega+i\eta=\omega_{i\sigma}} \end{aligned}$$

and define

$$\alpha_{i\sigma} = \left( 1 - \frac{dR_\sigma(k, \omega)}{d\omega} \Big|_{\omega=\omega_{i\sigma}} \right)^{-1}. \quad (13)$$

Substituting this to Eq. (11) the Green's function in the vicinity of resonance  $\omega_{i\sigma}$  is given by

$$G_{k\sigma}(\omega + i\eta) = \alpha_{i\sigma} \frac{1}{\omega - \omega_{i\sigma} + i[\eta + \alpha_{i\sigma}|I_\sigma^{(i)}(k)]} \quad (14)$$

and the spectral function obtains the form

$$A_{k\sigma}(\omega + i\eta) = \sum_i \frac{\alpha_{i\sigma}}{\pi} \frac{\eta + \alpha_{i\sigma}|I_\sigma^{(i)}(k)|}{(\omega - \omega_{i\sigma})^2 + [\eta + \alpha_{i\sigma}|I_\sigma^{(i)}(k)]^2}, \quad (15)$$

which corresponds to a sum of Lorentz distributions at the resonances  $\omega_{i\sigma}$  with a broadening of

$$B^{(i)}(\eta) = \eta + \alpha_{i\sigma}|I_\sigma^{(i)}(k)|. \quad (16)$$

Hence the broadening computed with the correction-vector method is essentially the sum of the natural broadening  $\alpha_{i\sigma}|I_\sigma^{(i)}(k)|$  and the artificially introduced broadening  $\eta$  and therefore  $B(\eta)$  linearly depends on  $\eta$ . Note that from the Lehmann representation of the spectral density one can find that  $I_\sigma^{(i)}(k) \leq 0$ .

The broadened spectral density  $A_{k\sigma}(\omega + i\eta)$  is a convolution of the nonbroadened spectral density  $A_{k\sigma}(\omega)$  with a Lorentzian of width  $\eta$ . We now assume that the spectral density consists of a sum of Lorentz distributions, which are separated by non-Lorentzian regions. The outcome of the convolution of two Lorentzians again is a Lorentzian, where the broadenings behave additively. As the broadening corresponds to an inverse lifetime, we can define the lifetime of the quasiparticle corresponding to the  $i$ th resonance as

$$\tau = \lim_{\eta \rightarrow 0} \frac{1}{\eta + \alpha_{i\sigma}|I_\sigma^{(i)}(k)|} = \frac{1}{\alpha_{i\sigma}|I_\sigma^{(i)}(k)|}. \quad (17)$$

The limitations of this method are obvious. First of all the excitation must cause a Lorentzian-shaped peak in the spectral density. To be able to extract the broadening of such a peak all other peaks must be energetically separated from this one. Thus we have to check whether the conditions of our theory are fulfilled or not. We can check whether the spectral density has a Lorentzian shape (see Fig. 8) and  $B(\eta)$  has to depend linearly on  $\eta$  (see Fig. 3). In Eq. (12) we only take zeroth order in  $\omega + i\eta$  of the imaginary part of the self-energy into account. The first order leads to a small additional linear  $\eta$ -dependent contribution in  $B(\eta)$  in Eq. (16),




FIG. 3. (Color online) The linear fit of  $B(\eta)$  vs  $\eta$  is shown for three different quasimomenta, for a system with 48 sites,  $n=0.125$ ,  $t=1$ , and Kondo coupling  $J=1$ . The data have been offset by  $\eta$ .

which is the reason for the small finite slope of  $B(\eta) - \eta$  in Fig. 3.

We have now two possible estimates for the lifetime of a quasiparticle:

(1) Lower estimate: use the inverse broadening  $B(\eta)^{-1}$  directly (without  $\eta \rightarrow 0$ ). With Eq. (17)  $B(\eta)^{-1}$  is smaller than  $B(0)^{-1}$ ; this is therefore a reliable lower estimate.

(2) Extrapolation: calculate the broadening for several different  $\eta$ . From this one can extract the linear dependence of  $B(\eta)$  on  $\eta$  and  $B(0)^{-1}$  gives the extrapolated lifetime. See also Fig. 3.

It turns out that due to long lifetimes only extrapolated lifetimes are meaningful. Therefore in the next section we will discuss the results obtained by the second method only. From the Lorentzian fit of a single resonance peak of the spectral function in Eq. (15) one can also estimate the spectral weight  $\alpha_{i\sigma}$  of the corresponding excitation.

### III. RESULTS

In this section we will present the results obtained using the methods we discussed in the last section. First we will show the calculated dispersion relations considering a Kondo lattice model at half-filling and at partial filling. Our half-filling results show a qualitative agreement with the results in Ref. 8. In the second part we show several spectral functions and the calculated lifetimes, which leads to the conclusion that we find a bound-polaron state. The hopping parameter  $t$  is set to  $t=1$  in all calculations.

#### A. Dispersion

The half-filled KLM serves as the touchstone of our method, where we can compare our results to those of Trebst *et al.*,<sup>8</sup> who did a strong-coupling expansion up to 11th order in  $t/J$ . We calculated the dispersion relation for different values of  $t/J$  (see Fig. 4) and used lattice sizes of 32 and 48 sites. The calculations have converged in the sense that we could not find any deviations between calculations of differ-




FIG. 4. (Color online) Dispersion relations of the half-filled KLM. The comparison to 11th-order perturbation theory is taken from Ref. 8. The line for  $t/J=0.8$  is meant only as a guide to the eyes.

ent system sizes. The dispersion relations are a compound of the single spectral densities or, in other words, the dispersion relation can be obtained by gluing all the spectral densities for all values of  $k$  together. Our results show very well agreement to the results in Ref. 8 for  $t/J=0.4$  with small deviations for small  $k$ . By strong-coupling expansion it is found that the band flattens out for  $k \rightarrow \pi$  around  $t/J_c \approx 0.50 \pm 0.02$  and therefore the effective quasiparticle mass diverges. This is also found by DMRG for a higher value of  $t/J_c \approx 0.576 \pm 0.002$ . As one can see, in Fig. 4, the strong-coupling expansion-dispersion relation has a pronounced minimum at  $k \approx 0.7\pi$  for  $t/J=0.6$ . This minimum is not visible by eyes only in the DMRG data; still it is there at  $k \approx 0.9\pi$ . The minimum becomes easily visible also in the DMRG data for  $t/J=0.8$  as shown in Fig. 4. Summarizing, the DMRG results match very well to the strong-coupling expansion for low  $t/J$  but the agreement becomes worse for  $t/J \geq 0.6$ . Qualitatively, the same things happen, but for a larger value of  $t/J$  in the DMRG calculations. DMRG is the more reliable method in that regime because it is nonperturbative and the error can be easily controlled by very small DMRG truncation errors. In this case it is easy to keep the truncation error reasonably low. We can confirm the physical picture established by Trebst *et al.*, namely, that the quasiparticles gain an enormously high mass, which is due to a growing correlation between the conduction and the  $f$  electrons. The quasiparticles with high momenta therefore behave like almost localized  $f$  electrons.

Now we consider the dispersion relation of the KLM for partial band filling (see Fig. 5). For the ground-state calculation of the KLM with 48 sites, a filling of  $n=0.125$  and  $J=1$  we used about 100 DMRG ansatz states. The calculation of the correction vectors needed 800 DMRG states to reach good convergence. In Fig. 5 and all other figures of dispersion relations, we neglected the chemical potential, which would shift the lower band edge to nearly zero. One can distinguish two different bands. The higher one behaves as  $-2t \cos k$  (up to a constant offset) and can therefore be attributed to free electrons, which do not form bound states




FIG. 5. (Color online) Dispersion of a KLM with 48 sites,  $n=0.125$ ,  $J=1$ , and  $t=1$  compounded of spectral densities for all values of  $k$ . The color scale corresponds to the height of the spectral density. The broadening is  $\eta=0.2$ . The lower band is the bound quasiparticles band, while the upper one is the scattering states band.

with the localized spins. From now on, this band will be referred to as the *scattering state band* in the sense that these excited states rapidly decay. The lower one of the two bands represents the states of the system which are formed by the conduction electrons bound to the localized spins, which from now on referred to as *quasiparticle or spin-polaron band*. Contrary to the scattering state band, this band consists only of one state, which is separated from the continuum (for large  $L$ , where  $L$  is the lattice size) of scattering states and has a Lorentzian shape from which the lifetime can be extracted, which is very long in most of the cases (see Sec. III B). Even on lowering  $\eta$  the excitation peak does not differ from its Lorentzian shape; therefore, we can be sure that only one excitational state can be responsible for this. In contrast to the spin-polaron band the scattering states band does not change its width linearly with  $\eta$ . Keeping the same filling  $n=0.125$ , but raising the Kondo coupling constant  $J$  to 3.5 (see Fig. 6), the quasiparticle band becomes more separated from the scattering states because the quasiparticle state is now energetically lowered. This can be understood by a simple physical picture. For that we rewrite the exchange




FIG. 6. (Color online) Dispersion relation (as in Fig. 5) of a KLM with  $n=0.125$ ,  $J=3.5$ , and  $t=1$ . The lower band is the bound quasiparticles band, while the upper one is the scattering states band.




FIG. 7. (Color online) Dispersion relations of a KLM with  $J=3.5$ ,  $t=1$ , and three different fillings,  $n=0.125$ ,  $n=0.25$ , and  $n=0.75$  extracted of the spectral densities.

coupling of the Hamiltonian as  $\sum_i^L [J^z S_i^z s_i^z + \frac{J_\perp}{2} (S_i^+ s_i^- + S_i^- s_i^+)]$  and we now set  $J_\perp=0$ . The quasiparticle state of the KLM almost only consists of an electron with spin antiparallel to the localized spins. With respect to our modified exchange coupling, this results in a lowered energy of  $J^z/4$  per electron. The scattering states also contain important contributions with an electron spin oriented parallel to the localized spins. This leads to a higher energy of  $J^z/4$ . Therefore the energy difference between quasiparticle and scattering states is  $J^z/2$  and scales with  $J^z$ . Taking also a finite  $J_\perp$  into account the quasiparticle energy is even lowered more due to spin-flip processes. The scattering states band has a similar shape as before, as expected. The weight of the quasiparticle states band is also increasing with  $J$ . This is also expected because the state becomes energetically more favorable with increasing  $J$ .

In Fig. 7 we show the dispersion relation of the quasiparticle of a system with  $J=3.5$  for three different fillings,  $n=0.125$ ,  $n=0.25$ , and  $n=0.75$ . The ground state is ferromagnetic in all cases. We conclude that even in the presence of many electrons the spin-polaron state can be clearly identified.

### B. Lifetime estimations from spectral functions

In a further step we take a look at single spectral densities for fixed quasimomentum  $k$ , which provides the possibility to calculate quasiparticle lifetimes of the bound quasiparticles and proves the existence of bound-polaron states. We consider only the calculation of the extrapolated lifetimes, as described in Sec. II C and whose extrapolation scheme is shown in Fig. 3. Calculating extrapolated lifetimes this way, we have to be very careful due to the assumption that the spectral density complies with a Lorentz distribution. For the spectral density being a Lorentzian the imaginary part of the self-energy has to be very small compared to the energy of the resonance and it should not vary too much in the vicinity of the resonance. The expansion of the self-energy leads then to a Lorentzian function. Thus the spectral density is not




FIG. 8. (Color online) Spectral function of a KLM with 48 Sites,  $n=0.125$ ,  $J=1$ ,  $t=1$ ,  $k=\frac{\pi}{49}$ , and  $\eta=0.02$ . The green line is obtained by fitting a Lorentzian to the DMRG data.

Lorentzian shaped in higher orders of the expansion and it has to be checked (see, for example, Fig. 8) whether it is good enough. Figure 8 shows a spectral density for a KLM with 48 sites,  $n=0.125$ ,  $J=1$ , and quasimomentum  $k=\frac{\pi}{49}$ . The artificial broadening is set to  $\eta=0.02$ . The number of data points obtained provides the possibility of a very precise fit of the Lorentz distribution. Figure 8 sharply supports the assumption made in Sec. II C that the spectral density has a Lorentzian shape, which is necessary to calculate quasiparticle lifetimes.

We would like to make a comment concerning the lifetimes in the paramagnetic phase, which is supposed to be of Luttinger liquid type. In a Luttinger liquid we would not expect to have well-defined quasiparticles. Therefore the approximation of Lorentzian-shaped excitations in the spectral density is crude in the paramagnetic regime of the KLM. Then it is even more surprising that this approximation fits the DMRG data relatively well. But we also find that those excitations in the paramagnetic phase decay fast compared to the ferromagnetic phase (where we do not expect a Luttinger liquid because of a finite spin gap) and this would be expected.

The extrapolated lifetimes are summarized in Fig. 9 and Table I. There we can see that the lifetime strongly depends on the parameters filling  $n$  and Kondo coupling constant  $J$  as well as on the quasimomentum  $k$ . The lifetimes in the ferromagnetic phase (this concerns the  $[n, J]$  pairs  $\{[0.125, 0.5], [0.125, 1], [0.25, 1], [0.29, 1], [0.75, 3.5]\}$ ) decrease by approaching the paramagnetic phase by either lowering  $J$  or increasing  $n$ . For fixed and low quasimomentum  $k$  it seems that the lifetime decreases by increasing  $n$  (even if  $J$  is increased at the same time, so that the distance to the paramagnetic phase is still large (compare, e.g., the pairs  $[0.125, 1]$  and  $[0.75, 3.5]$ ). This indicates that the lifetime is influenced by the presence of other quasiparticles, probably by an effective interaction between the quasiparticles mediated via the coupling to the localized spins. This is further substantiated by the dependence of the quasiparticle lifetime on the quasimomentum  $k$  in the ferromagnetic phase. For  $k$  approaching the Fermi level, the lifetime increases, which is consistent




FIG. 9. (Color online) Simplified phase diagram of the 1D Kondo lattice model taken from Ref. 24. The points mark the parameters at which extrapolated lifetimes have been calculated. The lifetimes for points 1 ( $J=1$ ), 2 ( $J=0.5$ ), and 3 ( $J=0.25$ ) and quasimomentum  $k=\frac{\pi}{49}$  are given directly in the picture by  $\tau_1$ ,  $\tau_2$ , and  $\tau_3$ ; the lifetimes for the other points are listed in Table I.

with the fact that the phase space for quasiparticle interaction becomes smaller close to the Fermi level. In contrast, electron-magnon interaction is expected to be more important for quasiparticles close to the Fermi level because the energy of the spin polaron comes closer to the scattering band. This effect can be seen in the paramagnetic phase for the pairs  $[0.375, 0.5]$  and  $[0.75, 2.5]$ , where the lifetime decreases with increasing quasimomentum. Thus, in the paramagnetic phase, we conclude that electron-magnon interaction limits the lifetime of the spin polaron. Deep inside the paramagnetic phase at  $[n=0.375, J=0.5]$  the lifetime is short for all determined values of  $k$ . Therefore, as predicted earlier in Refs. 21 and 22 there exist no persistent quasiparticles in this regime.

We also extracted the spectral weight of the spin-polaron excitation from the Lorentzian fit and summarized them in Table I in the second row of the respective  $k$  value. Considering the three numbered points of Fig. 9 we calculated the spectral weights: (1)  $0.818\,004 \pm 0.000\,001$  (2)  $0.871\,19 \pm 0.000\,03$ , and (3)  $0.588 \pm 0.001$ , which do fulfill the expectation that the spectral weight should be significantly lower in the paramagnetic phase. The calculated weights are independent of  $\eta$  within the error bounds. They show a strong dependence on the quasimomentum (decreasing for growing  $k$ ) in the ferromagnetic as well as in the paramagnetic phase. This is expected because the spin-polaron states with higher value of  $k$  have higher energy and come closer to the scattering states. However, it is unexpected that the spectral weight is large for  $[0.375, 0.5]$  and this still has to be explained.

#### IV. SUMMARY

We have studied the one-dimensional Kondo lattice model at half-filling and at partial band fillings for various Kondo

TABLE I. Summarization of extrapolated lifetimes (first row of the respective  $k$  value) and spectral weight of the spin-polaron excitation (second row of the respective  $k$  value) for different fillings of the system, different coupling constants and different quasimomenta. The  $k$  values in parentheses correspond to  $[0.75, 2.5]$ , only.

$k$	$[n, J]$				
	[0.25, 1] (fm)	[0.29, 1] (fm)	[0.375, 0.5] (pm)	[0.75, 3.5] (fm)	[0.75, 2.5] (pm)
$\frac{1\pi}{49} (\frac{1\pi}{33})$	$831 \pm 16$	$183 \pm 3$	$67 \pm 9$	$38.6 \pm 0.8$	$25.7 \pm 1.6$
	$0.8211 \pm 0.0004$	$0.8292 \pm 0.0008$	$0.951 \pm 0.006$	$0.714 \pm 0.008$	$0.674 \pm 0.005$
$\frac{3\pi}{49}$	$2540 \pm 84$				
$\frac{8\pi}{49}$			$14.7 \pm 1.5$		
			$0.82 \pm 0.01$		
$\frac{11\pi}{49}$	$\mathcal{O}(10^5)$	$2850 \pm 600$			
	$0.6244 \pm 0.0001$	$0.6375 \pm 0.0005$			
$\frac{18\pi}{49} (\frac{12\pi}{33})$				$\mathcal{O}(10^3)$	$5.29 \pm 0.14$
				$0.5580 \pm 0.0008$	$0.3391 \pm 0.0005$
$\frac{36\pi}{49} (\frac{24\pi}{33})$				$\mathcal{O}(10^3)$	$12.76 \pm 0.62$
				$0.3391 \pm 0.0005$	$0.190 \pm 0.006$

couplings  $J$ . At half-filling we could verify the results of Ref. 8. This includes the dispersion relation and the divergence in the effective quasiparticle mass. At partial band fillings we were able to show that in the case of ferromagnetism long living quasiparticle states exist and we have suggested that these are spin-polaron quasiparticles as used in Ref. 17. The lifetime exceeds the lifetime of quasiparticle excitations deep inside the paramagnetic phase by several orders of magnitude. From the dependence on the quasimomentum we conclude that the dominant decay processes are the spin-polaron–spin-polaron interaction in the ferromagnetic phase and the interaction between electrons and spin waves in the paramagnetic phase. The weight of the spin-polaron state is very close to one even for special points in the paramagnetic phase. The results motivate the speculation that spin coherence can be significantly enhanced by coupling of electrons to magnons in the ferromagnetic phase of the localized spins.

As we have seen in this paper, the spin excitations in the ferromagnetic phase can in turn form spin-polaron bound states with the itinerant electrons, increasing their lifetime considerably. This effect persists in the presence of many electrons and becomes more efficient for quasimomenta close to the Fermi level. It is an interesting question for future research to investigate the consequences for the spin relaxation and dephasing rates in the Kondo lattice model by studying the spin-spin correlation functions.

#### ACKNOWLEDGMENTS

S.S. is grateful for a discussion with Simon Trebst. S.S., U.S., and H.S. acknowledge the support from the DFG-Forschergruppe 912 on “coherence and relaxation properties of electron spins.”

\*smerat@physik.rwth-aachen.de

<sup>1</sup>P. Coleman, *Handbook of Magnetism and Advanced Magnetic Materials* (John Wiley & Sons, New York, 2007), Vol. 1.

<sup>2</sup>H. Tsunetsugu, M. Sigrist, and K. Ueda, *Rev. Mod. Phys.* **69**, 809 (1997).

<sup>3</sup>F. Reininghaus, T. Korb, and H. Schoeller, *Phys. Rev. Lett.* **97**, 026803 (2006).

<sup>4</sup>J. Schliemann, A. Khaetskii, and D. Loss, *J. Phys.: Condens. Matter* **15**, R1809 (2003).

<sup>5</sup>I. V. Krive, R. I. Shekhter, and M. Jonson, *Low Temp. Phys.* **32**, 887 (2006).

<sup>6</sup>S. R. White, *Phys. Rev. Lett.* **69**, 2863 (1992).

<sup>7</sup>U. Schollwöck, *Rev. Mod. Phys.* **77**, 259 (2005).

<sup>8</sup>S. Trebst, H. Monien, A. Grzesik, and M. Sigrist, *Phys. Rev. B*

**73**, 165101 (2006).

<sup>9</sup>N. Read and D. M. Newns, *J. Phys. C* **16**, 3273 (1983).

<sup>10</sup>N. Read, D. M. Newns, and S. Doniach, *Phys. Rev. B* **30**, 3841 (1984).

<sup>11</sup>T. M. Rice and K. Ueda, *Phys. Rev. B* **34**, 6420 (1986).

<sup>12</sup>T. M. Rice and K. Ueda, *Phys. Rev. Lett.* **55**, 995 (1985).

<sup>13</sup>C. C. Yu and S. R. White, *Phys. Rev. Lett.* **71**, 3866 (1993).

<sup>14</sup>H. Tsunetsugu, Y. Hatsugai, K. Ueda, and M. Sigrist, *Phys. Rev. B* **46**, 3175 (1992).

<sup>15</sup>R. M. Fye and D. J. Scalapino, *Phys. Rev. Lett.* **65**, 3177 (1990).

<sup>16</sup>J. R. Schrieffer and P. Wolff, *Phys. Rev.* **149**, 491 (1966).

<sup>17</sup>M. Sigrist, H. Tsunetsugu, and K. Ueda, *Phys. Rev. Lett.* **67**, 2211 (1991).

<sup>18</sup>M. Sigrist, K. Ueda, and H. Tsunetsugu, *Phys. Rev. B* **46**, 175



- (1992).
- <sup>19</sup>M. Sigrist, H. Tsunetsugu, K. Ueda, and T. M. Rice, *Phys. Rev. B* **46**, 13838 (1992).
- <sup>20</sup>H. Tsunetsugu, M. Sigrist, and K. Ueda, *Phys. Rev. B* **47**, 8345 (1993).
- <sup>21</sup>G. Honner and M. Gulácsi, *Phys. Rev. Lett.* **78**, 2180 (1997).
- <sup>22</sup>I. P. McCulloch, A. Juozapavicius, A. Rosengren, and M. Gulácsi, *Phys. Rev. B* **65**, 052410 (2002).
- <sup>23</sup>S. Methfessel and D. C. Mattis, *Handbuch der Physik*, edited by H. P. J. Wijn (Springer-Verlag, Berlin, 1968), Vol. 18.
- <sup>24</sup>I. P. McCulloch, A. Juozapavicius, A. Rosengren, and M. Gulácsi, *Philos. Mag. Lett.* **81**, 869 (2001).
- <sup>25</sup>J. C. Xavier, E. Novais, and E. Miranda, *Phys. Rev. B* **65**, 214406 (2002).
- <sup>26</sup>N. Shibata and K. Ueda, *J. Phys.: Condens. Matter* **11**, R1 (1999).
- <sup>27</sup>F. D. M. Haldane, *J. Phys. C* **14**, 2585 (1981).
- <sup>28</sup>J. C. Xavier and E. Miranda, *Phys. Rev. B* **70**, 075110 (2004).
- <sup>29</sup>Z. G. Soos and S. Ramasesha, *J. Chem. Phys.* **90**, 1067 (1989).
- <sup>30</sup>S. Ramasesha, S. Pati, H. Krishnamurthy, Z. Shuai, and J. L. Bredas, *Synth. Met.* **85**, 1019 (1997).
- <sup>31</sup>T. D. Kühner and S. R. White, *Phys. Rev. B* **60**, 335 (1999).
- <sup>32</sup>E. Jeckelmann, *Phys. Rev. B* **66**, 045114 (2002).
- <sup>33</sup>Y. Saad and M. H. Schultz, *SIAM J. Sci. Comput. (USA)* **7**, 856 (1986).