

Lorentz- and CPT-violating Chern-Simons term in the functional integral formalism

J.-M. Chung*

Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

(Received 21 April 1999; published 29 October 1999)

We show that in the functional integral formalism, with the weaker condition of gauge invariance alone, one cannot determine the (finite) coefficient of the induced, Lorentz- and CPT-violating Chern-Simons term, arising from the Lorentz- and CPT-violating fermion sector. [S0556-2821(99)03220-8]

PACS number(s): 11.30.Cp, 11.30.Er, 12.20.Ds

In recent work [1], the following question has been posed: is a Lorentz- and CPT-violating Chern-Simons term induced by the Lorentz- and CPT-violating term $\bar{\psi} b \gamma_5 \psi$ (b_μ is a constant four-vector) in the conventional Lagrangian of QED. In the paper by Jackiw and Kosteletsky [2], this problem has been discussed. Their results depend on whether one uses a nonperturbative formalism or a perturbative formalism. In a nonperturbative formalism, radiative corrections arising from the axial vector term in the fermion sector induce a definite and nonzero Chern-Simons term, while when a perturbative formalism is used, radiative corrections are finite but undetermined.

The purpose of this paper is to view this indeterminacy¹ of the finite radiative corrections in the functional integral formalism.

Let us consider the following functional integral

$$Z(A) = \int d\bar{\psi} d\psi \exp[iI(A)], \quad (1)$$

where

$$I(A) = \int d^4x [i\bar{\psi} \not{D} \psi - m\bar{\psi} \psi - b_\mu j_5^\mu],$$

with $j_5^\mu(x) = \bar{\psi}(x) \gamma^\mu \gamma_5 \psi(x) - [(\text{possibly finite}) \text{ local counterterms}]$. The gauge field A in the covariant derivative $\not{D}$ is external. By changing the field variables

$$\psi(x) \rightarrow \exp[i\alpha(x) \gamma_5] \psi(x),$$

$$\bar{\psi}(x) \rightarrow \bar{\psi}(x) \exp[i\alpha(x) \gamma_5],$$

we see that the integration measure of Eq. (1) changes by

$$d\bar{\psi} d\psi \rightarrow d\bar{\psi} d\psi \exp\left[-i \int \frac{\alpha(x)}{8\pi^2} {}^*F^{\mu\nu}(x) F_{\mu\nu}(x)\right],$$

where this form for the anomaly is obtained when $\bar{\psi} \not{D} \psi$ is defined in a gauge-invariant manner. The action changes into

$$I(A) \rightarrow \int d^4x [i\bar{\psi} \not{D} \psi - (\partial_\mu \alpha) \bar{\psi} \gamma^\mu \gamma_5 \psi - m\bar{\psi} e^{2i\alpha(x) \gamma_5} \psi - b_\mu j_5^\mu], \quad (2)$$

where the second term $[(\partial_\mu \alpha) \bar{\psi} \gamma^\mu \gamma_5 \psi]$ in the integrand of Eq. (2) comes from transformation of the gauge invariant kinetic term, hence the axial vector current $\bar{\psi} \gamma^\mu \gamma_5 \psi$ is gauge invariant, and does not necessarily equal to j_5^μ , which need not be gauge invariant. However, we must insist that $\int d^4x j_5^\mu$ is gauge invariant, i.e., the density need not be gauge invariant but its space-time integral—the action—is gauge invariant. Since the Chern-Simons term $\frac{1}{2} \epsilon^{\mu\alpha\beta\gamma} F_{\alpha\beta} A_\gamma = {}^*F^{\mu\nu} A_\nu$ behaves precisely in this same way under gauge transformation, we may use it to represent gauge noninvariant portion of j_5^μ . Thus we write

$$j_5^\mu(x) = \bar{\psi}(x) \gamma^\mu \gamma_5 \psi(x) - c {}^*F^{\mu\nu} A_\nu,$$

where c is an unknown constant. By choosing $\alpha(x) = -x^\mu b_\mu$, the gauge invariant current $\bar{\psi} \gamma^\mu \gamma_5 \psi$ cancels and we arrive at

$$\begin{aligned} Z(A) = & \exp\left(-i \int \frac{d^4x}{4\pi^2} b_\mu {}^*F^{\mu\nu} A_\nu\right) \\ & \times \exp\left(i c \int d^4x b_\mu {}^*F^{\mu\nu} A_\nu\right) \int d\bar{\psi} d\psi \\ & \times \exp\left(i \int d^4x [i\bar{\psi} \not{D} \psi - m\bar{\psi} e^{2i\gamma_5 x^\mu b_\mu} \psi]\right). \end{aligned}$$

Now let us calculate the vacuum polarization for the theory governed by the action in the functional integral. The propagator $G(x, y)$ satisfies the following equation:

$$(i \not{\partial}_x - m e^{2i\gamma_5 x \cdot b}) G(x, y) = i \delta(x - y)$$

and its formal solution can be expressed as

$$G(x, y) = \frac{i}{i \not{\partial}_x - m e^{2i\gamma_5 x \cdot b}} \delta(x - y),$$

which to first order in b becomes

*Electronic address: chung@ctpa03.mit.edu

¹See also Refs. [3,4].

$$\frac{i}{i\partial_x - m - 2im\gamma_5 x^\alpha b_\alpha} \delta(x-y)$$

$$\approx S(x-y) - i \int S(x-z) 2im\gamma_5 z^\alpha b_\alpha S(z-y) d^4z$$

$$\equiv S(x-y) + \Delta G(x,y),$$

where $S(x-y)$ is the free propagator

$$\frac{i}{i\partial_x - m} \delta(x-y).$$

This decomposition of the propagator splits the vacuum polarization tensor into three parts:

$$\Pi^{\mu\nu} = \Pi_0^{\mu\nu} + \Pi_b^{\mu\nu} + \Pi_{bb}^{\mu\nu}.$$

The first term $\Pi_0^{\mu\nu}$ is the usual lowest-order vacuum polarization tensor of QED, and the last term $\Pi_{bb}^{\mu\nu}$ is at least quadratic in b . Our main concern here is to calculate the middle term $\Pi_b^{\mu\nu}$, which is linear in b . It is expressed as

$$\Pi_b^{\mu\nu}(x,y) = \text{tr}[\gamma^\mu S(x-y) \gamma^\nu \Delta G(y,x)] + \text{tr}[\gamma^\mu \Delta G(x,y) \gamma^\nu S(y-x)] \equiv \Pi_{b1}^{\mu\nu}(x,y) + \Pi_{b2}^{\mu\nu}(x,y).$$

The Fourier transform of $\Pi_{b1}^{\mu\nu}(x,y)$ is readily obtained as

$$\begin{aligned} \Pi_{b1}^{\mu\nu}(p,q) &= \int d^4x d^4y e^{-ipx} e^{iqy} \Pi_{b1}^{\mu\nu}(x,y) = \int d^4x d^4y d^4z e^{-ipx} e^{iqy} \text{tr}[\gamma^\mu S(x-y) \gamma^\nu (-i) S(y-z) 2im\gamma_5 z^\alpha b_\alpha S(z-x)] \\ &= -2imb_\alpha \text{tr} \left[\int d^4x d^4y d^4z \frac{d^4l}{(2\pi)^4} \frac{d^4r}{(2\pi)^4} \frac{d^4k}{(2\pi)^4} e^{i(-p-l+k)x} e^{i(l-r+q)y} e^{i(r-k)z} \gamma^\mu \frac{1}{l-m} \gamma^\nu \frac{1}{r-m} \gamma_5 z^\alpha \frac{1}{k-m} \right]. \end{aligned}$$

After carrying out x, y, z, r , and k integrations, we are left with a simple momentum integration:

$$\begin{aligned} \Pi_{b1}^{\mu\nu}(p,q) &= -2imb_\alpha \text{tr} \left[\int \frac{d^4l}{(2\pi)^4} \partial_q^\alpha \delta(q-p) \gamma^\mu \frac{1}{l-m} \gamma^\nu \frac{1}{l+q-m} \gamma_5 \frac{1}{l+p-m} \right] \\ &= 2imb_\alpha \delta(q-p) \text{tr} \left[\int \frac{d^4l}{(2\pi)^4} \gamma^\mu \frac{1}{l-m} \gamma^\nu \partial_q^\alpha \frac{1}{l+q-m} \gamma_5 \frac{1}{l+p-m} \right]. \end{aligned}$$

Note that in spite of the explicit x dependence in the action, the vacuum polarization is translation invariant [proportional to $\delta(p-q)$ in momentum space].

Using the identity

$$\partial_q^\alpha \frac{1}{l+q-m} = -\frac{1}{l+q-m} \gamma^\alpha \frac{1}{l+q-m},$$

we have

$$\Pi_{b1}^{\mu\nu}(p,q) = -2imb_\alpha \delta(q-p) \text{tr} \left[\int \frac{d^4l}{(2\pi)^4} \gamma^\mu \frac{1}{l-m} \gamma^\nu \frac{1}{l+q-m} \gamma^\alpha \frac{1}{l+q-m} \gamma_5 \frac{1}{l+p-m} \right],$$

which can be written as

$$\Pi_{b1}^{\mu\nu}(p,q) = 2imb_\alpha \delta(q-p) \text{tr} \left[\int \frac{d^4l}{(2\pi)^4} \gamma^\mu \frac{1}{l-m} \gamma^\nu \frac{1}{l+p-m} \gamma^\alpha \gamma_5 \frac{1}{(l+p)^2 - m^2} \right] \equiv \delta(p-q) b_\alpha \tilde{\Pi}_{b1}^{\mu\nu\alpha}(p).$$

If we observe that $\Pi_{b2}^{\mu\nu}(x,y) = \Pi_{b1}^{\nu\mu}(y,x)$, then we readily obtain that the momentum conserving Fourier transform of $\Pi_{b2}^{\mu\nu}(x,y)$ is given by

$$\tilde{\Pi}_{b2}^{\mu\nu\alpha}(p) = \tilde{\Pi}_{b1}^{\nu\mu\alpha}(-p).$$

The calculation of the integral determining $\tilde{\Pi}_{b1}^{\mu\nu\alpha}(p)$ can be found in Ref. [2]:

$$\tilde{\Pi}_{b1}^{\mu\nu\alpha}(p) = \frac{-i\epsilon^{\mu\nu\alpha\beta} p_\beta}{4\pi^2} \frac{\theta}{\sin\theta},$$

where $\theta \equiv 2\sin^{-1}(\sqrt{p^2}/2m)$ and $p^2 < 4m^2$. Therefore we obtain the induced, parity violating, nonlocal action, which agrees with [2],

$$\frac{1}{4\pi^2} \int \frac{d^4p}{(2\pi)^4} b_\mu {}^*F^{\mu\nu}(p) \left[\frac{\theta}{\sin\theta} - 1 + 4\pi^2 c \right] A_\nu(-p).$$

Since $(\theta/\sin\theta)_{p^2=0}=1$, we obtain the Chern-Simons term with an undetermined strength c . This completes our argument.

We have shown in the functional integral formalism that the (finite) coefficient of the induced, Lorentz- and *CPT*-violating Chern-Simons term arising from the Lorentz-

and *CPT*-violating fermion sector is undetermined. Our result is basically a consequence of the weaker condition for gauge invariance: since $\bar{\psi}\gamma_\mu\gamma_5\psi$ does not couple to any other field, physical gauge invariance is maintained provided $\bar{\psi}\gamma_\mu\gamma_5\psi$ is gauge invariant at zero four-momentum [2].

I thank Professor R. Jackiw for enlightening discussions and acknowledge the Center for Theoretical Physics, MIT, for warm hospitality. This work was supported in part by the Korea Science and Engineering Foundation, and in part by the United States Department of Energy under grant number DF-FC02-94ER40818.

-
- [1] D. Colladay and V. A. Kostelecký, Phys. Rev. D **58**, 116002 (1998).
 [2] R. Jackiw and V. A. Kostelecký, Phys. Rev. Lett. **82**, 3572

- (1999).
 [3] R. Jackiw, Rajaramanfest lecture, hep-th/9903044.
 [4] W. F. Chen, Phys. Rev. D **60**, 085007 (1999).