

**Erratum: Spin asymmetries A_1 of the proton and the deuteron in the low x
and low Q^2 region from polarized high energy muon scattering
[Phys. Rev. D 60, 072004 (1999)]**

B. Adeva, E. Arik, A. Arvidson, B. Badelek, G. Baum, P. Berglund, L. Betev, R. Birsa, N. de Botton, F. Bradamante, A. Bravar, S. Bültmann, E. Burtin, D. Crabb, J. Cranshaw, T. Çuhadar, S. Dalla Torre, R. van Dantzig, B. Derro, A. Deshpande, S. Dhawan, C. Dulya, S. Eichblatt, D. Fasching, F. Feinstein, C. Fernandez, B. Froise, A. Gallas, J. A. Garzon, H. Gilly, M. Giorgi, S. Goertz, G. Garcia, N. de Groot, M. Grosse Perdekamp, K. Haft, D. von Harrach, T. Hasegawa, P. Hautle, N. Hayashi, C. A. Heusch, N. Horikawa, V. W. Hughes, G. Igo, S. Ishimoto, T. Iwata, E. M. Kabuŝ, A. Karev, H. J. Kessler, T. J. Ketel, J. Kiryluk, Yu. Kisselev, D. Krämer, W. Kröger, K. Kurek, J. Kynnäräinen, M. Lamanna, U. Landgraf, J. M. Le Goff, F. Lehar, A. de Lesquen, J. Lichtenstadt, M. Litmaath, A. Magnon, G. K. Mallot, F. Marie, A. Martin, J. Martino, T. Matsuda, B. Mayes, J. S. McCarthy, K. Medved, W. Meyer, G. van Middelkoop, D. Miller, Y. Miyachi, K. Mori, J. Nassalski, T. O. Niinikoski, J. E. J. Oberski, A. Ogawa, C. Ozben, H. Pereira, F. Perrot-Kunne, D. Peshekhonov, R. Piegaia, L. Pinsky, S. Platchkov, M. Plo, D. Pose, H. Postma, J. Pretz, R. Puntaferro, G. Rädcl, G. Reicherz, J. Roberts, M. Rodriguez, E. Rondio, I. Sabo, J. Saborido, A. Sandacz, I. Savin, P. Schiavon, E. P. Sichtermann, F. Simeoni, G. I. Smirnov, A. Staude, A. Steinmetz, U. Stiegler, H. Stuhmann, F. Tessarotto, D. Thers, W. Tłaczala, A. Tripet, G. Unel, M. Velasco, J. Vogt, R. Voss, C. Whitten, R. Willumeit, R. Windmolders, W. Wiŝlicki, A. Witzmann, A. M. Zanetti, K. Zaremba, and J. Zhao

(Spin Muon Collaboration)
(Published 21 August 2000)

PACS number(s): 13.60.Hb, 13.88.+e, 99.10.+g

The virtual photon-proton (-deuteron) asymmetries $A_1^{p,d}$ presented in [1] were measured in the kinematic region where the four-momentum transfer Q^2 extended down to 0.01 GeV². A full account of the formalism is given in [2]. In this kinematic region one cannot neglect m_μ^2/Q^2 terms in the expression for the cross section. These terms were correctly taken into account in the unpolarized part of the cross section, $\bar{\sigma}$ [cf. Eq. (2.2) in [2]]; they were however omitted in the polarized part, $\Delta\sigma$ [Eqs. (2.4)–(2.6) in [2]].

The cross sections $\Delta\sigma_\parallel$ and $\Delta\sigma_T$, corresponding to the two configurations where the nucleon spin is either along or orthogonal to the muon spin [cf., Eq. (2.4) in [2]] should be written as follows:

$$\frac{d^2\Delta\sigma_\parallel}{dx dQ^2} = \frac{16\pi\alpha^2 y}{Q^4} \left[\left(1 - \frac{y}{2} - \frac{\gamma^2 y^2}{4} - \frac{m_\mu^2 y^2}{Q^2} \right) g_1 - \frac{\gamma^2 y}{2} g_2 \right] \quad (1)$$

and

$$\frac{d^3\Delta\sigma_T}{dx dQ^2 d\phi} = -\cos\phi \frac{8\alpha^2 y}{Q^4} \gamma \sqrt{1 - y - \frac{\gamma^2 y^2}{4}} \left[\frac{y}{2} \left(1 + \frac{2m_\mu^2}{Q^2} y \right) g_1 + g_2 \right]. \quad (2)$$

The measured asymmetries $A_\parallel$ and $A_\perp$ are related to A_1 and A_2 [cf., Eqs (2.7)–(2.8) in [2]] through the depolarization factor D ,

$$D = \frac{y[(1 + \gamma^2 y/2)(2 - y) - 2y^2 m_\mu^2/Q^2]}{y^2(1 - 2m_\mu^2/Q^2)(1 + \gamma^2) + 2(1 + R)(1 - y - \gamma^2 y^2/4)}, \quad (3)$$

the factor d for the orthogonal spin configuration,

$$d = \frac{\sqrt{1 - y - \gamma^2 y^2/4}(1 + \gamma^2 y/2)}{(1 - y/2)(1 + \gamma^2 y/2) - y^2 m_\mu^2/Q^2} D, \quad (4)$$

and kinematic factors η and ξ ,

$$\eta = \frac{\gamma(1 - y - \gamma^2 y^2/4 - y^2 m_\mu^2/Q^2)}{(1 + \gamma^2 y/2)(1 - y/2) - y^2 m_\mu^2/Q^2}, \quad (5)$$

$$\xi = \frac{\gamma(1-y/2-y^2m_\mu^2/Q^2)}{1+\gamma^2y/2}. \quad (6)$$

The only approximation applied in these equations is in neglecting terms m_μ^2/E^2 which are of the order of 10^{-7} in our kinematic range. With the above definition the depolarization factor is always smaller than unity.

The missing m_μ^2/Q^2 terms in the polarized part of the cross section is most apparent at low Q^2 . Therefore our low x , low Q^2 data presented in [1] were reanalyzed using the corrected equations. The results for the reanalyzed proton and deuteron spin asymmetries $A_1^{p,d}$ and spin structure functions $g_1^{p,d}$ are given here in Tables I and II for newly accessed region at low Q^2 . The change in $A_1^{p,d}$ and in its statistical error is significant only in the two bins corresponding to the smallest values of x and Q^2 . The average values of x and Q^2 change in the first bin of x because D is used in the weight calculations. Changes at higher x are negligible and the physics conclusions given in [1] are unchanged.

TABLE I. Modifications to Tables III and V in Ref. [1]. The asymmetry $A_1^p(x)$ and the spin structure function $g_1^p(x)$ at the average Q^2 for newly accessed region at low x .

$\langle x \rangle$	$\langle Q^2 \rangle$ (GeV ²)	A_1^p	g_1^p
0.00011	0.03	$0.026 \pm 0.023 \pm 0.009$	$3.5 \pm 3.1 \pm 0.4$
0.00022	0.06	$0.019 \pm 0.019 \pm 0.005$	$2.5 \pm 2.5 \pm 0.6$
0.00039	0.10	$0.002 \pm 0.020 \pm 0.002$	$0.3 \pm 2.5 \pm 0.2$
0.00063	0.17	$-0.004 \pm 0.022 \pm 0.002$	$-0.4 \pm 2.3 \pm 0.2$

TABLE II. Modifications to Tables IV and VI in Ref. [1]. The asymmetry $A_1^d(x)$ and the spin dependent structure function $g_1^d(x)$ at the average Q^2 for newly accessed region at low x .

$\langle x \rangle$	$\langle Q^2 \rangle$ (GeV ²)	A_1^d	g_1^d
0.00011	0.03	$-0.013 \pm 0.050 \pm 0.006$	$-1.7 \pm 6.6 \pm 0.6$
0.00022	0.06	$0.056 \pm 0.040 \pm 0.015$	$7.6 \pm 5.3 \pm 1.6$
0.00039	0.10	$0.030 \pm 0.043 \pm 0.008$	$3.7 \pm 5.3 \pm 1.0$
0.00063	0.17	$0.047 \pm 0.046 \pm 0.012$	$5.0 \pm 4.9 \pm 1.2$

[1] SMC, B. Adeva *et al.*, Phys. Rev. D **60**, 072004 (1999).

[2] SMC, D. Adams *et al.*, Phys. Rev. D **56**, 5330 (1997).