

Magnetic Z_N symmetry in hot QCD and the spatial Wilson loop

C. P. Korthals Altes

Centre Physique Theorique au CNRS, Case 907, Luminy, F13288, Marseille Cedex, France

A. Kovner

Theoretical Physics, University of Oxford, 1 Keble Road, Oxford, OX1 3NP, United Kingdom

(Received 10 April 2000; revised manuscript received 30 May 2000; published 9 October 2000)

We discuss the relation between the deconfining phase transition in gauge theories and the realization of the magnetic Z_N symmetry. At low temperature the Z_N symmetry is spontaneously broken while above the phase transition it is restored. This is intimately related to the change of behavior of the spatial 't Hooft loop. We also point out that the realization of magnetic symmetry has a bearing on the behavior of the spatial Wilson loop. We give a physical argument to the effect that at zero temperature the spatial Wilson loop must have perimeter law behavior in the symmetric phase but area law behavior in the spontaneously broken phase. At high temperature the argument does not hold and the restoration of magnetic Z_N is consistent with the area law for the Wilson loop.

PACS number(s): 11.10.Wx

I. INTRODUCTION

This paper is devoted to further study of the theoretical aspects of the deconfining temperature phase transition in non-Abelian gauge theories. It is an immediate continuation of our earlier work [1]. In [1] we showed that the deconfining phase transition in the pure Yang-Mills (YM) theory is characterized by the change of behavior of the 't Hooft loop operator $V(C)$. In the “cold” phase the 't Hooft loop has a perimeter law behavior $\langle V(C) \rangle \propto \exp\{-aP(C)\}$, while in the “hot” phase it has an area law behavior $\langle V(C) \rangle \propto \exp\{-\alpha S(C)\}$.

In the present paper we want to sharpen somewhat this observation and further discuss related questions. We wish to point out that V is in fact an order parameter which probes the breaking of a physical symmetry of the Yang-Mills theory. The symmetry in question is the magnetic Z_N symmetry discussed by 't Hooft [2]. The deconfining phase transition is therefore characterized by the change in the mode of realization of a global Z_N symmetry: the symmetry is broken spontaneously in the “cold” phase while it is restored in the “hot” phase.

The previous two paragraphs may sound at first like a red herring. After all an order parameter for the deconfining phase transition and a related Z_N symmetry have been discussed for many years. The order parameter in question is the free energy of an external static color source in the fundamental representation: the Polyakov line $P = \text{Tr } P \exp\{ig \int_0^\beta dt A_0\}$. The Z_N symmetry is the transformation $P \rightarrow \exp\{i2\pi/N\}P$. We will refer to this transformation as the electric Z_N . There is however a great difference between the physical nature of P and V and the associated Z_N symmetries. The operator V is a canonical operator in the physical Hilbert space of the Yang-Mills theory. The magnetic Z_N symmetry similarly is a transformation that acts on quantum states in the physical Hilbert space. On the other hand P has a very different status. It is not an operator in the Hilbert space and as such not a canonical order parameter. It appears

as an auxiliary object when projecting onto the gauge invariant physical subspace of the Hilbert space. The “electric” Z_N —the operation that transforms P by multiplying it by a phase—similarly is not a canonical symmetry. There is no transformation of states in the physical Hilbert space that is related to this “symmetry,” although it is indeed a symmetry of the Euclidean path integral representing the statistical sum.

This is not to say of course that P and electric Z_N are useless concepts. The standard effective action, defined by the constrained path integral

$$\exp - S_{\text{eff}}(P) = \int DA_0 \delta(P - P(A_0)) \exp - S(A), \quad (1)$$

is gauge invariant. It is instrumental in computing the vortex expectation value. The way the electric $Z(N)$ symmetry is realized in S_{eff} is also related to the behavior of the order parameter of the magnetic $Z(N)$. We will discuss this in detail in Sec. III [1].

However, if one wants to describe the deconfinement phase transition in terms of a canonical order parameter in the same way as the Ising transition is described in terms of magnetization, one should zero in on V rather than on P and should study the magnetic Z_N symmetry rather than electric Z_N . This is what we intend to do in this paper.

The action of the magnetic Z_N symmetry is very different in the $(2+1)$ - and $(3+1)$ -dimensional cases. In $2+1$ dimensions it acts very much like usual global symmetry in a scalar theory with the order parameter being a scalar vortex field. In $3+1$ dimensions the symmetry acts not like a standard global symmetry—its “charge” is an integral over a two dimensional spacelike surface rather than over the whole of the three dimensional space.¹ As a consequence its order

¹These type of symmetries nowadays are frequently discussed in the context of “M theory” [3].

parameter is not a local field but rather a magnetic vortex stretching over macroscopic distances.

It is therefore convenient to start the discussion with the three dimensional gauge theories and to present all the arguments in this case. The generalization of appropriate aspects of this discussion to 3+1 dimensions will be given in the last part of every section.

The plan of this paper is the following. In Sec. II we recap the definition of the 't Hooft loop and its $(2+1)$ dimensional analog—the magnetic vortex operator. We formulate the arguments for the existence of the magnetic Z_N symmetry in theories without fundamental matter fields. We also show by explicit construction that the generator of this symmetry in the pure gluodynamics is none other than the spatial Wilson loop.

In Sec. III we discuss the relation between the behavior of the 't Hooft loop and the realization of the magnetic Z_N in the ground state of the theory. We demonstrate that the mode of the realization of the symmetry changes at the deconfining phase transition, while spontaneously broken at low temperature the symmetry is restored above the phase transition.

In Sec. IV we present in a toy model a simple physical picture explaining how the behavior of a spatial Wilson loop discriminates at zero temperature between the phases with broken and unbroken magnetic Z_N . In the phase where the Z_N symmetry is broken, W must have an area law while in the case of unbroken Z_N it must have a perimeter law. We explain why this argument does not generalize to the high temperature phase and thus why the area law behavior of the Wilson loop in the hot phase is consistent with restoration of the magnetic Z_N symmetry.

Finally in Sec. V we conclude with a short discussion.

II. THE MAGNETIC Z_N SYMMETRY AND THE 'T HOOFT LOOP OPERATOR

In this section we discuss the notion of the magnetic Z_N symmetry and its order parameter—the 't Hooft loop or magnetic vortex operator. Most of the material contained here is not new and, perhaps with the exception of explicit identification of the Z_N generator with the spatial Wilson loop, is contained in [2,4,5]. At the risk of being repetitive we have decided nevertheless to include this extended introductory part, since we feel that the concept of magnetic Z_N symmetry is not widely appreciated in the community. The Z_N symmetry structure is the basis for our discussion of the deconfining phase transition in the following sections.

Let us start by recalling the argument due to 't Hooft that a non-Abelian $SU(N)$ gauge theory with charged fields in adjoint representation possesses a global Z_N symmetry [2].

We discuss the $(2+1)$ -dimensional case first. Consider a theory with several adjoint Higgs fields so that varying parameters in the Higgs sector the $SU(N)$ gauge symmetry can be broken completely. In this phase the perturbative spectrum will contain the usual massive “gluons” and Higgs particles. However, in addition to that there will be heavy stable magnetic vortices. Those are the analogues of Abrikosov-Nielsen-Olesen vortices in the superconductors and they must be stable by virtue of the following topologi-

cal argument. The vortex configuration away from the vortex core has all the fields in the pure gauge configuration

$$H^\alpha(x) = U(x)h^\alpha, \quad A^\mu = iU\partial^\mu U^\dagger. \quad (2)$$

Here the index α labels the scalar fields in the theory, h^α are the constant vacuum expectation values of these fields, and $U(x)$ is a unitary matrix. As one goes around the location of the vortex in space, the matrix U winds nontrivially in the gauge group. This is possible, since the gauge group in the theory without fundamental fields is $SU(N)/Z_N$ and it has a nonvanishing first homotopy group $\Pi_1(SU(N)/Z_N) = Z_N$. Practically it means that when going around the vortex location full circle, U does not return to the same $SU(N)$ group element U_0 , but rather ends up at $\exp\{i2\pi/N\}U_0$. Adjoint fields do not feel this type of discontinuity in U and therefore the energy of such a configuration is finite. Since such a configuration cannot be smoothly deformed into a trivial one, a single vortex is stable. Processes involving annihilation of N such vortices into a vacuum are allowed since N -vortex configurations are topologically trivial. One can of course find explicit vortex solutions once the Higgs potential is specified. As with any other semiclassical solution in the weak coupling limit the energy of such a vortex is inversely proportional to the gauge coupling constant and therefore very large. One is therefore in a situation where the spectrum of the theory contains a stable particle even though its mass is much higher than masses of many other particles (gauge and Higgs bosons) and the phase space for its decay into these particles is enormous. The only possible reason for the existence of such a heavy stable particle is that it must carry a conserved quantum number. The theory therefore must possess a global symmetry which is unbroken in the completely Higgsed phase. The symmetry group must be Z_N since the number of vortices is only conserved modulo N .

Now imagine changing smoothly the parameters in the Higgs sector so that the expectation values of the Higgs fields become smaller and smaller, and finally the theory undergoes a phase transition into the confining phase. One can further change the parameters so that the adjoint scalars become heavy and eventually decouple completely from the glue. This limiting process does not change the topology of the gauge group and therefore does not change the symmetry content of the theory. We conclude that the pure Yang-Mills theory also possesses a Z_N symmetry. Of course, since the confining phase is separated from the completely Higgsed phase by a phase transition, one may expect that the Z_N symmetry in the confining phase is represented differently. In fact the original paper of 't Hooft as well as subsequent work [4] convincingly argued that in the confining phase the Z_N symmetry is spontaneously broken and this breaking is related to the confinement phenomenon.

The physical considerations given above can be put on firmer formal basis. In particular one can construct explicitly the generator of the Z_N as well as the order parameter associated with it—the operator that creates the magnetic vortex [5]. We will now describe this construction.

A. Abelian case

Consider first an Abelian gauge theory. In this case the homotopy group is Z and therefore we expect the $U(1)$ rather than Z_N magnetic symmetry. It is in fact absolutely straightforward to identify the relevant charge. It is none other than the magnetic flux through the equal time plane, with the associated conserved current being the dual of the electromagnetic field strength

$$\Phi = \int d^2x B(x), \quad \partial^\mu \tilde{F}_\mu = 0. \quad (3)$$

The current conservation is ensured by the Bianchi identity. A group element of the $U(1)$ magnetic symmetry group is $\exp\{i\alpha\Phi\}$ for any value of α . A local order parameter—a local field $V(x)$ which carries the magnetic charge—is also readily constructed. It has a form of the singular gauge transformation operator with the singularity at the point x ,

$$V(x) = \exp \frac{i}{g} \int d^2y \left[\epsilon_{ij} \frac{(x-y)_j}{(x-y)^2} E_i(y) + \Theta(x-y) J_0(y) \right] \quad (4)$$

where $\Theta(x-y)$ is the polar angle function and J_0 is the electric charge density of whatever matter fields are present in the theory. The cut discontinuity in the function Θ is not physical and can be chosen parallel to the horizontal axis. Using the Gauss' law constraint this can be cast in a different form, which we will find more convenient for our discussion,

$$V(x) = \exp \frac{2\pi i}{g} \int_C dy^i \epsilon_{ij} E_j(y), \quad (5)$$

where the integration goes along the cut of the function Θ which starts at the point x and goes to spatial infinity. The operator does not depend on where precisely one chooses the cut to lie. To see this, note that changing the position of the cut $C-C'$ adds to the phase $(2\pi/g) \int_S d^2x \partial_i E^i$ where S is the area bounded by $C-C'$. In the theory we consider only charged particles with charge multiples of g are present. Therefore the charge within any closed area is a multiple integer of the gauge coupling $\int_S d^2x \partial_i E^i = gn$ and the extra phase factor is always unity.

The meaning of the operator V is very simple. From the commutation relation

$$V(x)B(y)V^\dagger(x) = B(y) + \frac{2\pi}{g} \delta^2(x-y) \quad (6)$$

it is obvious that V creates a pointlike magnetic vortex of flux $2\pi/g$. Despite its nonlocal appearance the operator V can be proved to be a local Lorentz scalar field [6]. The locality is the consequence of the fact that $V(x)$ commutes with any local gauge invariant operator in the theory $O(y)$ except when $x=y$. This is due to the coefficient $2\pi/g$ in the exponential which ensures that the Aharonov-Bohm phase of the vortex created by V and any dynamical charged particle present in the theory vanishes. Equations (3), (5) formalize the physical arguments of 't Hooft in the Abelian case.

B. Non-Abelian case at weak coupling

Let us now move onto the analogous construction for non-Abelian theories. Ultimately we are interested in the pure Yang-Mills theory. It is however illuminating to start with the theory with an adjoint Higgs field and take the decoupling limit explicitly later. For simplicity we discuss the $SU(2)$ gauge theory. Consider the Georgi-Glashow model— $SU(2)$ gauge theory with an adjoint Higgs field:

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu} + \frac{1}{2} (\mathcal{D}_\mu^{ab} H^b)^2 + \tilde{\mu}^2 H^2 - \tilde{\lambda} (H^2)^2 \quad (7)$$

where

$$\mathcal{D}_\mu^{ab} H^b = \partial_\mu H^a - g f^{abc} A_\mu^b H^c. \quad (8)$$

At large and positive $\tilde{\mu}^2$ the model is weakly coupled. The $SU(2)$ gauge symmetry is broken down to $U(1)$ and the Higgs mechanism takes place. Two gauge bosons, $W^\pm$, acquire a mass, while the third one, the ‘‘photon,’’ remains massless to all orders in perturbation theory. The theory in this region of parameter space resembles very much electrodynamics with vector charged fields. The Abelian construction can therefore be repeated. The $SU(2)$ gauge invariant analogue of the conserved dual field strength is

$$\tilde{F}^\mu = \frac{1}{2} \left[\epsilon^{\mu\nu\lambda} F_{\nu\lambda}^a n^a - \frac{1}{g} \epsilon^{\mu\nu\lambda} \epsilon^{abc} n_a (\mathcal{D}_\nu n)^b (\mathcal{D}_\lambda n)^c \right] \quad (9)$$

where $n^a \equiv H^a/|H|$ is the unit vector in the direction of the Higgs field. Classically this current satisfies the conservation equation

$$\partial^\mu \tilde{F}_\mu = 0. \quad (10)$$

The easiest way to see this is to choose a unitary gauge of the form $H^a(x) = H(x) \delta^{a3}$. In this gauge $\tilde{F}$ is equal to the Abelian part of the dual field strength in the third direction in color space. Its conservation then follows by the Bianchi identity. Thus classically the theory has a conserved $U(1)$ magnetic charge $\Phi = \int d^2x \tilde{F}_0$ just like QED. However, the unitary gauge cannot be imposed at the points where H vanishes, which necessarily happens in the core of a 't Hooft–Polyakov monopole. It is well known of course [7] that the monopoles are the most important nonperturbative configurations in this model. Their presence leads to a nonvanishing small mass for the photon as well as to confinement of the charged gauge bosons with a tiny nonperturbative string tension. As far as the monopole effects on the magnetic flux are concerned, their presence leads to a quantum anomaly in the conservation equation (10). As a result only the discrete Z_2 subgroup of the transformation group generated by Φ remains unbroken in the quantum theory. The detailed discussion of this anomaly, the residual Z_2 symmetry and their relation to monopoles is given in [5].

The order parameter for the magnetic Z_2 symmetry is constructed analogously to QED as a singular gauge transformation generated by the gauge invariant electric charge operator

$$J^\mu = \epsilon^{\mu\nu\lambda} \partial_\nu (\tilde{F}_\lambda^a n^a), \quad Q = \int d^2x J_0(x). \quad (11)$$

Explicitly

$$\begin{aligned} V(x) &= \exp \frac{i}{g} \int d^2y \left[\epsilon_{ij} \frac{(x-y)_j}{(x-y)^2} n^a(y) E_i^a(y) \right. \\ &\quad \left. + \Theta(x-y) J_0(y) \right] \\ &= \exp \frac{2\pi i}{g} \int_C dy^i \epsilon_{ij} n^a E_j^a(y). \end{aligned} \quad (12)$$

One can think of it as a singular $SU(2)$ gauge transformation with the field dependent gauge function

$$\lambda^a(y) = \frac{1}{g} \Theta(x-y) n^a(\vec{y}). \quad (13)$$

This field dependence of the gauge function ensures the gauge invariance of the operator V . Just like in QED it can be shown [5,6] that the operator V is a local scalar field. Again, like in QED, the vortex operator V is a local eigenoperator of the Abelian magnetic field $B(x) = \tilde{F}_0$:

$$[V(x), B(y)] = -\frac{2\pi}{g} V(x) \delta^2(x-y). \quad (14)$$

That is to say, when acting on a state it creates a pointlike magnetic vortex which carries a quantized unit of magnetic flux. The Z_2 magnetic symmetry transformation is generated by the operator

$$U = \exp \left\{ i \frac{g}{2} \Phi \right\} \quad (15)$$

and acts on the vortex field V as a phase rotation by π :

$$e^{i(g/2)\Phi} V(x) e^{-i(g/2)\Phi} = -V(x). \quad (16)$$

An operator closely related to U and which will be of interest to us in the following is the generator of the magnetic Z_2 transformation only inside some closed contour C :

$$U(C) = \exp \left\{ i \frac{g}{2} \int_S d^2x B(x) \right\} \quad (17)$$

where the integration is over the area S bounded by C . The analogue of the commutator, Eq. (16), for this operator is

$$\begin{aligned} U_C V(x) U_C^\dagger &= -V(x), \quad x \in S, \\ V(x), \quad x \notin S. \end{aligned} \quad (18)$$

Taking the contour C to run at infinity U_C becomes the generator of Z_2 .

We now have the explicit realization of the magnetic Z_2 symmetry in the Georgi-Glashow model.

C. Pure gauge theory

Our next step is to move on to the pure Yang-Mills theory. This is achieved by smoothly varying the $\tilde{\mu}^2$ coefficient in the Lagrangian so that the coefficient of the mass term of the Higgs field becomes positive and eventually arbitrarily large. It is well known that in this model the weakly coupled Higgs regime and strongly coupled confining regime are not separated by a phase transition [8]. The pure Yang-Mills limit in this model is therefore smooth.

In the pure Yang-Mills limit expressions (9), (12), (17) have to be taken with care. When the mass of the Higgs field is very large, the configurations that dominate the path integral are those with very small value of the modulus of the Higgs field $|H| \propto 1/M$. The modulus of the Higgs field in turn controls the fluctuations of the unit vector n^a , since the kinetic term for n in the Lagrangian is $|H|^2 (D_\mu n)^2$. Thus as the mass of the Higgs field increases the fluctuations of n grow in both amplitude and frequency and the magnetic field operator B as defined in Eq. (9) fluctuates wildly. This situation is of course not unusual. It happens whenever one wants to consider in the effective low energy theory an operator which explicitly depends on fast, high energy variables. The standard way to deal with it is to integrate over the fast variables. There could be two possible outcomes of this procedure. Either the operator in question becomes trivial (if it depends strongly on the fast variables) or its reduced version is well defined and regular on the low energy Hilbert space. The ‘‘magnetic field’’ operator B in Eq. (9) is obviously of the first type. Since in the pure Yang-Mills limit all the orientations of n^a are equally probable, integrating over the Higgs field at fixed A_μ will lead to vanishing of B . However, what interests us is not so much the magnetic field but rather the generator of the magnetic Z_2 transformation U_C of Eq. (17). In the pure Yang-Mills limit we are thus lead to consider the operator

$$\begin{aligned} U_C &= \lim_{H \rightarrow 0} \int Dn^a \exp \{ -|H|^2 (\vec{D}n_a)^2 \} \\ &\quad \times \exp \left\{ i \frac{g}{4} \int_C d^2x \left(\epsilon_{ij} F_{ij}^a n^a \right. \right. \\ &\quad \left. \left. - \frac{1}{g} \epsilon^{ij} \epsilon^{abc} n_a (\mathcal{D}_i n)^b (\mathcal{D}_j n)^c \right) \right\}. \end{aligned} \quad (19)$$

The weight for the integration over n is the kinetic term for the isovector n_a . As was noted before the action does not depend on n^a in the YM limit. This term however regulates the integral and we keep it for this reason. This operator may look somewhat unfamiliar at first sight. However, in a re-

markable paper [9] Diakonov and Petrov showed that Eq. (19) is equal to the trace of the fundamental Wilson loop along the contour C :²

$$U_C = W_C \equiv \text{Tr} \mathcal{P} \exp \left\{ ig \int_C dl^i A^i \right\}. \quad (20)$$

We conclude that in the pure Yang-Mills theory the generator of the magnetic Z_2 symmetry is the fundamental spatial Wilson loop along the boundary of the spatial plane.

There is a slight subtlety here that may be worth mentioning. The generator of a unitary transformation should be a unitary operator. The trace of the fundamental Wilson loop on the other hand is not unitary. One should therefore strictly speaking consider instead a unitarized Wilson loop $\tilde{W} = W/\sqrt{WW^\dagger}$. However, the factor between the two operators $\sqrt{WW^\dagger}$ is an operator that is only sensitive to behavior of the fields at infinity. It commutes with all physical local operators $O(x)$ unless $x \rightarrow \infty$. In this it is very different from the Wilson loop itself, which has a nontrivial commutator with vortex operators $V(x)$ at all values of x . Since the correlators of all gauge invariant local fields in the pure Yang-Mills theory are massive and therefore short range, the operator $\sqrt{WW^\dagger}$ must be a constant operator on all finite energy states. The difference between W and $\tilde{W}$ is therefore a trivial constant factor and we will not bother with it in the following. Perhaps of more concern is the difference between W_C and $\tilde{W}_C$ when the contour C is not at infinity. However, here again the factor between the two operators, $\sqrt{W_C W_C^\dagger}$, is only sensitive to physical degrees of freedom on the contour C and not inside it. Because of its presence, the vacuum averages of W_C and $\tilde{W}_C$ may differ at most by a factor which has a perimeter behavior, $\langle W_C \rangle = \exp\{mP(C)\} \langle \tilde{W}_C \rangle$, where $P(C)$ is the perimeter of C . The question we will be interested in is whether $\langle W_C \rangle$ has a perimeter or area behavior. As far as the answer to this question is concerned W_C and $\tilde{W}_C$ are completely equivalent, and we will not make distinction between them. In the rest of this paper we will therefore refer to W as the generator of Z_2 , keeping this little caveat in mind.

Next we consider the vortex operator, Eq. (12). Again we have to integrate it over the orientations of the unit vector n^a . This integration in fact is equivalent to averaging over the gauge group. Following [9] one can write n_a in terms of the $SU(2)$ gauge transformation matrix Ω :

$$\vec{n} = \frac{1}{2} \text{Tr} \Omega \tau \Omega^\dagger \tau_3. \quad (21)$$

The vortex operator in the pure gluodynamics limit then becomes

²We note that Diakonov and Petrov had to introduce a regulator to define the path integral over n . The regulator they required was precisely of the same form as in Eq. (19). It is pleasing to see that this regulator appears naturally in our approach as the remnant of the kinetic term of the Higgs field.

$$\tilde{V}(x) = \int D\Omega \exp \frac{2\pi i}{g} \int_C dy^i \epsilon_{ij} \text{Tr} \Omega E_j \Omega^\dagger \tau_3. \quad (22)$$

This form makes it explicit that $\tilde{V}(x)$ is defined as the gauge singlet part of the following, apparently non-gauge-invariant operator:

$$V(x) = \exp \frac{2\pi i}{g} \int_C dy^i \epsilon_{ij} E_i^3(y). \quad (23)$$

The integration over Ω obviously projects out the gauge singlet part of V . In the present case however this projection is redundant. This is because even though V itself is not gauge invariant, when acting on a physical state it transforms it into another physical state.³ By physical states we mean the states which satisfy the Gauss' constraint in the pure Yang-Mills theory. This property of V was noticed by 't Hooft [2]. To show this let us consider $V(x)$ as defined in Eq. (23) and its gauge transform $V_\Omega = \Omega^\dagger V \Omega$ where Ω is an arbitrary nonsingular gauge transformation operator. The wave functional of any physical state depends only on gauge invariant characteristics of the vector potential, i.e. only on the values of Wilson loops over all possible contours:

$$\Psi[A_i] = \Psi[\{W(C)\}]. \quad (24)$$

Acting on this state by the operators V and V_Ω respectively we obtain

$$V|\Psi\rangle = \Psi_V[A_i] = \Psi[\{VW(C)V^\dagger\}]$$

$$V_\Omega|\Psi\rangle = \Psi_V^\Omega[A_i] = \Psi[\{V_\Omega W(C)V_\Omega^\dagger\}]. \quad (25)$$

It is however easy to see that the action of $V(x)$ and $V_\Omega(x)$ on the Wilson loop is identical—they both multiply it by the center-group phase (which stays unaffected by Ω) if x is inside C and do nothing otherwise. Therefore we see that

$$V|\Psi\rangle = V_\Omega|\Psi\rangle \quad (26)$$

for any physical state Ψ . Thus we have

$$\Omega V|\Psi\rangle = \Omega V \Omega^\dagger |\Psi\rangle = V|\Psi\rangle \quad (27)$$

where the first equality follows from the fact that a physical state is invariant under action of any gauge transformation Ω and the second equality follows from Eq. (26). But this equation is nothing but the statement that the state $V|\Psi\rangle$ is physical, i.e. invariant under any nonsingular gauge transformation.

We have therefore proved that when acting on a physical state the vortex operator creates another physical state. For an operator of this type the gauge invariant projection only affects its matrix elements between unphysical states. Since we are only interested in calculating correlators of V between

³This is not a trivial statement, since a generic non-gauge-invariant operator has nonvanishing matrix elements between the physical and an unphysical sector.

physical states, the gauge projection is redundant and we can freely use V rather than $\tilde{V}$ to represent the vortex operator.

It is instructive to note that this property is not shared by the Wilson loop. One can in fact represent the Wilson loop as a singlet gauge projection of a simple Abelian loop operator. The second exponential in Eq. (20) can be written as

$$\exp\left\{i\frac{g}{2}\int_C dl^i A_d^i n^a - \frac{i}{2}\int d^2x \epsilon_{ij} \epsilon^{abc} n_a \partial_i n_b \partial_j n_c\right\}. \quad (28)$$

Using Eq. (21) we can rewrite the integral in Eq. (20)—omitting the regulating kinetic piece—as

$$W_C = \int D\Omega \exp\left\{i\frac{g}{2}\int \text{Tr} \tau_3(\Omega A^i \Omega^\dagger + i\Omega \partial^i \Omega^\dagger) dl^i\right\}. \quad (29)$$

The Wilson loop is therefore the gauge singlet part of the Abelian loop:

$$U_C^A = \exp i\frac{g}{2}\int \text{Tr} A^i \tau_3 dl^i. \quad (30)$$

The matrix elements of W_C and U_C^A on physical subspace therefore are the same. However, U_C^A as opposed to V does have nonvanishing nondiagonal matrix elements, that is matrix elements between the physical and the unphysical sectors. It therefore *cannot* be used instead of W_C in gauge theory calculations. For example non-gauge-invariant states will contribute as intermediate states in the calculation of quantities like the correlation function $\langle U_{C1}^A U_{C2}^A \rangle$, while their contribution vanishes in similar correlators which involve the Wilson loop.

The generalization of the preceding discussion to $SU(N)$ gauge theories is straightforward. Once again one can start with the Georgi-Glashow-like model, where the $SU(N)$ is broken down to $U(1)^{(N-1)}$ by the Higgs mechanism.⁴ The construction of the vortex operator and the generator of Z_N in this case is very similar and the details are given in [5]. Taking the mass of the Higgs field to infinity again projects the generator onto the trace of the fundamental Wilson loop. The vortex operator can be taken as

$$V(x) = \exp\left\{\frac{4\pi i}{gN} \int_C dy^i \epsilon_{ij} \text{Tr}[YE_i(y)]\right\} \quad (31)$$

where the hypercharge generator Y is defined as

$$Y = \text{diag}(1, 1, \dots, -(N-1)) \quad (32)$$

⁴In $SU(N)$ theories with $N > 2$ there in principle can be phases separated from each other due to spontaneous breaking of some global symmetries. For instance the $SU(3)$ gauge theory with adjoint matter has a phase with spontaneously broken charge conjugation invariance [10]. Still even in this phase the confining properties are the same as in the strongly coupled pure Yang-Mills theory, with the Wilson loop having an area law.

and the electric field is taken in the matrix notation $E_i = \lambda^a E_i^a$ with λ^a —the $SU(N)$ generator matrices in the fundamental representation.

D. Generalization to 3+1 dimensions

To conclude this section we discuss how the magnetic symmetry structure generalizes to four dimensions. The conserved Z_N generator in the Georgi-Glashow model is defined through

$$U_S = \exp\left\{i\frac{g}{2}\int_S d^2S^i \left(B_i^a n^a - \frac{1}{g} \epsilon^{ijk} \epsilon^{abc} n_a (\mathcal{D}_j n)^b (\mathcal{D}_k n)^c\right)\right\}. \quad (33)$$

Although the definition of U contains explicitly the surface S through which the Abelian magnetic flux is integrated, the operator in fact does not depend on S but is specified completely by its boundary. This is because changing S changes the phase of U by the magnetic flux through the closed surface. The only dynamical objects that carry magnetic flux in the theory are 't Hooft–Polyakov monopoles. Since their flux is quantized in units of $4\pi/g$, the change in the phase is always a multiple integer of 2π . In the pure Yang-Mills limit the operator U_S again reduces to the trace of the fundamental Wilson loop along the boundary of S . Taking the contour to infinity defines the generator of magnetic Z_N . As we have already noted, this charge is a little unusual in that it is defined as a surface integral rather than a volume integral. As a result the order parameter for this symmetry transformation is not a local but rather a stringy field. This is of course just a restatement of the fact that magnetic vortices in 3+1 dimensions are stringlike objects. The operator that creates a vortex can still be defined in a way similar to 2+1 dimensions. Skipping the intermediate steps which we went through in the previous discussion we give the final result for the pure Yang-Mills $SU(N)$ gauge theory. The magnetic vortex along the curve C is created by the following operator of the “singular gauge transformation”⁵

$$\begin{aligned} V(C) &= \exp\left\{\frac{i}{gN} \int d^3x \text{Tr}(D^i \omega_C Y) E^i\right\} \\ &= \exp\left\{\frac{4\pi i}{gN} \int_S d^2S^i \text{Tr}(Y E^i)\right\} \end{aligned} \quad (34)$$

with $\omega_C(x)$, the singular gauge function which is equal to the solid angle subtended by C as seen from the point x . The function ω is continuous everywhere, except on a surface S bounded by C , where it jumps by 4π . Other than the fact that S is bounded by C , its location is arbitrary. The vortex loop and the spatial Wilson loop satisfy the 't Hooft algebra

$$V^\dagger(C) W(C') V(C) = e^{(2\pi i/N)n(C, C')} W(C') \quad (35)$$

⁵The derivative term $\partial^i \omega$ in this expression should be understood to contain only the smooth part of the derivative and to exclude the contribution due to the discontinuity of ω on the surface S .

where $n(C, C')$ is the linking number of the curves C and C' . One can consider closed contours C or infinite contours that run through the whole system. For an infinite contour C and the Wilson loop along the spatial boundary of the system the linking number is always unity. The $V(C)$ for an infinite loop is therefore an eigenoperator of the Z_N magnetic symmetry and is the analogue of the vortex operator $V(x)$ in 2+1 dimensions. Any closed vortex loop of fixed size commutes with the Wilson loop if the contour C' is very large. Such a closed loop is thus an analogue of the vortex-antivortex correlator $V(x)V^\dagger(y)$, which also commutes with the global symmetry generator, but has a nontrivial commutator with U_C if C encloses only one of the points x or y .

To summarize this section, we have shown that pure Yang-Mills theory in 2+1 and 3+1 dimensions has a global Z_N magnetic symmetry. The generator of the symmetry group in both cases is the trace of the fundamental Wilson loop along the spatial boundary of the system. The order parameter for this symmetry in 2+1 dimensions is a local scalar field $V(x)$, while in 3+1 dimensions a stringlike field $V(C)$. In both cases the field V is gauge invariant on physical states and is a *bona fide* canonical order parameter which distinguishes in gauge invariant way the phases of the theory. In the next section we discuss the realization of the magnetic symmetry in the confined and the deconfined phases.

III. HOT AND COLD REALIZATIONS OF THE MAGNETIC Z_N

As with any global symmetry, it is important to understand what is the mode of realization of magnetic Z_N in the ground state of the theory. This mode of realization depends of course on the parameters of the theory as well as on the temperature. The situation at zero temperature is well understood.

A. 2+1 dimensions

Again we start with three dimensions. There is a very general argument⁶ due to 't Hooft [2] stating that if the theory does not have zero mass excitations, the area law of the Wilson loop implies the nonvanishing expectation value of the vortex operator $V(x)$. Conversely, if the Wilson loop has a perimeter law, the expectation value of $V(x)$ must vanish and the correlation function $V(x)V^\dagger(y)$ must have an exponential falloff with $|x-y|$. It follows that in the pure Yang-Mills theory the vacuum expectation value of the vortex operator does not vanish and therefore the Z_N magnetic symmetry is spontaneously broken. The same is true in the partially broken Higgs phase of the Georgi-Glashow model. As mentioned in the last section the confining and the Higgs regimes in this model are analytically connected and therefore the realization of all global symmetries in the two regimes is the same.

In fact in the weakly coupled Higgs phase this can be verified by the direct calculation of the expectation value of V [5]. This calculation maps very simply into the classic monopole plasma calculation of Polyakov and was discussed in detail in [5]. One can also explicitly construct the low energy effective Lagrangian in terms of the field V which realizes the spontaneously broken Z_N symmetry and describes the low energy spectrum of the Georgi-Glashow vacuum:

$$\mathcal{L} = \partial_\mu V^* \partial^\mu V - \lambda (V^* V - \mu^2)^2 - \zeta (V^2 + V^{*2}). \quad (36)$$

A similar effective Lagrangian with some quantitative differences was argued to be valid also for the pure Yang-Mills theory in [11].

The application of the 't Hooft argument at finite temperature is somewhat less straightforward. Since at finite temperature the Lorentz invariance is broken, the temporal and spatial Wilson loops do not necessarily have the same behavior and one has to be more careful. The original argument relates the behavior of the vortex operator and the temporal Wilson loop. At finite temperature in the Euclidean formalism the extent of the system in the temporal direction is finite. As a result it is not possible to distinguish between the area and perimeter law for “asymptotically” large temporal loops. Instead the role of the temporal Wilson loop is taken over by the Polyakov line—the loop that winds around the total volume of the system in the temporal direction. Thus one expects that in the deconfining phase where the Polyakov line has a nonvanishing vacuum average, the vortex operator should have vanishing expectation value. Indeed this can be easily confirmed by the explicit calculation of the vacuum expectation value (VEV) of the vortex operator using the method of [1]. In [1] the calculation was performed in 3+1 dimensions, but adapting it to the (2+1)-dimensional case is trivial. We give below a brief outline.

Consider the equal time vortex-antivortex correlation function. At finite temperature it is given by the following expression:

$$\begin{aligned} \langle V(x)V^\dagger(y) \rangle &= \text{Tr} \exp \left(-\frac{\beta}{2} (E^2 + B^2) \right) \\ &\times \exp \left(i \frac{2\pi i}{g} \int_x^y \epsilon_{ij} dl^i E_3^j \right). \end{aligned} \quad (37)$$

The line integral can be taken along the straight line L connecting the points x and y . For definiteness we take x and y to be separated in the direction of the first axis. Introducing the imaginary time axis and the Lagrange multiplier field A_0 in the standard way this expression can be transformed to

$$\begin{aligned} \langle V(x)V^\dagger(y) \rangle &= \int DA_i DA_0 \exp \left\{ -\frac{1}{2} \int_0^\beta dt \int d^3x \right. \\ &\quad \times \left[\partial_0 A_i^a - (D_i A_0)^a - \delta(t) a_i^a \right]^2 + (B^a)^2 \left. \right\} \end{aligned} \quad (38)$$

⁶The original argument as stated in [2] is formulated for (3+1)-dimensional theories; however, its generalization to 2+1 dimensions requires only linguistic changes.

where the “external field” a^i is given by

$$a_i^a(\mathbf{x}) = \delta^{a3} \delta_{i2} \frac{2\pi}{g} \delta(\mathbf{x} - L). \quad (39)$$

The delta function in time in front of the external field a^i in Eq. (38) appears for the following reason. As we saw in Sec. II C the product of the vortex and an antivortex operator is gauge invariant. This is because it induces a singular gauge transformation which is continuous up to the center element. However, if we split it up in imaginary time into infinitesimal bits $V_{dt} = \exp[i dt T(2\pi i/g) \int_x^\gamma \epsilon_{ij} dl^j E_3^i]$, then any single such bit separately is not gauge invariant, since the transformation it induces is genuinely discontinuous across the line connecting the points x and y . The operators V_{dt} therefore do not commute with the projection operator on physical states. To obtain the path integral representation for the expectation value, Eq. (37), we should introduce the projection operator only at the last point in imaginary time and not at the intermediate points. In path integral language this corresponds to the gauge fixing $A_0 = 0$ everywhere except at one time, say $t = 0$. In this gauge it is straightforward to see that the Gaussian integration over the electric field leads to the usual path integral with $\partial_i A_0$ shifted by a_i . This can then be rotated to an arbitrary gauge with the result Eq. (38).⁷

To evaluate the path integral, Eq. (38), we follow the standard procedure and integrate out all modes except for the Polyakov loop in a saddle point approximation [13,14]. This leads to the effective action $S_{eff}(q, a_i)$, where $1/2 \text{Tr } P = \cos q$:

$$\langle V(x) V^\dagger(y) \rangle = \int Dq \exp - S_{eff}(q, a_i). \quad (40)$$

To one loop order the effective action is given by

$$S_{eff} = \int d^2x \left[\frac{2T}{g^2} \left(\partial_i q + \frac{g}{2} a_i \right)^2 + U(q) \right]. \quad (41)$$

The matrices τ^a in A_μ are the generators of $SU(2)$ in the fundamental representation and are normalized according to $\text{tr } \tau^a \tau^b = \frac{1}{2} \delta^{ab}$.

The one loop effective potential U is related to a Bernoulli polynomial and can be read off the expressions in [12,13]. The only property of U which is important to us is that it has two degenerate minima at $q = 0, \pi$.

To calculate the correlator we have to find the configuration of q which minimizes the action, Eq. (41). Qualitatively the form of the solution is clear. The considerations identical to those in [1] tell us that it must be the “broken” electric Z_2 domain wall: half a wall [$q \rightarrow_{x_2 \rightarrow \infty} 0$, $q(x_2 = 0) = \pi/2$] above

the line L and half a wall [$q(x_2 = 0) = -\pi/2$, $q \rightarrow_{x_2 \rightarrow -\infty} 0$] below the line L separated by a discontinuity $\delta q = \pi$. The action of such a configuration is $S_{eff} = \tilde{\alpha} |x - y|$ where $\tilde{\alpha}$ is the “ Z_2 domain wall tension.” The vortex correlator is thus given by

$$\langle V(x) V^\dagger(y) \rangle = \exp\{-\tilde{\alpha} |x - y|\}. \quad (42)$$

As $|x - y|$ becomes large the correlation function decreases exponentially, and thus the expectation value of the vortex operator vanishes. For the $SU(N)$ group this calculation trivially generalizes and gives the same result. The exponential decay is also obtained for the correlator of V^m with any power $m < N$.

Recall that the vortex operator is the order parameter for the magnetic Z_N symmetry. Moreover, the powers of V exhaust all possible local order parameters.⁸ Their vanishing is therefore an unambiguous indication that the magnetic Z_N is restored in the high temperature deconfined phase.

In hindsight this is not very surprising. Indeed, we are dealing with physical discrete symmetry which is spontaneously broken at zero temperature. When the system is heated it is unavoidable that entropy effects take over and at some sufficiently high temperature the symmetry must be restored. A good qualitative guide here is the effective Lagrangian, Eq. (36). It describes a simple Z_2 invariant scalar theory. There is very little doubt that a system described by this Lagrangian indeed undergoes a symmetry restoring phase transition at some T_c . Moreover, the effective Lagrangian approach also suggests that this phase transition has deconfining character. As shown in [5,6,11] the charged states in the effective theory, Eq. (36), are represented by solitonic configurations of the vortex field V with unit winding number. The energy of any such state is linearly divergent in the infrared. The reason is that due to the finite degeneracy of vacuum states, the minimum energy configuration looks like a quasi-one-dimensional strip across which the phase of V winds. The energy density inside this “electric flux tube” is proportional to the vacuum expectation value of V . When the VEV vanishes, so does the string tension. Stated in other words, when V vanishes, the phase fluctuations are large and the winding number is not a sharp observable. An external charge is thus screened easily by regions of space around it with vanishing V . The phase with $\langle V \rangle = 0$ is therefore not confining. In the theory with several Higgs fields this phase exists even at zero temperature and corresponds to a completely Higgsed phase—where the gauge group is broken completely. In such a Higgs phase indeed the color is screened rather than confined. In the pure Yang-Mills theory this phase is absent at zero temperature, but is realized as the deconfined phase at $T > T_c$. We thus see that the behavior of

⁷In this derivation we dropped commutator terms between the Hamiltonian and the exponent in the vortex-antivortex operator. These commutator terms only exist at $t = 0$ and therefore drop out in the continuum limit [i.e. are $O(dt)$]. In the lattice realization they are indeed present and complete the expression (38) to the twisted plaquette representation [12].

⁸The latter statement is correct modulo multiplication of V^m by local gauge invariant and Z_N invariant operators. These possible factors do not change the fact of the exponential decay of the correlators and are therefore unimportant for our discussion.

the vortex operator at high temperature does indeed match the simple intuition coming from a Z_N invariant effective Lagrangian very well.

B. Extension to 3+1 dimensions

The 't Hooft argument now states that the vanishing vacuum average of the Polyakov line is incompatible with the area law behavior of the spatial 't Hooft loop and vice versa. This means that in the confining phase the 't Hooft loop has perimeter law. In the high temperature deconfined phase the behavior of the spatial 't Hooft loop must become the area since the average of the Polyakov line is finite. Again this is confirmed by explicit calculation in [1].

A more subtle question is how the behavior of the 't Hooft loop relates to the realization of the magnetic Z_N symmetry. The Z_N symmetry does not have an order parameter which is a local field defined at a point. The only order parameters in the strict sense (an eigenoperator with a non-vanishing eigenvalue) is a 't Hooft line $V(C)$ which runs through the whole system [15].

In a system which is finite in the direction of the loop, but is infinite in the perpendicular directions, everything is clear cut. In this case there are two possibilities:

- (a) $\langle V \rangle \neq 0$ and the magnetic Z_N broken or
- (b) $\langle V \rangle = 0$ and the magnetic I_N restored.

In the system infinite in all directions C is necessarily an infinite line, and the expectation value $\langle V(C) \rangle$ clearly vanishes irrespective of whether Z_N is broken or not. The 't Hooft loop along a closed contour on the other hand is never zero, since it is globally invariant under the Z_N transformation. It is therefore impossible to find an operator whose VEV distinguishes between the two possible realizations of the magnetic symmetry by vanishing in one phase and not vanishing in the other. Nevertheless, the behavior of the closed loop does indeed reflect the realization mode of the symmetry, since it is qualitatively different in the two possible phases. Namely the vacuum expectation value of a large closed 't Hooft loop (by large, as usual we mean that the linear dimensions of the loop are much larger than the correlation length in the theory) has an area law decay if the magnetic symmetry is spontaneously broken and perimeter law decay if the vacuum state is invariant.

To understand the physics of this behavior it is useful to think of the 't Hooft line as built of "local" operators—little "magnetic dipoles." Consider Eq. (34) with the contour C running along the x axis and the surface S chosen as the (x, y) plane. Let us mentally divide the line into (short) segments of length 2Δ centered at x_i . Each one of these segments is a little magnetic dipole and the 't Hooft loop is a product of the operators that create these dipoles. The definition of these little dipole operators is somewhat ambiguous but since we only intend to use them here for the purpose of an intuitive argument any reasonable definition will do. It is convenient to define a single dipole operator in the following way:

$$D_\Delta(x) = \exp \left\{ i \int d^3y [a_i^+(x + \Delta - y) + a_i^-(x - \Delta - y)] \text{Tr}[YE^i(y)] \right\} \quad (43)$$

where $a_i^\pm(x - y)$ is the c-number vector potential of the Abelian magnetic monopole (antimonopole) of strength $4\pi/gN$. The monopole field corresponding to a_i contains both the smooth x_i/x^3 part as well as the Dirac string contribution. The Dirac string of the monopole-antimonopole pair in Eq. (43) is chosen so that it connects the points $x - \Delta$ and $x + \Delta$ along the straight line. The dipole operators obviously have the property

$$D_\Delta(x)D_\Delta(x + 2\Delta) = D_{2\Delta}(x + \Delta). \quad (44)$$

This is because in the product the smooth field contribution of the monopole in $D_\Delta(x)$ cancels the antimonopole contribution in $D_\Delta(x + 2\Delta)$, while the Dirac string now stretches between the points $(x - \Delta)$ and $(x + 3\Delta)$. When multiplied over the closed contour, the smooth fields cancel out completely, while the surviving Dirac string is precisely the magnetic vortex created by a closed 't Hooft loop operator. The 't Hooft loop can therefore be written as

$$V(C) = \prod_{x_i} D_\Delta(x_i). \quad (45)$$

The dipole operator $D(x_i)$ is an eigenoperator of the magnetic flux defined on a surface that crosses the segment $[x_i - \Delta, x_i + \Delta]$. Suppose the magnetic symmetry is broken. Then we expect the dipole operator to have a nonvanishing expectation value⁹ $\langle D \rangle = d(\Delta)$. If there are no massless excitations in the theory, the operators $D(x_i)$ and $D(x_j)$ should be decorrelated if the distance $x_i - x_j$ is greater than the correlation length l . Because of Eq. (45), the expectation value of the 't Hooft loop should therefore roughly behave as

$$\langle V(C) \rangle = d(l)^{L/l} = \exp \left\{ -\ln \left(\frac{1}{d(l)} \right) \frac{L}{l} \right\} \quad (46)$$

where L is the perimeter of the loop. In the system of finite length L_x , the vacuum expectation value of the vortex line which winds around the system in the x direction is therefore finite as in Eq. (46) with $L \rightarrow L_x$.

On the other hand in the unbroken phase the VEV of the dipole operator depends on the size of the system in the perpendicular plane L_y . For large L_y it must vanish exponentially as $d = \exp\{-aL_y\}$. So the expectation value of V behaves at finite L_y in the unbroken phase as

$$\langle V(C) \rangle = \exp\{-aL_y L_x\} \quad (47)$$

and vanishes as $L_y \rightarrow \infty$. Thus in a system which is finite in x direction, but infinite in y direction, the 't Hooft line in the x direction has a finite VEV in the broken phase and vanishing VEV in the unbroken phase.

In the limit of the infinite system size $L_x \rightarrow \infty$ the VEV obviously vanishes in both phases. This is of course due to

⁹The magnetic dipole operators defined above are strictly speaking not local, since they carry the long range magnetic field of a dipole. However, the dipole field falls off with distance very fast. Therefore even though this falloff is not exponential the slight non-locality of D should not affect the following qualitative discussion.

the fact that V is a product of infinite number of dipole operators, and this product vanishes even if individual dipole operators have finite VEV.¹⁰ However, one can avoid any reference to a finite size system and infinite vortex lines by considering closed 't Hooft loops.

For a closed loop with long sides along x axis at $y=0$ and $y=R$ the above argument leads to the conclusion that in the broken phase V must have a perimeter law, Eq. (46). In the unbroken phase the correlation between the dipoles at $y=0$ and dipoles at $y=R$ should decay exponentially $\langle D(0)D(R) \rangle \propto \exp\{-\alpha R/l\}$ and thus

$$\langle V(C) \rangle = \exp\left\{-\alpha \frac{LR}{l^2}\right\} = \exp\left\{-\alpha \frac{S}{l^2}\right\}. \quad (48)$$

Thus the perimeter behavior of $\langle V(C) \rangle$ indicates a vacuum state which breaks spontaneously the magnetic Z_N while the area behavior means that the magnetic Z_N is unbroken. The results of [1] then mean that in 3+1 dimensions as well as in 2+1 dimensions the magnetic symmetry is restored above the deconfining phase transition, in the sense of Eq. (47).

In the next section we discuss what is the implication of this conclusion on the behavior of the spatial Wilson loop.

IV. SPATIAL WILSON LOOP AT LOW AND HIGH TEMPERATURE

As we have shown in Sec. II the spatial Wilson loop is the generator of the magnetic Z_N symmetry. We expect therefore that the mode of realization of the magnetic Z_N is strongly linked to the behavior of W . The argument is simplest to state for a toy model which exemplifies the basic physics in a very simple setting.

Rather than talk about non-Abelian gauge theory, consider a scalar theory of a complex field ϕ with global Z_N theory in 2+1 dimensions:

$$\mathcal{L} = \partial_\mu \phi \partial_\mu \phi^* + \lambda (\phi^* \phi - \mu^2)^2 + \zeta (\phi^N + (\phi^*)^N). \quad (49)$$

The generator of the Z_N symmetry is given by

$$U = \exp\left\{i \frac{2\pi}{N} \int d^2x j_0(x)\right\} = \exp\left\{\frac{2\pi}{N} \int d^2x (\pi \phi - \pi^* \phi^*)\right\} \quad (50)$$

where $\pi = \partial_0 \phi^*$ is the momentum conjugate to the field ϕ . Obviously with the canonical commutation relations between π and ϕ one has

$$U \phi(x) U^\dagger = e^{i2\pi/N} \phi(x). \quad (51)$$

We will be interested in the behavior of the operator which generates the Z_N transformation only inside some region S of the two dimensional plane:

$$U(S) = \exp\left\{\frac{2\pi}{N} \int_S d^2x (\pi \phi - \pi^* \phi^*)\right\} \quad (52)$$

$$U(S) \phi(x) U^\dagger(S) = e^{i2\pi/N} \phi(x) \quad (x \in S) \\ = \phi(x) \quad (x \notin S).$$

We will refer to this operator as the U loop. Throughout this discussion we assume that there are no massless excitations in the spectrum of the theory and that the linear dimensions of the area S are much larger than the correlation length.

The statement we are aiming at is that at zero temperature in the phase with broken Z_N the U loop has an area law behavior while in the phase with unbroken Z_N this changes into the perimeter law behavior.

A. U loop in the broken phase

Consider the broken phase first. We are interested in the vacuum expectation value of $U(S)$. This is nothing but the overlap of the vacuum state $\langle 0|$ and the state which is obtained by acting with $U(S)$ on the vacuum state $|S\rangle = U|0\rangle$. If the symmetry is broken, the field ϕ in the vacuum state is pointing in some fixed direction in the internal space. In the state $|S\rangle$ on the other hand its direction in the internal space is different—rotated by $2\pi/N$ —at points inside the area S . In the local theory with finite correlation length the overlap between the two states approximately factorizes into the product of the overlaps taken over the region of space of linear dimension of order of the correlation length l :

$$\langle 0|S\rangle = \prod_x \langle 0_x|S_x\rangle \quad (53)$$

where the label x is the coordinate of the point in the center of a given small region of space. For x outside the area S the two states $|0_x\rangle$ and $|S_x\rangle$ are identical and therefore the overlap is unity. However, for x inside S the states are different and the overlap is therefore some number $e^{-\gamma}$ smaller than unity. The number of such regions inside the area is obviously of order S/l^2 and we thus

$$\langle U(S) \rangle = \exp\left\{-\gamma \frac{S}{l^2}\right\}. \quad (54)$$

In a weakly coupled theory this argument is confirmed by explicit calculation. The expectation value of the U loop in the theory, Eq. (49), is given by the following path integral:

$$\langle U(C) \rangle = \int d\phi d\phi^* \exp\left\{-\int (\partial_\mu \phi + i\phi \chi_\mu)(\partial_\mu \phi^* - i\phi^* \chi_\mu) + \lambda (\phi^* \phi - \mu^2)^2 + \zeta (\phi^N + (\phi^*)^N)\right\} \quad (55)$$

with

$$\chi_\mu(x) = \frac{2\pi}{N} \delta^{\mu 0} \delta(x_0) \quad (x \in S) \quad (56)$$

¹⁰The VEV of the dipole D must be smaller than one since D is defined as a unitary operator.

$$=0, \quad (x \notin S). \quad (57)$$

This expression directly follows from Eq. (52) and integration over the canonical momentum in the phase space path integral. At weak coupling this path integral is dominated by a simple classical configuration. First, it is clear that the solution must be such that the phase of the field ϕ has a discontinuity of $2\pi/N$ when crossing the surface S since otherwise the action is UV divergent due to singular χ . Asymptotically at large distance from the surface the field should approach its vacuum expectation value. Since the source term χ vanishes outside S , everywhere where ϕ is continuous it has to solve classical equations of motion. Also, for values of x_1 and x_2 which are well inside S the profile ϕ should not depend on these coordinates, but should only depend on x_0 . It is easy to see that a solution with these properties exists: it is given by the “broken” domain wall solution. Recall that the vacuum is degenerate and so there certainly exists a classical solution of the equations of motion which interpolates between two adjacent vacuum states $\phi \rightarrow_{x_0 \rightarrow \infty} \phi_0$ and $\phi \rightarrow_{x_0 \rightarrow -\infty} e^{i2\pi/N} \phi_0$. Breaking this classical solution along the plane $x_0=0$ and rotating the piece $x_0 < 0$ by $2\pi/N$ produces precisely the configuration with the correct boundary conditions and the discontinuity structure. The path integral in Eq. (55) is therefore dominated by this classical configuration. Its action (up to corrections associated with the boundary effects of S) is αS where α is the classical wall tension of the domain wall which separates two adjacent Z_N vacua. Thus we find that the expectation value of the U loop is related to the domain wall tension of the Z_N domain wall by

$$\langle U(S) \rangle = \exp\{-\alpha S\}. \quad (58)$$

B. U loop in the unbroken phase

Now consider the unbroken phase. Again the U loop average has the form of the overlap of two states which factorize as in Eq. (53). Now however all observables noninvariant under Z_N vanish in the vacuum. The action of the symmetry generator does not affect the state $|0\rangle$. The state $|S\rangle$ is therefore locally exactly the same as the state $|0\rangle$ except along the boundary of the area S . Therefore the only regions of space which contribute to the overlap are those which lay within one correlation length from the boundary. Thus

$$\langle U(S) \rangle = \exp\{-\gamma P(S)\} \quad (59)$$

where $P(S)$ is the perimeter of the boundary of S . The absence of the area law is again easily verified by a perturbative calculation. In the unbroken phase the fluctuations of the field ϕ as well as the current density $j_0 = i(\pi\phi - \pi^*\phi^*)$ are small. To leading order in the coupling constant

$$\langle U(S) \rangle = \exp\left\{-\frac{1}{2} \int_{x,y \in S} d^2x d^2y \langle j_0(x) j_0(y) \rangle\right\}. \quad (60)$$

The possible area law contribution in the exponent is

$$S \int d^2x \langle j_0(0) j_0(x) \rangle = S \lim_{p \rightarrow 0} G(p) \quad (61)$$

where $G(p)$ is the Fourier transform of the charge density correlation function. The correlator of the charge densities however vanishes at zero momentum. This is because in the leading perturbative order the symmetry of the theory is actually $U(1)$ and not just Z_N as seen in Eq. (55). Since the vacuum state is invariant, it follows that the total charge $Q = \int d^2x j_0(x) = j_0(p=0)$ on this state vanishes, and so does any correlation function that involves zero momentum component of the charge density. So the area contribution in Eq. (61) is zero. Strictly speaking in the leading order in perturbation theory, Eq. (60) is not the complete result. The exact expression contains in the exponential also higher point correlators of the current density. Again however the possible area law contribution contains correlators of the total charge Q with powers of j_0 and therefore vanishes.

C. U loop at high temperature

Let us see now how the argument changes at high temperature. The important difference is that the vacuum is not a pure state but rather a statistical ensemble. The average of the U loop is therefore not given by a single matrix element but rather by

$$\langle U \rangle = \sum_i e^{-E_i T} \langle i | U | i \rangle. \quad (62)$$

Let us consider the theory in which the Z_N symmetry is broken at zero temperature. For concreteness we will think about Z_2 symmetric theory, although qualitatively the discussion does not change for any N . The two degenerate vacuum states are characterized by the value of the condensate $\langle \phi \rangle = \pm \mu$.

In order to understand the behavior of the U loop we have to figure out what types of states contribute to the thermal ensemble. At zero temperature the only states that are of interest are those with finite energy. There are two towers of such states $|n\rangle_\mu$ and $|n\rangle_{-\mu}$ —constructed above each one of the degenerate vacua. These two towers of states do not talk to each other, not only because their overlap is zero, but also because they cannot be connected to each other by action of any local (or semilocal) operator ${}_\mu \langle n | O | n' \rangle_{-\mu} = 0$. An immediate corollary of this is that a superposition of the type $|\alpha, \beta\rangle = \alpha |n\rangle_\mu + \beta |n\rangle_{-\mu}$ violates clustering property of the correlators of local operators. For this reason at zero temperature in a spontaneously broken theory we are never interested in states which carry sharp quantum numbers of the broken symmetry.

At finite temperature however we are also asking after states with finite energy density and, therefore, infinite energy. This part of the spectrum looks rather different if the energy density involved is high enough. The two vacuum configurations of the potential in Eq. (49) are separated by a finite barrier. Let us call the height of this barrier H . The states with energy density lower than H still separate into two towers. We will denote these states by $|I\rangle$. However,

higher energy density states, with $\epsilon \gg H$, have a different nature. Their wave function is not localized in the field space to the vicinity of one of the vacua, but rather is spread over distances larger than the distance between the two vacuum states, 2μ . These states therefore naturally carry sharp quantum numbers with respect to the broken Z_2 symmetry. These states we will denote by $|h\rangle$. In fact one expects that the higher the energy density the more these states look like the multiparticle states of a symmetric phase. That is to say as long as $\epsilon \gg H$ it does not matter whether the potential has the double well structure or a single vacuum. These highly excited states should look like states with finite density of particles which carry the Z_2 charge.

At low temperatures, when the entropy effects are not important the contribution to the thermal ensemble comes only from the $|l\rangle$ sector since the Boltzmann factor for any of the $|h\rangle$ states vanishes exponentially in the infinite volume limit. As we have argued earlier, the average of U in each one of these states has an area law behavior and so does the whole temperature average of U .

When the temperature reaches T_c the phase transition occurs. The reason for the onset of the phase transition is that when the equilibrium energy density reaches critical threshold value, the $|h\rangle$ sector states start contributing to the thermal ensemble. The sudden change in the entropy due to these new channels drives the phase transition. Above the phase transition therefore there are two kinds of states that contribute to thermal averages. One can then write

$$\langle U \rangle_{T>T_c} = \sum_n e^{-E_n^l/T} \langle n, l | U | n, l \rangle + \sum_h e^{-E_n^h/T} \langle n, h | U | n, h \rangle \quad (63)$$

where n stands for all other quantum numbers. In fact once the entropy effects become important enough to excite the $|h\rangle$ sector, the contribution of $|l\rangle$ states to any physical observable becomes negligible. As discussed earlier each state in the second term gives a perimeter contribution $\exp\{\gamma_s P\}$ to the average of the U loop, so one could be tempted to conclude that the loop must have a perimeter law just like in zero temperature vacuum of a symmetric phase. This however is not necessarily the case. The reason is that $\langle n, h | U | n, h \rangle$ is not positive definite. In fact the number of states in which it is positive is roughly equal to the number of states on which it is negative. It is therefore very likely that the leading perimeter behavior will cancel and the net result will again be an area law for $\langle U \rangle$.

Indeed, if the ensemble can be thought of as an ensemble of Z_2 charged free particles, the area law for the U loop follows immediately.¹¹ The U loop in such an ensemble is

$$\langle U(S) \rangle = \sum_{x_i, n} \frac{1}{n!} \mu^n (-1)^n \quad (64)$$

where μ is the fugacity of a single particle, and the summation goes over all coordinates x_i of the particles inside the area S and over all possible numbers of particles n . Assuming that particles have a finite size Δ , so that there are S/Δ possibilities to place one particle inside the area S in the dilute gas approximation, the sum gives

$$\langle U(S) \rangle = \exp \left\{ -\frac{\mu}{\Delta} S \right\}. \quad (65)$$

We stress that the thermal ensemble of particles is Z_2 invariant. The density matrix for such an ensemble can be written in the particle basis as

$$\rho = \sum_{n(x)} \frac{1}{n!} \mu^n |n\rangle \langle n|. \quad (66)$$

The operator of the Z_2 transformation acts on the n -particle states as

$$U|n\rangle = (-1)^n |n\rangle \quad (67)$$

and so

$$U \rho U^\dagger = \rho. \quad (68)$$

The explicit simple formula (65) is derived in the dilute gas approximation. We expect that the physics will be similar as long as the interaction between the particles is short range. Whenever the interaction is long range the behavior of $\langle U(S) \rangle$ can be different. For instance one does not expect area law behavior if the particles are bound into pairs since in this case only Z_2 invariant states contribute to the thermal ensemble.

Our conclusion is that at finite temperature the behavior of the U loop is not strongly related to the mode of the realization of the Z_N symmetry. It is rather more likely to have an area behavior.

To reiterate, the physics involved is very simple. At zero temperature when acting on a state, the U loop performs the Z_N transformation inside the loop. The only degrees of freedom that are changed by this operation inside the loop are the Z_N -noninvariant fields. If the vacuum wave function depends on the configuration of the noninvariant degrees of freedom (the state in question is not Z_N invariant), the action of the U loop affects the state everywhere inside the loop. The VEV of the U loop then falls off as an area. If the vacuum is Z_N invariant, the wave function does not depend on the configuration of the noninvariant degrees of freedom. The action of the U loop then perturbs the state only along the perimeter, hence the perimeter law in the unbroken phase.

At finite temperature however the thermal ensemble may even in the symmetric phase contain significant contributions from states with nonvanishing Z_N charges. The U loop therefore perturbs the thermal ensemble very significantly everywhere inside the area, and the natural outcome is an area law.

The argument is quite general and does not depend on the exact form of the Z_N invariant potential and more generally on the field content of the theory—we could have added any

¹¹This argument is borrowed from [16]. We thank Mike Teper and Biaggio Lucini for discussions of this point.

number of extra fields to the theory, Eq. (49), without changing the conclusions. The same relation must exist between the mode of realization of the magnetic Z_N symmetry and the behavior of the Wilson loop in the pure Yang-Mills theory. The direct analogues of the scalar field $\phi(x)$ in Eq. (49) and the U loop of the scalar theory are correspondingly the vortex field $V(x)$ and the spatial Wilson loop $W(C)$.

As we have shown in the previous section, the magnetic Z_N is restored at high temperature. The Wilson loop is nevertheless likely to have an area law as is indeed indicated by all existing lattice data. In this context we note that analytic strong coupling results also give area behavior [17].

D. Wilson loop in 3+1 dimensions

The previous considerations generalize to 3+1 dimensions. At zero temperature in the broken phase when acting with the Wilson loop $W(C)$ on the vacuum one changes the state of those magnetic vortices which loop through C . The number of such vortices which are present in a generic configuration in the broken phase is proportional to the minimal area subtending C . The number of degrees of freedom that is changed by the action of W is thus proportional to the area S . Each of these degrees of freedom contributes a factor smaller than unity to the overlap with the vacuum state and so the VEV of W scales with the exponential of the area. In the unbroken phase the vacuum does not contain vortices of arbitrarily large size. The size of the vortices present in the vacuum is cut off by the relevant correlation length. This is the case if the gauge group is completely broken by the Higgs mechanism. Therefore, for contours C of linear dimension much larger than this length, the action of $W(C)$ only disturbs degrees of freedom close to the contour C itself and the VEV must have the perimeter behavior.

At high temperature even the symmetric thermal ensemble is populated by vortices. These vortices are not free since apart from them the ensemble also contains “free” charges. However, unless this background of charges induces long range interactions between the vortices, the most probable result for the Wilson loop is the area law. A more detailed knowledge of vortex dynamics is necessary to draw a firm conclusion.

To close this section we note that the present considerations do not apply to Abelian theories. The magnetic symmetry does exist in this case too, but here it is the continuous $U(1)$ group and the spectrum is massless. In this case there is no reason to expect the local factorization of the overlap and generically therefore the arguments of this section do not hold. In particular in the presence of long range correlations it is perfectly possible that the Wilson loop has a perimeter law even though the state is perturbed everywhere inside the area bounded by the loop.¹²

¹²In 2+1 dimensions it is actually only the noncompact Abelian theories that are excluded from the consideration. Compact theories are massive and therefore should behave in the same way as the non-Abelian Yang-Mills theory.

V. DISCUSSION

In this paper our aim was to point out two facts. First that the calculation of the VEV of the ’t Hooft loop [1] implies the restoration of the magnetic Z_N symmetry above the deconfinement transition. Second, that the mode of realization of magnetic symmetry is closely related to the behavior of the spatial Wilson loop.

At zero temperature this relation is very rigid: spontaneous breaking of Z_N implies the area law behavior for W while unbroken Z_N leads to perimeter law behavior. At high temperature however even though the Z_N symmetry is restored, the Wilson loop may have an area law. This is the consequence of the fact that even a Z_N -invariant thermal ensemble can contain a significant contribution of Z_N non-singlet states. The area law is particularly simply understood if the thermal ensemble at high temperature is well approximated by an ensemble of weakly interacting magnetic vortices. The vortex gas argument has been previously brought up in favor of the area behavior of the spatial Wilson loop in [18]. This behavior is also confirmed by several lattice gauge theory calculations [19].

An alternative possibility is that due to the as yet unknown vortex dynamics, there is vortex-antivortex binding. To explore such a possibility it would be very interesting to measure on the lattice the free energy of a magnetic vortex. In Ref. [15] the behavior of the free energy of magnetic and electric fluxes has been discussed in the *low* temperature phase. To be able to do it in the lattice framework one has to define the theory in a finite volume. As discussed by ’t Hooft this can be achieved by imposing on the potentials periodic boundary conditions modulo a gauge transformation. As discussed in Ref. [15] this admits the presence of vortices in 2+1 and of the vortex lines in 3+1 dimensions. ’t Hooft’s discussion was based on a Euclidean rotation identity for the twisted 4D path integrals valid for any temperature, and a factorization property of magnetic and electric fluxes. In the notation of Ref. [15],

$$F(\vec{e}, \vec{m}) = F_e(\vec{e}) + F_m(\vec{m}). \quad (69)$$

Its validity at low T is very reasonable, but is inconsistent with the Euclidean rotation identity at high T . Based on this ’t Hooft could prove ($N \leq 3$) that in the confining phase, where the free energy of an electric flux is linear with the length (with the string tension ρ), the free energy of magnetic flux vanishes exponentially in the infinite volume limit. For a magnetic flux in, say the z direction it is $\exp(-\rho L_x L_y)$. Thus the free energy of a magnetic flux is related to the behavior of the Wilson loop. The free energy of an electric flux in the z direction in the *hot* phase vanishes exponentially like $\exp(-\alpha L_x L_y)$ where α is the surface tension found in Ref. [1]. So the next obvious question is how the magnetic flux free energy behaves in the hot phase. In a vortex gas picture this free energy should vanish exponentially in the infinite volume limit. Such a calculation in 2+1 dimensions has been performed and (modulo some uncertainty related to imperfect measurement of global vorticity) results are consistent with this expectation [16]. It will be

instructive to see how in the hot phase the additivity of electric and magnetic fluxes is broken.

We note that a recent lattice calculation [20] measures the monopole-antimonopole correlation. The results of [20] point to the screened behavior of this correlation function for *all* temperatures. So in the hot phase it behaves like its electric partner, the correlator of Polyakov loops.¹³ This, via 't Hooft's argument, is consistent with the measured area behavior of Wilson loops [19] and would imply that the magnetic flux energy would fall off with an area law for *all* temperatures.

It is interesting to note that the vortex gas picture is equally applicable in high temperature confining and non-confining gauge theory. In particular one can consider an $SU(N)$ gauge theory with sufficient number of adjoint Higgs

fields, so that the gauge theory at zero temperature is broken completely. In this situation the magnetic Z_N symmetry is unbroken in the vacuum and the Wilson loop has a perimeter law. Magnetic vortices are finite energy excitations with the mass of order $M = M_v^2/g^2$, where M_v is the vector boson mass. When the system is heated one expects that the thermal ensemble will contain a dilute gas of these vortices at any temperature. Therefore at any finite temperature the spatial string tension should be nonzero, although at low temperatures it will be exponentially suppressed if the theory is weakly coupled: $\sigma \propto M_v^2 \exp\{-M_v^2/g^2 T\}$.

ACKNOWLEDGMENTS

The work of A.K. is supported by PPARC. The work of C.P.K.A. and A.K. is supported by a joint CNRS-Royal Society project. We are indebted to Frithjof Karsch for bringing Ref. [17] to our attention. We thank Ian Kogan, Biaggio Lucini and Mike Teper for very interesting and useful discussions, which in particular helped to clarify a fundamental flaw in the arguments in the first version of this paper.

¹³We note however that this simulation [20] also points to the Coulomb behavior for the spatial 't Hooft loop in the hot phase, in contradiction to analytic results [1,21] and early lattice results [22]. We feel that here more work should be done to clarify the situation.

-
- [1] C.P. Korthals Altes, A. Kovner, and M. Stephanov, Phys. Lett. B **469**, 205 (1999).
 - [2] G. 't Hooft, Nucl. Phys. **B138**, 1 (1978).
 - [3] O. Aharony, S.S. Gubser, J. Maldacena, H. Ooguri, and Y. Oz, Phys. Rep. **323**, 183 (2000).
 - [4] S. Samuel, Nucl. Phys. **B154**, 62 (1979); N.J. Snyderman, *ibid.* **B218**, 381 (1983).
 - [5] A. Kovner and B. Rosenstein, Int. J. Mod. Phys. A **7**, 7419 (1992).
 - [6] A. Kovner, B. Rosenstein, and D. Eliezer, Mod. Phys. Lett. A **5**, 2661 (1990); Nucl. Phys. **B350**, 325 (1991); A. Kovner and B. Rosenstein, Phys. Lett. B **266**, 443 (1991).
 - [7] A.M. Polyakov, Nucl. Phys. **B120**, 429 (1977).
 - [8] E. Fradkin and S.H. Shenker, Phys. Rev. D **19**, 3682 (1979).
 - [9] D. Diakonov and V. Petrov, Phys. Lett. B **224**, 131 (1989); hep-th/9606104.
 - [10] S. Bronoff and C.P. Korthals Altes, Phys. Lett. B **448**, 85 (1999).
 - [11] A. Kovner and B. Rosenstein, J. High Energy Phys. **03**, 9809 (1998).
 - [12] C. P. Korthals Altes, A. Michels, M. Stephanov, and M. Teper, Phys. Rev. D **55**, 1047 (1997).
 - [13] C.P. Korthals Altes, Nucl. Phys. **B420**, 637 (1994).
 - [14] N. Weiss, Phys. Rev. D **24**, 475 (1981); **25**, 2667 (1982); D. Gross, R.D. Pisarski, and L.G. Yaffe, Rev. Mod. Phys. **53**, 497 (1981).
 - [15] G. 't Hooft, Nucl. Phys. **B153**, 141 (1979).
 - [16] A. Hart, B. Lucini, Z. Schram, and M. Teper, J. High Energy Phys. **06**, 040 (2000).
 - [17] C. Borgs, Nucl. Phys. **B261**, 455 (1985).
 - [18] R. Bertle, M. Faber, J. Greensite, and S. Olejnik, J. High Energy Phys. **03**, 019 (1999); K. Langfeld, O. Tennert, M. Engelhardt, and H. Reinhard, Phys. Lett. B **452**, 301 (1999); M.N. Chernodub, M.I. Polikarpov, A.I. Veselov, and M.A. Zubkov, Nucl. Phys. B (Proc. Suppl.) **73**, 575 (1999).
 - [19] F. Karsch, E. Laermann, and M. Lutgemeier, Phys. Lett. B **346**, 94 (1995).
 - [20] Ch. Hoebbling, C. Rebbi, and V.A. Rubakov, Nucl. Phys. B (Proc. Suppl.) **83-84**, 485 (2000); hep-lat/0003010.
 - [21] T. Bhattacharya, A. Gocksch, C.P. Korthals Altes, and R.D. Pisarski, Phys. Rev. Lett. **66**, 998 (1991); Nucl. Phys. **B383**, 497 (1992).
 - [22] K. Kajantie and L. Kärkkäinen, Phys. Lett. B **214**, 595 (1988); K. Kajantie, L. Kärkkäinen, and K. Rummukainen, Nucl. Phys. **B333**, 100 (1990); **B357**, 693 (1991); Phys. Lett. B **286**, 125 (1992); S. Huang, J. Potvin, C. Rebbi, and S. Sanieleveci, Phys. Rev. D **42**, 2864 (1990); **43**, 2056(E) (1991); R. Brower, S. Huang, J. Potvin, and C. Rebbi, *ibid.* **46**, 2703 (1992); R. Brower, S. Huang, J. Potvin, C. Rebbi, and J. Ross, *ibid.* **46**, 4736 (1992).