

Low-scale seesaw mechanisms for light neutrinos

Francesca Borzumati

Theory Group, KEK, Tsukuba, Ibaraki 305-0801, Japan

and Scuola Internazionale Superiore di Studi Avanzati (SISSA), I-34013 Trieste, Italy

Yasunori Nomura

Department of Physics, University of Tokyo, Tokyo 113-0033, Japan

(Received 12 September 2000; published 25 July 2001)

Alternatives to the seesaw mechanism are explored in supersymmetric models with three right-handed or sterile neutrinos. Tree-level Yukawa couplings can be drastically suppressed in a natural way to give sub-eV Dirac neutrino masses. If, in addition, a $B-L$ gauge symmetry broken at a large scale M_G is introduced, a wider range of possibilities opens up. The value of the right-handed neutrino mass M_R can be easily disentangled from that of M_G . Dirac and Majorana neutrino masses at the eV scale can be generated radiatively through the exchange of sneutrinos and neutralinos. Dirac masses m_D owe their smallness to the pattern of light-heavy scales in the neutralino mass matrix. The smallness of the Majorana masses m_L is linked to a similar seesaw pattern in the sneutrino mass matrix. Two distinct scenarios emerge. In the first, with very small or vanishing M_R , the physical neutrino eigenstates are, for each generation, either two light Majorana states with a mixing angle ranging from very small to maximal, depending on the ratio m_D/M_R , or one light Dirac state. In the second scenario, with a large value of M_R , the physical eigenstates are two nearly unmixed Majorana states with masses $\sim m_L$ and $\sim M_R$. In both cases, the $(B-L)$ -breaking scale M_G is, in general, much smaller than that in the traditional seesaw mechanism.

DOI: 10.1103/PhysRevD.64.053005

PACS number(s): 14.60.Pq, 12.60.Jv

I. INTRODUCTION

It has long been known that neutrinos are very light. Whether they are only light or completely massless, however, is a question which has remained unanswered up until recently.

No direct measurement of neutrino masses exists as yet. Nevertheless, overwhelming evidence has been gathering in the past years of the fact that neutrinos of different generations oscillate one into the other [1]. Oscillations have been observed by atmospheric [3], solar [4], and accelerator [5] neutrino experiments. Quantum mechanics relates oscillations among neutrinos of different flavor, ν_e , ν_μ , ν_τ , to nonvanishing values of mass for the neutrino mass eigenstates, ν_1 , ν_2 , and ν_3 , which are linear combinations of ν_e , ν_μ , and ν_τ . The profusion of experimental results on neutrino oscillations turns, therefore, into a strong indication that neutrinos have mass.

This fact has profound consequences for particle physics, astrophysics, and cosmology. Having mass, neutrinos may be a crucial ingredient of the hot dark matter component that contributes to the mass density of the Universe [6]. Moreover, if their mass does not exceed the eV range, they may have played an important role in the formation of the observed structure of galaxies [6]. Far from being a tested result, this value is a constraint to be kept into account by model building, more stringent than the direct bounds from collider experiments: $m_{\nu_e} \lesssim 5$ eV, $m_{\nu_\mu} \lesssim 170$ keV, and $m_{\nu_\tau} \lesssim 18$ MeV [2]. It is also possible, however, that some or all neutrinos are even lighter than a few eV. Indeed, oscillation experiments give only an indication of the splitting of neutrino masses squared, but leave the absolute size of neutrino

masses still open to speculation. They are not yet conclusive even on the number of neutrino species undergoing oscillations.

The experimental situation can be schematically summarized as follows. Atmospheric neutrino experiments from Super-Kamiokande [3,7] indicate that muon neutrinos ν_μ 's oscillate into different neutrinos, ν_x 's. The mass-squared splitting of the two mass eigenstate neutrinos (whose main components are ν_μ and ν_x) is $10^{-3} \text{ eV}^2 < \Delta m^2 < 10^{-2} \text{ eV}^2$. The neutrino ν_x seems to be predominantly a tau neutrino ν_τ [7]. Results from the CHOOZ reactor experiment [8] strongly suggest that ν_x is unlikely to be the electron neutrino ν_e , although this possibility cannot be excluded by the Super-Kamiokande Collaboration [7]. Solar neutrino experiments [4] indicate that ν_e 's oscillate into some other type of neutrinos ν_y 's. If the MSW mechanism [9] is used to explain the deficit in the solar neutrino flux, the relevant mass-squared splitting is $\Delta m^2 \sim 10^{-5} \text{ eV}^2$ for the large and small mixing angle solution, or $\Delta m^2 \sim 10^{-7} \text{ eV}^2$ for the LOW solution [10]. It is $\Delta m^2 \sim 10^{-10} \text{ eV}^2$ if oscillations in the vacuum between Sun and Earth [11] are invoked. Finally, the Liquid Scintillator Neutrino Detector (LSND) experiment [5] seems to support the hypothesis of a $\nu_\mu \rightarrow \nu_e$ oscillation with corresponding mass-squared splitting $\Delta m^2 \sim 1 \text{ eV}^2$.

Barring experimental systematic errors which may lead to inconsistent results, these data cannot be explained by assuming the existence of only three light neutrinos and that each of these oscillations is described by a single Δm^2 [1]. One simple way out is to assume that there exist additional neutrinos, also light. Accurate measurements of the invisible width of the Z boson at the CERN e^+e^- collider LEP have unequivocally shown that there are only three light neutrinos coupling to the Z boson [12]. Other possible light states,

therefore, must have as main constituent neutrinos invisible for the Z boson, i.e., right-handed neutrinos. When discussing neutrino oscillations, these are often called sterile neutrinos, in contrast to the left-handed components of $SU(2)$ doublets, or active neutrinos. If sterile neutrinos exist, the results of solar neutrino experiments may be viewed in terms of oscillations between electron and sterile neutrinos. Up until recently, these were preferably accommodated in a four-neutrino spectrum of the “ $2+2$ ” variety. This spectrum consists of two pairs of neutrinos separated by the LSND Δm^2 , with the component of each pair split, respectively, by the atmospheric and solar Δm^2 . Recent measurements of the solar neutrino flux by the Super-Kamiokande Collaboration, however, seem to disfavor the possibility of ν_γ 's being sterile neutrinos at the 95% C.L. [13]. (A different conclusion is reached in [14].) This fact, together with a lower oscillation probability recently claimed by LSND [15], resurrects the “ $3+1$ ” neutrino spectrum [16] previously excluded by the incompatibility between LSND and negative searches for ν_e and ν_μ disappearance. In this spectrum, one mainly-sterile neutrino is separated by the LSND mass gap from the other three, which are mainly active, roughly degenerate, and sufficient to explain atmospheric and solar neutrino data. The “ $3+1$ ” spectrum assumes a small mixing between the active and sterile content of the four mass eigenstates. For a model in which this spectrum can be easily implemented, see Ref. [17].

The challenge, which particle physics is then faced with, is twofold. On one side there is the longstanding issue of the smallness of neutrino masses. Although clear experimental evidence that neutrinos are massive is a recent acquisition, there has always been a strong theoretical prejudice in this direction. The exact vanishing of neutrino masses, which is consistent with the standard-model (SM) particle content, would nevertheless demand for a deeper explanation, possibly through a symmetry principle, in an underlying theory of which the SM is an effective limit. Rejecting the possibility of vanishing neutrino masses on theoretical ground, there has been a tremendous effort to explain why these masses are much smaller than those of quarks and charged leptons.

Technically, two ways are known to obtain light neutrinos. The first follows the conventional mechanism for generating quark and charged-lepton masses. The SM is enlarged to include three right-handed neutrinos. These couple to the neutral components of the left-handed leptonic doublets through Yukawa interaction terms. After electroweak symmetry breaking, there remain three light mass eigenstates ν_1 , ν_2 , and ν_3 , which are Dirac fermions. For these neutrinos to be at the eV scale, Yukawa couplings as small as $\sim 10^{-11}$ are needed. The second known way of generating small neutrino masses is through the seesaw mechanism [18,19]. This invokes the existence of right-handed neutrinos with a very large mass $M_R \sim 10^{12}$ GeV, whereas the couplings of the Yukawa neutrino interactions are unsuppressed. The smallness of the light neutrino masses is then explained in terms of the seesaw pattern of light-heavy scales in the neutrino mass matrix. The three light mass eigenstates ν_1 , ν_2 , and ν_3 are now Majorana fermions, with mass $\sim m_{\text{weak}}^2/M_R$. The other three mass eigenstates n_1 , n_2 , and

n_3 , also of Majorana type, remain at the high scale M_R . (On a different line of thought is the recent proposal of Ref. [20], where the smallness of neutrino masses is explained by embedding the SM and right-handed neutrinos in a higher-dimensional spacetime. In a scenario of Dirac neutrino masses, for example, the Dirac Yukawa couplings between left-handed neutrinos localized on the SM three-brane and the right-handed ones, residing in the bulk, are suppressed by the volume factor of compactified dimensions.)

The other challenge for theoretical particle physics has been unexpectedly kindled by recent experimental results. The interpretation of data coming from solar neutrino experiments in terms of oscillations between active and sterile neutrinos requires extensions of the SM that explain the existence of light mass eigenstates, other than ν_1 , ν_2 , ν_3 , mainly corresponding to additional $SU(2) \times U(1)$ singlets. It is possible that the experimental situation will evolve in the future in such a way that (i) the preliminary findings of the Super-Kamiokande Collaboration on solar neutrino oscillations disfavoring the sterile neutrino option are confirmed, and (ii) the LSND results are clearly excluded. The SNO experiment [21] will soon be able to resolve the issue as to whether the electron neutrinos produced in the Sun oscillate to active or sterile neutrinos. In the meantime, the Super-Kamiokande Collaboration will keep collecting more data on solar neutrino oscillations that may possibly strengthen their preliminary results. Moreover, there is a number of already operating [22] and planned [23] accelerator experiments to seek for oscillations in the LSND region. If no signal is found in these searches, the two Δm^2 of solar and atmospheric neutrino oscillations can be easily accommodated in a scenario of three light neutrinos. This, of course, does not mean that sterile neutrinos do not exist. It may only mean that low-energy physics is blind to them, either because they are heavy or because, even if light, they are very weakly mixed with the active ones. Ideal detectors may turn out to be astronomical ones, such as supernovas. Sterile neutrinos with a mass of a few keV (or tens of keV) have been proposed to explain the motion of pulsars born in supernova explosions [24]. Light sterile neutrinos, with mass above the eV scale, may also be of interest in explaining supernova nucleosynthesis and big bang nucleosynthesis [25]. If in the mass range of 0.1 keV, they may also give rise to “cool” dark matter, as recently discussed in [26].

The natural setting for incorporating sterile neutrinos is that in which six final Majorana states are seesaw generated. Nevertheless, no satisfactory theoretical model has been found so far to induce the suppression factor $\sim m_{\text{weak}}^2/M_R$ needed to have light ν_1 , ν_2 , and ν_3 states, while simultaneously keeping very low the scale for some or all of the right-handed neutrinos, with resulting light states among n_1 , n_2 , and n_3 . (Infinitely many light sterile neutrinos, however, are obtained in extra-dimensional mechanisms for generating light neutrino masses [20].) In a typical model, the seesaw right-handed neutrinos may participate in an additional gauge interaction spontaneously broken at some scale M_G . In the simplest known model, free from gauge anomalies and mixed gauge and gravitational anomalies, this new interaction is $U(1)_{B-L}$ and the SM particle content is mini-

mally enlarged by a neutral gauge boson, three additional electroweak singlets, and a neutral $(B-L)$ -breaking Higgs boson. The mass of these new particles is closely related to the spontaneous violation of $B-L$ and all these particles are at the large scale $M_G(\sim M_R)$.

It is in general assumed that similar features hold also in the supersymmetric version of such a model. In this paper we explore other possibilities that naturally present themselves in supersymmetric models, in addition to the two outlined above, of generating small neutrino masses. Two mechanisms are described. In both, lowest-order Yukawa operators are forbidden by imposing some horizontal symmetries. Small neutrino masses can then be generated at the tree level, through higher-dimensional nonrenormalizable operators, in the first mechanism, or radiatively, through the virtual exchange of superpartners, in the second one. The first mechanism, described in Sec. II, does not require an additional $U(1)_{B-L}$ gauge interaction. The gauging of $U(1)_{B-L}$ is not needed either for the radiative generation of Majorana masses for active neutrinos, but it is, on the contrary, necessary to obtain radiative Dirac neutrino masses. For the sake of a simpler and more coherent discussion, however, the radiative mechanism is presented in the case of a gauged $U(1)_{B-L}$ symmetry. The modifications to the case without a $U(1)_{B-L}$ symmetry, when applicable, are obvious and will be briefly commented upon. The radiative mechanism is introduced in Sec. III and worked out in Secs. IV, V, and VI. The patterns of physical neutrino masses that can arise through both mechanisms are classified in Sec. VII. A possible embedding of these two mechanisms in full-fledged models is outlined in Sec. VIII. Up to this section, it is implicitly or explicitly assumed that the signal of supersymmetry breaking is transmitted from the hidden to the visible sector through supergravity mediation. Comments will be made in Sec. VIII on possible changes induced when this transmission is realized through gauge mediation. The paper ends with some final conclusions and remarks in Sec. IX.

II. SUPERSYMMETRIC TREE-LEVEL MECHANISM

The class of models considered in this paper¹ has three $SU(2) \times U(1)$ singlet superfields \bar{N} in addition to the three lepton doublets L , with neutral components N . The lowest-order operators giving rise to neutrino masses are

$$W = \frac{1}{4} \frac{h_\nu}{M_P} LHLH + y_\nu \bar{N}LH + \frac{1}{2} M_R \bar{N}\bar{N}, \quad (1)$$

with the scale M_R naturally given by the Planck mass M_P . All these operators can be forbidden by a horizontal discrete symmetry Z_n in conjunction with a continuous or discrete symmetry such as lepton number. As for the first symmetry,

it is sufficient to assign vanishing Z_n charges to the superfields L and H and $Z_n(\bar{N}) = +1$ to the superfields \bar{N} . This assignment forbids the lowest-order Yukawa operators, but leaves allowed higher-dimensional operators of Yukawa type:

$$W = \frac{k_\nu}{M_P} Z \bar{N} L H, \quad (2)$$

where Z is a spurion superfield with charge $Z_n(Z) = -1$. The second symmetry (different from Z_2) assigns the following charges to the lepton superfields:

$$L(+1), \quad \bar{N}(-1), \quad \bar{E}(-1), \quad (3)$$

and zero charge to all the other superfields. Therefore, it forbids both Majorana mass operators in Eq. (1) as well as the higher-dimensional operators:

$$W = \frac{1}{4} \frac{z_\nu}{M_P} Z Z \bar{N} \bar{N}. \quad (4)$$

In this scenario, the emerging picture is that of three very light neutrino states of Dirac type. The spurion field Z acquires, in general, a supersymmetry-conserving vacuum expectation value (VEV) \mathcal{A}_Z and a supersymmetry-violating one \mathcal{F}_Z . Through operators analogous to that in Eq. (2), the VEV \mathcal{A}_Z gives contributions to Yukawa couplings for quarks and charged leptons that are negligibly small, unless \mathcal{A}_Z is at some large scale $\gtrsim 10^{13}$ GeV. Acceptable Yukawa couplings may, however, be generated for neutrinos. Indeed, in a class of O'Raifeartaigh-type models discussed in Sec. VIII, \mathcal{A}_Z is vanishing in the supersymmetric limit, but is shifted to a value

$$\mathcal{A}_Z \simeq \frac{16\pi^2}{\lambda^3} m_{\text{weak}} \quad (5)$$

by supersymmetry-breaking effects, where λ is a dimensionless coupling. The induced Yukawa couplings

$$y_\nu = 16\pi^2 \frac{k_\nu}{\lambda^3} \frac{m_{\text{weak}}}{M_P} \quad (6)$$

give rise to Dirac neutrino masses of order 10^{-2} eV if λ and k_ν are of $\mathcal{O}(1)$,² i.e., in the range of the Δm^2 required by the MSW solution to the solar neutrino deficit. Smaller values of λ allow also neutrino masses in the range of atmospheric neutrino experiments. No room is left for sterile neutrinos and the observed oscillation patterns can be explained through flavor oscillations. This tree-level mechanism for generating small neutrino Yukawa couplings is rather ge-

¹Tree-level mechanisms were also advocated in [27] in the context of superstring theories, in [28] in the context of supersymmetric models with gauge mediation of supersymmetry breaking, and in [29] in the context of supergravity models.

²We thank T. Yanagida for pointing this out to us.

neric, in the sense that it can be easily incorporated in any supersymmetric extension of the SM model.

It is also possible to forbid the renormalizable operators in Eq. (1) leaving, however, allowed both operators in Eqs. (2) and (4), as well as the nonrenormalizable operator in Eq. (1). One way to achieve this is to couple the above Z_n symmetry with an R -parity symmetry under which \bar{N} is odd and Z is even. Dangerous terms of type $M_P Z \bar{N}$ are then forbidden, while Majorana mass terms $\sim M_P \bar{N} \bar{N}$ are avoided by requiring that Z_n be not Z_2 . In this case, both Dirac and Majorana masses become possible at the tree level: $\sim \mathcal{A}_Z v / M_P$ is obtained for Dirac masses; $\sim \mathcal{A}_Z^2 / M_P$ and $\sim v^2 / M_P$, for Majorana masses for right- and left-handed neutrinos, respectively, where v is the VEV of the neutral component of H . The physical neutrino states are six, of Majorana type: the three n_i states have masses $\sim \mathcal{A}_Z^2 / M_P$, the three states ν_i are lighter, i.e., of order of the Majorana mass for left-handed neutrinos, $\sim v^2 / M_P$. In the class of models in which supersymmetry breaking induces the value of \mathcal{A}_Z in Eq. (5), these masses range from the keV to the sub-eV region, depending on the value of λ and of the other dimensionless couplings z_ν , h_ν , and k_ν . Thus, light sterile neutrinos can be easily accommodated in such a scenario.

It should be noticed that the operator (4) gives rise to bilinear soft supersymmetry-breaking terms with large massive couplings, $BM_R \sim \mathcal{A}_Z \mathcal{F}_Z / M_P$. Such terms may induce VEV's for the scalar components of the superfields \bar{N} if \mathcal{F}_Z is of the same order of the largest supersymmetry-breaking VEV $\sim m_{\text{weak}} M_P$. A value of \mathcal{F}_Z low enough to avoid the spontaneous breaking of lepton number decreases also the value of the trilinear parameters $A \sim \mathcal{F}_Z / M_P$ generated by the operator (2). Nevertheless, large radiative contributions to the Majorana mass for active neutrinos arise. We discuss these contributions explicitly in the case of a gauged $U(1)_{B-L}$, when also potentially large contributions to Dirac masses are induced.

III. SUPERSYMMETRIC RADIATIVE MECHANISM

If the gauge sector is enlarged as in a typical model that leads to the seesaw mechanism, the phenomenology of the neutrino sector becomes much richer. The phenomenology of other sectors becomes also more interesting. The presence of the heavy scale M_G , typical of seesaw models, for example, is felt also in other sectors. Some neutralinos become very heavy and, depending on the details of the specific realization of the model, also some supersymmetric partners of neutrinos, or sneutrinos, can be at the same large scale of the $B-L$ violation.

A suppression factor for small neutrino masses, therefore, does not necessarily need to be induced by the right-handed neutrinos. Indeed, there is at least one and, possibly, two mass matrices (neutralino and sneutrino mass matrices) with the same seesaw pattern of light-heavy scales that was only present in the neutrino mass matrix in the nonsupersymmetric case. It is clear that the transmission to the neutrino sector of a suppression factor induced in the neutralino and/or

sneutrino sector may only happen at the quantum level.³ The advantage of such a procedure is obvious: the mechanism of mass generation for the light neutrinos can be disentangled from the scale of the sterile neutrinos, which does not need to be anymore that of the $B-L$ violation.

If a program of radiative generation of neutrino mass operators is implemented, the Yukawa couplings between active and sterile neutrinos become redundant. As in the generic supersymmetric scenario outlined above, they can be altogether forbidden. Higher-dimensional operators of Yukawa type [see Eq. (2)] remain in general allowed and provide Dirac neutrino masses naturally of order 10^{-2} eV. If larger physical masses are aimed at—for example, in the eV range—then the largest contribution to the Dirac neutrino mass can be of radiative origin if the scale of $B-L$ breaking is not too large. If this is not the case, both tree-level and radiative mechanisms complement each other, providing comparable contributions. The radiative contribution to the Majorana mass for active neutrinos is, on the contrary, always larger than the tree-level contribution arising in the second tree-level scenario described in Sec. II. The ingredients necessary for generating radiatively neutrino mass operators can be supplied by soft supersymmetry-breaking terms. The chiral-flavor violation needed for a Dirac mass operator can be provided by trilinear soft terms [33,34]. For a Majorana mass operator, the correct $SU(2)$ structure can also be induced by the same trilinear terms, whereas the explicit lepton-number violation can come from bilinear soft terms in the scalar components of the singlets \bar{N} . In both cases, the correct number of R charges for Dirac and Majorana mass operators is provided by neutralino mass terms.

The supersymmetric models that we consider here contain a $B-L$ symmetry broken at some large scale M_G and three additional SM singlets \bar{N} with tree-level mass M_R . Dirac mass operators mixing active and sterile neutrinos as well as Majorana mass operators for active neutrinos are generated at the quantum level via loops with virtual exchange of neutralinos and sneutrinos. Explicit expressions for the corresponding couplings m_D and m_L are given in Sec. V after a detailed description of the neutralino and sneutrino mass matrices in Sec. IV. For unsuppressed soft parameters A and B ($A \sim B \sim m_{\text{weak}}$), Dirac masses m_D are of order $\sim (m_{\text{weak}}/8\pi^2)(m_{\text{weak}}/M_G)^2$, whereas Majorana ones are $\sim (m_{\text{weak}}/8\pi^2)(m_{\text{weak}}/M_R)^3$ for $M_R \gg m_{\text{weak}}$ and $\sim (BM_R/m_{\text{weak}}/8\pi^2)$ for $M_R \ll m_{\text{weak}}$. (The qualitative behavior of the radiative contribution to Majorana masses obtained in the second scenario of Sec. II is discussed in Sec. V.) Of the resulting six eigenvalues, three are certainly light.

³Neutrino masses can be obtained radiatively also in other models. Among them, the Zee model and its supersymmetric version have long been known. For recent discussions on this possibility and references to other radiative mechanism scenarios, see Ref. [30]. See also Ref. [31] where Dirac masses are obtained at the quantum level in nonsupersymmetric models with extended gauge groups. Very popular are also R_p -violating models in which two neutrino masses are generated radiatively [32].

The other three are light or heavy depending on the value of the tree-level Majorana masses M_R for the singlets \bar{N} .

Thus, two classes of possible scenarios emerge, discriminated by the value of the mass parameter M_R . In the first one M_R is large, say, $M_R \sim M_G$, and the resulting three states n_i are of $\mathcal{O}(M_G)$. The states ν_i can be easily of $\mathcal{O}(\text{eV})$ or less. The main contribution to the corresponding eigenvalues comes from the Majorana masses m_L , which dominate over m_D^2/M_R . Therefore, the heavy $U(1)_{B-L}$ scale can be as light as a few hundred TeV. [Notice that, if the lack of a dynamical explanation for such a large scale M_R is tolerated, the same neutrino spectrum can be obtained without introducing a $U(1)_{B-L}$ symmetry.]

In the second class, M_R is very light and Dirac masses play a much more relevant role. The n_i states can be as light as, or lighter than, the ν_i 's, which are at the eV scale if M_G is $\sim 10^7$ GeV. They can be heavier, as, for example, up to the scale required to explain the motion of pulsars. They can also be degenerate with the ν_i states in the particular case of the exact vanishing of M_R : in this case, for each generation i , the two Majorana states ν_i and n_i are equivalent to one Dirac neutrino. The key ingredients for this second class of scenarios are the lightness of M_R and the seesaw suppression of m_D originating in the neutralino mass matrix. Ways to render M_R much smaller than M_G are discussed in Sec. IV.

From the phenomenological point of view, $U(1)_{B-L}$ scales as low as those needed for these two classes of scenarios are either too heavy or marginally suited to give clear collider signatures due to the additional neutral gauge boson or, possibly, heavy \bar{N} states. Drastic deviations from the phenomenology of the minimal supersymmetric standard model may, however, be observed in the scalar sector if $M_R \lesssim m_{\text{weak}}$. Phenomenological consequences as well as cosmological implications of these two scenarios are postponed to later work. Finally, in Sec. VI, we recall that in a scenario of radiative neutrino masses, also Yukawa couplings are generated radiatively. These are now *naturally* very small: i.e., $\sim m_D/m_{\text{weak}}$ for the dimensionless Dirac coupling and m_L/m_{weak}^2 for the dimensionful Majorana coupling. For Dirac and Majorana masses in the eV range, they are in general larger than the coupling induced by Yukawa operators of higher dimensionality.

IV. RADIATIVE SEESAW MECHANISMS

Six massive Majorana neutrino states are usually obtained in a three-family scenario with three left-handed neutrinos ν_L and three right-handed ones ν_R with mass operators

$$-\mathcal{L}_{\text{mass}} = -\frac{1}{2} \nu_L^T C m_L \nu_L + \bar{\nu}_R m_D \nu_L - \frac{1}{2} \nu_R^T C M_R^* \nu_R + \text{H.c.}, \quad (7)$$

where C is the charge conjugation matrix. Here and in the following, all family indices are omitted for simplicity. In the conventional seesaw mechanism, with $m_D \simeq m_{\text{weak}} \ll M_R$, the six Majorana states are split into three light ones, $\nu \simeq \nu_L$

+ ν_L^c , with mass $m_\nu \simeq -m_D(1/M_R)m_D^T$ and three heavy ones, $n \simeq \nu_R + \nu_R^c$, with mass $m_n \simeq M_R$.

In supersymmetric models with violation of $B-L$ at some large scale M_G , both Dirac and Majorana mass operators can be generated radiatively through neutralino-sneutrino loops. Suitable suppression factors, with the task of keeping m_D and m_L well below the electroweak scale, can be induced by the seesaw pattern of high-low scales in the neutralino and/or sneutrino mass matrix.

The SM fields are charged under the $B-L$ symmetry, which, for simplicity, is assumed to be a $U(1)$ symmetry. Nevertheless, the following discussion can be easily generalized to larger gauge groups. Under $U(1)_{B-L}$, quark, lepton, and Higgs chiral superfields are charged in such a way that all Yukawa interaction terms are invariant under $U(1)_{B-L}$. A charge assignment that leads to vanishing gauge anomalies is, however, unique [35]. It is given by

$$\begin{aligned} X_Q &= +1, & X_{\bar{U}} &= +1, & X_{\bar{D}} &= -3, & X_H &= -2, & X_{\bar{H}} &= +2, \\ X_L &= -3, & X_{\bar{E}} &= +1, & X_{\bar{N}} &= +5. \end{aligned} \quad (8)$$

Neutrino Yukawa terms such as those in Eq. (1) can be forbidden at the tree level by the discrete Z_n symmetry discussed in Sec. II. On the other hand, soft supersymmetry-breaking trilinear terms involving sneutrino fields can be induced by superpotential terms such as those in Eq. (2) when the spurion field Z , which is neutral under $U(1)_{B-L}$ and has Z_n charge $Z_n(Z) = -1$, acquires the supersymmetry-violating VEV \mathcal{F}_Z . As already explained in Sec. II, the same operator gives rise to tree-level Yukawa couplings when Z acquires the supersymmetry-conserving VEV \mathcal{A}_Z . In order to avoid too large neutrino masses, a hierarchy between \mathcal{A}_Z and \mathcal{F}_Z must exist. The possibility of implementing such a hierarchy, which was already assumed in Sec. II, will be discussed in Sec. VIII. The tree-level contributions can then be naturally suppressed to give neutrino Yukawa couplings as small as $\sim 10^{-13}$ and Dirac masses well below the eV range.

Majorana mass terms $(1/2)M_R \bar{N} \bar{N}$ are assumed to exist at the tree level. It is easy to see how M_R can be completely disentangled from M_G . Majorana mass terms, indeed, are induced by higher-dimensional operators in the superpotential [different from those considered in the case without a $U(1)_{B-L}$ symmetry discussed in Sec. II] that are neutral under Z_n and $U(1)_{B-L}$. For a generic Z_n symmetry, these operators are

$$W = \frac{1}{4} \frac{1}{(M_P)^{m+1}} \bar{\Phi}^m Z Z \bar{N} \bar{N}, \quad (9)$$

whereas, if the discrete symmetry Z_n is Z_2 , the lowest-order operators are

$$W = \frac{1}{2} \frac{1}{(M_P)^{m-1}} \bar{\Phi}^m \bar{N} \bar{N}. \quad (10)$$

In both cases, the field $\bar{\Phi}$ is a SM singlet that breaks $U(1)_{B-L}$ and gets a VEV of $\mathcal{O}(M_G)$. It is charged under $U(1)_{B-L}$ with charge $X_{\bar{\Phi}}$. The power m in Eqs. (9) and (10) depends on $X_{\bar{\Phi}}$, once the charge assignment for \bar{N} is fixed as in Eq. (8). It is clear, then, that the Majorana mass M_R does not need to be of the same order of M_G , and it can get values over a wide range of scales below M_G . Indeed, if the discrete symmetry that forbids tree-level Yukawa terms is Z_2 , it may be of $\mathcal{O}(M_G)$ if $m=1$. If $m=2$, depending on the value of M_G , it may be a few orders of magnitude above the electroweak scale or at the eV scale (see possible values of M_G discussed in Sec. VII). It can also be extraordinarily suppressed if a generic Z_n symmetry is assumed and if $\bar{\Phi}$ has the $U(1)_{B-L}$ charge $X_{\bar{\Phi}} = -1$. For the charge assignment in Eq. (8), it is, for example, $M_R \sim M_G^{10} \mathcal{A}_Z^2 / (M_P)^{11}$. Finally, if we impose the symmetry in Eq. (3), the higher-dimensional operators in Eqs. (9) and (10) are forbidden and the exact vanishing of M_R is guaranteed.⁴

Notice how the presence of a gauged $U(1)_{B-L}$ symmetry prevents, in general, unwanted vacua with nonvanishing lepton number. Indeed, in contrast to the situation described in the second scenario of Sec. II, no dangerously large bilinear soft supersymmetry-breaking terms BM_R are induced by the same operators that generate M_R .

Tree-level Majorana mass terms for left-handed neutrinos are strongly suppressed. They may arise from superpotential operators:

$$W = \frac{1}{4} \frac{1}{(M_P)^{m+1}} \Phi^m LHLH, \quad (11)$$

where Φ is another SM singlet needed in addition to $\bar{\Phi}$ to cancel $U(1)_{B-L}$ anomalies. It has $U(1)_{B-L}$ charge opposite to that of $\bar{\Phi}$. As in the case of the operators in Eqs. (9) and (10), also these Majorana mass terms can be forbidden by the symmetry in Eq. (3).

A. Sneutrino mass matrix

As long as M_R is different from zero, a bilinear and lepton number-violating soft supersymmetry-breaking term $(1/2)BM_R\bar{N}\bar{N}$ is always induced. Since the parameter B is, in general, at the electroweak scale, this term may be very large or very small, depending on the value of M_R . The complete sneutrino mass potential is given by

$$-\mathcal{L}_{\text{mass}} = m_{\tilde{l}}^2 \tilde{N}^* \tilde{N} + m_{\tilde{\nu}}^2 \tilde{N}^* \tilde{N} + M_R^2 \tilde{N}^* \tilde{N} + \frac{1}{2} M_R (B \tilde{N} \tilde{N} + B^* \tilde{N}^* \tilde{N}^*) + (A v \tilde{N} \tilde{N} + A^* v \tilde{N}^* \tilde{N}^*), \quad (12)$$

where v is the VEV of the relevant Higgs doublet. When redefining the sneutrino fields as $\tilde{\nu}_L \equiv \tilde{N}$ and $\tilde{\nu}_R \equiv \tilde{N}^*$, the above expression for $-\mathcal{L}_{\text{mass}}$ becomes

$$-\mathcal{L}_{\text{mass}} = m_{\tilde{l}}^2 \tilde{\nu}_L^* \tilde{\nu}_L + m_{\tilde{\nu}}^2 \tilde{\nu}_R^* \tilde{\nu}_R + M_R^2 \tilde{\nu}_R^* \tilde{\nu}_R + \frac{1}{2} M_R (B \tilde{\nu}_R^* \tilde{\nu}_R + B^* \tilde{\nu}_R \tilde{\nu}_R) + (A v \tilde{\nu}_R^* \tilde{\nu}_L + A^* v \tilde{\nu}_L^* \tilde{\nu}_R). \quad (13)$$

A nonvanishing soft parameter B in Eq. (13) induces a splitting in the CP -even and CP -odd sneutrino states. By decomposing $\tilde{\nu}_L$ and $\tilde{\nu}_R$ as $\tilde{\nu}_L \equiv (\tilde{\nu}_{L+} + i \tilde{\nu}_{L-})/\sqrt{2}$ and $\tilde{\nu}_R \equiv (\tilde{\nu}_{R+} + i \tilde{\nu}_{R-})/\sqrt{2}$, the sneutrino mass potential can, then, be expressed as

$$-\mathcal{L}_{\text{mass}} = \frac{1}{2} \begin{pmatrix} \tilde{\nu}_{L+} & \tilde{\nu}_{R+} & \tilde{\nu}_{L-} & \tilde{\nu}_{R-} \end{pmatrix} \mathcal{M}_{\text{sneut}}^2 \begin{pmatrix} \tilde{\nu}_{L+} \\ \tilde{\nu}_{R+} \\ \tilde{\nu}_{L-} \\ \tilde{\nu}_{R-} \end{pmatrix}. \quad (14)$$

The matrix $\mathcal{M}_{\text{sneut}}^2$ is, in general, a rather complicated matrix. For one-generation neutrino fields, i.e., ignoring the flavor-changing structure of A , B , $m_{\tilde{l}}$, and $m_{\tilde{\nu}}$, $\mathcal{M}_{\text{sneut}}^2$ is the following 4×4 matrix:

$$\begin{pmatrix} m_{\tilde{l}}^2 & \frac{1}{2}(A+A^*)v & 0 & \frac{i}{2}(A^*-A)v \\ \frac{1}{2}(A+A^*)v & \left(M_R^2 + \frac{1}{2}(B+B^*)M_R + m_{\tilde{\nu}}^2 \right) & \frac{i}{2}(A-A^*)v & \frac{i}{2}(B^*-B)M_R \\ 0 & \frac{i}{2}(A-A^*)v & m_{\tilde{l}}^2 & \frac{1}{2}(A+A^*)v \\ \frac{i}{2}(A^*-A)v & \frac{i}{2}(B^*-B)M_R & \frac{1}{2}(A+A^*)v & \left(M_R^2 - \frac{1}{2}(B+B^*)M_R + m_{\tilde{\nu}}^2 \right) \end{pmatrix}. \quad (15)$$

⁴Depending on $X_{\bar{\Phi}}$, it is also possible that no Majorana mass term for \bar{N} is allowed due to a residual unbroken discrete subgroup of $U(1)_{B-L}$, even if no other symmetry as that in Eq. (3) is imposed.

This matrix simplifies drastically when A and B are made real using phase rotations of the N and \bar{N} superfields: it reduces, in this case, to the block diagonal matrix

$$\mathcal{M}_{\text{sneut}}^2 = \begin{pmatrix} \mathcal{M}_+^2 & 0_2 \\ 0_2 & \mathcal{M}_-^2 \end{pmatrix}, \quad \text{with}$$

$$\mathcal{M}_\pm^2 = \begin{pmatrix} m_{\tilde{l}}^2 & Av \\ Av & M_R^2 \pm BM_R + m_{\tilde{\nu}}^2 \end{pmatrix}. \quad (16)$$

(The same decomposition of $\mathcal{M}_{\text{sneut}}^2$ is found in Ref. [36]; see also [37].)

For large M_R ($\sim M_G$), the two matrices \mathcal{M}_\pm^2 have the typical pattern of light-heavy scales present in the neutrino mass matrix in the conventional seesaw mechanism. Of the four eigenstates $\tilde{\nu}_{+,1}$, $\tilde{\nu}_{+,2}$, $\tilde{\nu}_{-,1}$, and $\tilde{\nu}_{-,2}$, two ($\tilde{\nu}_{\pm,1}$) are at the electroweak scale and two ($\tilde{\nu}_{\pm,2}$) are heavy, i.e., at the scale M_R . The two states in the light pair and those in the heavy pair have masses that are split by terms proportional to B . These splitting terms are suppressed by powers of (m_{weak}/M_R) . It is, indeed,

$$m_{\tilde{\nu}_{\pm,1}}^2 = m_{\tilde{l}}^2 \{1 - (ac)^2 \beta^2 \pm (ac)^2 b \beta^3 + \mathcal{O}(\beta^4)\},$$

$$m_{\tilde{\nu}_{\pm,2}}^2 = M_R^2 \{1 \pm b \beta + d^2 \beta^2 + \mathcal{O}(\beta^4)\}, \quad (17)$$

where the definitions

$$\beta \equiv \frac{m_{\tilde{l}}}{M_R}, \quad a \equiv \frac{A}{m_{\tilde{l}}}, \quad b \equiv \frac{B}{m_{\tilde{l}}}, \quad c \equiv \frac{v}{m_{\tilde{l}}}, \quad d \equiv \frac{m_{\tilde{\nu}}}{m_{\tilde{l}}}, \quad (18)$$

were used. At the same level of approximation, i.e., up to terms of $\mathcal{O}(\beta^4)$, the two diagonalization matrices of \mathcal{M}_\pm^2 , U_\pm , are given by

$$\begin{pmatrix} \tilde{\nu}_{L\pm} \\ \tilde{\nu}_{R\pm} \end{pmatrix} = U_\pm \begin{pmatrix} \tilde{\nu}_{\pm,1} \\ \tilde{\nu}_{\pm,2} \end{pmatrix}$$

$$\equiv \begin{pmatrix} 1 & -ac(-\beta^2 \pm b\beta^3) \\ ac(-\beta^2 \pm b\beta^3) & 1 \end{pmatrix} \begin{pmatrix} \tilde{\nu}_{\pm,1} \\ \tilde{\nu}_{\pm,2} \end{pmatrix}. \quad (19)$$

A Majorana mass operator radiatively induced by sneutrino-neutralino loops must have a coupling proportional to the mass splitting of the light or heavy sneutrino pairs, since this is induced by the lepton-violating B parameter. Notice that the splitting of the two light eigenvalues comes only at order $(m_{\text{weak}}/M_R)^3$. It is, therefore, conceivable that a Majorana mass term can be sufficiently suppressed by several powers of (m_{weak}/M_R) .

For very tiny M_R and small (BM_R/m_{weak}^2) , the correct expansion parameter needed to calculate eigenvalues and eigenstates of \mathcal{M}_\pm^2 is the inverse of β , which we indicate by γ . Since eigenvalues and eigenvectors are more involved in this case, we express them in terms of those obtained at the

zeroth order in γ . For $M_R=0$, \mathcal{M}_+^2 and \mathcal{M}_-^2 coincide, and the sneutrino mass potential has the well-known form

$$-\mathcal{L}_{\text{mass}} = \begin{pmatrix} \tilde{\nu}_L^* & \tilde{\nu}_R^* \end{pmatrix} \begin{pmatrix} m_{\tilde{l}}^2 & Av \\ Av & m_{\tilde{\nu}}^2 \end{pmatrix} \begin{pmatrix} \tilde{\nu}_L \\ \tilde{\nu}_R \end{pmatrix}, \quad (20)$$

once the phase of the A parameter is rotated away. Then, the eigenvalues are given by

$$m_{\tilde{\nu}_{1,2}}^2 = \frac{1}{2} \{m_{\tilde{l}}^2 + m_{\tilde{\nu}}^2 \pm \sqrt{(m_{\tilde{l}}^2 - m_{\tilde{\nu}}^2)^2 + 4(Av)^2}\}, \quad (21)$$

and the diagonalization matrix takes the following form:

$$U = \sqrt{\frac{m_{\tilde{\nu}}^2 - m_{\tilde{\nu}_1}^2}{m_{\tilde{\nu}_2}^2 - m_{\tilde{\nu}_1}^2}} \begin{pmatrix} 1 & \frac{Av}{m_{\tilde{\nu}}^2 - m_{\tilde{\nu}_1}^2} \\ -\frac{Av}{m_{\tilde{\nu}}^2 - m_{\tilde{\nu}_1}^2} & 1 \end{pmatrix}. \quad (22)$$

If γ is nonzero, the masses for the CP -even and CP -odd sneutrino states split, and the eigenvalues of \mathcal{M}_\pm^2 now become

$$m_{\tilde{\nu}_{\pm,1}}^2 = m_{\tilde{\nu}_1}^2 \pm \frac{m_{\tilde{\nu}_2}^2 - m_{\tilde{\nu}}^2}{m_{\tilde{\nu}_2}^2 - m_{\tilde{\nu}_1}^2} m_{\tilde{l}}^2 b \gamma + \mathcal{O}(m_{\tilde{l}}^2 \gamma^2),$$

$$m_{\tilde{\nu}_{\pm,2}}^2 = m_{\tilde{\nu}_2}^2 \pm \frac{m_{\tilde{\nu}}^2 - m_{\tilde{\nu}_1}^2}{m_{\tilde{\nu}_2}^2 - m_{\tilde{\nu}_1}^2} m_{\tilde{l}}^2 b \gamma + \mathcal{O}(m_{\tilde{l}}^2 \gamma^2), \quad (23)$$

with b defined in Eq. (18). At the same level of approximation, the corresponding diagonalization matrices U_\pm are

$$U_\pm = \sqrt{\frac{m_{\tilde{\nu}}^2 - m_{\tilde{\nu}_1}^2}{m_{\tilde{\nu}_2}^2 - m_{\tilde{\nu}_1}^2}} \begin{pmatrix} 1 + x_\pm & \frac{Av}{m_{\tilde{\nu}}^2 - m_{\tilde{\nu}_1}^2} - y_\pm \\ -\frac{Av}{m_{\tilde{\nu}}^2 - m_{\tilde{\nu}_1}^2} + y_\pm & 1 + x_\pm \end{pmatrix}, \quad (24)$$

where x_\pm and y_\pm are given by

$$x_\pm = \pm \frac{(Av)^2}{(m_{\tilde{\nu}}^2 - m_{\tilde{\nu}_1}^2)(m_{\tilde{\nu}_2}^2 - m_{\tilde{\nu}_1}^2)} m_{\tilde{l}}^2 b \gamma,$$

$$y_\pm = x_\pm \left\{ \frac{Av}{m_{\tilde{\nu}}^2 - m_{\tilde{\nu}_1}^2} + \frac{m_{\tilde{\nu}}^2 - m_{\tilde{l}}^2}{Av} \right\}. \quad (25)$$

Differently than in the case with opposite hierarchy, i.e., $M_R \gg m_{\text{weak}}$, here the first nonvanishing splitting of the two pairs of sneutrino eigenvalues is linear in the suppression parameter (M_R/m_{weak}).

In the second scenario discussed in Sec. II radiative contributions to the Majorana mass for left-handed neutrinos are possible and potentially large. In this scenario, M_R is small, i.e., $M_R \ll m_{\text{weak}}$, but BM_R , induced by the operator (4), is larger than all other entries in the sneutrino mass matrix if \mathcal{F}_Z is of the order of the maximal VEV inducing supersymmetry breaking. For coefficients z_ν of $\mathcal{O}(1)$, a value of \mathcal{F}_Z smaller than the typical VEV $\sim m_{\text{weak}} M_P$ is required in order to avoid a tachyonic sneutrino state. When \mathcal{F}_Z is minimally reduced, the splittings in the two pairs of eigenvalues of the two matrices \mathcal{M}_\pm^2 is of $\lesssim \mathcal{O}(m_{\text{weak}}^2)$. However, it is easy to convince oneself, at least *a posteriori*, that the requirement $m_{\nu_i} \lesssim 1$ eV implies $BM_R < m_{\text{weak}}^2$. Therefore, the approximated expressions in Eqs. (23), (24), and (25) still hold in this case.

B. Neutralino mass matrix

The neutralino sector of this model is also more involved than in the minimal supersymmetric extension of the SM. The spontaneous breaking of $U(1)_{B-L}$ is encapsulated in the superpotential terms

$$W = \frac{1}{\sqrt{2}} Y (\Phi \bar{\Phi} - v_\Phi^2). \quad (26)$$

Here, Y is a chiral superfield neutral under $U(1)_{B-L}$ that forces Φ and $\bar{\Phi}$ to acquire VEV's $\langle \Phi \rangle = \langle \bar{\Phi} \rangle = v_\Phi$, which can be assumed real and positive. The two fields Φ and $\bar{\Phi}$ can be reexpressed in terms of a Goldstone chiral multiplet Ψ , which is absorbed by the $U(1)_{B-L}$ gauge multiplet X , and a chiral field K . The effect of this super-Higgs mechanism is that the gauge multiplet acquires mass

$$M_G = 2 g_X |X_\Phi| v_\Phi, \quad (27)$$

where g_X is the $U(1)_{B-L}$ gauge coupling and X_Φ the $U(1)_{B-L}$ charge of the field Φ . The superpotential in Eq. (26) reduces to

$$W = \frac{1}{2\sqrt{2}} Y K^2 + v_\Phi Y K. \quad (28)$$

(Notice that a potentially dangerous coupling $Y H \bar{H}$, which could shift the scale of the SM breaking up to M_G , can be easily forbidden by making use of an R symmetry.)

The implementation of a spontaneous breaking of $U(1)_{B-L}$ requires therefore the inclusion of at least three SM neutral singlet superfields. As a consequence, new neutralino states are present in this model in addition to the usual \tilde{B} , \tilde{W}_3 , \tilde{H}_0 , and $\tilde{\tilde{H}}_0$. For the three new singlet superfields Φ , $\bar{\Phi}$, and Y , the new neutralino states are $\tilde{\Psi}$, \tilde{K} , \tilde{Y} , and

the $U(1)_{B-L}$ gaugino \tilde{X} . On the basis $(\tilde{B} \tilde{X} \tilde{\Psi} | \tilde{K} \tilde{Y} | \tilde{W}_3 \tilde{H}_0 \tilde{\tilde{H}}_0)^T$, the neutralino mass matrix, when normalized as

$$-\mathcal{L}_{\text{mass}} = 1/2 (\tilde{B} \tilde{X} \tilde{\Psi} | \tilde{K} \tilde{Y} | \tilde{W}_3 \tilde{H}_0 \tilde{\tilde{H}}_0) \times \mathcal{M}_{\text{neutr}} (\tilde{B} \tilde{X} \tilde{\Psi} | \tilde{K} \tilde{Y} | \tilde{W}_3 \tilde{H}_0 \tilde{\tilde{H}}_0)^T, \quad (29)$$

acquires the form

$$\mathcal{M}_{\text{neutr}} = \begin{pmatrix} m_{\tilde{B}} & m_{\text{mix}} & 0 & 0 & 0 & 0 & gv & gv \\ m_{\text{mix}} & m_{\tilde{X}} & M_G & 0 & 0 & 0 & gv & gv \\ 0 & M_G & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & v_\Phi & 0 & 0 & 0 \\ 0 & 0 & 0 & v_\Phi & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & m_{\tilde{W}} & gv & gv \\ gv & gv & 0 & 0 & 0 & gv & 0 & \mu \\ gv & gv & 0 & 0 & 0 & gv & \mu & 0 \end{pmatrix}. \quad (30)$$

In Eq. (30), the two-component notation for the spinor fields is used. The entry M_G is the Dirac-type mass that couples \tilde{X} to $\tilde{\Psi}$, whereas the Majorana-type mass $m_{\tilde{X}}$ for the gaugino \tilde{X} is a soft supersymmetry-breaking parameter of the order of the weak scale. The entry m_{mix} , also at the electroweak scale, is a mass terms that mixes the $U(1)_Y$ and $U(1)_{B-L}$ gauginos. The generic symbol gv stays here for the product of a $SU(2) \times U(1)$ gauge coupling times one of the two SM VEV's and some normalization factor, and it is not always the same in the different entries of $\mathcal{M}_{\text{neutr}}$. The diagonalization of this matrix through a unitary transformation ($D^\dagger \mathcal{M}_{\text{neutr}} D$) leads to eight mass eigenstates $\tilde{\chi}_i^0$, related to the current eigenstates through

$$(\tilde{B} \tilde{X} \tilde{\Psi} | \tilde{K} \tilde{Y} | \tilde{W}_3 \tilde{H}_0 \tilde{\tilde{H}}_0)^T = D (\tilde{\chi}_1^0 \tilde{\chi}_2^0 \tilde{\chi}_3^0 \tilde{\chi}_4^0 \tilde{\chi}_5^0 \tilde{\chi}_6^0 \tilde{\chi}_7^0 \tilde{\chi}_8^0)^T. \quad (31)$$

Of these mass eigenstates, two are degenerate with mass $\pm v_\Phi$, and two are almost degenerate, at the scale M_G . Their masses differ by an overall sign; the splitting in the absolute value of these masses is of order m_{weak} . The remaining four eigenstates are at the electroweak scale.

In order to visualize the relevance of these states for the radiative generation of neutrino masses, in particular of Dirac type, it is convenient to take the limit $v \rightarrow 0$ in the

neutralino mass matrix. Mixing terms due to SU(2) Higgsino-gaugino couplings as well as SU(2)-gaugino-U(1) gaugino couplings, which are of order $(m_Z/\tilde{m})^2$, with \tilde{m} a soft supersymmetry breaking parameter, are in this case neglected. In this limit, $\mathcal{M}_{\text{neutr}}$ reduces to a block-diagonal matrix. The lower 3×3 block has no relevance for the radiative generation of neutrino masses, since no SU(2) Higgsino-neutrino-sneutrino couplings are possible in the absence of tree-level neutrino Yukawa couplings. The central 2×2 block turns out to give a vanishing contribution to neutrino masses. Therefore, in this limit, the 3×3 upper block is the only one of interest to induce Dirac neutrino mass operators. Fermionic and scalar components of the singlets \tilde{N} , indeed, couple only to the gaugino \tilde{X} and the field \tilde{K} . However, once the rotation (31) is performed, all mass eigenstates $\tilde{\chi}_i^0$ participate in the vertex $\tilde{\nu}_R \tilde{\chi}_i^0 \tilde{\nu}_R$, with couplings that are larger for the heavy $\tilde{\chi}_i^0$ states. Two of these $\tilde{\chi}_i^0$ states are a linear combination of \tilde{K} and \tilde{Y} only, obtained through a rotation at 45° of \tilde{K} and \tilde{Y} . These states have exactly opposite masses $\pm v_\Phi$ and their contributions to Dirac mass operators cancel identically. A similar feature holds for the contribution coming from the two states that are mainly \tilde{X} and $\tilde{\Psi}$. In this case, however, the cancellation is not exact, but it is spoiled by powers of the suppression factor (m_{weak}/M_G) , where m_{weak}

can be any of the remaining parameters in the upper 3×3 block of $\mathcal{M}_{\text{neutr}}$, i.e., $m_{\tilde{B}}$, m_{mix} , or $m_{\tilde{X}}$. It turns out that the lowest-order suppression factor comes with power 2. As will be stressed later, this fact has important phenomenological consequences.

We close this section by giving explicitly eigenvalues and eigenvectors of the 3×3 upper block in the neutralino mass matrix (30), assuming that its elements are real. The three eigenvalues are

$$m_{\tilde{\chi}_0^0} = m_{\tilde{B}}, \quad m_{\tilde{\chi}_\pm^0} = M_G \left\{ \pm 1 + \alpha \frac{x}{2} \pm \alpha^2 \left(\frac{x^2}{8} + \frac{z^2}{2} \right) \right\}, \quad (32)$$

where α , z , and x are

$$\alpha = \frac{m_{\tilde{B}}}{M_G}, \quad z = \frac{m_{\text{mix}}}{m_{\tilde{B}}}, \quad x = \frac{m_{\tilde{X}}}{m_{\tilde{B}}}. \quad (33)$$

At the same level of precision in the small parameter α , the reduced 3×3 diagonalization matrix, defined by

$$(\tilde{B} \tilde{X} \tilde{\Psi})^T = D_{(3)} (\tilde{\chi}_0^0 \tilde{\chi}_+^0 \tilde{\chi}_-^0)^T, \quad (34)$$

is

$$D_{(3)} = \begin{pmatrix} 1 - \alpha^2 \frac{z^2}{2} & \frac{z}{\sqrt{2}} \left[\alpha + \alpha^2 \left(1 - \frac{x}{4} \right) \right] & \frac{z}{\sqrt{2}} \left[\alpha - \alpha^2 \left(1 - \frac{x}{4} \right) \right] \\ -\alpha^2 z & \frac{1}{\sqrt{2}} \left[1 + \alpha \frac{x}{4} - \alpha^2 \frac{x^2}{32} \right] & \frac{1}{\sqrt{2}} \left[-1 + \alpha \frac{x}{4} + \alpha^2 \frac{x^2}{32} \right] \\ -\alpha z & \frac{1}{\sqrt{2}} \left[1 - \alpha \frac{x}{4} - \alpha^2 \left(\frac{x^2}{32} + \frac{z^2}{2} \right) \right] & \frac{1}{\sqrt{2}} \left[1 + \alpha \frac{x}{4} - \alpha^2 \left(\frac{x^2}{32} + \frac{z^2}{2} \right) \right] \end{pmatrix}. \quad (35)$$

V. EFFECTIVE DIRAC AND MAJORANA NEUTRINO MASSES

We are now in a position to evaluate Dirac and Majorana mass couplings m_D and m_L . Since Majorana masses violate lepton number, they can only be generated for nonvanishing BM_R . In contrast, Dirac masses can be obtained also for $M_R = 0$. In the following, we give expressions for m_D and m_L classified according to the value of M_R . As in the previous section, also here the flavor-changing structure of the soft parameters in the sneutrino potential is ignored. All the following expressions for m_D and m_L hold for each generation. A generalization to the case in which A , B , $m_{\tilde{L}}$, and $m_{\tilde{\nu}}$, have a nontrivial matrix structure is technically straightforward but leads to more involved formulas.

A. Dirac neutrino mass m_D , $M_R \approx 0$

The radiatively generated Dirac mass m_D is evaluated in the mass eigenbasis for neutralinos and sneutrinos. The cor-

responding diagram is shown in Fig. 1, where the sneutrino soft parameter A , which breaks chiral symmetries, and the neutralino mass $m_{\tilde{\chi}_j^0}$, which connects left- and right-chiral fermions, are shown as mass insertions.

The Majorana mass M_R , which couples the singlets \tilde{N} , is assumed to exist at the tree level, but to be small. Therefore, the approximation in Eq. (20) is sufficient for the determina-




FIG. 1. Diagram contributing to the Dirac neutrino mass m_D .

tion of the sneutrino mass eigenstates. They are, in this case, only two, $\tilde{\nu}_1$ and $\tilde{\nu}_2$, and are both at the electroweak scale. The current eigenstate vertex $\tilde{\nu}_R \tilde{X} \tilde{\nu}_R$ has a coupling $-i\sqrt{2}g_X X_{\tilde{N}}$ where $X_{\tilde{N}}$ is the charge of the singlet superfield \tilde{N} under $U(1)_{B-L}$. The gaugino \tilde{X} couples also to the left-handed neutrinos and sneutrinos with a similar coupling $-i\sqrt{2}g_X X_L$. A straightforward calculation of the Dirac neutrino mass, in the one-generation case, then, yields

$$m_D = \frac{(Av)}{8\pi^2} \sum_{j=1}^6 (g_X X_{\tilde{N}} D_{2j}) m_{\tilde{\chi}_j^0} \{g_Y Y_L D_{1j} + g_X X_L D_{2j} + g_2 T_{3L} D_{6j}\} I(m_{\tilde{\nu}_1}^2, m_{\tilde{\nu}_2}^2, m_{\tilde{\chi}_j^0}^2), \quad (36)$$

where the function $I(m_1^2, m_2^2, m_3^2)$ is defined in [31], Y_L is the hypercharge of the $SU(2)$ leptonic doublet, T_{3L} is the isospin of the neutral component of this doublet, and D is the neutralino diagonalization matrix defined in Eq. (31).

This result becomes particularly transparent when the limit $v \rightarrow 0$ is taken in the neutralino mass matrix. In this limit, the mixed contribution of \tilde{W}_3 and heavy neutralinos vanishes, and the third term in the curly bracket of Eq. (36) drops out. Thus, in the approximation of vanishing phases in the neutralino mass matrix, the Dirac neutrino mass becomes

$$m_D \simeq \frac{g_Y g_X}{8\pi^2} Y_L X_{\tilde{N}} (Av) m_{\text{mix}} \left\{ -\left(\frac{m_{\tilde{B}}}{M_G}\right)^2 I(m_{\tilde{\nu}_1}^2, m_{\tilde{\nu}_2}^2, m_{\tilde{B}}^2) + I(m_{\tilde{\nu}_1}^2, m_{\tilde{\nu}_2}^2, M_G^2) \right\} + \frac{g_X^2}{8\pi^2} X_L X_{\tilde{N}} \left(\frac{Av}{m_{\tilde{B}}}\right) \left(\frac{m_{\tilde{X}}}{m_{\tilde{B}}}\right) \left(\frac{m_{\tilde{B}}}{M_G}\right)^2, \quad (37)$$

where higher orders in the expansion parameter $(m_{\tilde{B}}/M_G)$ have been neglected. Notice that in the absence of a mass term mixing the $U(1)_Y$ and $U(1)_{B-L}$ gauginos, a contribution to the Dirac mass m_D comes from pure \tilde{X} - \tilde{X} exchange. Finally, in the approximation $m_{\tilde{\nu}_1}^2 \simeq m_{\tilde{\nu}_2}^2 \simeq m_{\tilde{B}}^2 \ll M_G^2$, the above result can be cast in the simple form

$$m_D \sim \frac{g_X}{8\pi^2} X_{\tilde{N}} \left(\frac{Av}{m_{\tilde{B}}}\right) \left(\frac{m_{\tilde{B}}}{M_G}\right)^2 \left\{ g_Y Y_L \left(\frac{m_{\text{mix}}}{m_{\tilde{B}}}\right) \times \left[2 \log\left(\frac{M_G}{m_{\tilde{B}}}\right) - \frac{3}{2} \right] + g_X X_L \left(\frac{m_{\tilde{X}}}{m_{\tilde{B}}}\right) \right\}. \quad (38)$$

For soft supersymmetry-breaking terms at the electroweak scale, the Dirac neutrino mass m_D , besides being suppressed by a loop factor, is moved away from m_{weak} by the ratio $(m_{\text{weak}}/M_G)^2$. The behavior $m_D \simeq (m_{\text{weak}}/8\pi^2)(m_{\text{weak}}/M_G)^2$ is quite general and it is not limited to the choice of gauge group $U(1)_{B-L}$ made here. It has to be observed, however, that the suppression factor $(m_{\text{weak}}/M_G)^2$ is actually $(\tilde{m}/M_G)^2$, where \tilde{m} is a generic soft supersymmetry-breaking parameter. Therefore, because of its dependence on a fixed scale M_G , m_D tends to grow in the superpartner

decoupling limit. The effectiveness of the suppression factor $(\tilde{m}/M_G)^2$ is recovered only when also M_G is increased and/or the trilinear parameters A are decreased.

B. Dirac neutrino mass m_D , $M_R \gg m_{\text{weak}}$

If the Majorana mass M_R is non-negligible, then CP -even and CP -odd components of the sneutrino states are split by lepton flavor-violating mass terms. Therefore, when calculating m_D , the general form in Eq. (15) for the sneutrino mass matrix has to be used or, at least, that in Eq. (16) obtained in the basis in which the A and B parameters are real. In this basis, m_D gets the form

$$m_D = \frac{(Av)}{16\pi^2} \sum_{j=1}^6 (g_X X_{\tilde{N}} D_{2j}) m_{\tilde{\chi}_j^0} \{g_Y Y_L D_{1j} + g_X X_L D_{2j} + g_2 T_{3L} D_{6j}\} [I(m_{\tilde{\nu}_{+1}}^2, m_{\tilde{\nu}_{+2}}^2, m_{\tilde{\chi}_j^0}^2) + I(m_{\tilde{\nu}_{-1}}^2, m_{\tilde{\nu}_{-2}}^2, m_{\tilde{\chi}_j^0}^2)]. \quad (39)$$

When taking the limit $v \rightarrow 0$ in the neutralino mass matrix, also in this case, the mixed contribution of \tilde{W}_3 and heavy neutralinos vanishes. Moreover, if $M_R \simeq M_G$, much above the electroweak scale, while $m_{\tilde{\nu}_{-1}} \simeq m_{\tilde{\nu}_{+1}} \simeq m_{\tilde{l}} \simeq m_{\tilde{B}}$, the above expression reduces to

$$m_D \sim \frac{g_X}{8\pi^2} X_{\tilde{N}} \left(\frac{Av}{m_{\tilde{B}}}\right) \left(\frac{m_{\tilde{B}}}{M_R}\right)^2 \left\{ g_Y Y_L \left(\frac{m_{\text{mix}}}{m_{\tilde{B}}}\right) \left[2 \log\left(\frac{M_R}{M_G}\right) \right] + g_X X_L \left(\frac{m_{\tilde{X}}}{m_{\tilde{B}}}\right) \left[2 \log\left(\frac{M_R}{M_G}\right) - 1 \right] \right\}, \quad (40)$$

in the approximation of vanishing phases in the neutralino mass matrix. Although different in some details, this expression for m_D is qualitatively very similar to that obtained when $M_R \simeq 0$. The double suppression $(m_{\text{weak}}/M_G)^2$ seems, therefore, a typical feature of radiatively induced Dirac masses. It holds, indeed, for any value of M_R .

C. Majorana neutrino mass m_L , $M_R \gg m_{\text{weak}}$

The Majorana neutrino mass m_L violates lepton number and must have a direct dependence on the soft parameter B , which is the only lepton-number-violating parameter in the sneutrino potential.⁵ The diagram needed to obtain m_L is shown in Fig. 2. The parameters A and B as well as the relevant gaugino mass are shown as mass insertions, although the calculation is done using the mass eigenstate formalism. The neutrino mass is directly proportional to the

⁵A different possibility was considered in the last reference in [33]. In the absence of additional singlets \tilde{N} , a lepton-number-violating soft supersymmetry-breaking trilinear term was allowed. A mass splitting among left-handed sneutrinos was induced at the one-loop level and a Majorana neutrino mass m_L at the two-loop level.




FIG. 2. Diagram contributing to the Majorana neutrino mass m_L .

splitting of each of the two pairs of eigenvalues of the sneutrino mass matrix (16), induced by B . The largest contribution to the Majorana mass comes from the splitting in the lightest eigenvalues. The splitting in the two heavy eigenvalues, which appears at a lower order in the expansion parameter $m_{\tilde{l}}/M_R$ than the splitting in the two light eigenvalues, gives a much more suppressed contribution. The resulting expression is

$$m_L = \frac{1}{4\pi^2} \sum_{j=1}^6 (g_Y Y_L D_{1j} + g_X X_L D_{2j} + g_2 T_{3L} D_{6j}) m_{\tilde{\chi}_j^0} (g_Y Y_L D_{1j} + g_X X_L D_{2j} + g_2 T_{3L} D_{6j}) \times \left(\frac{Av}{m_{\tilde{l}}} \right)^2 \left(\frac{B}{m_{\tilde{l}}} \right) \left(\frac{m_{\tilde{l}}}{M_R} \right)^3 I(m_{\tilde{\nu}_{+1}}^2, m_{\tilde{\nu}_{-1}}^2, m_{\tilde{\chi}_j^0}^2). \quad (41)$$

When taking the limit $v \rightarrow 0$ in the neutralino mass matrix, the contribution from mixed propagators $\tilde{B}-\tilde{W}_3$ drops out and only the pure $\tilde{B}-\tilde{B}$, $\tilde{X}-\tilde{X}$, and $\tilde{W}_3-\tilde{W}_3$ propagators survive. Furthermore, assuming a real neutralino mass matrix, in the limit $m_{\tilde{\nu}_{+1}}^2 \simeq m_{\tilde{\nu}_{-1}}^2 \simeq m_{\tilde{l}}^2 \simeq m_{\tilde{B}}^2 \simeq m_{\tilde{W}}^2$, as well as $M_G \simeq M_R$ for the functions $I(\dots)$, the previous expression becomes

$$m_L \simeq \frac{1}{8\pi^2} \left(\frac{Av}{m_{\tilde{l}}^2} \right)^2 \left(\frac{B}{m_{\tilde{l}}} \right) \left(\frac{m_{\tilde{l}}}{M_R} \right)^3 \{ (g_Y Y_L)^2 m_{\tilde{B}} + (g_2 T_{3L})^2 m_{\tilde{W}} \}. \quad (42)$$

It is interesting to see that the suppression (m_{weak}/M_R) comes with a higher power than in the expression for the Dirac mass m_D . It is, indeed, $m_L \sim (m_{\text{weak}}/8\pi^2)(m_{\text{weak}}^2/M_R^3)$. Also in this case, m_L tends to increase in the superpartner decoupling limit and its smallness can be retrieved by tuning the parameters A and B to be very small or raising the value of M_R .

D. Majorana neutrino mass m_L , $M_R \ll m_{\text{weak}}$

If M_R is much smaller than the electroweak scale, the Majorana neutrino mass m_L is not suppressed by the large scale $\sim M_G$. In this case, all particles exchanged in the loop in Fig. 2 have masses of the order of the electroweak scale, and they all contribute to m_L . Thus, the suppression factor only comes from (M_R/m_{weak}) when all the supersymmetry-breaking parameters are around the weak scale. The Majorana neutrino mass is given by

$$m_L = \frac{1}{4\pi^2} \sum_{j=1}^6 (g_Y Y_L D_{1j} + g_X X_L D_{2j} + g_2 T_{3L} D_{6j}) m_{\tilde{\chi}_j^0} (g_Y Y_L D_{1j} + g_X X_L D_{2j} + g_2 T_{3L} D_{6j}) \times (BM_R) \frac{(Av)^2}{(m_{\tilde{\nu}_2}^2 - m_{\tilde{\nu}_1}^2)^2} [I(m_{\tilde{\nu}_1}^2, m_{\tilde{\nu}_1}^2, m_{\tilde{\chi}_j^0}^2) + I(m_{\tilde{\nu}_2}^2, m_{\tilde{\nu}_2}^2, m_{\tilde{\chi}_j^0}^2) - 2I(m_{\tilde{\nu}_1}^2, m_{\tilde{\nu}_2}^2, m_{\tilde{\chi}_j^0}^2)], \quad (43)$$

where $m_{\tilde{\nu}_{1,2}}^2$ are defined in Eq. (21). When taking the limit $v \rightarrow 0$ in the neutralino mass matrix, the contribution from mixed propagators $\tilde{B}-\tilde{W}_3$ drops out and the dominant contributions come from the $\tilde{B}-\tilde{B}$ and $\tilde{W}_3-\tilde{W}_3$ propagators. It is easy to see from Eq. (43) that the Majorana neutrino mass m_L generated when $M_R \ll m_{\text{weak}}$ is of order $(1/8\pi^2)(BM_R/m_{\text{weak}})$.

The expression (43) is valid also for the radiative contribution to m_L obtained in the second (tree-level) scenario of Sec. II, in which no $U(1)_{B-L}$ symmetry is imposed, and generically for the contributions obtained when $M_R \ll m_{\text{weak}}$ and $BM_R \lesssim m_{\text{weak}}^2$. As was already argued in Sec. IV A, acceptable values of m_L arise, in general, when BM_R is not too close to m_{weak}^2 . Although the supersymmetry-breaking parameter B is very large, the approximations made in Sec. IV A, as well as the resulting expression (43), are still applicable.

VI. EFFECTIVE YUKAWA NEUTRINO AND SNEUTRINO INTERACTIONS

The Yukawa interaction terms Higgs boson-lepton doublet right-handed neutrino are also induced radiatively. The diagrams for the neutral-Higgs-boson-left-handed-neutrino-right-handed neutrino interaction term are similar to that in Fig. 1, with the external scalar line being that of a physical neutral Higgs boson. Analogous diagrams generate also the vertex of charged Higgs bosons, charged leptons, and right-handed neutrinos. The interaction terms $\nu_L \nu_L H^0 H^0$ as well as eeH^+H^+ are obtained from variations of the diagram in Fig. 2. The main characteristic of the couplings associated with these terms is that they are momentum dependent with, in general, complicated form factors. Although different from m_D/v and m_L/v^2 , they have, however, the dimensionality and order of magnitude of these ratios.

For a Dirac mass in the eV range, for example, the effective Yukawa coupling is $\sim 10^{-11}$. The smallness of this coupling is now *natural* and, as the smallness of m_D , owes its origin to the seesaw pattern of widely different scales for the neutralinos entering in the loop calculation. In the SM, when the Higgs mechanism is the only way to obtain small neutrino masses, such a small value can only be imposed by hand. In the traditional seesaw mechanism, the couplings for Yukawa interaction terms are of $\mathcal{O}(1)$. While the phenomenological relevance of these terms for possible collider signals is marred by the presence of right-handed fields at scales $\sim 10^{12}$ GeV, these large couplings have subtle consequences

in other sectors. For example, they alter the pattern of Yukawa couplings unification [38]. Moreover, they lift the predictions for lepton-flavor-violating processes involving charged leptons, such as the decay $\mu \rightarrow e \gamma$, by inducing sizable splittings among charged sleptons of different generations. Indeed, these processes, which are realized at the quantum level, may be detectable if the mass of the charged sleptons exchanged in the loops is not too heavy [39].

In the scenarios presented here, with intrinsically small effective Yukawa couplings, large rates for lepton-flavor-violating processes are obtained if a large splitting among the charged-slepton masses of different generations exists already at the Planck scale or at the scale of $U(1)_{B-L}$ breaking. Notice that, although small, these effective Yukawa couplings, obtained radiatively, are, in general, much larger than those obtained at the tree level from the operators in Eq. (2). Indeed, these tree-level Yukawa couplings are naturally $\sim 10^{-13}$; see the discussion in Secs. II and VIII. In some specific cases, however, they may be larger and even comparable to those obtained radiatively. In this case, the actual couplings of neutrino Yukawa interactions are of mixed origin. They remain, however, very small.

The dimensionful coupling m_L/v^2 for the interaction terms $\nu_L \nu_L H^0 H^0$ and eeH^+H^+ is $\sim (10^{13} \text{ GeV})^{-1}$ for $m_L \sim 1 \text{ eV}$. This is the same dimensionful suppression factor that can be obtained by integrating out right-handed neutrinos of mass $M_R \sim 10^{13} \text{ GeV}$ if these couple at the tree level to Higgs and lepton doublets with couplings of $\mathcal{O}(1)$. Couplings that may arise from operators such as those in Eq. (11), in the case of a gauged $U(1)_{B-L}$ symmetry or from operators such as the first in Eq. (1) when $U(1)_{B-L}$ is not introduced, are much smaller than the effective couplings obtained radiatively.

Similar considerations hold also for Higgsino-sneutrino-neutrino operators, which also arise at the quantum level [31]. Their couplings are, in general, so small as to be irrelevant for phenomenology.

Large are, on the contrary, the trilinear sneutrino soft terms A , which can be as large as the electroweak scale. In this case, relatively large interaction terms $AH^0 \tilde{\nu}_R^* \tilde{\nu}_L$ and $AH^+ \tilde{\nu}_R^* \tilde{e}_L$ are present. They may possibly affect searches for neutral and charged Higgs bosons. Particularly interesting is, in this respect, the situation in which the right-handed sneutrino $\tilde{\nu}_R$ is very light. (For the effect of light sneutrinos in searches of the charged Higgs boson, within a generic supersymmetric model, see [40].) A dedicated study of this possibility is certainly important and is left for future work.

VII. PHYSICAL NEUTRINO MASSES

It was shown in the previous sections that small neutrino masses can be naturally obtained in supersymmetric models with right-handed neutrinos, in which (i) the lowest-order neutrino Yukawa operators are forbidden, (ii) effective Yukawa couplings are induced through higher-dimensional operators by a spurion field Z with a supersymmetry-conserving VEV much smaller than the supersymmetry-violating one, and (iii) a $U(1)_{B-L}$ gauge symmetry may be

introduced. In this class of models, contributions to Dirac and Majorana neutrino masses m_D and m_L entering the neutrino mass matrix

$$\begin{pmatrix} m_L & m_D \\ m_D & M_R \end{pmatrix} \quad (44)$$

are generated at the tree level and/or at the quantum level. The overall results obtained for neutrino masses can be classified according to the value of the right-handed neutrino mass M_R .

(1) If M_R is large enough to be hierarchically split from m_{weak} , $M_R \gg m_{\text{weak}}$, the complete or largest contributions to m_L and m_D are, naturally, of radiative origin if the light neutrino states are assumed to be $\sim 1 \text{ eV}$. This case is naturally realized when a $U(1)_{B-L}$ gauge symmetry broken at a large scale M_G is introduced. If $M_R \approx M_G$, the Majorana mass m_L , with its dependence $m_L \sim (m_{\text{weak}}/8\pi^2)(m_{\text{weak}}/M_R)^3$, is much smaller than the Dirac mass m_D , which is $\sim (m_{\text{weak}}/8\pi^2)(m_{\text{weak}}/M_G)^2$. Nevertheless, the eigenstates ν_i have mass dominated by m_L . The eigenstates n_i are, as in the usual seesaw mechanism, at the large scale M_R . The six eigenstates are Majorana fermions. The mixing angle θ_ν between light and heavy states is $|\sin 2\theta_\nu| \sim 2|m_D|/M_R$. Therefore, the states ν_i and n_i are, respectively, almost pure active and sterile neutrinos. Light states ν_i at the eV scale require $M_R(\sim M_G) \sim 100 \text{ TeV}$. This scale, however, can be decreased if the soft parameters A and B are suppressed with respect to m_{weak} . Notice that for light neutrinos in the sub-eV range, the tree-level contribution to m_D may be non-negligible with respect to the radiative one and affect the mixing angle.

(2) If M_R is around the electroweak scale, m_L is, in general, suppressed with respect to m_{weak} only by numerical loop factors. A drastic tuning of A and/or B is then required to reduce m_L to the eV scale. [Small values of these parameters may naturally occur in a gauge-mediated supersymmetry-breaking scenario (see Sec. VIII).] Also this case, as the previous one of large M_R , can be realized when a $U(1)_{B-L}$ gauge symmetry is introduced.

(3) If M_R is much smaller than the electroweak scale, the radiative contribution to m_L is given by $\sim (1/8\pi^2)(BM_R/m_{\text{weak}})$. The Dirac mass m_D has a tree-level contribution $\sim \mathcal{A}_Z v/M_P$ with v an electroweak VEV. In addition, if no $U(1)_{B-L}$ symmetry is introduced, the left- and right-handed Majorana masses $m_L \sim v^2/M_P$ and $M_R \sim \mathcal{A}_Z^2/M_P$ are also generated at the tree level as discussed in the second scenario of Sec. II. If a $U(1)_{B-L}$ gauge symmetry is introduced, on the other hand, we have a radiative contribution to the Dirac mass of $\sim (m_{\text{weak}}/8\pi^2)(m_{\text{weak}}/M_G)^2$. The relative size of M_R with respect to that of m_D has actually rather important physical consequences. Therefore, we further distinguish the following two cases.

(3a) For $M_R \geq m_D$, the six Majorana eigenstates (ν_i, n_i) have masses $\sim m_L + m_D^2/M_R$, M_R , if there is a hierarchy between M_R and m_D , or $(1/2)\{M_R \pm \sqrt{M_R^2 - 4m_D^2}\}$, if M_R is comparable to m_D . The first case can be realized through the

tree-level or the radiative mechanism, or both; the second is typical of the radiative generation of neutrino masses with a $U(1)_{B-L}$ gauge symmetry. The mixing angle between ν_i and n_i ranges from small, for $M_R \gg m_D$, to maximal, for $M_R \sim m_D$. Notice that $M_R \sim 10^{-3}$ eV and $m_D \sim 10^{-5}$ eV could explain the solar neutrino problem with an oscillation $\nu_e \rightarrow \nu_s$ and a small mixing angle. In order to avoid the same pattern of oscillation also in the other two generations, a particular flavor structure of the trilinear A terms and/or of the different M_R has to be implemented.

(3b) The case $M_R \ll m_D$ is typically realized in the radiative mechanism of neutrino Dirac mass generation. In this case, almost degenerate Majorana states (ν_i, n_i) are obtained with a nearly maximal mixing. Their masses are of order $m_D \sim (m_{\text{weak}}/8\pi^2)(m_{\text{weak}}/M_G)^2$, so that a scale of $B-L$ violation of $\sim 10^7$ GeV is required to obtain neutrino masses in the eV range.

(4) Finally, there is the possibility $M_R = 0$, for which the Majorana states (ν_i, n_i) have masses $\pm m_D$ and combine into a Dirac neutrino, for each family. As discussed in the previous sections, in both mechanisms of neutrino-mass generation, this possibility can be realized by imposing an additional symmetry, such as a lepton-number symmetry or its discrete subgroup given in Eq. (3). Notice that purely Dirac neutrinos emerge in this case, even when a $U(1)_{B-L}$ gauge symmetry is imposed. From the phenomenological point of view this case is very different from all the others discussed above. The observed oscillation structure of atmospheric and solar oscillations can then be explained only as a flavor oscillation. It is obviously very difficult to accommodate in such a scenario the results from the LSND experiment. Naturally, no signal is expected from the $\beta\beta$ decay. Moreover, relatively large neutrino dipole moments are predicted, i.e., dipole moments which are not loop suppressed with respect to neutrino masses (see the discussion on quark and lepton dipole moments in a scenario of radiative generation of fermion masses [33]).

Both mechanisms for generating small neutrino masses proposed in this paper offer a very fertile ground in which widely different scenarios of neutrino masses and mixing can be implemented. If the lowest-order tree-level Yukawa couplings exist, then the loops calculated above provide corrections to the tree-level neutrino masses. For unsuppressed Yukawa couplings, i.e., for couplings of $\mathcal{O}(1)$, these corrections are negligibly small once the scale M_R is raised, as in the usual seesaw case, to get tree-level masses of the correct order of magnitude. If tree-level Yukawa couplings are, however, suppressed by some other mechanism, as explained in Secs. II and VIII, then a large variety of neutrino spectra is possible. The fact that the lowest-order neutrino Yukawa operators are vanishing is therefore a key ingredient to unravel a much wider range of possibilities than that offered by the traditional seesaw mechanism.

VIII. MODEL EMBEDDING

In this section, an explicit model that generates the desired hierarchy between the supersymmetry-conserving and the supersymmetry-violating VEV of the Z field, i.e., \mathcal{A}_Z

$\ll \mathcal{F}_Z$, is presented. We recall that \mathcal{A}_Z is responsible for the tree-level generation of neutrino masses, whereas \mathcal{F}_Z gives rise to the sneutrino soft trilinear A parameter. This, together with a neutralino mass, induces quantum contributions to the Dirac mass m_D . The same parameter A , together with the soft bilinear B parameter and a neutralino mass, induces a radiative contribution to the Majorana mass m_L . Comments on the size of the parameters A and B in various supersymmetry-breaking scenarios are also made.

The model is based on a supersymmetric $SU(2)$ gauge theory with four doublet chiral superfields Q_i and six singlet superfields $Z^{ij} = -Z^{ji}$ where i, j are flavor indices ($i, j = 1, \dots, 4$) [41]. All the Q_i and Z^{ij} fields are singlets under the SM gauge groups. Without a superpotential, this theory has a flavor $SU(4)_F$ symmetry. This is explicitly broken to a global $SP(4)_F$ subgroup if the tree-level superpotential [42]

$$W_{\text{tree}} = \lambda Z(QQ) + \lambda_Z Z^a (QQ)_a \quad (45)$$

is introduced. Here, (QQ) and Z are singlets of $SP(4)_F$, while $(QQ)_a$ and Z^a are five-dimensional representations of $SP(4)_F$ ($a = 1, \dots, 5$). [Z and Z^a are certain linear combinations of the Z^{ij} fields, (QQ) and $(QQ)_a$ combinations of the gauge-invariant operators $(Q_i Q_j)$.] The model has a discrete symmetry Z_n that assigns the following charges to the Q_i and Z^{ij} fields:

$$Z_n(Q_i) = \frac{1}{2}, \quad Z_n(Z^{ij}) = -1, \quad (46)$$

and that is identified with the symmetry used in Secs. II and IV to forbid tree-level neutrino Yukawa terms $\bar{N}LH$.⁶ Then, the Z field can couple to the SM fields through the superpotential in Eq. (2).

The dynamics of the $SU(2)$ gauge theory causes the condensation of the Q_i fields through nonperturbative effects [43]. By integrating out the $SU(2)$ gauge fields together with Q_i and Z^a , the low-energy effective superpotential

$$W_{\text{eff}} \simeq \lambda \Lambda^2 Z \quad (47)$$

is obtained for $\lambda_Z \gg \lambda$, where Λ is a dynamical scale of the $SU(2)$ gauge interaction. [Notice that there is a mixed Z_n - $SU(2)$ anomaly and that Λ has therefore a Z_n charge.] Thus, the Z field acquires a nonvanishing F term, $\mathcal{F}_Z \simeq \lambda \Lambda^2 \neq 0$ [41], which induces a sneutrino A parameter of order \mathcal{F}_Z/M_P .

The Z direction is flat at the tree level, but it is lifted by loop corrections. The corrections to the Kähler potential from the strong $SU(2)$ gauge interaction are noncalculable and can only be estimated. They give rise to a quadratic term in Z in the scalar potential. If positive, this term, together with a linear term due to supergravity effects, induces a tiny

⁶Here, it is assumed that quadratic (if $n=2$) and cubic (if $n=3$) terms in Z are absent.

VEV for the scalar component of Z [44]. However, there are also contributions to the mass squared for the Z field near the origin coming from loops of light particles. These are calculable and larger than the noncalculable ones as long as λ is in a perturbative regime [45]. The sign of the mass squared is shown to be positive [45], so that $Z=0$ is a local minimum of the potential in the limit of a global supersymmetry. Thus, the scalar potential for the Z field has the form

$$V \simeq \frac{|\lambda|^4}{16\pi^2} \Lambda^2 |Z|^2 + (\lambda \Lambda^2 m_{3/2} Z + \text{H.c.}), \quad (48)$$

and the nonvanishing VEV for the scalar component of the Z field is

$$A_Z \simeq \frac{16\pi^2}{\lambda^3} m_{3/2}. \quad (49)$$

This VEV, in turn, induces effective tree-level neutrino Yukawa couplings of order A_Z/M_P through the superpotential term in Eq. (2). If $m_{3/2} \simeq 1$ TeV and $\lambda = O(1)$, the resulting Dirac neutrino masses $m_D \simeq 10^{-2}$ eV are sufficiently large to explain the solar neutrino deficit by the MSW mechanism. If λ is somewhat smaller than 1, it is even possible to generate neutrino masses in the range needed by LSND and atmospheric neutrino experiments. Note that this tree-level generation of neutrino masses generically occurs if a chiral superfield Z , which couples to SM fields through the superpotential term in Eq. (2), acquires a nonvanishing auxiliary component at the tree level, but has a vanishing scalar component in the global supersymmetry limit.

Thus, the above model can yield the desired size of the A parameter and of the effective tree-level Yukawa couplings through the supersymmetry-violating and supersymmetry-conserving VEV's of the Z field. On the other hand, a B parameter of the order of the gravitino mass $m_{3/2}$ is always generated by supergravity effects. In addition, if the Z_n symmetry needed to forbid the renormalizable neutrino Yukawa operators is generic (i.e., $n \neq 2$), the B parameter also has a contribution of order \mathcal{F}_Z/A_Z from Eq. (9), which is generically much larger than $m_{3/2}$. (In this case, the value of M_R is, however, rather small.) To summarize, in the present model, we obtain the A and B parameters

$$A \simeq \frac{\Lambda^2}{M_P}, \quad B \simeq \begin{cases} m_{3/2} + \frac{\Lambda^2}{16\pi^2 m_{3/2}} & [\text{for } Z_n \ (n \neq 2)], \\ m_{3/2} & (\text{for } Z_2), \end{cases} \quad (50)$$

and the tree-level neutrino Yukawa couplings y_ν :

$$y_\nu \simeq \frac{16\pi^2 m_{3/2}}{M_P}. \quad (51)$$

Here, we have set λ and the coupling of the superpotential in Eq. (2) to be of order 1. For simplicity, only the case of $B \simeq m_{3/2}$ will be considered in the following discussion.⁷

It is interesting to compare the size that the above parameters acquire in supergravity-mediated [46] and gauge-mediated [47,48] supersymmetry-breaking scenarios. In the first, the gravitino mass $m_{3/2}$, as well as the B parameter, is at the electroweak scale ~ 1 TeV. The effective neutrino Yukawa couplings are of order 10^{-13} and induce Dirac neutrino masses of order $m_D \simeq 10^{-2}$ eV. On the other hand, the A parameter is less constrained. Since the Z field is charged under the Z_n symmetry needed to forbid the lowest-order Yukawa operators, its VEV \mathcal{F}_Z cannot give sizable gaugino masses. Therefore, an additional singlet S with a supersymmetry-violating VEV $\mathcal{F}_S \simeq m_{3/2} M_P$ is needed to give soft supersymmetry-breaking mass to all superparticles. Thus, the phenomenological constraint on the supersymmetry-violating VEV of the Z field, \mathcal{F}_Z , is only $\mathcal{F}_Z \lesssim \mathcal{F}_S$, and this translates into $A \lesssim m_{3/2}$. Although it is logically possible that the A parameter is much smaller than the electroweak scale, we may naturally expect that \mathcal{F}_Z and \mathcal{F}_S are somehow related to each other. If this is the case, the A parameter is also of order of the gravitino mass. This situation was assumed throughout the previous sections.

In a gauge-mediated scenario, the VEV \mathcal{F}_Z of the Z field can be the primary source of supersymmetry breaking. The supersymmetry-breaking effect in the Z field can be transmitted to all the superparticles through gauge interactions [47–49], and no additional singlet is necessary. In this case, both A and B parameters are of the order of the gravitino mass (i.e., $\Lambda^2 \simeq m_{3/2} M_P$). Since in the gauge-mediated scenario $m_{3/2}$ is, in general, much smaller than that in the supergravity-mediated one, the A and B parameters are certainly below the electroweak scale. Thus, the neutrino Yukawa coupling in Eq. (51) cannot give neutrino masses in the range of the various Δm^2 suggested by different experiments, except perhaps for the vacuum-oscillation region of the solar neutrino experiments. A substantial contribution to neutrino masses may come from the radiative mechanism. Notice that all the expressions for m_L and m_D derived in Sec. V still apply to the gauge-mediated case. However, since the values of A and B are now smaller, the values of M_R and of the $(B-L)$ -breaking scale M_G needed to obtain sizable neutrino masses are also lower than those needed in the supergravity-mediated case.

It should be noticed here that, if the Z_n charge -1 is assigned not only to the \bar{N} field but also to all SU(2)-singlet SM fields \bar{U} , \bar{D} , and \bar{E} , then all the SM Yukawa couplings are forbidden and effective Yukawa operators are generated also for quark and charged-lepton fields through higher-

⁷If the discrete symmetry in Sec. IV is generic, i.e., Z_n with $n \neq 2$, it is possible to have $M_R \simeq 0$ and $M_R B \simeq m_{\text{weak}}^2$. Then, the neutrino mass matrix takes the form $\sim ((m_L, m_D)^T, (m_D, 0)^T)$, where m_L is generated by the diagram shown in Fig. 2, and it is suppressed with respect to m_{weak} only by a numerical loop factor if $A^2 \simeq (M_R B) \simeq m_{\text{weak}}^2$.

dimensional operators such as those in Eq. (2). The tree-level contributions that these operators give to Yukawa couplings of quarks and charged leptons are, however, completely negligible. Thus, their masses, as well as Yukawa couplings, are generated radiatively, as in the framework of Ref. [33], through trilinear supersymmetry-breaking parameters induced by \mathcal{F}_Z and gluino and/or neutralino masses. In this case, a unified radiative generation picture for quark and lepton masses emerges. The smallness of neutrino masses with respect to the masses of all other fermions is then easily explained by the fact that neutrinos feel the presence of neutralinos and sneutrinos at the large scales M_G and M_R of $U(1)_{B-L}$ violation. In contrast, the leading contribution to quark and charged-lepton masses comes from the exchange of sfermions and gluinos or sfermions and neutralinos at the electroweak scale.

IX. CONCLUSIONS

In this paper, two different mechanisms to obtain light physical neutrino states in supersymmetric models with three right-handed neutrinos are proposed. It is observed that if the lowest-order neutrino Yukawa operators are forbidden by means of a horizontal discrete symmetry, Yukawa couplings are still induced by higher-dimensional operators suppressed by the Planck mass. In these operators, the three fields that participate in a Yukawa interaction are coupled to a spurion field Z , which acquire a supersymmetry-conserving and a supersymmetry-violating VEV, respectively, \mathcal{A}_Z and \mathcal{F}_Z . If a large hierarchy between the two VEV's is possible, then naturally small Yukawa couplings can be induced at the tree level.

Models where such a hierarchy can be implemented exist and one is explicitly described in this paper. In this model, the Z direction is flat at the tree level. This flatness is lifted by loop corrections, and a supersymmetry-conserving VEV $\mathcal{A}_Z \sim 16\pi^2 m_{3/2}$ is induced by supersymmetry breaking. The strong suppression of neutrino Yukawa couplings y_ν gets naturally linked to the hierarchy between $m_{3/2}$ and M_P : $y_\nu \sim \mathcal{A}_Z/M_P \sim 16\pi^2(m_{3/2}/M_P)$. If Majorana masses for active and sterile neutrinos are forbidden by an additional symmetry such as, for example, lepton number, only Dirac neutrino masses are generated. They are of order $\sim \mathcal{A}_Z v/M_P$, with v an electroweak VEV, i.e., they can reach $\sim 10^{-2}$ eV if all interaction couplings are assumed of $\mathcal{O}(1)$. Intergenerational mass splittings and related oscillations rely on specific textures of these couplings.

Nevertheless, it is possible to forbid only the lowest-order mass term for right-handed neutrinos, but leave allowed higher-dimensional mass operators for left- as well as right-handed neutrinos (see the second tree-level scenario discussed in Sec. II). Besides the Dirac mass m_D given above, also Majorana masses for active and sterile neutrino m_L and M_R are present, respectively of order $\sim v^2/M_P$ and $\sim \mathcal{A}_Z^2/M_P$. Six physical states are generated of Majorana type: ν_i and n_i ($i=1,2,3$). The states n_i correspond mainly to sterile neutrinos and are the heaviest ones. For all interaction couplings of $\mathcal{O}(1)$, their mass is 10^4 – 10^6 times larger than the mass of the states ν_i , which is of order $\sim 10^{-5}$ eV.

Different patterns for different generations may be induced by allowing a nontrivial structure of the interaction couplings.

The second mechanism proposed is that of the radiative generation of neutrino masses via sneutrino-neutralino loops. It is illustrated in the supersymmetrized version of a typical model that leads to the traditional seesaw mechanism: i.e., a supersymmetric model with three right-handed neutrinos and an additional gauge interaction broken at a large scale M_G . Nevertheless, large radiative contributions to the Majorana mass m_L may arise also in models without any additional gauge interaction, but with an explicit violation of lepton number. Also for this mechanism, as in the tree-level one, the chief assumption is the vanishing of the lowest-order neutrino Yukawa operators.

It is shown that (i) the Majorana neutrino mass M_R for right-handed neutrinos is not necessarily of order $\mathcal{O}(M_G)$ and can even be exactly vanishing, and (ii) a seesaw pattern of light-heavy scales is present in the neutralino and (possibly) the sneutrino mass matrix. Dirac and Majorana masses m_D and m_L are induced by trilinear and bilinear soft sneutrino terms as well as neutralino mass terms, which provide, respectively, information on chirality breaking, lepton-number violation, and the correct number of R charges. Apart from a loop suppression factor, ratios of light over heavy scales, present in the neutralino and sneutrino mass matrices, steer m_D and m_L away from the electroweak scale. Since these ratios enter in the expression for m_D and m_L with powers larger than 1, their effectiveness in suppressing m_D and m_L is stronger than in the usual seesaw mechanism, and the scale M_G can be much lower.

Such a mechanism allows a large variety of possible neutrino spectra. In the case of exact vanishing of M_R , three Dirac neutrino states are possible, as in the first scenario of the tree-level mechanism or in a traditional Higgs mechanism with highly tuned Yukawa couplings. These states can have now arbitrary mass depending on the value of M_G . In this case, a texture of neutrino masses consistent with experimental observations finds its origin in the texture of the trilinear A terms for sneutrinos.

In general, for nonvanishing M_R , six Majorana states ν_i and n_i are generated. For heavy M_R , the three states n_i are heavy and no space is left for light sterile neutrinos. For light M_R , the situation is very rich. The six states are all light. The three n_i neutrinos, however, can still be much heavier than the ν_i 's, in which case the mixing angles between n_i 's and ν_i 's are very small. There are finally the other interesting cases in which the ν_i and n_i states are roughly at the same scale or nearly degenerate, with mixing angles ranging from small to maximal. Flavor oscillations can be accommodated in this class of scenarios, as before, i.e., relying on specific textures of the trilinear and bilinear neutrino soft parameters. Oscillations among active and sterile neutrinos are, however, also possible. Whether this is the answer to the puzzle posed by the present results of solar, atmospheric, and reactor neutrino experiments is a question which new experimental data will answer, hopefully, soon. Irrespective of this, it is nontrivial that such oscillations, as well as those between active and much heavier sterile neutrinos advocated in supernova

physics, can be easily accommodated in this class of scenarios.

Since neutrino masses are much smaller than all other masses, it is plausible to assume that the lowest-order neutrino Yukawa operators inducing neutrino masses are vanishing and that the origin of neutrino masses is linked to operators of higher dimensionality or to radiative generation. We have shown that both possibilities are easily implemented in supersymmetric models. The resulting mechanisms encompass the typical Higgs and seesaw mechanisms, although with different realizations, and give also rise to various interesting neutrino spectra.

Note added. While this work was being completed, we received Ref. [50], which makes another proposal for small neutrino masses in supersymmetric models.

ACKNOWLEDGMENTS

The authors thank T. Yanagida for discussions and interest in this work. They also acknowledge discussions with S. Davidson and K. Hagiwara. F.B. thanks the KEK theory group, in particular Y. Okada, for hospitality. The work of Y.N. was supported by the Japanese Society for the Promotion of Science.

-
- [1] For reviews, see B. Kayser, in Particle Data Group [2], p. 307; P. Fisher, B. Kayser, and K. S. McFarland, *Annu. Rev. Nucl. Part. Sci.* **49**, 481 (1999); V. Barger and K. Whisnant, hep-ph/0006235.
 - [2] Particle Data Group, C. Caso *et al.*, *Eur. Phys. J. C* **3**, 1 (1998).
 - [3] Super-Kamiokande Collaboration, Y. Fukuda *et al.*, *Phys. Rev. Lett.* **81**, 1562 (1998).
 - [4] B. T. Cleveland *et al.*, *Astrophys. J.* **496**, 505 (1998); SAGE Collaboration, J. Abdurashitov *et al.*, *Phys. Rev. Lett.* **77**, 4708 (1996); GALLEX Collaboration, W. Hampel *et al.*, *Phys. Lett. B* **447**, 127 (1999); Super-Kamiokande Collaboration, Y. Fukuda *et al.*, *Phys. Rev. Lett.* **82**, 1810 (1999); **82**, 2430 (1999).
 - [5] LSND Collaboration, C. Athanassopoulos *et al.*, *Phys. Rev. Lett.* **77**, 3082 (1996); **81**, 1774 (1998).
 - [6] J. R. Bond, G. Efstathiou, and J. Silk, *Phys. Rev. Lett.* **45**, 1980 (1980); E. Gawiser and J. Silk, *Science* **280**, 1405 (1998); see also P. J. E. Peebles, *Principles of Physical Cosmology* (Princeton University Press, Princeton, 1993).
 - [7] Super-Kamiokande Collaboration, H. Sobel, talk at the “Neutrino 2000” Conference, Sudbury, Canada, 2000.
 - [8] M. Apollonio *et al.*, *Phys. Lett. B* **420**, 397 (1998).
 - [9] L. Wolfenstein, *Phys. Rev. D* **17**, 2369 (1978); **20**, 2634 (1979); S. Mikheyev and A. Yu. Smirnov, *Yad. Fiz.* **42**, 1441 (1985) [*Sov. J. Nucl. Phys.* **42**, 913 (1985)].
 - [10] M. C. Gonzalez-Garcia, *Nucl. Phys. B (Proc. Suppl.)* **95**, 100 (2001).
 - [11] V. N. Gribov and B. M. Pontecorvo, *Phys. Lett.* **28B**, 493 (1969).
 - [12] D. Karlen, in Particle Data Group [2], p. 319.
 - [13] Super-Kamiokande Collaboration, Y. Suzuki, talk at the “Neutrino 2000” Conference, Sudbury, Canada, 2000.
 - [14] M. C. Gonzalez-Garcia and C. Peña-Garay, *Nucl. Phys. B (Proc. Suppl.)* **91**, 80 (2000).
 - [15] G. Mills, talk at the “Neutrino 2000” Conference, Sudbury, Canada, 2000.
 - [16] V. Barger, B. Kayser, J. Learned, T. Weiler, and K. Whisnant, *Phys. Lett. B* **489**, 345 (2000); B. Kayser, hep-ph/0010206; O. L. Peres and A. Y. Smirnov, *Nucl. Phys.* **B599**, 3 (2001).
 - [17] F. Borzumati, K. Hamaguchi, and T. Yanagida, *Phys. Lett. B* **497**, 259 (2001).
 - [18] T. Yanagida, in *Proceedings of the Workshop on the Unified Theory and Baryon Number in the Universe*, edited by O. Sawada and A. Sugamoto, KEK Report No. 79-18, 1979, p. 95; M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, edited by P. van Nieuwenhuizen and D. Z. Freedman (North-Holland, Amsterdam, 1979), p. 315.
 - [19] R. Barbieri, D. V. Nanopoulos, G. Morchio, and F. Strocchi, *Phys. Lett.* **90B**, 91 (1980); R. N. Mohapatra and G. Senjanovic, *Phys. Rev. Lett.* **44**, 912 (1980); T. Yanagida, *Prog. Theor. Phys.* **64**, 1103 (1980); T. Chikashige, G. Gelmini, R. D. Peccei, and M. Roncadelli, *Phys. Lett.* **94B**, 499 (1980).
 - [20] N. Arkani-Hamed, S. Dimopoulos, G. Dvali, and J. March-Russel, hep-ph/9811448; K. R. Dienes, E. Dudas, and T. Gherghetta, *Nucl. Phys.* **B557**, 25 (1999).
 - [21] SNO Collaboration, J. Boger *et al.*, *Nucl. Instrum. Methods Phys. Res. A* **449**, 172 (2000).
 - [22] KARMEN Collaboration, K. Eithel *et al.*, in *ICHEP-98*, Vancouver, edited by A. Astbury, D. Axen, and J. Robinson (World Scientific, Singapore, 1999).
 - [23] E898 Collaboration, E. Church *et al.*, FNAL Report No. Fermilab-P-0898, 1997; N. Armenise *et al.*, CERN Report No. CERN-SPSC/97-21, 1997.
 - [24] A. Kusenko and G. Segre, *Phys. Lett. B* **396**, 197 (1997); *Phys. Rev. D* **59**, 061302 (1999); E. Nardi and J. I. Zuluaga, hep-ph/0006285.
 - [25] X. Shi and G. M. Fuller, *Phys. Rev. D* **59**, 063006 (1999); astro-ph/9812232; D. O. Caldwell, G. M. Fuller, and Y.-Z. Qian, *Phys. Rev. D* **61**, 123005 (2000).
 - [26] S. Dodelson and L. M. Widrow, *Phys. Rev. Lett.* **72**, 17 (1994); X.-D. Shi and G. M. Fuller, *ibid.* **82**, 2832 (1999); M. Drees and D. Wright, hep-ph/0006274.
 - [27] K. Benakli and A. Yu. Smirnov, *Phys. Rev. Lett.* **79**, 4314 (1997).
 - [28] G. Dvali and Y. Nir, *J. High Energy Phys.* **10**, 014 (1998).
 - [29] K. S. Babu and T. Yanagida, *Phys. Lett. B* **491**, 148 (2000); F. Borzumati, K. Hamaguchi, Y. Nomura, and T. Yanagida, in the *Proceedings of the Post Summer Institute Workshop on Neutrino Physics*, Fujiyoshida, Japan, hep-ph/0012118.
 - [30] L. Lavoura, *Phys. Rev. D* **62**, 093011 (2000); T. Kitabayashi and M. Yasue, *Phys. Lett. B* **490**, 236 (2000); hep-ph/0006040.
 - [31] P. Q. Hung, *Phys. Rev. D* **62**, 053015 (2000).
 - [32] See, for example, R. Hempfling, *Nucl. Phys.* **B478**, 3 (1996); F. Borzumati, Y. Grossman, E. Nardi, and Y. Nir, *Phys. Lett. B* **384**, 123 (1996); H.-P. Nilles and N. Polonsky, *Nucl. Phys.* **B484**, 33 (1997); additional references can be found in T.

- Hambye, E. Ma, and U. Sarkar, *ibid.* **B590**, 429 (2000).
- [33] F. Borzumati, G. R. Farrar, N. Polonsky, and S. Thomas, hep-ph/9712428; hep-ph/9805314; Nucl. Phys. **B555**, 53 (1999).
- [34] N. Polonsky and S. Su, Phys. Rev. D **63**, 035007 (2001).
- [35] S. Weinberg, *The Quantum Theory of Fields: Modern Applications* (Cambridge University Press, Cambridge, England, 1996).
- [36] Y. Grossman and H. Haber, Phys. Rev. Lett. **78**, 3438 (1997).
- [37] S. Davidson and S. King, Phys. Lett. B **445**, 191 (1998).
- [38] F. Vissani and A. Yu. Smirnov, Phys. Lett. B **341**, 173 (1994); A. Brignole, H. Murayama, and R. Rattazzi, *ibid.* **335**, 345 (1994); G. K. Leontaris, S. Lola, and G. G. Ross, Nucl. Phys. **B454**, 25 (1995); M. Carena, J. Ellis, S. Lola, and C. E. M. Wagner, Eur. Phys. J. C **12**, 507 (2000).
- [39] F. Borzumati and A. Masiero, Phys. Rev. Lett. **57**, 961 (1986); J. Hisano, T. Moroi, K. Tobe, M. Yamaguchi, and T. Yanagida, Phys. Lett. B **357**, 579 (1995); J. Hisano, T. Moroi, K. Tobe, and M. Yamaguchi, Phys. Rev. D **53**, 2442 (1996); J. Hisano, D. Nomura, and T. Yanagida, Phys. Lett. B **437**, 351 (1998); J. Hisano and D. Nomura, Phys. Rev. D **59**, 116005 (1999); W. Buchmüller, D. Delepine, and F. Vissani, Phys. Lett. B **459**, 171 (1999); J. Ellis, M. Gomez, G. Leontaris, S. Lola, and D. Nanopoulos, Eur. Phys. J. C **14**, 319 (2000); W. Buchmüller, D. Delepine, and L. T. Handoko, Nucl. Phys. **B576**, 445 (2000); G. Barenboim, K. Huitu, and M. Raidal, Phys. Rev. D **63**, 055006 (2001).
- [40] F. Borzumati and A. Djouadi, hep-ph/9806301.
- [41] K. I. Izawa and T. Yanagida, Prog. Theor. Phys. **95**, 829 (1996); K. Intriligator and S. Thomas, Nucl. Phys. **B473**, 121 (1996).
- [42] K. I. Izawa, Y. Nomura, K. Tobe, and T. Yanagida, Phys. Rev. D **56**, 2886 (1997).
- [43] For a review, K. Intriligator and N. Seiberg, Nucl. Phys. B (Proc. Suppl.) **45BC**, 1 (1996).
- [44] K. I. Izawa and T. Yanagida, Prog. Theor. Phys. **94**, 1105 (1995).
- [45] Z. Chacko, M. A. Luty, and E. Ponton, J. High Energy Phys. **12**, 016 (1998).
- [46] A. H. Chamseddine, R. Arnowitt, and P. Nath, Phys. Rev. Lett. **49**, 970 (1982); R. Barbieri, S. Ferrara, and C. A. Savoy, Phys. Lett. **119B**, 343 (1982); L. J. Hall, J. Lykken, and S. Weinberg, Phys. Rev. D **27**, 2359 (1983).
- [47] M. Dine, W. Fischler, and M. Srednicki, Nucl. Phys. **B189**, 575 (1982); S. Dimopoulos and S. Raby, *ibid.* **B192**, 353 (1981); L. Alvarez-Gaumé, M. Claudson, and M. Wise, *ibid.* **B207**, 96 (1982).
- [48] M. Dine and A. E. Nelson, Phys. Rev. D **48**, 1277 (1993); M. Dine, A. E. Nelson, and Y. Shirman, *ibid.* **51**, 1362 (1995); M. Dine, A. E. Nelson, Y. Nir, and Y. Shirman, *ibid.* **53**, 2658 (1996).
- [49] Y. Nomura, K. Tobe, and T. Yanagida, Phys. Lett. B **425**, 107 (1998).
- [50] N. Arkani-Hamed, L. Hall, H. Murayama, D. Smith, and N. Weiner, hep-ph/0006312.