

Hydrostatic pressure of the $O(N)$ ϕ^4 theory in the large N limit

Petr Jizba*

DAMTP, University of Cambridge, Silver Street, Cambridge, CB3 9EW, United Kingdom
and Institute for Theoretical Physics, University of Tsukuba, Ibaraki 305-8571, Japan[†]

(Received 18 November 2002; revised manuscript received 8 September 2003; published 27 April 2004)

With nonequilibrium applications in mind we present in this paper (the first in a series of three) a self-contained calculation of the hydrostatic pressure of the $O(N)$ $\lambda\phi^4$ theory at finite temperature. By combining the Keldysh-Schwinger closed-time path formalism with thermal Dyson-Schwinger equations we compute in the large N limit the hydrostatic pressure in a fully resummed form. We also calculate the high-temperature expansion for the pressure (in $D=4$) using the Mellin transform technique. The result obtained extends the results found by Drummond *et al.* [Nucl. Phys. **B524**, 579 (1998)] and Amelino-Camelia and Pi [Phys. Rev. D **47**, 2356 (1993)]. The latter are reproduced in the limits $m_r(0) \rightarrow 0$, $T \rightarrow \infty$, and $T \rightarrow \infty$, respectively. Important issues of renormalizability of composite operators at finite temperature are addressed and the improved energy-momentum tensor is constructed. The utility of the hydrostatic pressure in the nonequilibrium quantum systems is discussed.

DOI: 10.1103/PhysRevD.69.085011

PACS number(s): 11.10.Wx, 11.10.Gh, 11.15.Pg

I. INTRODUCTION

In order to give a theoretical description of the properties of matter under extreme conditions (such as neutron stars, the early universe or heavy-ion collisions) one is often forced to use statistical quantum field theory (QFT). The latter is due to the inherent quantum nature of these processes and due to an overwhelming number of degrees of freedom involved. In recent years, considerable effort has been devoted to the understanding of both equilibrium and nonequilibrium behavior of such systems (see e.g. [1,2] and citations therein). In fact, the equilibrium description is worked out relatively well and a number of methodologies for doing quantum field theory on systems at or near (local) equilibrium are available. On this level two modes of description have been formulated: the imaginary-time (or Matsubara) approach [3–6] and real-time approach [3–5,7]. In contrast with equilibrium, the theoretical understanding of nonequilibrium quantum field theories is still very rudimentary. The complications involved are essentially twofold. The first is related to the appropriate choice of the nonequilibrium initial-time conditions and their implementation into a quantum description [2,8]. The second problem is to construct the density matrix pertinent to the level of description one aims at. The latter requires usually some sort of coarse-graining (e.g., truncation of higher point Wigner functions in the infinite tower of Schwinger-Dyson equations [9]) or projecting over irrelevant subsystems (incorporated, e.g., via projection operator method [10] or maximal entropy—MaXent—prescription [11]). However, when the density matrix is known one may, in principle, apply the cumulant expansion to convert the calculations into those mimicking usual equilibrium techniques [9,12]. Yet, the boundary problem prohibits *per se* many of equilibrium approaches. Imaginary-

time approach is clearly not applicable due to its lack of the explicit time dependence and build-in equilibrium (Kubo-Martin-Schwinger) boundary conditions. Among the real-time formalisms only the Schwinger-Keldysh or closed-time path formalism (CTP) [3,4,9,13,14] and thermofield dynamics (TFD) [3,15,16] has found a wider utility in nonequilibrium computations. The CTP formulation was conveniently applied, for instance, in the study of nonequilibrium gluon matter [17], cosmological back reaction problem [18] or in the time evolution of a nonequilibrium chiral phase transition [19]. On the other hand, the nonequilibrium TFD was recently used in deriving the transport equations for dense quantum systems [20], or in a study of transport properties of quantum fields with continuous mass spectrum [16].

To extract information on the underlying field dynamics or on nonequilibrium transport characteristics one needs to specify an appropriate set of observables (be it conductivity, damping rates, edge temperature jumps, viscosity, etc.). Pressure is often one of the key parameters used in the diagnostics of off-equilibrium quantum media. Hydrostatic pressure measurements in superfluid He^4 (i.e., in He II phase) [21] and in superconductors [22] provide examples. It is thus clear that an extension of the pressure calculations to nonequilibrium systems could enhance our predicative ability in such areas as (realistic) phase transitions, early universe cosmology or hot fusion dynamics. However, the usual procedure known from equilibrium QFT, i.e., calculations based on the partition function or effective potential [5,23–25] cannot be employed here. This is because the (grand)-canonical potential from which the *thermodynamic* pressure is derived does not exist away from equilibrium. Fortunately, more general definition of pressure, not hinging on existence of (grand)-canonical potential, exists. This is the so called *hydrostatic* pressure and its form is deduced from the expectation value of the energy-momentum tensor. It might be shown that in thermal equilibrium the (classical) thermodynamic and (classical) hydrostatic pressures are identical on account of the (classical) virial theorem [26].

In this and two companion papers we aim at clarifying the calculation of the hydrostatic pressure away from equilibrium and at studying its bearings to various nonequilibrium

*Email address: petr@cm.ph.tsukuba.ac.jp

and p.jizba@damtp.cam.ac.uk

[†]Present address.

situations. Calculation of the expectation value of the energy-momentum tensor is, however, quite delicate task even in thermal equilibrium as computations involved are qualitatively very different from those known, for instance, from the effective action approach. This is because the energy-momentum tensor is a composite operator and as such it requires a different methodology of treatment including a different approach to renormalization issues [3,27]. It should then come as no surprise that in thermal QFT the equivalence between hydrostatic and thermodynamic pressure (or effective action) is more fragile than in corresponding classical statistical systems. In fact, the validity of the *quantum* virial theorem is by no means established conclusively, and it is conjectured that it could break down, for instance, in gauge theories [3]. Besides, there is clearly no virial theorem away from equilibrium (not even classically) and so in such a case one must expect disparity between hydrostatic pressure and effective action.

In order to understand the difficulties involved we concentrate in the present paper on the calculation of the hydrostatic pressure in thermal equilibrium. To this end, we utilize the CTP approach which both in spirit and in many technical details mimics the realistic nonequilibrium calculations [9,11,12,19]. Presented CTP formalism in addition to its theoretical structure which is interesting in its own right, is important because it can be with minor changes directly applied to translationally invariant nonequilibrium QFT systems [11]. In order to keep the discussion as simple as possible we illustrate our reasonings on $O(N)$ symmetric scalar $\lambda\phi^4$ theory. The model is sufficiently simple yet complex enough to serve as an illustration of basic characteristics of the presented method in contrast to other ones in use. The latter has the undeniable merit of being exactly solvable in the large- N limit both at zero and finite temperature [23,28–33]. It might be shown that the leading order approximation in $1/N$ is closely related to the Hartree-Fock mean field approximation which has been much studied in nuclear, many-body, atomic and molecular chemistry applications [23,34]. In addition, in the case of a pure state it corresponds to a Gaussian ansatz for the Schrödinger wave functional [35]. We will amplify some of these points in later papers. We should also emphasize that although the $O(N)$ ϕ^4 theory frequently serves as a useful playground for study of finite-temperature phase transitions with a scalar order parameter, this point is not objective of this work and hence we will not pursue it here.

The setup of the paper is the following: In Sec. II we briefly review the derivation of the thermodynamic and hydrostatic pressures. In Sec. III we lay down the mathematical framework needed for the finite-temperature renormalization of the energy-momentum tensor (for an extensive review on renormalization of composite operators the reader may consult e.g., Refs. [3,36,37]). The latter is discussed on the $O(N)$ ϕ^4 theory. It is a common wisdom that the zero temperature renormalization takes care also of the UV divergences of the corresponding finite temperature theory [3,5,38]. The situation with energy-momentum tensor is, however, more complicated as there is no well defined expectation value of the stress tensor at $T=0$ [3,27]. We show how this problem can be amended at finite temperature. The

key original results obtained here is the prescription for the improved energy-momentum tensor of the $O(N)$ ϕ^4 theory. The latter is achieved by means of the Zimmerman forest formula. With the help of the improved stress tensor we are able to find the corresponding QFT extension of hydrostatic pressure and hence obtain the prescription for the renormalized pressure. This latter result is also original finding. As a byproduct we renormalize ϕ_a^2 and $\phi_a\phi_b$ operators.

Resumed form for the pressure in the large- N limit, together with the discussion of both coupling constant and mass renormalization is presented in Sec. IV. Calculations are substantially simplified by use of the thermal Dyson-Schwinger equations. For simplicity's sake our analysis is confined to the part of the parameter space where the ground state at large N has the $O(N)$ symmetry of the original Lagrangian and the spontaneous symmetry breakdown and Goldstone phenomena are not possible (Bardeen and Moshe's parameter space [31]).

In Sec. V we end up with the high-temperature expansion of the pressure. Calculations are performed in $D=4$ both for massive and massless fields, and the result is expressed in terms of the renormalized mass $m_r(T)$ and the thermal mass shift $\delta m^2(T)$. The expansion is done by means of the Mellin transform technique. In appropriate limits we recover the results of Drummond *et al.* [39] and Amelino-Camelia and Pi [40] for thermodynamic pressure (effective action).

The paper is furnished with two appendixes. In Appendix A we clarify some mathematical manipulations needed in Sec. IV. For the completeness' sake we compute in Appendix B the high-temperature expansion of the thermal-mass shift $\delta m^2(T)$ which will prove useful in Sec. V.

II. HYDROSTATIC PRESSURE

In thermal quantum field theory where one deals with systems in thermal equilibrium there is an easy prescription for a pressure calculation. The latter is based on the observation that for thermally equilibrated systems the grand canonical partition function Z is given as

$$Z = e^{-\beta\Omega} = \text{Tr}(e^{-\beta(H - \mu_i N_i)}), \quad (2.1)$$

where Ω is the grand canonical potential, H is the Hamiltonian, N_i are conserved charges, μ_i are corresponding chemical potentials, and β is the inverse temperature: $\beta = 1/T$ ($k_B=1$). Using identity $\beta(\partial/\partial\beta) = -T(\partial/\partial T)$ together with (2.1) one gets

$$T \left(\frac{\partial \Omega}{\partial T} \right)_{\mu_i, V} = \Omega - E + \mu_i N_i, \quad (2.2)$$

with E and V being the averaged energy and volume of the system, respectively. A comparison of (2.2) with a corresponding thermodynamic expression for the grand canonical potential [3,41–43] requires that entropy $S = -(\partial\Omega/\partial T)_{\mu_i, V}$, so that

$$d\Omega = -S dT - p dV - N_i d\mu_i \Rightarrow p = - \left(\frac{\partial \Omega}{\partial V} \right)_{\mu_i, T}. \quad (2.3)$$

For large systems one can usually neglect surface effects so E and N_i become extensive quantities. Equation (2.1) then immediately implies that Ω is extensive quantity as well, so Eq. (2.3) simplifies to

$$p = - \frac{\Omega}{V} = \frac{\ln Z}{\beta V}. \quad (2.4)$$

The pressure defined by Eq. (2.4) is so called thermodynamic pressure.

Since $\ln Z$ can be systematically calculated summing up all connected closed diagrams (i.e., bubble diagrams) [3,44,45], the pressure calculated via Eq. (2.4) enjoys a considerable popularity [39,3,4,46]. Unfortunately, the latter procedure can not be extended to out of equilibrium as there is, in general, no definition of the partition function Z nor grand-canonical potential Ω away from an equilibrium.

Yet another, alternative definition of a pressure not hinging on thermodynamics can be provided; namely the hydrostatic pressure which is formulated through the energy-momentum tensor $\Theta^{\mu\nu}$. The formal argument leading to the hydrostatic pressure in D space-time dimensions is based on the observation that $\langle \Theta^{0j}(x) \rangle$ is the mean (or macroscopic) density of momenta \mathbf{p}^j in the point x^μ . Let \mathbf{P} be the mean total $(D-1)$ -momentum of an infinitesimal volume $V^{(D-1)}$ centered at \mathbf{x} , then the rate of change of j -component of \mathbf{P} reads

$$-\frac{d\mathbf{P}^j(x)}{dt} = \int_{V^{(D-1)}} d^{D-1}\mathbf{x}' \frac{\partial}{\partial x^0} \langle \Theta^{0j}(x^0, \mathbf{x}') \rangle = \sum_{i=1}^{D-1} \int_{\partial V^{(D-1)}} ds^i \langle \Theta^{ij} \rangle. \quad (2.5)$$

In the second equality we have exploited the continuity equation for $\langle \Theta^{\mu j} \rangle$ and successively we have used Gauss' theorem.¹ The $\partial V^{(D-1)}$ corresponds to the surface of $V^{(D-1)}$.

Anticipating a system out of equilibrium, we must assume a nontrivial distribution of the mean particle four-velocity $U^\mu(x)$ (hydrodynamic velocity). Now, a pressure is by definition a scalar quantity. This particularly means that it should not depend on the hydrodynamic velocity. We must thus go to the local rest frame and evaluate pressure there. However, in the local rest frame, unlike the equilibrium, the notion of a pressure acting equally in all directions is lost. In order to retain the scalar character of pressure, one customarily defines the *pressure at a point* [in the following denoted as

$p(x)$] [48], which is simply the “averaged pressure”² over all directions at a given point. In the local rest frame Eq. (2.5) describes j -component of the force exerted by the medium on the infinitesimal volume $V^{(D-1)}$. [By definition there is no contribution to $d\mathbf{P}^j(x)/dt$ caused by the particle convection through $V^{(D-1)}$.] Averaging the left-hand side (LHS) of Eq. (2.5) over all directions of the normal $\mathbf{n}(\mathbf{x})$, we get³

$$\begin{aligned} & \frac{1}{(S_1^{D-2})} \sum_{j=1}^{D-1} \int \frac{d\mathbf{P}^j(x)}{dt} \mathbf{n}^j d\Omega(\mathbf{n}) \\ &= \frac{1}{(S_1^{D-2})} \sum_{j,i=1}^{D-1} \int_{\partial V^{(D-1)}} ds \langle \Theta^{ij}(x') \rangle \int d\Omega(\mathbf{n}) \mathbf{n}^i \mathbf{n}^j \\ &= - \frac{1}{(D-1)} \sum_{i=1}^{D-1} \int_{\partial V^{(D-1)}} ds \langle \Theta_i^i(x') \rangle, \end{aligned} \quad (2.6)$$

where $d\Omega(\mathbf{n})$ is an element of solid angle about \mathbf{n} and S_1^{D-2} is the surface of $(D-2)$ -sphere with unit radius ($\int d\Omega(\mathbf{n}) = S_1^{D-2} = 2\pi^{(D-1)/2}/\Gamma[(D-1)/2]$). On the other hand, from the definition of the pressure at a point x^μ we might write

$$(S_1^{D-2})^{-1} \sum_{j=1}^{D-1} \int \frac{d\mathbf{P}^j(x)}{dt} \mathbf{n}^j d\Omega(\mathbf{n}) = -p(x) \int_{\partial V^{(D-1)}} ds, \quad (2.7)$$

here the minus sign reflects that the force responsible for a compression (conventionally assigned as a positive pressure) has reversed orientation than the surface normals \mathbf{n} (pointing outward). In order to keep track with the standard textbook definition of a sign of a pressure [42,48] we have used in Eq. (2.7) the normal \mathbf{n} in a contravariant notation (note, $\mathbf{n}^i = -\mathbf{n}_i$). Comparing Eq. (2.6) with Eq. (2.7) we can write for a sufficiently small volume $V^{(D-1)}$,

$$p(x) = - \frac{1}{(D-1)} \sum_{i=1}^{D-1} \langle \Theta_i^i(x) \rangle. \quad (2.8)$$

We should point out that in equilibrium the thermodynamic pressure is usually identified with the hydrostatic one via the virial theorem [3,49]. In the remainder of this note we shall deal with the hydrostatic pressure at equilibrium. We shall denote the foregoing as $\mathcal{P}(T)$, where T stands for temperature. We consider the nonequilibrium case in a future paper.

III. RENORMALIZATION

If we proceed with Eq. (2.8) to QFT this leads to the notorious difficulties connected with the fact that $\Theta^{\mu\nu}$ is a

¹The macroscopic conservation law for $\langle \Theta^{\mu\nu} \rangle$ (i.e., the continuity equation) has to be postulated. For some systems, however, the later can be directly derived from the corresponding microscopic conservation law [47].

²To be precise, we should talk about averaging the normal components of stress [48].

³The angular average is standardly defined for scalars (say, A) as $\int A d\Omega(\mathbf{n}) / \int d\Omega(\mathbf{n})$, and for vectors (say, \mathbf{A}^i) as $\Sigma_j \int \mathbf{A}^j \mathbf{n}^j d\Omega(\mathbf{n}) / \int d\Omega(\mathbf{n})$. Similarly we might write the angular averages for tensors of a higher rank.

(local) composite operator. If only a free theory would be in question then the normal ordering prescription would be sufficient to render $\langle \Theta^{\mu\nu} \rangle$ finite. In the general case, when the interacting theory is of interest, one must work with the Zimmermann *normal* ordering prescription instead. Let us demonstrate the latter on the $O(N)$ ϕ^4 theory. (In this section we keep N arbitrary.) Such a theory is defined by the bare Lagrange function

$$\mathcal{L} = \frac{1}{2} \sum_{a=1}^N ((\partial \phi_a)^2 - m_0^2 \phi_a^2) - \frac{\lambda_0}{8N} \left(\sum_{a=1}^N (\phi_a)^2 \right)^2, \quad (3.1)$$

and we assume that $m_0^2 > 0$. The corresponding *canonical* energy-momentum tensor is given by

$$\Theta_c^{\mu\nu} = \sum_a \partial^\mu \phi_a \partial^\nu \phi_a - g^{\mu\nu} \mathcal{L}. \quad (3.2)$$

The Feynman rules for Green's functions with the energy-momentum insertion can be easily explained in momentum space. In the reasonings to follow we shall need the (thermal) composite Green's function⁴

$$D^{\mu\nu}(x^n|y) = \langle T^* \{ \phi_r(x_1) \dots \phi_r(x_n) \Theta_c^{\mu\nu}(y) \} \rangle. \quad (3.3)$$

Here the subscript r denotes the renormalized fields in the Heisenberg picture (the internal indices are suppressed) and T^* is the so-called T^* product (or covariant T product) [24,50–52]. The T^* product is defined in such a way that it is simply the T product with all differential operators \mathcal{D}_{μ_i} pulled out of the T -ordering symbol, i.e.,

$$\begin{aligned} T^* \{ \mathcal{D}_{\mu_1}^{x_1} \phi_r(x_1) \dots \mathcal{D}_{\mu_n}^{x_n} \phi_r(x_n) \} \\ = \mathcal{D}(i\partial_{\{\mu\}}) T \{ \phi_r(x_1) \dots \phi_r(x_n) \}, \end{aligned} \quad (3.4)$$

where $\mathcal{D}(i\partial_{\{\mu\}})$ is just a useful short-hand notation for $\mathcal{D}_{\mu_1}^{x_1} \mathcal{D}_{\mu_2}^{x_2} \dots \mathcal{D}_{\mu_n}^{x_n}$. In the case of thermal Green's functions, T represents a contour ordering symbol [3–5]. It is the mean value of the T^* ordered fields rather than the T ones, which corresponds at $T=0$ and at equilibrium to the Feynman path integral representation of Green's functions [52,53].

A typical contribution to $\Theta_c^{\mu\nu}(y)$ can be written as

$$\mathcal{D}_{\mu_1} \phi(y) \mathcal{D}_{\mu_2} \phi(y) \dots \mathcal{D}_{\mu_n} \phi(y), \quad (3.5)$$

so the typical term in Eq. (3.3) is

$$\begin{aligned} \mathcal{D}(i\partial_{\{\mu\}}) \langle T^* \{ \phi_r(x_1) \dots \phi_r(x_n) \phi(y_1) \dots \phi(y_k) \} \rangle \Big|_{y_i=y} \\ \equiv D_{\{\mu\}}(x^n|y^k) \Big|_{y_i=y}. \end{aligned}$$

⁴By ϕ we shall mean the field in the Heisenberg picture. The subscript H will be introduced in cases when a possible ambiguity could occur.




FIG. 1. The graphical representation of $D^{\mu\nu}(p^n|p)$.

Performing the Fourier transform in Eq. (3.3) we get

$$\begin{aligned} D^{\mu\nu}(p^n|p) = \sum_{k=\{2,4\}} \int \left(\prod_{i=1}^k \frac{d^D q_i}{(2\pi)^D} \right) (2\pi)^D \delta^D \left(p - \sum_{j=1}^k q_j \right) \\ \times \mathcal{D}_{(k)}^{\mu\nu}(q_{\{\mu\}}) D(p^n|q^k), \end{aligned} \quad (3.6)$$

where $\mathcal{D}_{(k)}^{\mu\nu}(\dots)$ is a Fourier transformed differential operator corresponding to the quadratic ($k=2$) and quartic ($k=4$) terms in $\Theta_c^{\mu\nu}$. Denoting the new vertex corresponding to $\mathcal{D}_{(k)}^{\mu\nu}(\dots)$ as \otimes , we can graphically represent Eqs. (3.3) through (3.6) as Fig. 1.

For the case at hand one can easily read off from Eq. (3.2) an explicit form of the bare composite vertices, the foregoing are

$$\begin{aligned} \text{Diagram 1: } \otimes \sim \mathcal{D}_{(2)}^{\mu\nu}(q_{\{\mu\}}) = \frac{1}{2} \delta_{ab} \{ 2(q_1 - p)^\mu q_1^\nu \\ - g^{\mu\nu} ((q_1 - p)_\lambda q_1^\lambda - m_0^2) \}, \end{aligned}$$

$$\begin{aligned} \text{Diagram 2: } \otimes \sim \mathcal{D}_{(4)}^{\mu\nu}(q_{\{\mu\}}) = \frac{g^{\mu\nu} \lambda_0}{8N} \{ 2(\delta_{ab} \delta_{cd} + \delta_{ac} \delta_{bd} \\ + \delta_{ad} \delta_{bc}) - 5 \delta_{ab} \delta_{cd} \delta_{ac} \}. \end{aligned}$$

(For the internal indices we do not adopt Einstein's summation convention.) The blobs in Fig. 1 comprise the sum of all $n+2$ - and $n+4$ - (not necessarily connected) Green functions. As usual, the disjoint bubble diagrams in Green functions (blobs) can be divided out from the very beginning. We have also implicitly assumed that the summation over internal indices is understood.

In case when we deal with finite temperature, we choose the contour ordering in Eq. (3.3) to run along the time contour depicted in Fig. 3. It is possible to show that for Green's function calculations only horizontal paths contribute [14,54,55]. In addition, the “physical” fields occurring on the external lines of Green's functions have time arguments on the upper horizontal path (type-1 fields) while the “ghost” fields have time arguments on the lower horizontal path (type-2 fields). The latter modify the Feynman rules in a

nontrivial fashion [4,5,14]. From the foregoing discussion should be clear that in the case of thermal composite Green's function, the new (composite) vertices are of type-1 as the fields from which they are deduced are all physical.⁵

A. Renormalization of $\phi_a(x)\phi_b(x)$

Now, if there would be no $\Theta_c^{\mu\nu}$ insertion in Eq. (3.3), the latter would be finite, and so it is natural to define the renormalized energy-momentum tensor [$\Theta_c^{\mu\nu}$] (or Zimmermann normal ordering) in such a way that

$$D_r^{\mu\nu}(x^n|y) = \langle T^* \{ \phi_r(x_1) \dots \phi_r(x_n) [\Theta_c^{\mu\nu}] \} \rangle,$$

is finite for any $n > 0$. To see what is involved, we illustrate the mechanism of the composite operator renormalization on $\phi_a(x)\phi_b(x)$ first, the energy-momentum tensor case will be postponed to Sec. III B. In the following we shall use the mass-independent renormalization, and for definiteness we chose the minimal subtraction scheme (MS). In MS we can expand the bare parameters into the Laurent series which has a simple form [24,37,53], namely

$$\lambda_0 = \mu^{4-D} \lambda_r \left(1 + \sum_{k=1}^{\infty} \frac{a_k(\lambda_r; D)}{(D-4)^k} \right), \quad (3.7)$$

$$m_0^2 = m_r^2 \left(1 + \sum_{k=1}^{\infty} \frac{b_k(\lambda_r; D)}{(D-4)^k} \right). \quad (3.8)$$

Here a_0 and b_0 are analytic in $D=4$. The parameter μ is the scale introduced by the renormalization in order to keep λ_r dimensionless. An important point is that both a_k 's and b_k 's are mass, temperature, and momentum independent.

It was Zimmermann who first realized that the forest formula known from the ordinary Green's function renormalization [24,36] can be also utilized for the composite Green's functions rendering them finite [36,56]. That is, we start with Feynman diagrams expressed in terms of physical (i.e., finite) coupling constants and masses. As we calculate diagrams to a given order, we meet UV divergences which might be cancelled by adding counterterm diagrams. The forest formula then prescribes how to systematically cancel all the UV loop divergences by counterterms to all orders. However, in contrast to the coupling constant renormalization, the composite vertex need not to be renormalized multiplicatively. We shall illustrate this fact in the sequel. Let us also observe that in the lowest order (no loop) the renormalized composite vertex equals to the bare one, and so to that order $A = [A]$, for any composite operator A .

Now, from Eqs. (3.7) and (3.8) follows that for any function $F = F(m_r, \lambda_r)$ we have

$$\frac{\partial F}{\partial m_r^2} = \frac{\partial m_0^2}{\partial m_r^2} \frac{\partial F}{\partial m_0^2} = \frac{m_0^2}{m_r^2} \frac{\partial F}{\partial m_0^2}.$$

⁵For a brief introduction to the real-time formalism in thermal QFT see, for example, Refs. [3,4,7].

So particularly for

$$F = D(x_1, \dots, x_n) = \langle T^* \{ \phi_r(x_1) \dots \phi_r(x_n) \} \rangle,$$

one reads

$$\begin{aligned} m_r^2 \frac{\partial}{\partial m_r^2} D(x_1, \dots, x_n) &= m_0^2 \frac{\partial}{\partial m_0^2} D(x_1, \dots, x_n) \\ &= \left(-\frac{i}{2} \right) \mathcal{N} \int d^D x \sum_{a=1}^N \int \mathcal{D} \phi \phi_r(x_1) \dots \\ &\quad \times \phi_r(x_n) m_0^2 \phi_a^2(x) \exp(iS[\phi, T]) \\ &= \left(-\frac{i}{2} \right) \int d^D x \sum_{a=1}^N D_a(x_1, \dots, x_n | x; m_0^2). \end{aligned} \quad (3.9)$$

Here \mathcal{N}^{-1} is the standard denominator of the path integral representation of Green's function. We should apply the derivative also on \mathcal{N} but this would produce disconnected graphs with bubble diagrams. The former precisely cancel the very same disconnected graphs in the first term, so we are finally left with no bubble diagrams in Eq. (3.9). In the Fourier space Eq. (3.9) reads

$$m_r^2 \frac{\partial}{\partial m_r^2} D(p_1, \dots, p_n) = \left(-\frac{i}{2} \right) \sum_{a=1}^N D_a(p_1, \dots, p_n | 0; m_0^2). \quad (3.10)$$

As the LHS is finite there cannot be any pole terms on the right-hand side (RHS) either, and so $\sum_a m_0^2 \phi_a^2$ is by itself a renormalized composite operator. We see that m_0^2 precisely compensates the singularity of $\sum_{a=1}^N \phi_a^2$.

Now, it is well known that any second-rank tensor (say M_{ab}) can be generally decomposed into three irreducible tensors; an antisymmetric tensor, a symmetric traceless tensor and an invariant tensor. Let us set $M_{ab} = \phi_a \phi_b$, so the symmetric traceless tensor K_{ab} reads

$$K_{ab}(x) = \phi_a(x) \phi_b(x) - (\delta_{ab}/N) \sum_{c=1}^n \phi_c^2(x), \quad (3.11)$$

while the invariant tensor I_{ab} is

$$I_{ab}(x) = (\delta_{ab}/N) \sum_{c=1}^N \phi_c^2(x).$$

Because the renormalized composite operators have to preserve a tensorial structure of the bare ones, we immediately have that

$$K_{ab} = A_1 [K_{ab}] \quad \text{and} \quad I_{ab} = A_2 [I_{ab}], \quad (3.12)$$


where both A_1 and A_2 must have structure $(1 + \Sigma(\text{poles}))$. The foregoing guarantees that to the lowest order K_{ab}

$=[K_{ab}]$ and $I_{ab}=[I_{ab}]$. As we saw in Eq. (3.10), $m_0^2 I_{ab}$ is renormalized, and so from Eq. (3.12) follows that $m_0^2 I_{ab} = C[I_{ab}]$. Here C has dimension $[m^2]$ and is analytic in $D=4$. We can uniquely set $C=m_r^2$ because only this choice fulfills the lowest order condition $I_{ab}=[I_{ab}]$ [cf. Eq. (3.8)]. Collecting our results together we might write

$$\sum_c \phi_c^2 = Z_{\Sigma\phi^2} \left[\sum_c \phi_c^2 \right] = Z_{\Sigma\phi^2} \sum_c [\phi_c^2], \quad (3.13)$$

with $Z_{\Sigma\phi^2} = A_2 = m_r^2/m_0^2$. In the second equality we have used an obvious linearity [36] of $[\dots]$. From Eqs. (3.11) and (3.13) follows that

$$\phi_a(x)\phi_b(x) = A_1[\phi_a(x)\phi_b(x)] - \frac{\delta_{ab}}{N}(A_1 - Z_{\Sigma\phi^2}) \sum_{c=1}^N [\phi_c^2(x)]. \quad (3.14)$$



where

$$\textcircled{1} \textcircled{1} = \left\{ -\frac{1}{4} \textcircled{1} \textcircled{1} - \frac{1}{4} \textcircled{1} \textcircled{1} \right\} \quad (3.16)$$

$$\textcircled{2} \textcircled{2} = \left\{ -\frac{1}{12} \textcircled{1} \textcircled{1} - \frac{1}{12} \textcircled{1} \textcircled{1} \right\}$$

Here cross-hatched blobs refer to (renormalized) 1PI $(n+2)$ -point Green's function, circled indices mark a type of the field propagated on the indicated line, and uncircled numbers refer to thermal indices (we explicitly indicate only relevant thermal indices). The counterterms, symbolized by a heavy dot, are extracted from the boxed diagrams (elementary Zimmermann forests). In MS scheme one gets the following results:

$$\textcircled{1} \textcircled{1} = \frac{i\lambda_r \mu^{4-D}}{4} \int \frac{d^D q}{(2\pi i)^D} \{ \mathbb{D}_{11}(q) \mathbb{D}_{11}(-q) - \mathbb{D}_{12}(q) \mathbb{D}_{12}(-q) \} |_{\text{MS pole term}}$$

So particularly for ϕ_a^2 one reads

$$\phi_a^2 = \frac{1}{N} ((N-1)A_1 + Z_{\Sigma\phi^2}) [\phi_a^2] - \frac{1}{N} (A_1 - Z_{\Sigma\phi^2}) \sum_{c \neq a} [\phi_c^2]. \quad (3.15)$$

From the discussion above it does not seem to be possible to obtain more information about A_1 without doing an explicit perturbative calculations, however it is easy to demonstrate that $A_1 \neq Z_{\Sigma\phi^2}$. To show this, let us consider the simplest nontrivial case; i.e., $N=2$, and calculate A_1 to order λ_r . For that we need to discuss the renormalization of the n -point composite Green's function with, say, ϕ_1^2 insertion. To do that, it suffices to discuss the renormalization of the corresponding 1PI n -point Green's function. The perturbative expansion for the composite vertex to order λ_r can be easily generated via the Dyson-Schwinger (DS) equation [57] and it reads⁶

$$= -\frac{1}{4} \partial_{m_r^2} \left(\frac{\Gamma(1-\frac{D}{2})}{(4\pi)^{D/2}} \lambda_r \mu^{4-D} m_r^{D-2} \right) \Big|_{\text{MS}} = -\lambda_r \mu^{4-D} / 2(D-4)(4\pi)^2$$

$$\textcircled{2} \textcircled{2} = -\lambda_r \mu^{4-D} / 6(D-4)(6\pi)^2.$$

Here \mathbb{D}_{11} and \mathbb{D}_{12} are the usual thermal propagators in the real-time formalism [3,4,7] (see also Sec. IV). From Eq. (3.16) we can directly read off that

$$[\phi_1^2] = \left(1 - \frac{\lambda_r \mu^{4-D}}{2(D-4)(4\pi)^2} + \mathcal{O}(\lambda_r^2) \right) \phi_1^2 + \left(-\frac{\lambda_r \mu^{4-D}}{6(D-4)(4\pi)^2} + \mathcal{O}(\lambda_r^2) \right) \phi_2^2.$$

⁶Throughout the paper we accept the usual convention: Ordinary (not necessarily connected) N -point Green's functions are represented with dotted blobs with N external legs, connected N -point Green's functions are represented with hatched blobs with N external legs and 1PI N -point Green's functions are represented with cross hatched blobs with N truncated legs (represented by solid circles in vertices).

FIG. 2. Counterterm renormalization of the last two diagrams in Eq. (3.17). (Cut legs indicate amputations.)

As the coefficient before ϕ_2^2 is not zero, we conclude that $A_1 \neq Z_\Sigma \phi_2$. It is not a great challenge to repeat the previous calculations for the $\phi_1 \phi_2$ insertion. The latter gives

$$A_1 = 1 - \frac{\lambda_r \mu^{4-D}}{3(D-4)(4\pi)^2} + \mathcal{O}(\lambda_r^2).$$

Equation (3.15) exhibits the so-called operator mixing [24]; the renormalization of ϕ_a^2 cannot be considered independently of the renormalization of ϕ_c^2 ($c \neq a$). The latter is a general feature of composite operator renormalization. Note, however, that $\phi_a \phi_b$ ($a \neq b$) do not mix by renormal-

ization, i.e., they renormalize multiplicatively. It can be shown that composite operators mix under renormalization only with those composite operators which have dimension less or equal [24,36,56].

Unfortunately, if we apply the previous arguments to $nn=0$, the result is not finite; another additional renormalization must be performed. The fact that the expectation values of $[\dots]$ are generally UV divergent, in spite of being finite for the composite Green's functions,⁷ can be nicely illustrated with the composite operator $[\phi^2]$ in the $N=1$ theory. Taking the diagrams for $D(0|0)$ and applying successively the (unrenormalized) DS equation [3,57] we get

Equation (3.17) might be rewritten as

$$\begin{aligned} D(0|0) &= D(0|0)|_{\lambda_r^0} + \frac{1}{2} \int \frac{d^D q_1}{(2\pi)^D} \frac{d^D q_2}{(2\pi)^D} \delta^D(q_1 + q_2) \\ &\quad \times D^{\text{amp}}(q^2|0)|_{\lambda_r} D(q^2) \\ &\quad + \frac{1}{36} \int \prod_{i=1}^6 \frac{d^D q_i}{(2\pi)^D} \delta^D\left(\sum_{j=1}^6 q_j\right) \\ &\quad \times D^{\text{amp}}(q^6|0)|_{\lambda_r} D(q^6), \end{aligned} \quad (3.18)$$

where $D^{\text{amp}}(q^m|0)|_{\lambda_r^k}$ is the m -point amputated composite

Green's function to order λ_r^k , and $D(q^m)$ is the full m -point Green's function. The crucial point is that we can write $D(0|0)$ as a sum of terms, which, apart from the first (free field) diagram, are factorized to the product of the composite Green's function with $n > 0$ and the full Green's function. [The factorization is represented in Eq. (3.17) by the dashed lines.] Note that the expansion Eq. (3.17) is not unique as various other ways of pulling vertices out of Green's function may be utilized but this particular form will prove to be important in the next section [see Eq. (3.26)].

Now, utilizing the counterterm renormalization to the last two diagrams in Eq. (3.17) we get situation depicted in Fig. 2. Terms inside of the parentheses are finite, this is because

⁷Also called the matrix elements of $[\dots]$.

both the composite Green's functions ($n \geq 2$!) and the full Green's functions are finite after renormalization. The counterterm diagrams, which appear on the RHS of the parentheses, precisely cancel the UV divergences coming from the loop integrations over momenta $q_1 \dots q_i$ which must be finally performed. The heavy dots schematically indicates the corresponding counterterms. In the spirit of the counterterm renormalization we should finally subtract the counterterm associated with the overall superficial divergence⁸ related to the diagrams in question. But as we saw this is not necessary; individual counterterm diagrams (Zimmermann forests) mutually cancel their divergences leaving behind a finite result.

So the only UV divergence in Eq. (3.17) which cannot be cured by existing counterterms is that coming from the first (i.e., free field or ring) diagram. The foregoing divergence is evidently temperature independent (to see that, simply use an explicit form of the free thermal propagator D_{11}). Hence, if we define

$$\langle \phi^2 \rangle_{\text{renorm}} = \langle [\phi^2] \rangle - \langle 0 | [\phi^2] | 0 \rangle, \quad (3.19)$$

or, alternatively

$$\langle \phi^2 \rangle_{\text{renorm}} = \langle [\phi^2] \rangle - \langle [\phi^2] \rangle|_{\text{free fields}}, \quad (3.20)$$

we get finite quantities, as desired. On the other hand, we should emphasize that

$$\begin{aligned} \langle \phi^2 \rangle - \langle 0 | \phi^2 | 0 \rangle &= Z_{\phi^2} \{ \langle [\phi^2] \rangle - \langle 0 | [\phi^2] | 0 \rangle \} \\ &\neq \text{finite in } D=4. \end{aligned} \quad (3.21)$$

An extension of the previous reasonings to any $N > 1$ is straightforward, only difference is that we must deal with operator mixing which makes Eqs. (3.19) and (3.20) less trivial.

The important lesson which we have learned here is that the naive “double dotted” normal product (i.e., subtraction of the vacuum expectation value from a given operator) does not generally give a finite result. The former is perfectly suited for the free theory ($Z_{\Sigma \phi^2} = 1$) but in the interacting case we must resort to the prescription Eq. (3.19) or (3.20) instead.

B. Renormalization of the energy-momentum tensor

In order to calculate the hydrostatic pressure, we need to find such $\langle \Theta_c^{\mu\nu} \rangle|_{\text{renorm}}$ which apart from being finite is also consistent with our derivation of the hydrostatic pressure introduced in the introductory section. In view of the previous treatment, we however cannot, however, expect that $\Theta_c^{\mu\nu}$ will be renormalized multiplicatively. Instead, new terms with a different structure than $\Theta_c^{\mu\nu}$ itself will be generated

during renormalization. The latter must add up to $\Theta_c^{\mu\nu}$ in order to render $D^{\mu\nu}(x^n|y)$ finite.⁹

Now, the key ingredient exploited in Eq. (2.5) is the conservation law (continuity equation). It is well known that one can “modify” $\Theta_c^{\mu\nu}$ in such a way that the new tensor $\Theta^{\mu\nu}$ preserves the convergence properties of $\Theta_c^{\mu\nu}$. Such a modification (the Pauli transformation) reads

$$\begin{aligned} \Theta^{\mu\nu} &= \Theta_c^{\mu\nu} + \partial_\lambda X^{\lambda\mu\nu}, \\ X^{\lambda\mu\nu} &= -X^{\mu\lambda\nu}. \end{aligned} \quad (3.22)$$

For scalar fields Eq. (3.22) is the only transformation which neither changes the divergence properties of $\Theta_c^{\mu\nu}$ nor the generators of the Poincare group constructed out of $\Theta_c^{\mu\nu}$ [3,24,47,51]. Because the renormalized (or improved) energy momentum tensor must be conserved (otherwise theory would be anomalous), it has to mix with $\Theta_c^{\mu\nu}$ under renormalization only via the Pauli transformation, i.e.,

$$[\Theta_c^{\mu\nu}] = \Theta_c^{\mu\nu} + \partial_\lambda X^{\lambda\mu\nu}. \quad (3.23)$$

In order to determine $X^{\lambda\mu\nu}$, we should realize that its role is to cancel divergences present in $\Theta_c^{\mu\nu}$. Such a cancellation can be, however, performed only by means of composite operators which are even in the number of fields (note that $\Theta_c^{\mu\nu}$ is even in fields and Green's functions with the odd number of fields vanish). Recalling the condition that renormalization can mix only operators with dimension less or equal, we see that the dimension of $X^{\lambda\mu\nu}$ must be $D-1$, and that $X^{\lambda\mu\nu}$ must be quadratic in fields. The only possible form which is compatible with tensorial structure Eq. (3.22) is then

$$X^{\lambda\mu\nu} = \sum_{a,b=1}^N c_{ab}(\lambda_r; D) (\partial^\mu g^{\lambda\nu} - \partial^\lambda g^{\mu\nu}) \phi_a \phi_b. \quad (3.24)$$

From the fact that $\Theta_c^{\mu\nu}$ and $[\Theta_c^{\mu\nu}]$ are $O(N)$ invariant [see Eq. (3.2)], $\partial_\lambda X^{\lambda\mu\nu}$ must be also $O(N)$ invariant, so $c_{ab} = \delta_{ab}c$. Thus, finally we can write

$$[\Theta_c^{\mu\nu}] = \Theta_c^{\mu\nu} + c(\lambda_r; D) \sum_{a=1}^N (\partial^\mu \partial^\nu - g^{\mu\nu} \partial^2) \phi_a^2, \quad (3.25)$$

with $c = c_0 + \Sigma(\text{poles})$, here c_0 is analytic in D . Structure of $c(\lambda_r; D)$ could be further determined, similarly as in the $N=1$ theory, employing a renormalization group equation [37]. We do not intend to do that as the detailed structure of

⁸A simple power counting in the ϕ^4 theory reveals [24] that for a composite operator A with dimension ω_A the superficial degree of divergence ω corresponding to an n -point diagram is $\omega = \omega_A - n$.

⁹In fact it can be shown [3,36] that the Noether currents corresponding to a given internal symmetry are renormalized, i.e., $J^a = [J^a]$, however, this is not the case for the Noether currents corresponding to external symmetries (like $\Theta_c^{\mu\nu}$ is).

c will show totally irrelevant for the following discussion, however, it turns out to be important in nonequilibrium cases.

Now, similarly as before, $[\Theta_c^{\mu\nu}]$ gives the finite compos-

ite Green's functions if $n > 0$ but the expectation value $\langle [\Theta_c^{\mu\nu}] \rangle$ is divergent (for discussion of the $N=1$ theory see, e.g., Brown [37]). The unrenormalized DS equation for $D^{\mu\nu}(0|0)$ reads [3]

$$\begin{aligned}
 & \text{Diagram 1} + \text{Diagram 2} = \text{Diagram 3} + \left(\frac{1}{2} \text{Diagram 4} + \frac{1}{2} \text{Diagram 5} \right) \\
 & + \left(\frac{1}{36} \text{Diagram 6} + \frac{1}{48} \text{Diagram 7} \right). \quad (3.26)
 \end{aligned}$$

The structure of the composite vertices in Eq. (3.26) is that described at the beginning of this section. Note that the amputated composite Green's functions in individual parentheses are of the same order in λ_r . Performing the counterterm renormalization as in the case of $\langle [\phi^2] \rangle$, we factorize the graphs inside of parentheses into the product of the renormalized 2- (and 6-) point composite Green's function and the renormalized full 2- (and 6-) point Green's function. The latter are finite. The UV divergences arisen during the integrations over momenta connecting both composite and full Green's functions are precisely cancelled by the remaining counterterm diagrams. Only divergence comes from the free-field contribution, more precisely from the $T=0$ ring diagram. Defining

$$\langle \Theta_c^{\mu\nu} \rangle|_{\text{renorm}} = \langle [\Theta_c^{\mu\nu}] \rangle - \langle 0 | [\Theta_c^{\mu\nu}] | 0 \rangle, \quad (3.27)$$

or

$$\langle \Theta_c^{\mu\nu} \rangle|_{\text{renorm}} = \langle [\Theta_c^{\mu\nu}] \rangle - \langle [\Theta_c^{\mu\nu}] \rangle|_{\text{free field}}, \quad (3.28)$$

we get the finite expressions. Note that the conservation law is manifest in both cases. In equilibrium (and in $T=0$) we can, due to space-time translational invariance of $\langle \dots \rangle$, write

$$\langle [\Theta_c^{\mu\nu}] \rangle = \langle \Theta_c^{\mu\nu} \rangle + \partial_\lambda \langle X^{\lambda\mu\nu} \rangle = \langle \Theta_c^{\mu\nu} \rangle. \quad (3.29)$$

Using Eq. (3.27) or (3.28) we get either the thermal interaction pressure or the interaction pressure, respectively. This can be explicitly written as

$$\begin{aligned}
 \mathcal{P}_{\text{th.int.}}(T) &= \mathcal{P}(T) - \mathcal{P}(0) \\
 &= -\frac{1}{(D-1)} \sum_{i=1}^{D-1} \{ \langle \Theta_{ci}^i \rangle - \langle 0 | \Theta_{ci}^i | 0 \rangle \}, \quad (3.30)
 \end{aligned}$$

or

$$\begin{aligned}
 \mathcal{P}_{\text{int.}}(T) &= \mathcal{P}(T) - \mathcal{P}_{\text{free field}}(T) \\
 &= -\frac{1}{(D-1)} \sum_{i=1}^{D-1} \{ \langle \Theta_{ci}^i \rangle - \langle \Theta_{ci}^i \rangle|_{\text{free field}} \}. \quad (3.31)
 \end{aligned}$$

In order to keep connection with calculations done by Drummond *et al.* in [39] we shall in the sequel deal with the thermal interaction pressure only. If instead of an equilibrium, a nonequilibrium medium would be in question, translational invariance of $\langle \dots \rangle$ might be lost, in that case either prescription Eq. (3.27) or (3.28) is obligatory, and consequently $c(\lambda_r; D)$ in Eq. (3.25) must be further specified.

IV. HYDROSTATIC PRESSURE CALCULATION

In the preceding section we have prepared ground for a hydrostatic pressure calculation. In this section we aim to apply the previous results to the massive $O(N)\phi^4$ theory in the large- N limit. Anticipating an out of equilibrium application, we shall use the real-time formalism even if the imaginary-time one is more natural in the equilibrium context. As we aim to evaluate the hydrostatic pressure in 4 dimensions, we use here, similarly as in the preceding section, the usual dimensional regularization to regulate the theory (i.e., here and throughout we keep D slightly away from the physical value $D=4$).

In order to actually pursue the pressure calculation we feel it is necessary to briefly review the mass and coupling renormalization of the model at hand. This will also help to clarify the notation used. While we hope to provide all essentials requisite for our task, good discussion of alternative approaches and renormalization prescriptions may be obtained for instance in [29,30,33].

A. Mass renormalization

In the Dyson multiplicative renormalization the fact that the complete propagator has a pole at the physical mass leads to the usual mass renormalization prescription [24]:

$$m_r^2 = m_0^2 + \Sigma(m_r^2), \quad (4.1)$$

where m_r is renormalized mass and $\Sigma(m_r^2)$ is the proper self-energy evaluated at the mass shell; $p^2 = m_r^2$. In fact, Eq. (4.1) is nothing but the statement that 2-point vertex function $\Gamma_r^{(2)}$ evaluated at the mass-shell must vanish. The Dyson-Schwinger equation corresponding to the proper self-energy reads [29,38,57,58]:

$$\begin{aligned} \Sigma^{aa} = & \frac{1}{2} \sum_{b=1}^N \left(\text{tadpole loop with blob } b \right) + \text{setting sun diagram} \\ & + \frac{i}{2} \sum_{b=1}^N \left(\text{superdaisy diagram with hatched blobs} \right) \end{aligned} \quad (4.2)$$

$$\Sigma^{aa}|_{a \neq c} = 0; \quad \text{blob} \sim \lambda_o / N$$

where hatched blobs represent 2-point connected Green's functions while cross-hatched blobs represent proper vertices $\Gamma_r^{(4)}$ (i.e., 1PI 4-point Green's function). As Σ^{aa} are the same for all a , we shall simplify notation and write Σ instead. In the sequel the following convention is accepted.

The second term in Eq. (4.2) actually does not contribute in the large- N limit. It is easy to see that the third term does not contribute either. This is because each hatched blob behaves at most as N^0 while $\Gamma^{(4)}$ goes maximally¹⁰ as N^{-1} . Consequently, various contributions from the first graph in Eq. (4.2) contribute at most N^0 , whereas in the second graph the contributions

$$\text{cross-hatched blob} = \Gamma_r^{(n)} \text{abcd} \dots$$

contribute up to order N^{-1} . So the first diagram dominates, provided we retain only such 2-point connected Green's functions which are proportional to N^0 (as mentioned in the footnote, these are comprised only of tadpole loops). After neglecting the “setting sun” graph, Eq. (4.2) generates upon iterating the so-called superdaisy diagrams [39,25,38].

Let us now define $\Sigma(m_r^2) = \lambda_0 \mathcal{M}(m_r^2)$. Because the tadpole diagram in Eq. (4.2) can be easily resummed we observe that

¹⁰In the ϕ^4 theory there is a simple relation between the number of loops (L), vertices (V) and external lines (E); $4V = 2I + E$. Together with the Euler relation for connected graphs; $L = I - V + 1$ (here I is the number of internal lines), we have $L - V = (2 - E)/2$. As each loop carries maximally a factor of N (this is saturated only for “tadpole” loops) and each vertex carries a factor of N^{-1} , the overall blob contribution behaves at most as $N^{L-V} = N^{(2-E)/2}$.

$$\begin{aligned} \mathcal{M}(m_r^2) &= \frac{1}{2} \int \frac{d^D q}{(2\pi)^D} \frac{i}{q^2 - m_0^2 - \Sigma(m_r^2) + i\epsilon} \\ &= \frac{1}{2} \int \frac{d^D q}{(2\pi)^D} \frac{i}{q^2 - m_r^2 + i\epsilon}, \end{aligned} \quad (4.3)$$

hence we see that Σ is external-momentum independent. If we had started with the renormalization prescription: $i\Gamma_r^{(2)}(p^2=0) = -m_r^2$, we would have arrived at Eq. (4.1) as well (this is not the case for $N=1$!).

At finite temperature the strategy is analogous. Due to a doubling of degrees of freedom, the full propagator is a 2×2 matrix. The latter satisfies, similarly as at $T=0$, Dyson's equation

$$D = D_F + D_F(-i\Sigma)D. \quad (4.4)$$

An important point is that there exists a real, nonsingular matrix \mathbb{M} (Bogoliubov matrix) [3,4,7] having a property that

$$D_F = \mathbb{M} \begin{pmatrix} i\Delta_F & 0 \\ 0 & -i\Delta_F^* \end{pmatrix} \mathbb{M} \quad \text{and} \quad \Sigma = \mathbb{M}^{-1} \begin{pmatrix} \Sigma_T & 0 \\ 0 & -\Sigma_T^* \end{pmatrix} \mathbb{M}^{-1}. \quad (4.5)$$

Here Δ_F is the standard Feynman propagator and $*$ denotes the complex conjugation. Consequently, the full matrix propagator may be written as

$$D = \mathbb{M} \begin{pmatrix} \frac{i}{p^2 - m_0^2 - \Sigma_T + i\epsilon} & 0 \\ 0 & \frac{-i}{p^2 - m_0^2 - \Sigma_T^* - i\epsilon} \end{pmatrix} \mathbb{M}. \quad (4.6)$$

Similarly as in many body systems, the position of the (real) pole of D in p^2 fixes the temperature-dependent effective mass $m_r(T)$ [4,59]. The latter is determined by the equation

$$m_r^2(T) = m_0^2 + \text{Re}(\Sigma_T(m_r^2(T))). \quad (4.7)$$

From the explicit form of \mathbb{M} it is possible to show [3,4] that $\text{Re} \Sigma_{11} = \text{Re} \Sigma_T$. As before, the structure of the proper self-energy can be deduced from the corresponding Dyson-Schwinger equation. Following the usual real-time formalism convention (type-1 vertex $\sim -i\lambda_0$, type-2 vertex $\sim i\lambda_0$), the former reads:

$$-i\Sigma_{11} = \frac{i}{2} \left(\text{tadpole with blob} \right) - i\Sigma_{22} = \frac{i}{2} \left(\text{tadpole with blob} \right), \quad (4.8)$$

where

and similarly for D_{22} . In Eq. (4.8) we have omitted diagrams which are of order $\mathcal{O}(1/N)$ or less. Note that the fact that no setting sun diagrams are present implies that the off-diagonal elements of Σ are zero. Inspection of Eq. (4.8) reveals that

$$\Sigma_{11} = \frac{\lambda_0}{2} \int \frac{d^D q}{(2\pi)^D} D_{11}(q; T)$$

and

$$\Sigma_{22} = -\frac{\lambda_0}{2} \int \frac{d^D q}{(2\pi)^D} D_{22}(q; T). \quad (4.9)$$

It directly follows from Eq. (4.9) that both Σ_{11} and Σ_{22} are external-momentum independent and real.¹¹ If we define $\Sigma_T(m_r^2(T)) = \lambda_0 \mathcal{M}_T(m_r^2(T))$, then Eq. (4.7) through Eq. (4.9) implies that

$$m_r^2(T) = m_0^2 + \lambda_0 \mathcal{M}_T(m_r^2(T)). \quad (4.10)$$

A resummed version of D_{11} is easily obtainable from Eq. (4.6) [3,4] and consequently Eq. (4.9) yields

$$\begin{aligned} \mathcal{M}_T(m_r^2(T)) &= \frac{1}{2} \int \frac{d^D q}{(2\pi)^D} \left\{ \frac{i}{q^2 - m_r^2(T) + i\epsilon} \right. \\ &\quad \left. + (4\pi) \delta^+(q^2 - m_r^2(T)) \frac{1}{e^{q_0\beta} - 1} \right\} \\ &= - \int \frac{d^D q}{(2\pi)^D} \frac{\varepsilon(q_0)}{e^{q_0\beta} - 1} \text{Im} \frac{1}{q^2 - m_r^2(T) + i\epsilon}. \end{aligned} \quad (4.11)$$

Let us remark that Eq. (4.11) is manifestly independent of any particular real-time formalism version.

In passing it may be mentioned that because $\Sigma_{11}(m_r^2)$ is momentum independent, the wave function renormalization $Z_\phi = 1$. [The Källen-Lehmann representation requires the renormalized propagator to have a pole of residue i at p^2

$= m_r^2$. The former in turn implies that $Z_\phi = (1 - \Sigma'_{11}(p^2)|_{p^2=m_r^2})^{-1} = 1$.] Trivial consequence of the foregoing fact is that $\Gamma_r^{(2)} = \Gamma^{(2)}$ and $\Gamma_r^{(4)} = \Gamma^{(4)}$.

B. Coupling constant renormalization

Let us choose the coupling constant to be defined at $T = 0$. This will have the advantage that the high temperature expansion of the pressure (see Sec. V) will become more transparent. In addition, such a choice will allow us to select safely the part of the parameter space in which spontaneous symmetry breakdown is not possible. An alternative renormalization procedure based on the effective action is presented in [29].

By assumption the fields ϕ_a have nonvanishing masses, so we can safely choose the renormalization prescription for λ_r at $s=0$ (s is the standard Mandelstam variable). For example, one may require that for the scattering $aa \rightarrow bb$,

$$\Gamma^{(4)}(s=0) = -\lambda_r/N \quad (b \neq a). \quad (4.12)$$

The formula Eq. (4.12) clearly agrees with the tree level value $\Gamma_{tree}^{(4)aaab}(s=0) = -\lambda_0/N$. Let us also mention that Ward's identities corresponding to the internal $O(N)$ symmetry enforce $\Gamma^{(4)aaaa}$ to obey the constraint¹²

$$\begin{aligned} \Gamma^{(4)aaaa}(p_1; p_2; p_3; p_4) &= \Gamma^{(4)bbba}(p_1; p_2; p_3; p_4) \\ &\quad + \Gamma^{(4)baba}(p_1; p_2; p_3; p_4) \\ &\quad + \Gamma^{(4)baab}(p_1; p_2; p_3; p_4), \end{aligned} \quad (4.13)$$

for any $b \neq a$. The structure of $\Gamma^{(4)}$ is encoded in the following Dyson-Schwinger equation (see also [24,57]).

¹¹Reality of Σ_{11} can be most easily seen from the largest-time equation [58]. The LTE states that $\Sigma_{11} + \Sigma_{22} + \Sigma_{12} + \Sigma_{21} = 0$. Because no setting sun graphs are present, $\Sigma_{12} + \Sigma_{21} = 0$, on the other hand $\Sigma_{11} + \Sigma_{22} = 2i \text{Im} \Sigma_{11}$ [see Eq. (4.5)].

¹²Actually, Ward's identities read [57] $\int d^D x [\delta\Gamma[\phi]/\delta\phi_a(x)] \times \phi_b(x) = \int d^D x [\delta\Gamma[\phi]/\delta\phi_b(x)] \phi_a(x)$ [here $\phi_a = (\delta W/\delta J_a)$; W is the generating functional of connected Green's functions]. Performing successive variations with respect to $\phi_a(v), \phi_a(z), \phi_a(y)$, and $\phi_b(w)$, taking the Fourier transform, and setting the physical condition $\phi_c = 0$, we get directly Eq. (4.13).

$$\begin{aligned}
 & \text{Diagram} = - \frac{i}{2} \sum_{i=1}^3 \text{Diagram} - \frac{1}{6} \text{Diagram} \\
 & + \frac{1}{2} \sum_{i=1}^3 \text{Diagram} ; \quad \times \sim \lambda_o/N
 \end{aligned} \tag{4.14}$$

In the latter the sum $\sum_{i=1}^3$ schematically represents a summation over s, t and u scattering channels. For clarity's sake the internal indices are suppressed. Similarly as before, we can argue that both the third and fourth graphs contribute at most N^{-2} , while the second (“fish”) graph may contribute up to order N^{-1} . So in the large- N limit the last three diagrams may be neglected, provided we keep in the 4-point vertex function only graphs proportional to N^{-1} . However, the former can be only fulfilled if we retain such a “fish” graph where summation over internal index on the loop is allowed. Remaining two graphs in the sum $\sum_{i=1}^3$ (i.e., t and u scattering channels) are suppressed by the factor N^{-1} as the internal index on the loop is fixed. In this way we are left with the relation

$$\begin{aligned}
 \Gamma^{(4) aabb}(s=0) &= -\frac{\lambda_0}{N} - \frac{i\lambda_0}{2N} \sum_{c \neq b} \int \frac{d^D q}{(2\pi)^D} \Gamma^{(4) bccc}(s) \\
 &\times \frac{i}{(q^2 - m_r^2 + i\epsilon)} \frac{i}{((q-Q)^2 - m_r^2 + i\epsilon)} \Big|_{s=0} \\
 &= -\frac{\lambda_0}{N} - \frac{\lambda_0 \lambda_r (N-1)}{2N^2} \int_0^1 dx \int \frac{d^D q}{(2\pi)^D} \\
 &\times \frac{i}{(q^2 - m_r^2 + x(1-x)s + i\epsilon)^2} \Big|_{s=0}, \tag{4.15}
 \end{aligned}$$

with $Q = p_1 + p_2$ and $s = Q^2$, p_1, p_2 are the external momenta. To leading order in $1/N$ we may equivalently write

$$\lambda_r = \lambda_0 + \lambda_0 \lambda_r \mathcal{M}'(m_r^2), \tag{4.16}$$

the prime means differentiation with respect to m_r^2 ; $\mathcal{M}(m_r^2)$ is defined by Eq. (4.3). Evaluating explicitly $\mathcal{M}'(m_r^2)$, we get from Eq. (4.16),

$$\lambda_0 = \frac{\lambda_r}{1 - \lambda_r \Gamma\left(2 - \frac{D}{2}\right) (m_r)^{D-4}/2(4\pi)^{D/2}}. \tag{4.17}$$

Assuming that both $\lambda_0 \geq 0$ and $\lambda_r \geq 0$ (note $\lambda_r < 0$ would be incompatible with $\lambda_0 \geq 0$ and $m_r^2 > 0$), we can infer from Eq. (4.17) that

$$0 \leq \lambda_r \leq \frac{2(4\pi)^{D/2} (m_r)^{4-D}}{\Gamma\left(2 - \frac{D}{2}\right)}, \tag{4.18}$$

and so for $D=4$ we inevitably get that $\lambda_r = 0$. The latter indicates that the theory is trivial [39,23,31], or, in other words, the $O(N)\phi^4$ theory is a renormalized free theory in the large- N limit. This conclusion is also consistent with the observation that the theory does not possess any nontrivial UV fixed point in the large- N limit [23,31,32,60].

On the other hand, if we were assuming that $\lambda_0 < 0$, we would get indeed a nontrivial renormalized field theory in $D=4$ [actually, from Eq. (4.17) we see that $\lambda_0 \rightarrow 0_-$, provided that λ_r is fixed and positive and $D \rightarrow 4_-$]. However, as it was pointed out in Refs. [39,23,31,33], such a theory is intrinsically unstable as the ground-state energy is unbounded from below. This is reflected, for instance, in the existence of tachyons in the theory [39,31,33,61], therefore the case with negative λ_0 is clearly inconsistent.

The straightforward remedy for this situation was suggested by Bardeen and Moshe [31]. They showed that the only meaningful (stable) $O(N)\phi^4$ theory in the large- N limit is that with $\lambda_r, \lambda_0 \geq 0$. This is provided that we view it as an effective field theory at momenta scale small compared to a fixed UV cutoff Λ . The cutoff itself is further determined by Eq. (4.16) because in that case (assuming $m_r \ll \Lambda$)

$$\lambda_0 = \frac{\lambda_r}{1 - \frac{\lambda_r}{32\pi^2} \ln\left(\frac{\Lambda^2}{m_r^2}\right)}, \tag{4.19}$$

which implies that for $\lambda_r, \lambda_0 \geq 0$ we have $\Lambda^2 < m_r^2 \exp(32\pi^2/\lambda_r)$. The case $\Lambda^2 = m_r^2 \exp(32\pi^2/\lambda_r)$ corresponds to the Landau ghost [62] (tachyon pole [39,31]). For reasonably small λ_r , Λ is truly huge¹³ and so it does not represent any significant restriction.

¹³For example, if $\lambda_r = 1$ and $m_r \approx 100$ MeV, we get $\Lambda < 10^{141}$ MeV or equivalently $\Lambda < 10^{131}$ K (this is far beyond the Planck temperature $\sim 10^{32}$ K).




FIG. 3. The Keldysh-Schwinger time path.

In passing we may observe that from Eqs. (4.1) and (4.19),

$$\frac{m_r^2}{\lambda_r} = \frac{m_0^2}{\lambda_0} + \frac{1}{23\pi^2} \Lambda^2, \quad (4.20)$$

and so the fraction m_r^2/λ_r is renormalization invariant. It was argued in [31] that for the part of the parameter space where $\lambda_0 > 0$ and $m_r^2/\lambda_r > 0$ the ground state is $O(N)$ symmetric, Goldstone phenomena cannot materialize and hence the expectation value of the field is zero. The latter fact has been implicitly used, for instance, in the derivation our Dyson-Schwinger equations. To avoid a delicate discussion of the phase structure of the $O(N)$ ϕ^4 theory and to emphasize our primary objective, i.e., hydrostatic pressure calculation, we confine ourselves to the parameter space defined above. Such an effective theory will provide a suitable playground to explore all the basic salient points involved in the hydrostatic pressure calculation. Furthermore, because the mass-shift equation (gap equation) has a particularly simple form in this case the high- T analysis of the hydrostatic pressure will be easy to perform.

C. Resumed hydrostatic pressure

The partition function Z has a well-known path-integral representation at finite temperature, namely,

$$Z[T] = \exp(\Phi[T]) = \int \mathcal{D}\phi \exp(iS[\phi; T]),$$

$$S[\phi; T] = \int_C d^D x \mathcal{L}(x). \quad (4.21)$$

Here $\Phi = -\beta\Omega$ is a Massieu function (the Legendre transform of the entropy) [3,41–43] and $\int_C d^D x = \int_C dx_0 \int_V d^{D-1} \mathbf{x}$ with the subscript C suggesting that the time runs along some contour in the complex plane. In the real-time formalism, which we adopt throughout, the most natural version is the so-called Keldysh-Schwinger one [3,4], which is represented by the contour in Fig. 3. Let us mention that the fields within the path-integral Eq. (4.21) are further restricted by the periodic boundary condition (KMS condition) [3,4,7] which in our case reads

$$\phi_a(t_i - i\beta, \mathbf{x}) = \phi_a(t_i, \mathbf{x}).$$

As explained in Sec. III, we can use for a pressure calculation the canonical energy-momentum tensor $\Theta_c^{\mu\nu}$. Employing for $\Theta_c^{\mu\nu}(x)$ its explicit form Eq. (3.2) together with Eq. (3.6), one may write

$$\langle \Theta_c^{\mu\nu} \rangle = \frac{N}{2} \int \frac{d^D q}{(2\pi)^D} (2q^\mu q^\nu - g^{\mu\nu}(q^2 - m_0^2)) D_{11}(q; T)$$

$$+ \frac{\lambda_0}{8N} g^{\mu\nu} \left\langle \left(\sum_{a=1}^N \phi_a^2(0) \right)^2 \right\rangle, \quad (4.22)$$

where D_{11} is the Dyson-resumed thermal propagator [3,4], i.e.,

$$D_{11}(q; T) = \frac{i}{q^2 - m_r^2(T) + i\epsilon}$$

$$+ (2\pi) \delta(q^2 - m_r^2(T)) \frac{1}{e^{|q_0|\beta} - 1}. \quad (4.23)$$

Note that we have exploited in Eq. (4.22) the fact that the expectation value of $\Theta_c^{\mu\nu}(x)$ is x independent. On the other hand, in Eq. (4.23) we have used the fact that m_r^2 is q independent. In order to calculate the expectation value of the quartic term in Eq. (4.22), let us observe [cf. Eq. (4.21)] that the derivative of Φ with respect to the bare coupling λ_0 (taken at fixed m_0) gives

$$\frac{\partial \Phi[T]}{\partial \lambda_0} = -\frac{i}{8N} \int_C d^D x \left\langle \left(\sum_{a=1}^N \phi_a^2(0) \right)^2 \right\rangle, \quad (4.24)$$

which implies that

$$\left\langle \left(\sum_{a=1}^N \phi_a^2(0) \right)^2 \right\rangle = -\frac{N8}{\beta V} \frac{\partial \Phi[T]}{\partial \lambda_0}. \quad (4.25)$$

The key point now is that we can calculate $\Phi[T]$ in a non-perturbative form. The latter is based on the fact that we know the Dyson-resumed propagator $D_{11}(q; T)$ [see Eq. (4.23)]. Indeed, taking derivative of Φ with respect to m_0^2 (keeping λ_0 fixed) we obtain

$$\frac{\partial \Phi[T]}{\partial m_0^2} = -\frac{iN}{2} \int_C d^D x \langle \phi^2(0) \rangle$$

$$= -\frac{\beta V N}{2} \int \frac{d^D q}{(2\pi)^D} D_{11}(q; T)$$

$$= -\beta V N \mathcal{M}_T(m_r^2(T)), \quad (4.26)$$

thus

$$\Phi[T; \lambda_0; m_0^2] = \beta V N \int_{m_0^2}^{\infty} d\hat{m}_0^2 \mathcal{M}_T(\hat{m}_r^2(T)) + \Phi[T; \lambda_0; \infty]. \quad (4.27)$$

Let us note that $\Phi[T; \lambda_0; \infty]$ is actually zero¹⁴ because $\Phi[T; \lambda_0; m_0^2]$ has the standard loop expansion [3,57] depicted in Fig. 4. It is worth mentioning that in the latter expansion one must always have at least one type-1 vertex [39]. The RHS of Fig. 4 clearly tends to zero for $m_0 \rightarrow \infty$ as all the (free) thermal propagators from which the individual diagrams are constructed tend to zero in this limit. The former result can be also deduced from the CJT effective action formalism [25] or from a heuristic argumentation based on a thermodynamic pressure [39]. Note that in the large- N limit the fourth and fifth diagrams in Fig. 4 must be omitted.

The expectation value Eq. (4.25) can be now explicitly written as

$$\left\langle \left(\sum_{a=1}^N \phi_a^2(0) \right)^2 \right\rangle = 8N^2 \int_{m_0^2}^{\infty} d\hat{m}_0^2 \int \frac{d^D q}{(2\pi)^D} \frac{\varepsilon(q_0)}{e^{q_0\beta} - 1} \times \text{Im} \left(\frac{\partial \Sigma_T(\hat{m}_r^2(T))}{\partial \lambda_0} \right) \frac{1}{(q^2 - \hat{m}_r^2 + i\epsilon)^2}. \quad (4.28)$$

In fact, the differentiation of the proper self-energy in Eq. (4.28) can be carried out easily. Using Eq. (4.10), we get

$$\frac{\partial \Sigma_T}{\partial \lambda_0} = \frac{\Sigma_T}{\lambda_0} + \lambda_0 \mathcal{M}'_T \frac{\partial \Sigma_T}{\partial \lambda_0} \Rightarrow \frac{\partial \Sigma_T}{\partial \lambda_0} = \frac{\Sigma_T}{\lambda_0(1 - \lambda_0 \mathcal{M}'_T)}.$$

From Eq. (4.10) it directly follows that

$$\frac{dm_r^2(T)}{dm_0^2} = \frac{1}{(1 - \lambda_0 \mathcal{M}'_T)},$$

which, together with the definition of \mathcal{M}_T , gives

$$\begin{aligned} \left\langle \left(\sum_{a=1}^N \phi_a^2(0) \right)^2 \right\rangle &= 8N^2 \int_{m_r^2(T)}^{\infty} d\hat{m}_r^2 \int \frac{d^D q}{(2\pi)^D} \frac{\varepsilon(q_0)}{e^{q_0\beta} - 1} \\ &\times \text{Im} \frac{\mathcal{M}_T(\hat{m}_r^2)}{(q^2 - \hat{m}_r^2 + i\epsilon)^2} \\ &= -8N^2 \int_{m_r^2(T)}^{\infty} d\hat{m}_r^2 \mathcal{M}_T(\hat{m}_r^2) \frac{\partial \mathcal{M}_T(\hat{m}_r^2)}{\partial \hat{m}_r^2} \\ &= 4N^2 \mathcal{M}_T^2(m_r^2(T)), \end{aligned} \quad (4.29)$$

where we have exploited in the last line the fact that $\mathcal{M}_T(m_r^2 \rightarrow \infty) = 0$. Let us mention that the crucial point in the previous manipulations was that m_r is both real and momentum independent. Collecting our results together, we can write for the hydrostatic pressure per particle [cf. Eq. (3.30)]

¹⁴To be precise, we should also include in Fig. 4 an (infinite) circle diagram corresponding to the free pressure [38,39]. However, the latter is λ_0 independent (although m_0 dependent) and so it is irrelevant for the successive discussion [cf. Eq. (4.25)].




FIG. 4. First few bubble diagrams in the Φ expansion.

$$\begin{aligned} \mathcal{P}(T) - \mathcal{P}(0) &= -\frac{1}{(D-1)N} (\langle \Theta_{ci}^i \rangle - \langle 0 | \Theta_{ci}^i | 0 \rangle) \\ &= +\frac{1}{2} \int \frac{d^D q}{(2\pi)^{D-1}} \left(\frac{2\mathbf{q}^2}{(D-1)} \right) \frac{\varepsilon(q_0)}{e^{q_0\beta} - 1} \\ &\times \delta(q^2 - m_r^2(T)) \\ &- \frac{1}{2} \int \frac{d^D q}{(2\pi)^{D-1}} \left(\frac{2\mathbf{q}^2}{(D-1)} \right) \delta^+(q^2 - m_r^2(0)) \\ &+ \frac{1}{2\lambda_0} (\Sigma_T^2(m_r^2(T)) - \Sigma^2(m_r^2(0))). \end{aligned} \quad (4.30)$$

Applying Green's theorem to the last two integrals and eliminating the surface terms (for details see Appendix A) we find

$$\begin{aligned} \mathcal{P}(T) - \mathcal{P}(0) &= \frac{1}{2} \int \frac{d^D q}{(2\pi)^{D-1}} \frac{\varepsilon(q_0)}{e^{q_0\beta} - 1} \theta(q^2 - m_r^2(T)) \\ &- \frac{1}{2} \int \frac{d^D q}{(2\pi)^{D-1}} \theta(q_0) \theta(q^2 - m_r^2(0)) \\ &+ \frac{1}{2\lambda_0} (\Sigma_T^2(m_r^2(T)) - \Sigma^2(m_r^2(0))) \\ &= \mathcal{N}_T(m_r^2(T)) - \mathcal{N}(m_r^2(0)) \\ &+ \frac{1}{2\lambda_0} (\Sigma_T^2(m_r^2(T)) - \Sigma^2(m_r^2(0))), \end{aligned} \quad (4.31)$$

where we have introduced new functions $\mathcal{N}_T(m_r^2(T))$ and $\mathcal{N}(m_r^2)$;

$$\mathcal{N}_T(m_r^2(T)) = \frac{1}{2} \int \frac{d^D q}{(2\pi)^{D-1}} \frac{\varepsilon(q_0)}{e^{q_0\beta} - 1} \theta(q^2 - m_r^2(T))$$

$$\mathcal{N}(m_r^2) = \lim_{T \rightarrow 0} \mathcal{N}_T(m_r^2(T)). \quad (4.32)$$

Equation (4.31) can be rephrased into a form which exhibits an explicit independence of bar quantities. Using the trivial identity:

$$\begin{aligned}
& \frac{1}{2\lambda_0}(\Sigma_T^2(m_r^2(T)) - \Sigma^2(m_r(0))) \\
&= \frac{1}{2\lambda_0}(\Sigma_T(m_r^2(T)) - \Sigma(m_r^2(0))) \\
&\quad \times (\Sigma_T(m_r^2(T)) + \Sigma(m_r^2(0))) \\
&= \frac{\delta m^2(T)}{2}(\mathcal{M}_T(m_r^2(T)) + \mathcal{M}(m_r^2(0))), \quad (4.33)
\end{aligned}$$

we get

$$\begin{aligned}
\mathcal{P}(T) - \mathcal{P}(0) &= \mathcal{N}_T(m_r^2(T)) - \mathcal{N}(m_r^2(0)) \\
&\quad + \frac{\delta m^2(T)}{2}(\mathcal{M}_T(m_r^2(T)) + \mathcal{M}(m_r^2(0))), \quad (4.34)
\end{aligned}$$

where $\delta m^2(T) = m_r^2(T) - m_r^2(0)$. Let us finally mention that the finding Eq. (4.30) is an original result of this paper. The result Eq. (4.34) has been previously obtained by authors [39] in the purely thermodynamic pressure framework.

V. HIGH-TEMPERATURE EXPANSION OF THE HYDROSTATIC PRESSURE IN $D=4$

In order to obtain the high-temperature expansion of the pressure in $D=4$, it is presumably the easiest to go back to Eq. (4.30) and employ identity Eq. (4.33). Let us split this task into two parts. We first evaluate the integrals with potentially UV divergent parts using the dimensional regularization. The remaining integrals, with the Bose-Einstein distribution insertion, are safe of UV singularities and can be computed by means of the Mellin transform technique.

Inspecting Eqs. (4.30) and (4.33), we observe that the only UV divergent contributions come from the integrals:

$$\begin{aligned}
& + \frac{1}{(D-1)} \int \frac{d^D q}{(2\pi)^{D-1}} \mathbf{q}^2 (\delta^+(q^2 - m_r^2(T)) \\
& \quad - \delta^+(q^2 - m_r^2(0))) + \frac{\delta m^2(T)}{4} \int \frac{d^D q}{(2\pi)^D} \\
& \quad \times \left(\frac{i}{q^2 - m_r^2(T) + i\epsilon} + \frac{i}{q^2 - m_r^2(0) + i\epsilon} \right), \quad (5.1)
\end{aligned}$$

which, if integrated over, give

$$\begin{aligned}
(5.1) &= + \frac{\Gamma\left(\frac{-D}{2}\right) \Gamma\left(\frac{D}{2} + \frac{1}{2}\right)}{(D-1) \Gamma\left(\frac{D-1}{2}\right) (4\pi)^{D/2}} \\
&\quad \times ((m_r^2(T))^{D/2} - (m_r^2(0))^{D/2}) + \frac{\delta m^2(T) \Gamma\left(1 - \frac{D}{2}\right)}{4(4\pi)^{D/2}} \\
&\quad \times ((m_r^2(T))^{(D/2)-1} + (m_r^2(0))^{(D/2)-1}). \quad (5.2)
\end{aligned}$$

Taking the limit $D=4-2\epsilon \rightarrow 4$ and using expansions

$$\Gamma(-n+\epsilon) = \frac{(-1)^n}{n!} \left(\frac{1}{\epsilon} + \sum_{k=1}^n \frac{1}{k} - \gamma + \mathcal{O}(\epsilon) \right),$$

$$a^{x+\epsilon} = a^x (1 + \epsilon \ln a + \mathcal{O}(\epsilon^2)),$$

(γ is the Euler-Mascheroni constant) we are finally left with

$$\begin{aligned}
(5.1)|_{D \rightarrow 4} &= - \frac{m_r^2(0)m_r^2(T)}{64\pi^2} \ln \left(\frac{m_r^2(T)}{m_r^2(0)} \right) \\
&\quad + \delta m^2(T)(m_r^2(T) + m_r^2(0)) \frac{1}{128\pi^2}. \quad (5.3)
\end{aligned}$$

The fact that we get the finite result should not be surprising as the entire analysis of Sec. III was made to show that $\mathcal{P}(T) - \mathcal{P}(0)$ defined via $\Theta_c^{\mu\nu}$ is finite in $D=4$.

We may now concentrate on the remaining terms in Eq. (4.30), the latter read (we might, and we shall, from now on work in $D=4$)

$$\begin{aligned}
& \frac{1}{3} \int \frac{d^4 q}{(2\pi)^3} \mathbf{q}^2 \frac{1}{e^{|q_0|\beta} - 1} \delta(q^2 - m_r^2(T)) \\
& + \frac{\delta m^2(T)}{4} \int \frac{d^4 q}{(2\pi)^3} \frac{1}{e^{|q_0|\beta} - 1} \delta(q^2 - m_r^2(T)). \quad (5.4)
\end{aligned}$$

Our following strategy is based on the observation that the previous integrals have generic form:

$$\begin{aligned}
I_{2\nu}(m_r) &= \int \frac{d^4 q}{(2\pi)^3} \mathbf{q}^{2\nu} \frac{1}{e^{|q_0|\beta} - 1} \delta(q^2 - m_r^2) \\
&= \frac{m_r^{2+2\nu}}{2\pi^2} \int_1^\infty dx (x^2 - 1)^{(1+2\nu)/2} \frac{1}{e^{xy} - 1}, \quad (5.5)
\end{aligned}$$

with $\nu=0,1$ and $y=m_r\beta$. Unfortunately, the integral Eq. (5.5) cannot be evaluated exactly, however, its small y (i.e., high-temperature) behavior can be successfully analyzed by means of the Mellin transform technique [3,38]. Before going further, let us briefly outline the basic steps needed for such a small y expansion.

The Mellin transform $\hat{f}(s)$ is done by the prescription [3,38,63–66]:

$$\hat{f}(s) = \int_0^\infty dx x^{s-1} f(x), \quad (5.6)$$

with s being a complex number. One can easily check that the inverse Mellin transform reads

$$f(x) = \frac{1}{i(2\pi)} \int_{-i\infty+a}^{i\infty+a} ds x^{-s} \hat{f}(s), \quad (5.7)$$

where the real constant a is chosen in such a way that $\hat{f}(s)$ is convergent in the neighborhood of a straight line $(-i\infty + a, i\infty + a)$. So particularly if $f(x) = 1/[e^{xy} - 1]$ one can find ([65]; formula I.3.19) that

$$\hat{f}(s) = \Gamma(s) \zeta(s) y^{-s} \quad (\text{Re } s > 1), \quad (5.8)$$

where ζ is the Riemann zeta function ($\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$). Now we insert the Mellin transform of $f(x) = 1/[e^{xy} - 1]$ to Eq. (5.5) and interchange integrals (this is legitimate only if the integrals are convergent before the interchange). As a result we have

$$\int_0^{\infty} dx g(x) \frac{1}{e^{xy} - 1} = \int_{-i\infty+a}^{i\infty+a} \frac{ds}{i(2\pi)} \Gamma(s) \zeta(s) y^{-s} \hat{g}(1-s), \quad (5.9)$$

with $g(x) = \theta(x-1)(x^2-1)^{(1+2\nu)/2}$. Using the tabulated result ([66]; formula 6.2.32) we find

$$\hat{g}(1-s) = \frac{1}{2} B(-\nu-1 + \frac{1}{2}s; \frac{3}{2} + \nu) \quad (\text{Re } s > 2+2\nu), \quad (5.10)$$

with $B(\cdot)$ being the beta function. Because the integrand on the RHS of Eq. (5.9) is analytic for $\text{Re } s > 2+2\nu$ and the LHS is finite, we must choose such a that the integration is defined. The foregoing is achieved choosing $a > 2+2\nu$. Other useful expressions for $\hat{g}(1-s)$ are ([66]; formula I.2.34 or I.2.37)

$$\begin{aligned} \hat{g}(1-s) &= B(\frac{3}{2} + \nu; -2-2\nu+s) \\ &\times {}_2F_1[-\frac{1}{2} - \nu; -2-2\nu+s; -\frac{1}{2} - \nu+s; -1] \end{aligned}$$

$$\begin{aligned} &= 2^{\frac{1}{2}+\nu} B(\frac{3}{2} + \nu; -2-2\nu+s) \\ &\times {}_2F_1[-\frac{1}{2} - \nu; \frac{3}{2} + \nu; -\frac{1}{2} - \nu+s; \frac{1}{2}], \end{aligned}$$

where ${}_2F_1$ is the (Gauss) hypergeometric function [66]. Using identity

$$\Gamma(2x) = \frac{2^{2x-1}}{\sqrt{\pi}} \Gamma(x) \Gamma\left(x + \frac{1}{2}\right),$$

we can write

$$\begin{aligned} (5.9) &= \frac{\Gamma\left(\frac{3}{2} + \nu\right)}{4\sqrt{\pi}} \int_{-i\infty+a}^{i\infty+a} \frac{ds}{i(2\pi)} \Gamma\left(\frac{1}{2}s\right) \\ &\times \zeta(s) \left(\frac{1}{2}y\right)^{-s} \Gamma\left(-\nu-1 + \frac{1}{2}s\right). \quad (5.11) \end{aligned}$$

The integrand of Eq. (5.11) has simple poles in $s = -2n$ ($n = 1, 2, \dots$), $s = 1$, $s = -2n+2\nu+2$ ($n = 0, 1, \dots, \nu$) and double pole in $s = 0$. An important point in the former pole analysis was the fact that $\zeta(s)$ has simple zeros in $-2m$ ($m > 0$) and only one simple pole in $s = 1$. The former together with identity

$$\Gamma\left(\frac{x}{2}\right) \pi^{-x/2} \zeta(x) = \Gamma\left(\frac{1-x}{2}\right) \pi^{(x-1)/2} \zeta(1-x),$$

shows that no double pole except for $s = 0$ is present in Eq. (5.11). Now, we can close the contour to the left as the value of the contour integral around the large arc is zero in the limit of infinite radius (cf. [65] and [67]; formula 8.328.1). Using successively the Cauchy theorem we obtain

$$\begin{aligned} \frac{4\sqrt{\pi}(5.9)}{\Gamma\left(\frac{3}{2} + \nu\right)} &= \sum_{n=0}^{\nu} y^{2n-2\nu-2} \frac{\pi^{-2n+2\nu+2} (-n+\nu)! (-1)^n |B_{-2n+2\nu+2}|}{n! (-2n+2\nu+2)! 2^{4n-4\nu-4}} + \sum_{n=1}^{\infty} y^{2n} \frac{\pi^{-2n} (2n)! \zeta(1+2n) (-1)^{n+\nu+1}}{n! (n+1+\nu)! 2^{4n-1}} \\ &+ y^{-1} \frac{\pi (-1)^{\nu+1} (\nu+1)! 2^{2\nu+3}}{(2\nu+2)!} + \frac{2(-1)^{\nu+1}}{(\nu+1)!} \left\{ \ln\left(\frac{y}{4\pi}\right) + \gamma - \frac{1}{2} \sum_{k=1}^{\nu+1} \frac{1}{k} \right\}, \quad (5.12) \end{aligned}$$

where B_n 's are the Bernoulli numbers. Let us mention that for $\zeta(2n+1)$ only numerical values are available.

Inserting Eq. (5.12) back to Eq. (5.4), we get for $\mathcal{P}(T) - \mathcal{P}(0)$,

$$\begin{aligned} \mathcal{P}(T) - \mathcal{P}(0) &= (5.3) + \frac{1}{3} I_2(m_r(T)) + \frac{\delta m^2(T)}{4} I_0(m_r(T)) \\ &= \frac{T^4 \pi^2}{90} - \frac{T^2}{24} \left(m_r^2(T) - \frac{\delta m^2(T)}{2} \right) + \frac{T m_r(T)}{4\pi} \left(\frac{m_r^2(T)}{3} - \frac{\delta m^2(T)}{4} \right) + \frac{m_r^2(T) m_r^2(0)}{32\pi^2} \left(\ln\left(\frac{m_r(0)}{T4\pi}\right) + \gamma - \frac{1}{2} \right) - \frac{m_r^4(0)}{128\pi^2} \\ &- \sum_{n=1}^{\infty} \left(m_r^2(T) - \frac{(n+2)}{2} \delta m^2(T) \right) \frac{m_r^{2n+2}(T) \pi^{-2n-2} (2n)! \zeta(1+2n) (-1)^{1+n}}{T^{2n} n! (n+2)! 2^{4n+4}}. \quad (5.13) \end{aligned}$$

Note that Eq. (5.3) cancelled against the same term in $\frac{1}{3}I_2(m_r(T)) + [\delta m^2(T)/4]I_0(m_r(T))$. One can see that Eq. (5.13) rapidly converges for large T , so that only first four terms dominate at sufficiently high temperature. The aforementioned terms come from the poles nearby the straight line $(-i\infty + a, i\infty + a)$ (the more dominant contribution the closer pole). It is a typical feature of the Mellin transform technique that integrals of type

$$\int_0^\infty dx g(x) \frac{1}{e^{xy} - 1},$$

can be expressed as an expansion which rapidly converges for small y (high-temperature expansion) or large y (low-temperature expansion).¹⁵

Expansion (5.13) is the sought result. To check its consistency we will apply it to two important cases: high T case and $m_r(0)=0$ case. Concerning the first case, note that for a sufficiently large T we can use the high-temperature expansion of $\delta m^2(T)$ found in Appendix B. Inserting Eq. (B6) to Eq. (5.13) we obtain

$$\begin{aligned} \mathcal{P}(T) - \mathcal{P}(0) = & \frac{T^4 \pi^2}{90} - \frac{T^2 m_r^2(T)}{24} + \frac{T^3 m_r(T)}{12\pi} \\ & + \frac{\lambda_r}{8} \left(\frac{T^4}{144} - \frac{T^3 m_r(T)}{24\pi} + \frac{T^2 m_r^2(T)}{16\pi^2} \right) \\ & + \mathcal{O} \left(m_r^4(T) \ln \left(\frac{m_r(T)}{T^4 \pi} \right) \right). \end{aligned} \quad (5.14)$$

Up to a sign, the result Eq. (5.14) coincides with that found by Amelino-Camelia and Pi [40] for the effective potential.¹⁶ Actually, they used instead of the $N \rightarrow \infty$ limit the Hartree-Fock approximation which is supposed to give the same V_{eff} as the leading $1/N$ approximation [62].

As for the second case, we may observe that our discussion of the mass renormalization in Sec. III A can be directly extended to the case when $m_r(0)=0$ (this does not apply to our discussion of λ_r !). Latter can be also seen from the fact that Eq. (5.13) is continuous in $m_r(0)=0$ (however not analytic). The foregoing implies that the original massless scalar particles acquire the thermal mass $m_r^2(T) = \delta m^2(T)$. From Eq. (5.13) one then may immediately deduce the pressure for massless fields ϕ_a in terms of $\delta m(T)$. The latter reads

$$\begin{aligned} \mathcal{P}(T) - \mathcal{P}(0) = & \frac{T^4 \pi^2}{90} - \frac{T^2 (\delta m(T))^2}{48} + \frac{T (\delta m(T))^3}{48\pi} \\ & + \sum_{n=1}^{\infty} \frac{(\delta m(T))^{2n+4} \pi^{-2n-2} (2n)! \zeta(1+2n) (-1)^{n+1}}{T^{2n} (n-1)! (n+2)! 2^{4n+5}}. \end{aligned} \quad (5.15)$$

This result is identical to that found by Drummond *et al.* in [39].

A noteworthy observation is that when the energy of a thermal motion is much higher than the mass of particles in the rest, then the massive theory approaches the massless one. This is justified in the first (high-temperature dominant) term of Eqs. (5.13) and (5.15). This term is nothing but a half of the black-body radiation pressure for photons [41,42] (photons have two degrees of freedom connected with two transverse polarizations). One could also obtain the temperature dominant contributions directly from the Stefan-Boltzmann law [3,41,42] for the density energy (i.e., $\langle \Theta^{00} \rangle$). The formal argument leading to this statement is based on the noticing that at high energy (temperature) the scalar field theory is approximately conformally invariant, which in turn implies that the energy-momentum tensor is traceless [52]. Taking into account the definition of the hydrostatic pressure Eq. (2.8), we can with a little effort recover the leading high-temperature contributions for the massive case.

VI. CONCLUSIONS

In the present paper we have clarified the status of hydrostatic pressure in (equilibrium) thermal QFT. The former is explained in terms of the thermal expectation value of the “weighted” space-like trace of the energy-momentum tensor $\Theta^{\mu\nu}$. In classical field theory there is a clear microscopic picture of the hydrostatic pressure which is further enhanced by a mathematical connection (through the virial theorem) with the thermodynamic pressure. In addition, it is the hydrostatic pressure which can be naturally extended to a non-equilibrium medium. Quantum theoretic treatment of the hydrostatic pressure is however pretty delicate. In order to get a sensible, finite answer we must give up the idea of total hydrostatic pressure. Instead, thermal interaction pressure or/and interaction pressure must be used [see Eqs. (3.30) and (3.31)]. We have established this result for a special case when the theory in question is the scalar ϕ^4 theory with $O(N)$ internal symmetry; but it can be easily extended to more complex situations. Moreover, due to a lucky interplay between the conservation of $\Theta^{\mu\nu}$ and the space-time translational invariance of an equilibrium (and $T=0$) expectation value we can use the simple canonical (i.e., unrenormalized) energy-momentum tensor. In the course of our treatment in Sec. III we heavily relied on the counterterm renormalization, which seems to be the most natural when one discusses renormalization of composite Green’s functions. To be specific, we have resorted to the minimal subtraction scheme which has proved useful in several technical points.

¹⁵By the same token we get the low-temperature expansion if the integral contour must be closed to the right.

¹⁶Let us remind [40,25,29] that from the definition of V_{eff} the thermodynamic pressure is $-V_{\text{eff}}$. In order to obtain Eq. (5.14) from V_{eff} in [40], one must subtract the zero temperature value of V_{eff} and restrict oneself to vanishing field expectation value and positive bare mass squared.

We have applied the prescriptions obtained for the QFT hydrostatic pressure to ϕ^4 theory in the large- N limit. The former has the undeniable advantage of being exactly soluble. This is because of the fact that the large- N limit eliminates “nasty” classes of diagrams in the thermal Dyson-Schwinger expansion. The survived class of diagrams (superdaisy diagrams) can be exactly resumed, because the (thermal) proper self-energy Σ , as well as the renormalized coupling constant λ_r are momentum independent. We have also stressed that the $O(N)\phi^4$ theory in the large- N limit is consistent only if we view it as an effective field theory. Fortunately, the upper bound on the UV cutoff is truly huge, and it does not represent any significant restriction. For the model at hand the resummed form of the pressure with $m_r(0)=0$ was first derived (in the purely thermodynamic pressure context) by Drummond *et al.* in [39]. We have checked, using the prescription Eq. (3.30) for the thermal interaction pressure, that their results are in agreement with ours. The former is a nice vindication of the validity of the virial theorem for the QFT system at hand. In this connection we should perhaps mention that the latter is by no means obvious. For example, for quantized gauge fields the conformal (trace) anomaly may even invalidate the virial theorem [3]. The fact that this point is indeed nontrivial is illustrated on the QCD case in [68].

The expression for the pressure obtained was in a suitable form which allowed us to take advantage of the Mellin transform technique. We were then able to write down the high-temperature expansion for the pressure in $D=4$ (both for massive and massless fields) in terms of renormalized masses $m_r(T)$ and $m_r(0)$. We have explicitly checked that all UV divergences present in the individual thermal diagrams “miraculously” cancel in accordance with our analysis of the composite operators in Sec. III.

ACKNOWLEDGMENTS

I am indebted to P.V. Landshoff for reading the manuscript and for invaluable discussion. I am also grateful to N.P. Landsman, H. Osborn, R. Jackiw and H.J. Schnitzer for useful discussions. Finally I would like to thank the Fitzwilliam College of Cambridge and the Japanese Society for Promotion of Science for financial supports.

APPENDIX A

In this appendix we give some details of the derivation of Eq. (4.31). We particularly show that the surface integrals arisen during the transition from Eq. (4.30) to Eq. (4.31) mutually cancel among themselves. As usual, the integrals will be evaluated for integer values of D and corresponding results then analytically continued to a desired (generally complex) D .

The key quantity in question is

$$\begin{aligned} & + \frac{1}{2} \int \frac{d^D q}{(2\pi)^{D-1}} \left(\frac{2\mathbf{q}^2}{(D-1)} \right) \frac{\varepsilon(q_0)}{e^{\beta q_0} - 1} \delta(q^2 - m_r^2(T)) \\ & - \frac{1}{2} \int \frac{d^D q}{(2\pi)^{D-1}} \left(\frac{2\mathbf{q}^2}{(D-1)} \right) \delta^+(q^2 - m_r^2(0)). \quad (\text{A1}) \end{aligned}$$

Applying Green’s theorem (i.e., integrating by parts with respect to \mathbf{q}) on Eq. (A1) one finds

$$\begin{aligned} (\text{A1}) &= \mathcal{N}_T(m_r^2(T)) - \mathcal{N}(m_r^2(0)) \\ &+ \lim_{R \rightarrow \infty} \frac{1}{2(D-1)} \int \frac{dq_0}{(2\pi)^{D-1}} \int_{\partial S_R^{D-2}} d\mathbf{s} \mathbf{q} \\ &\times \theta(q^2 - m_r^2(T)) \theta(q_0) \left(\frac{2}{e^{\beta q_0} - 1} + 1 \right) \\ &- \lim_{R \rightarrow \infty} \frac{1}{2(D-1)} \int \frac{dq_0}{(2\pi)^{D-1}} \int_{\partial S_R^{D-2}} d\mathbf{s} \mathbf{q} \\ &\times \theta(q^2 - m_r^2(0)) \theta(q_0). \quad (\text{A2}) \end{aligned}$$

As usual, $\mathbf{ab} = \sum_{i=1}^{D-1} \mathbf{a}_i \mathbf{b}_i$ and S_R^{D-2} is a $(D-2)$ -sphere with the radius R . The expressions for \mathcal{N}_T and \mathcal{N} are done by Eq. (4.32).

With the relation Eq. (A3) we can show that the surface terms cancel in the large R limit. Let us first observe that

$$\begin{aligned} & \lim_{R \rightarrow \infty} \int \frac{dq_0}{(2\pi)^{D-1}} \int_{\partial S_R^{D-2}} d\mathbf{s} \mathbf{q} \theta(q^2 - m_r^2(T)) \frac{2\theta(q_0)}{e^{\beta q_0} - 1} \\ &= \lim_{R \rightarrow \infty} \frac{2\pi^{(D-1/2)} R^{D-1}}{\Gamma\left(\frac{D-1}{2}\right)} \int \frac{dq_0}{(2\pi)^{D-1}} \\ &\times \theta(q_0^2 - R^2 - m_r^2(T)) \frac{2\theta(q_0)}{e^{\beta q_0} - 1} \\ &= \lim_{R \rightarrow \infty} \frac{\pi^{(1-D/2)} R^{D-1}}{2^{D-2} \Gamma\left(\frac{D-1}{2}\right)} \int_{\sqrt{R^2 + m_r^2(T)}}^{\infty} dq_0 \frac{2}{e^{\beta q_0} - 1} = 0. \quad (\text{A3}) \end{aligned}$$

In the second line we have exploited Gauss’ theorem and in the last line we have used L’Hôpital’s rule as the expression is in the indeterminate form 0/0. The remaining surface terms in Eq. (A3) read

$$\begin{aligned} & \lim_{R \rightarrow \infty} \int \frac{dq_0}{(2\pi)^{D-1}} \int_{\partial S_R^{D-2}} d\mathbf{s} \mathbf{q} \\ &\times \{ \theta(q^2 - m_r^2(T)) - \theta(q^2 - m_r^2(0)) \} \theta(q_0) \\ &= \lim_{R \rightarrow \infty} \frac{\pi^{[(1-D)/2]} R^{D-1}}{2^{D-2} \Gamma\left(\frac{D-1}{2}\right)} \left\{ \int_{\sqrt{R^2 + m_r^2(T)}}^{\infty} - \int_{\sqrt{R^2 + m_r^2(0)}}^{\infty} \right\} \\ &\times dq_0 = 0. \quad (\text{A4}) \end{aligned}$$

The last identity follows either by applying L’Hôpital’s rule or by a simple transformation of variables which renders

both integrals inside of $\{ \dots \}$ equal. Expressions on the last lines in Eqs. (A3) and (A4) can be clearly (single-valuedly) continued to the region $\text{Re } D > 1$ as they are analytic there. We thus end up with the statement that

$$(A1) = \mathcal{N}_T(m_r^2(T)) - \mathcal{N}(m_r^2(0)).$$

APPENDIX B

In this appendix we shall derive the high-temperature expansion of the mass shift $\delta m^2(T)$ in the case when fields ϕ_a are massive [i.e., $m_r^2(0) \neq 0$].

Consider Eqs. (4.1) and (4.10). If we combine them together, we easily obtain the following transcendental equation for $\delta m^2(T)$:

$$\delta m^2(T) = \lambda_0 \left\{ \mathcal{M}(m_r^2(T)) - \mathcal{M}(m_r^2(0)) + \frac{1}{2} I_0(m_r^2(0) + \delta m^2(T)) \right\}. \quad (B1)$$

Here \mathcal{M} and I_0 are done by Eqs. (4.3) and (5.5), respectively.

Now, both λ_0 and \mathcal{M} are divergent in $D=4$. If we reexpress λ_0 in terms of λ_r , divergences must cancel, as $\delta m^2(T)$ is finite in $D=4$. The latter can be easily seen if we Taylor expand \mathcal{M} , i.e.,

$$\mathcal{M}(m_r^2(T)) = \mathcal{M}(m_r^2(0)) + \delta m^2(T) \mathcal{M}'(m_r^2(0)) + \hat{\mathcal{M}}(m_r^2(0); \delta m^2(T)). \quad (B2)$$

Obviously, $\hat{\mathcal{M}}$ is finite in $D=4$ as \mathcal{M} is quadratically divergent. Inserting Eq. (B2) to Eq. (B1) and employing Eq. (4.16) we get

$$\delta m^2(T) = \lambda_r \left\{ \hat{\mathcal{M}}(m_r^2(0); \delta m^2(T)) + \frac{1}{2} I_0(m_r^2(0) + \delta m^2(T)) \right\}. \quad (B3)$$

This is sometimes referred to as the renormalized gap equation. In order to determine $\hat{\mathcal{M}}$ we must go back to Eq. (B2). From the former we read off that

$$\begin{aligned} \hat{\mathcal{M}}(m_r^2(T); \delta m^2(T)) &= \mathcal{M}(m_r^2(T)) - \mathcal{M}(m_r^2(0)) - \delta m^2(T) \mathcal{M}'(m_r^2(0)) \\ &= \frac{\Gamma\left(1 - \frac{D}{2}\right)}{2(4\pi)^{D/2}} \left\{ (m_r^2(T))^{(D/2)-1} - (m_r^2(0))^{(D/2)-1} - \delta m^2(T) \left(\frac{D}{2} - 1\right) (m_r^2(0))^{(D/2)-2} \right\} \\ &\stackrel{D \rightarrow 4}{=} \frac{1}{32\pi^2} \left\{ m_r^2(T) \ln\left(\frac{m_r^2(T)}{m_r^2(0)}\right) - \delta m^2(T) \right\}. \end{aligned} \quad (B4)$$

So

$$\delta m^2(T) = \lambda_r \left\{ \frac{(m_r^2(0) + \delta m^2(T)) \ln\left(1 + \frac{\delta m^2(T)}{m_r^2(0)}\right) - \delta m^2(T)}{32\pi^2} + \frac{1}{2} I_0 \right\}. \quad (B5)$$

Analogous relation was also derived in [30] where authors used finite temperature renormalization group. In the latter the zero-momentum renormalization prescription was utilized. Equation (B5) was first obtained and numerically solved in [39]. It was shown that the solution is double valued. The former behavior was also observed in the effective action approach. Namely by Abbott *et al.* [33] at $T=0$, and by Bardeen and Moshe [31] at both $T=0$ and $T \neq 0$. The relevant solution is only that which fulfills the consistency condition $\delta m^2(T) \rightarrow 0$ when $T \rightarrow 0$. For such a solution it can be shown (cf. [39], Fig. 3) that $\delta m^2(T)/m_r^2(0) \ll 1$ for a sufficiently high T . So the high-temperature expansion of Eq. (B5) reads

$$\begin{aligned} \delta m^2(T) &= \lambda_r \left\{ \frac{\frac{(\delta m^2(T))^2}{2m_r^2(0)} - \frac{(\delta m^2(T))^3}{6m_r^4(0)} + \frac{(\delta m^2(T))^4}{12m_r^6(0)} + \dots}{32\pi^2} + \frac{1}{2} I_0 \right\} \\ &\doteq \frac{\lambda_r}{2} I_0 = \frac{\lambda_r T^2}{24} - \frac{\lambda_r m_r(T)}{8\pi} T + \mathcal{O}\left(m_r^2(T) \ln\left(\frac{m_r(T)}{T4\pi}\right)\right). \end{aligned} \quad (B6)$$

- [1] S. Habib, C. Molina-Paris, and E. Mottola, Phys. Rev. D **61**, 024010 (2000); L.M.A. Bettencourt, F. Cooper, and K. Pao, Phys. Rev. Lett. **89**, 112301 (2002); S.-Y. Wang, D. Boyanovsky, and Kin-Wang Ng, Nucl. Phys. **A699**, 819 (2002); S.-Y. Wang and D. Boyanovsky, Phys. Rev. D **63**, 051702 (2001).
- [2] F. Cooper, E. Mottola, and G.C. Nayak, Phys. Lett. B **555**, 181 (2003).
- [3] N.P. Landsman and Ch.G. van Weert, Phys. Rep. **145**, 141 (1987).
- [4] M. LeBellac, *Thermal Field Theory* (Cambridge University Press, Cambridge, 1996).
- [5] A. Das, *Finite Temperature Field Theory* (World Scientific, London, 1997).
- [6] J.I. Kapusta, *Finite-Temperature Field Theory* (Cambridge University Press, Cambridge, 1993).
- [7] T. Altherr, Int. J. Mod. Phys. A **8**, 5605 (1993).
- [8] A.L. Licht, Ann. Phys. (N.Y.) **34**, 161 (1965); N.P. Landsman, *ibid.* **34**, 141 (1981); P.A. Henning, Nucl. Phys. **B337**, 337 (1990); P.A. Henning, GSI Report No. 96-(1996), hep-ph/9605372.
- [9] E. Calzetta and B.L. Hu, Phys. Rev. D **37**, 2878 (1988).
- [10] R. Zwanzig, J. Chem. Phys. **33**, 138 (1960); *Nonequilibrium Statistical Mechanics* (Oxford University Press, Oxford, 2001); S. Nakajima, Prog. Theor. Phys. **20**, 948 (1958); D. Zubarev, V. Morozov, and G. Ropke, *Statistical Mechanics of Nonequilibrium Processes, Vol. 1, Basic Concepts, Kinetic Theory* (Wiley, Berlin, 1996).
- [11] P. Jizba and E.S. Tututi, Phys. Rev. D **60**, 105013 (1999).
- [12] K.C. Chou, Z.B. Su, B.L. Hao, and L. Yu, Phys. Rep. **118**, 1 (1985).
- [13] F. Cooper, S. Habib, Y. Kluger, E. Mottola, J.P. Paz, and P.R. Anderson, Phys. Rev. D **50**, 2848 (1994).
- [14] T.S. Evans and A.C. Pearson, Phys. Rev. D **52**, 4652 (1995).
- [15] L. Lapple, F. Mancini, and H. Umezawa, Phys. Rep. **10**, 151 (1974); T. Arimitsu and H. Umezawa, Prog. Theor. Phys. **77**, 32 (1987); **77**, 53 (1987); T. Arimitsu, Phys. Lett. A **153**, 163 (1991); T. Arimitsu and N. Arimitsu, Phys. Rev. E **50**, 121 (1994); T. Arimitsu, Condens. Matter Phys. **4**, 26 (1994); H. Umezawa, *Advanced Field Theory: Micro, Macro and Thermal Physics* (IOP, New York, 1993); I. Ojima, Ann. Phys. (N.Y.) **137**, 1 (1981).
- [16] P.A. Henning, Phys. Rep. **253**, 235 (1995).
- [17] C.-W. Kao, G.C. Nayak, and W. Greiner, Phys. Rev. D **66**, 034017 (2002).
- [18] E. Calzetta and B.L. Hu, Phys. Rev. D **35**, 495 (1987).
- [19] F. Cooper, hep-th/9504073.
- [20] M.V. Tokarchuk, T. Arimitsu, and A.E. Kobryn, Condens. Matter Phys. **1**, 605 (1998).
- [21] T. Haruyama, N. Kimura, and T. Nakamoto, Adv. Cryog. Eng. **43**, 781 (1998).
- [22] T. Tomita, J.J. Hamlin, J.S. Schilling, D.G. Hinks, and J.D. Jorgensen, Phys. Rev. B **64**, 092505 (2001); D. Andres, M.V. Kartsovnik, W. Biberacher, K. Neumaier, and H. Müller, cond-mat/0210062.
- [23] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena* (Oxford Science Publisher, Oxford, 2001).
- [24] C. Itzykson and J.B. Zuber, *Quantum Field Theory* (McGraw-Hill, New York, 1980).
- [25] J.M. Cornwall, R. Jackiw, and E. Tomboulis, Phys. Rev. D **10**, 2428 (1974).
- [26] E. Papp, Phys. Rep. **136**, 103 (1986).
- [27] P. Bucci and M. Pietroni, Phys. Rev. D **63**, 045026 (2001).
- [28] H. Kleinert and V. Schulte-Frohlinde, *Critical Properties of ϕ^4 -Theories* (World Scientific, Singapore, 2001).
- [29] H.J. Schnitzer, Phys. Rev. D **10**, 1800 (1974).
- [30] K. Funakubo and M. Sakamoto, Phys. Lett. B **186**, 205 (1987).
- [31] W.A. Bardeen and M. Moshe, Phys. Rev. D **28**, 1372 (1983).
- [32] J. Fröhlich, Nucl. Phys. **B200**, 281 (1982).
- [33] L.F. Abbot, J.S. Kang, and H.J. Schnitzer, Phys. Rev. D **13**, 2212 (1976).
- [34] S. Epstein, *The Variational Method in Quantum Chemistry* (Academic, New York, 1974); F. Cooper, S.-Y. Pi, and P.N. Stancioff, Phys. Rev. D **34**, 3831 (1986).
- [35] O. Éboli, R. Jackiw, and S.-Y. Pi, Phys. Rev. D **37**, 3557 (1988).
- [36] J. Collins, *Renormalization; An Introduction to Renormalization, the Renormalization Group, and the Operator-Product Expansion* (Cambridge University Press, Cambridge, 1984).
- [37] L.S. Brown, Ann. Phys. (N.Y.) **126**, 135 (1979).
- [38] M. van Eijck, "Thermal field theory and the finite-temperature renormalization group," Thesis, University of Amsterdam, 1995.
- [39] I.T. Drummond, R.R. Horgan, P.V. Landshoff, and A. Rebhan, Nucl. Phys. **B524**, 579 (1998).
- [40] G. Amelino-Camelia and S.-Y. Pi, Phys. Rev. D **47**, 2356 (1993).
- [41] G. Morandi, *Statistical Mechanics, An Intermediate Course* (World Scientific, London, 1995).
- [42] R. Kubo, *Statistical Mechanics* (North-Holland, Oxford, 1965).
- [43] H.B. Callen, *Thermodynamics and Introduction to Thermostatistics* (Wiley, New York, 1985).
- [44] M.C.J. Lermakers and Ch.G. van Weert, Nucl. Phys. **B248**, 671 (1984).
- [45] L. Dolan and R. Jackiw, Phys. Rev. D **9**, 3320 (1974).
- [46] I.T. Drummond, R.R. Horgan, P.V. Landshoff, and A. Rebhan, Phys. Lett. B **398**, 326 (1997).
- [47] S.R. de Groot, W.A. van Leeuwen, and Ch.G. van Weert, *Relativistic Kinetic Theory. Principles and Applications* (North-Holland, Oxford, 1980).
- [48] G.K. Batchelor, *An Introduction to Fluid Dynamics* (Cambridge University Press, London, 1967).
- [49] D.N. Zubarev, *Nonequilibrium Statistical Thermodynamics* (Consultants Bureau, London, 1974).
- [50] Y. Nambu, Prog. Theor. Phys. **7**, 131 (1952).
- [51] R. Jackiw, in *Current Algebra and Anomalies*, edited by S.B. Treiman, R. Jackiew, B. Zumino, and E. Witten (World Scientific, Singapore, 1985).
- [52] C.G. Callan, S. Coleman, and R. Jackiw, Ann. Phys. (N.Y.) **59**, 42 (1970).
- [53] G. Serman, *An Introduction to Quantum Field Theory* (Cambridge University Press, Cambridge, 1993).
- [54] M. Le Bellac and H. Mabilat, Phys. Lett. B **381**, 262 (1996).
- [55] F. Gelis, Phys. Lett. B **455**, 205 (1999); Z. Phys. C **70**, 321 (1996).

- [56] W. Zimmermann, in *Lectures on Elementary Particles and Quantum Field Theory*, edited by S. Deser *et al.* (M.I.T. Press, Cambridge, MA, 1970).
- [57] P. Cvitanovic, *Field Theory, Lecture Notes* (Nordita, Copenhagen, 1983); <http://www.cns.gatech.edu/FieldTheory/>
- [58] P. Jizba, Phys. Rev. D **57**, 3634 (1998).
- [59] A.L. Fetter and J.D. Walecka, *Quantum Theory of Many Particle Systems* (McGraw-Hill, New York, 1971).
- [60] B. Freeman, P. Smolensky, and D. Weingarten, Phys. Lett. B **113B**, 481 (1982).
- [61] R.G. Root, Phys. Rev. D **10**, 3322 (1974).
- [62] G. Amelino-Camelia, Phys. Lett. B **407**, 268 (1997).
- [63] H.W. Braden, Phys. Rev. D **25**, 1028 (1982).
- [64] H.E. Haber and E.A. Weldon, J. Math. Phys. **23**, 1852 (1982).
- [65] F. Oberhettinger, *Tables of Mellin Transforms* (Springer, Berlin, 1974).
- [66] A. Erdélyi, *Tables of Integral Transformations I* (McGraw-Hill, London, 1954).
- [67] I.S. Gradshteyn and I.M. Ryzhik, *Tables of Integrals, Series, and Products* (Academic, London, 1980).
- [68] I.T. Drummond, R.R. Horgan, P.V. Landshoff, and A. Rebhan, Phys. Lett. B **460**, 197 (1999).