

Measurement of the branching fractions for inclusive B^- and $\bar{B}^0$ decays to flavor-tagged D , D_s , and Λ_c

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ J.-M. Gaillard,¹ A. Hicheur,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ V. Tisserand,¹ A. Zghiche,¹ A. Palano,² A. Pompili,² J. C. Chen,³ N. D. Qi,³ G. Rong,³ P. Wang,³ Y. S. Zhu,³ G. Eigen,⁴ I. Ofte,⁴ B. Stugu,⁴ G. S. Abrams,⁵ A. W. Borgland,⁵ A. B. Breon,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵ R. N. Cahn,⁵ E. Charles,⁵ C. T. Day,⁵ M. S. Gill,⁵ A. V. Gritsan,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵ R. W. Kadel,⁵ J. Kadyk,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ G. Kukartsev,⁵ G. Lynch,⁵ L. M. Mir,⁵ P. J. Oddone,⁵ T. J. Orimoto,⁵ M. Pripstein,⁵ N. A. Roe,⁵ M. T. Ronan,⁵ V. G. Shelkov,⁵ W. A. Wenzel,⁵ M. Barrett,⁶ K. E. Ford,⁶ T. J. Harrison,⁶ A. J. Hart,⁶ C. M. Hawkes,⁶ S. E. Morgan,⁶ A. T. Watson,⁶ M. Fritsch,⁷ K. Goetzen,⁷ T. Held,⁷ H. Koch,⁷ B. Lewandowski,⁷ M. Pelizaeus,⁷ M. Steinke,⁷ J. T. Boyd,⁸ N. Chevalier,⁸ W. N. Cottingham,⁸ M. P. Kelly,⁸ T. E. Latham,⁸ F. F. Wilson,⁸ T. Cuhadar-Donszelmann,⁹ C. Hearty,⁹ N. S. Knecht,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹ D. Thiessen,⁹ A. Khan,¹⁰ P. Kyberd,¹⁰ L. Teodorescu,¹⁰ A. E. Blinov,¹¹ V. E. Blinov,¹¹ V. P. Druzhinin,¹¹ V. B. Golubev,¹¹ V. N. Ivanchenko,¹¹ E. A. Kravchenko,¹¹ A. P. Onuchin,¹¹ S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ A. N. Yushkov,¹¹ D. Best,¹² M. Bruinsma,¹² M. Chao,¹² I. Eschrich,¹² D. Kirkby,¹² A. J. Lankford,¹² M. Mandelkern,¹² R. K. Mommsen,¹² W. Roethel,¹² D. P. Stoker,¹² C. Buchanan,¹³ B. L. Hartfiel,¹³ S. D. Foulkes,¹⁴ J. W. Gary,¹⁴ B. C. Shen,¹⁴ K. Wang,¹⁴ D. del Re,¹⁵ H. K. Hadavand,¹⁵ E. J. Hill,¹⁵ D. B. MacFarlane,¹⁵ H. P. Paar,¹⁵ Sh. Rahatlou,¹⁵ V. Sharma,¹⁵ J. W. Berryhill,¹⁶ C. Campagnari,¹⁶ B. Dahmes,¹⁶ O. Long,¹⁶ A. Lu,¹⁶ M. A. Mazur,¹⁶ J. D. Richman,¹⁶ W. Verkerke,¹⁶ T. W. Beck,¹⁷ A. M. Eisner,¹⁷ C. A. Heusch,¹⁷ J. Kroseberg,¹⁷ W. S. Lockman,¹⁷ G. Nesom,¹⁷ T. Schalk,¹⁷ B. A. Schumm,¹⁷ A. Seiden,¹⁷ P. Spradlin,¹⁷ D. C. Williams,¹⁷ M. G. Wilson,¹⁷ J. Albert,¹⁸ E. Chen,¹⁸ G. P. Dubois-Felsmann,¹⁸ A. Dvoretzki,¹⁸ D. G. Hitlin,¹⁸ I. Narsky,¹⁸ T. Piatenko,¹⁸ F. C. Porter,¹⁸ A. Ryd,¹⁸ A. Samuel,¹⁸ S. Yang,¹⁸ S. Jayatilake,¹⁹ G. Mancinelli,¹⁹ B. T. Meadows,¹⁹ M. D. Sokoloff,¹⁹ T. Abe,²⁰ F. Blanc,²⁰ P. Bloom,²⁰ S. Chen,²⁰ W. T. Ford,²⁰ U. Nauenberg,²⁰ A. Olivas,²⁰ P. Rankin,²⁰ J. G. Smith,²⁰ J. Zhang,²⁰ L. Zhang,²⁰ A. Chen,²¹ J. L. Harton,²¹ A. Soffer,²¹ W. H. Toki,²¹ R. J. Wilson,²¹ Q. Zeng,²¹ D. Altenburg,²² T. Brandt,²² J. Brose,²² M. Dickopp,²² E. Feltresi,²² A. Hauke,²² H. M. Lacker,²² R. Müller-Pfefferkorn,²² R. Nogowski,²² S. Otto,²² A. Petzold,²² J. Schubert,²² K. R. Schubert,²² R. Schwierz,²² B. Spaan,²² J. E. Sundermann,²² D. Bernard,²³ G. R. Bonneaud,²³ F. Brochard,²³ P. Grenier,²³ S. Schrenk,²³ Ch. Thiebaux,²³ G. Vasileiadis,²³ M. Verderi,²³ D. J. Bard,²⁴ P. J. Clark,²⁴ D. Lavin,²⁴ F. Muheim,²⁴ S. Playfer,²⁴ Y. Xie,²⁴ M. Andreotti,²⁵ V. Azzolini,²⁵ D. Bettoni,²⁵ C. Bozzi,²⁵ R. Calabrese,²⁵ G. Cibinetto,²⁵ E. Luppi,²⁵ M. Negrini,²⁵ L. Piemontese,²⁵ A. Sarti,²⁵ E. Treadwell,²⁶ F. Anulli,²⁷ R. Baldini-Ferrolì,²⁷ A. Calcaterra,²⁷ R. de Sangro,²⁷ G. Finocchiaro,²⁷ P. Patteri,²⁷ I. M. Peruzzi,²⁷ M. Piccolo,²⁷ A. Zallo,²⁷ A. Buzzo,²⁸ R. Capra,²⁸ R. Contri,²⁸ G. Crosetti,²⁸ M. Lo Vetere,²⁸ M. Macri,²⁸ M. R. Monge,²⁸ S. Passaggio,²⁸ C. Patrignani,²⁸ E. Robutti,²⁸ A. Santroni,²⁸ S. Tosi,²⁸ S. Bailey,²⁹ G. Brandenburg,²⁹ K. S. Chaisanguanthum,²⁹ M. Morii,²⁹ E. Won,²⁹ R. S. Dubitzky,³⁰ U. Langenegger,³⁰ W. Bhimji,³¹ D. A. Bowerman,³¹ P. D. Dauncey,³¹ U. Egede,³¹ J. R. Gaillard,³¹ G. W. Morton,³¹ J. A. Nash,³¹ M. B. Nikolich,³¹ G. P. Taylor,³¹ M. J. Charles,³² G. J. Grenier,³² U. Mallik,³² J. Cochran,³³ H. B. Crawley,³³ J. Lamsa,³³ W. T. Meyer,³³ S. Prell,³³ E. I. Rosenberg,³³ A. E. Rubin,³³ J. Yi,³³ M. Biasini,³⁴ R. Covarelli,³⁴ M. Pioppi,³⁴ M. Davies,³⁵ X. Giroux,³⁵ G. Grosdidier,³⁵ A. Höcker,³⁵ S. Laplace,³⁵ F. Le Diberder,³⁵ V. Lepeltier,³⁵ A. M. Lutz,³⁵ T. C. Petersen,³⁵ S. Plaszczyński,³⁵ M. H. Schune,³⁵ L. Tantot,³⁵ G. Wormser,³⁵ C. H. Cheng,³⁶ D. J. Lange,³⁶ M. C. Simani,³⁶ D. M. Wright,³⁶ A. J. Bevan,³⁷ C. A. Chavez,³⁷ J. P. Coleman,³⁷ I. J. Forster,³⁷ J. R. Fry,³⁷ E. Gabathuler,³⁷ R. Gamet,³⁷ D. E. Hutchcroft,³⁷ R. J. Parry,³⁷ D. J. Payne,³⁷ R. J. Sloane,³⁷ C. Touramanis,³⁷ J. J. Back,^{38,*} C. M. Cormack,³⁸ P. F. Harrison,^{38,*} F. Di Lodovico,³⁸ G. B. Mohanty,^{38,*} C. L. Brown,³⁹ G. Cowan,³⁹ R. L. Flack,³⁹ H. U. Flaecher,³⁹ M. G. Green,³⁹ P. S. Jackson,³⁹ T. R. McMahon,³⁹ S. Ricciardi,³⁹ F. Salvatore,³⁹ M. A. Winter,³⁹ D. Brown,⁴⁰ C. L. Davis,⁴⁰ J. Allison,⁴¹ N. R. Barlow,⁴¹ R. J. Barlow,⁴¹ P. A. Hart,⁴¹ M. C. Hodgkinson,⁴¹ G. D. Lafferty,⁴¹ A. J. Lyon,⁴¹ J. C. Williams,⁴¹ A. Farbin,⁴² W. D. Hulsbergen,⁴² A. Jawahery,⁴² D. Kovalskyi,⁴² C. K. Lae,⁴² V. Lillard,⁴² D. A. Roberts,⁴² G. Blaylock,⁴³ C. Dallapiccola,⁴³ K. T. Flood,⁴³ S. S. Hertzbach,⁴³ R. Kofler,⁴³ V. B. Koptchev,⁴³ T. B. Moore,⁴³ S. Saremi,⁴³ H. Staengle,⁴³ S. Willocq,⁴³ R. Cowan,⁴⁴ G. Sciolla,⁴⁴ S. J. Sekula,⁴⁴ F. Taylor,⁴⁴ R. K. Yamamoto,⁴⁴ D. J. J. Mangeol,⁴⁵ P. M. Patel,⁴⁵ S. H. Robertson,⁴⁵ A. Lazzaro,⁴⁶ V. Lombardo,⁴⁶ F. Palombo,⁴⁶ J. M. Bauer,⁴⁷ L. Cremaldi,⁴⁷ V. Eschenburg,⁴⁷ R. Godang,⁴⁷ R. Kroeger,⁴⁷ J. Reidy,⁴⁷ D. A. Sanders,⁴⁷ D. J. Summers,⁴⁷ H. W. Zhao,⁴⁷ S. Brunet,⁴⁸ D. Côté,⁴⁸ P. Taras,⁴⁸ H. Nicholson,⁴⁹ N. Cavallo,^{50,†} F. Fabozzi,^{50,†} C. Gatto,⁵⁰ L. Lista,⁵⁰ D. Monorchio,⁵⁰ P. Paolucci,⁵⁰ D. Piccolo,⁵⁰ C. Sciacca,⁵⁰ M. Baak,⁵¹ H. Bulten,⁵¹ G. Raven,⁵¹ H. L. Snoek,⁵¹ L. Wilden,⁵¹ C. P. Jessop,⁵² J. M. LoSecco,⁵² T. Allmendinger,⁵³ K. K. Gan,⁵³ K. Honscheid,⁵³ D. Hufnagel,⁵³ H. Kagan,⁵³ R. Kass,⁵³ T. Pulliam,⁵³ A. M. Rahimi,⁵³ R. Ter-Antonyan,⁵³

Q. K. Wong,⁵³ J. Brau,⁵⁴ R. Frey,⁵⁴ O. Igonkina,⁵⁴ C. T. Potter,⁵⁴ N. B. Sinev,⁵⁴ D. Strom,⁵⁴ E. Torrence,⁵⁴ F. Colecchia,⁵⁵ A. Dorigo,⁵⁵ F. Galeazzi,⁵⁵ M. Margoni,⁵⁵ M. Morandin,⁵⁵ M. Posocco,⁵⁵ M. Rotondo,⁵⁵ F. Simonetto,⁵⁵ R. Stroili,⁵⁵ G. Tiozzo,⁵⁵ C. Voci,⁵⁵ M. Benayoun,⁵⁶ H. Briand,⁵⁶ J. Chauveau,⁵⁶ P. David,⁵⁶ Ch. de la Vaissière,⁵⁶ L. Del Buono,⁵⁶ O. Hamon,⁵⁶ M. J. J. John,⁵⁶ Ph. Leruste,⁵⁶ J. Malcles,⁵⁶ J. Ocariz,⁵⁶ M. Pivk,⁵⁶ L. Roos,⁵⁶ S. T'Jampens,⁵⁶ G. Therin,⁵⁶ P. F. Manfredi,⁵⁷ V. Re,⁵⁷ P. K. Behera,⁵⁸ L. Gladney,⁵⁸ Q. H. Guo,⁵⁸ J. Panetta,⁵⁸ C. Angelini,⁵⁹ G. Batignani,⁵⁹ S. Bettarini,⁵⁹ M. Bondioli,⁵⁹ F. Bucci,⁵⁹ G. Calderini,⁵⁹ M. Carpinelli,⁵⁹ F. Forti,⁵⁹ M. A. Giorgi,⁵⁹ A. Lusiani,⁵⁹ G. Marchiori,⁵⁹ F. Martinez-Vidal,^{59,‡} M. Morganti,⁵⁹ N. Neri,⁵⁹ E. Paoloni,⁵⁹ M. Rama,⁵⁹ G. Rizzo,⁵⁹ F. Sandrelli,⁵⁹ J. Walsh,⁵⁹ M. Haire,⁶⁰ D. Judd,⁶⁰ K. Paick,⁶⁰ D. E. Wagoner,⁶⁰ N. Danielson,⁶¹ P. Elmer,⁶¹ Y. P. Lau,⁶¹ C. Lu,⁶¹ V. Miftakov,⁶¹ J. Olsen,⁶¹ A. J. S. Smith,⁶¹ A. V. Telnov,⁶¹ F. Bellini,⁶² G. Cavoto,^{61,62} R. Faccini,⁶² F. Ferrarotto,⁶² F. Ferroni,⁶² M. Gaspero,⁶² L. Li Gioi,⁶² M. A. Mazzoni,⁶² S. Morganti,⁶² M. Pierini,⁶² G. Piredda,⁶² F. Safai Tehrani,⁶² C. Voena,⁶² S. Christ,⁶³ G. Wagner,⁶³ R. Walldi,⁶³ T. Adye,⁶⁴ N. De Groot,⁶⁴ B. Franek,⁶⁴ N. I. Geddes,⁶⁴ G. P. Gopal,⁶⁴ E. O. Olaiya,⁶⁴ R. Aleksan,⁶⁵ S. Emery,⁶⁵ A. Gaidot,⁶⁵ S. F. Ganzhur,⁶⁵ P.-F. Giraud,⁶⁵ G. Hamel de Monchenault,⁶⁵ W. Kozanecki,⁶⁵ M. Legendre,⁶⁵ G. W. London,⁶⁵ B. Mayer,⁶⁵ G. Schott,⁶⁵ G. Vasseur,⁶⁵ Ch. Yèche,⁶⁵ M. Zito,⁶⁵ M. V. Purohit,⁶⁶ A. W. Weidemann,⁶⁶ J. R. Wilson,⁶⁶ F. X. Yumiceva,⁶⁶ D. Aston,⁶⁷ R. Bartoldus,⁶⁷ N. Berger,⁶⁷ A. M. Boyarski,⁶⁷ O. L. Buchmueller,⁶⁷ R. Claus,⁶⁷ M. R. Convery,⁶⁷ M. Cristinziani,⁶⁷ G. De Nardo,⁶⁷ D. Dong,⁶⁷ J. Dorfan,⁶⁷ D. Dujmic,⁶⁷ W. Dunwoodie,⁶⁷ E. E. Elsen,⁶⁷ S. Fan,⁶⁷ R. C. Field,⁶⁷ T. Glanzman,⁶⁷ S. J. Gowdy,⁶⁷ T. Hadig,⁶⁷ V. Halyo,⁶⁷ C. Hast,⁶⁷ T. Hryn'ova,⁶⁷ W. R. Innes,⁶⁷ M. H. Kelsey,⁶⁷ P. Kim,⁶⁷ M. L. Kocian,⁶⁷ D. W. G. S. Leith,⁶⁷ J. Libby,⁶⁷ S. Luitz,⁶⁷ V. Luth,⁶⁷ H. L. Lynch,⁶⁷ H. Marsiske,⁶⁷ R. Messner,⁶⁷ D. R. Muller,⁶⁷ C. P. O'Grady,⁶⁷ V. E. Ozcan,⁶⁷ A. Perazzo,⁶⁷ M. Perl,⁶⁷ S. Petrak,⁶⁷ B. N. Ratcliff,⁶⁷ A. Roodman,⁶⁷ A. A. Salnikov,⁶⁷ R. H. Schindler,⁶⁷ J. Schwiening,⁶⁷ G. Simi,⁶⁷ A. Snyder,⁶⁷ A. Soha,⁶⁷ J. Stelzer,⁶⁷ D. Su,⁶⁷ M. K. Sullivan,⁶⁷ J. Va'vra,⁶⁷ S. R. Wagner,⁶⁷ M. Weaver,⁶⁷ A. J. R. Weinstein,⁶⁷ W. J. Wisniewski,⁶⁷ M. Wittgen,⁶⁷ D. H. Wright,⁶⁷ A. K. Yarritu,⁶⁷ C. C. Young,⁶⁷ P. R. Burchat,⁶⁸ A. J. Edwards,⁶⁸ T. I. Meyer,⁶⁸ B. A. Petersen,⁶⁸ C. Roat,⁶⁸ S. Ahmed,⁶⁹ M. S. Alam,⁶⁹ J. A. Ernst,⁶⁹ M. A. Saeed,⁶⁹ M. Saleem,⁶⁹ F. R. Wappler,⁶⁹ W. Bugg,⁷⁰ M. Krishnamurthy,⁷⁰ S. M. Spanier,⁷⁰ R. Eckmann,⁷¹ H. Kim,⁷¹ J. L. Ritchie,⁷¹ A. Satpathy,⁷¹ R. F. Schwitters,⁷¹ J. M. Izen,⁷² I. Kitayama,⁷² X. C. Lou,⁷² S. Ye,⁷² F. Bianchi,⁷³ M. Bona,⁷³ F. Gallo,⁷³ D. Gamba,⁷³ L. Bosisio,⁷⁴ C. Cartaro,⁷⁴ F. Cossutti,⁷⁴ G. Della Ricca,⁷⁴ S. Dittongo,⁷⁴ S. Grancagnolo,⁷⁴ L. Lanceri,⁷⁴ P. Poropat,^{74,§} L. Vitale,⁷⁴ G. Vuagnin,⁷⁴ R. S. Panvini,⁷⁵ Sw. Banerjee,⁷⁶ C. M. Brown,⁷⁶ D. Fortin,⁷⁶ P. D. Jackson,⁷⁶ R. Kowalewski,⁷⁶ J. M. Roney,⁷⁶ R. J. Sobie,⁷⁶ H. R. Band,⁷⁷ B. Cheng,⁷⁷ S. Dasu,⁷⁷ M. Datta,⁷⁷ A. M. Eichenbaum,⁷⁷ M. Graham,⁷⁷ J. J. Hollar,⁷⁷ J. R. Johnson,⁷⁷ P. E. Kutter,⁷⁷ H. Li,⁷⁷ R. Liu,⁷⁷ A. Mihalys,⁷⁷ A. K. Mohapatra,⁷⁷ Y. Pan,⁷⁷ R. Prepost,⁷⁷ P. Tan,⁷⁷ J. H. von Wimmersperg-Toeller,⁷⁷ J. Wu,⁷⁷ S. L. Wu,⁷⁷ Z. Yu,⁷⁷ M. G. Greene,⁷⁸ and H. Neal⁷⁸

(BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

³Institute of High Energy Physics, Beijing 100039, China

⁴Institute of Physics, University of Bergen, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁷Institut für Experimentalphysik I, Ruhr Universität Bochum, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 1TL, United Kingdom

⁹University of British Columbia, Vancouver, BC, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, California 92697, USA

¹³University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁴University of California at Riverside, Riverside, California 92521, USA

¹⁵University of California at San Diego, La Jolla, California 92093, USA

¹⁶University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁷Institute for Particle Physics, University of California at Santa Cruz, Santa Cruz, California 95064, USA

¹⁸California Institute of Technology, Pasadena, California 91125, USA

¹⁹University of Cincinnati, Cincinnati, Ohio 45221, USA

²⁰University of Colorado, Boulder, Colorado 80309, USA

²¹Colorado State University, Fort Collins, Colorado 80523, USA

²²Institut für Kern- und Teilchenphysik, Technische Universität Dresden, D-01062 Dresden, Germany

²³Ecole Polytechnique, LLR, F-91128 Palaiseau, France

- ²⁴University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
- ²⁵Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
- ²⁶Florida A&M University, Tallahassee, Florida 32307, USA
- ²⁷Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
- ²⁸Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
- ²⁹Harvard University, Cambridge, Massachusetts 02138, USA
- ³⁰Physikalisches Institut, Universität Heidelberg, Philosophenweg 12, D-69120 Heidelberg, Germany
- ³¹Imperial College London, London, SW7 2AZ, United Kingdom
- ³²University of Iowa, Iowa City, Iowa 52242, USA
- ³³Iowa State University, Ames, Iowa 50011-3160, USA
- ³⁴Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
- ³⁵Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
- ³⁶Lawrence Livermore National Laboratory, Livermore, California 94550, USA
- ³⁷University of Liverpool, Liverpool L69 7ZE, United Kingdom
- ³⁸Queen Mary, University of London, E1 4NS, United Kingdom
- ³⁹University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
- ⁴⁰University of Louisville, Louisville, Kentucky 40292, USA
- ⁴¹University of Manchester, Manchester M13 9PL, United Kingdom
- ⁴²University of Maryland, College Park, Maryland 20742, USA
- ⁴³University of Massachusetts, Amherst, Massachusetts 01003, USA
- ⁴⁴Laboratory for Nuclear Science, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA
- ⁴⁵McGill University, Montréal, QC, Canada H3A 2T8
- ⁴⁶Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
- ⁴⁷University of Mississippi, University, Mississippi 38677, USA
- ⁴⁸Laboratoire René J. A. Lévesque, Université de Montréal, Montréal, QC, Canada H3C 3J7
- ⁴⁹Mount Holyoke College, South Hadley, Massachusetts 01075, USA
- ⁵⁰Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
- ⁵¹NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
- ⁵²University of Notre Dame, Notre Dame, Indiana 46556, USA
- ⁵³The Ohio State University, Columbus, Ohio 43210, USA
- ⁵⁴University of Oregon, Eugene, Oregon 97403, USA
- ⁵⁵Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
- ⁵⁶Laboratoire de Physique Nucléaire et de Hautes Energies, Universités Paris VI et VII, F-75252 Paris, France
- ⁵⁷Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy
- ⁵⁸University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
- ⁵⁹Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
- ⁶⁰Prairie View A&M University, Prairie View, Texas 77446, USA
- ⁶¹Princeton University, Princeton, New Jersey 08544, USA
- ⁶²Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
- ⁶³Universität Rostock, D-18051 Rostock, Germany
- ⁶⁴Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
- ⁶⁵DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
- ⁶⁶University of South Carolina, Columbia, South Carolina 29208, USA
- ⁶⁷Stanford Linear Accelerator Center, Stanford, California 94309, USA
- ⁶⁸Stanford University, Stanford, California 94305-4060, USA
- ⁶⁹State University of New York, Albany, New York 12222, USA
- ⁷⁰University of Tennessee, Knoxville, Tennessee 37996, USA
- ⁷¹University of Texas at Austin, Austin, Texas 78712, USA
- ⁷²University of Texas at Dallas, Richardson, Texas 75083, USA
- ⁷³Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
- ⁷⁴Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
- ⁷⁵Vanderbilt University, Nashville, Tennessee 37235, USA
- ⁷⁶University of Victoria, Victoria, BC, Canada V8W 3P6
- ⁷⁷University of Wisconsin, Madison, Wisconsin 53706, USA

*Now at Department of Physics, University of Warwick, Coventry, United Kingdom.

†Also with Università della Basilicata, Potenza, Italy.

‡Also with IFIC, Instituto de Física Corpuscular, CSIC-Universidad de Valencia, Valencia, Spain.

§Deceased.

⁷⁸*Yale University, New Haven, Connecticut 06511, USA*
(Received 23 August 2004; published 29 November 2004)

We report on the inclusive branching fractions of B^- and of $\bar{B}^0$ mesons decaying to D^0X , $\bar{D}^0X$, D^+X , D^-X , D_s^+X , D_s^-X , Λ_c^+X , $\bar{\Lambda}_c^-X$, based on a sample of 88.9×10^6 $B\bar{B}$ events recorded with the *BABAR* detector at the $\Upsilon(4S)$ resonance. Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. We measure the number of charmed and of anticharmed particles per B decay and derive the total charm yield per B^- decay $n_c^- = 1.313 \pm 0.037 \pm 0.062^{+0.063}_{-0.042}$ and per $\bar{B}^0$ decay $n_c^0 = 1.276 \pm 0.062 \pm 0.058^{+0.066}_{-0.046}$, where the first uncertainty is statistical, the second is systematic, and the third reflects the charm branching-fraction uncertainties.

DOI: 10.1103/PhysRevD.70.091106

PACS numbers: 13.25.Hw, 11.30.Er, 12.15.Hh

The dominant process for the decay of a b quark is $b \rightarrow cW^{*-}$ [1], resulting in a (flavor) correlated c quark and a virtual W . In the decay of the W , the production of a $\bar{u}d$ or a $\bar{c}s$ pair are both Cabibbo-allowed and should be equal, the latter being suppressed only by a phase-space factor. The first process dominates hadronic b decays, while the second can be easily distinguished as it will produce a (flavor) anticorrelated $\bar{c}$ quark. Experimentally, correlated and anticorrelated charm production can be investigated through the measurement of the inclusive B -decay rates to flavor-tagged charmed mesons or baryons. Current measurements [2–4] of these rates have statistically limited precision and do not distinguish among the different B parent states.

Most of the charged and neutral D mesons produced in $\bar{B}$ decays come from correlated production $\bar{B} \rightarrow DX$. However, a significant number of $\bar{B} \rightarrow \bar{D}X$ decays are expected through $b \rightarrow c\bar{c}s$ transitions, such as $\bar{B} \rightarrow D^{(*)}\bar{D}^{(*)}\bar{K}^{(*)}(n\pi)$. Although the branching fractions of the three-body decays $\bar{B} \rightarrow D^{(*)}\bar{D}^{(*)}\bar{K}$ have been measured [5,6], it is not clear whether they saturate $\bar{B} \rightarrow \bar{D}X$ transitions. It is therefore important to improve the precision on the branching fraction $\mathcal{B}(\bar{B} \rightarrow \bar{D}X)$.

By contrast, the anticorrelated D_s^- production $\bar{B} \rightarrow D_s^-D(n\pi)$ is expected to dominate $\bar{B}$ decays to D_s mesons, since correlated production needs an extra $s\bar{s}$ pair created from the vacuum to give $\bar{B} \rightarrow D_s^+K^-(n\pi)$. There is no prior published measurement of $\mathcal{B}(\bar{B} \rightarrow D_s^+X)$.

All strangeless charmed baryons decay to Λ_c . Correlated Λ_c are produced in decays such as $B^- \rightarrow \Lambda_c^+\bar{p}\pi^-(\pi)$, while anticorrelated $\bar{\Lambda}_c^-$ should originate from $B^- \rightarrow \Xi_c^-\bar{\Lambda}_c^-(\pi)$. Another possibility is $B^- \rightarrow \Lambda_c^+\bar{\Lambda}_c^-K^-$, the baryonic analogue of the $D\bar{D}K$ decay. The rates for Ξ_c production in B decays [7] are unknown, because there is no absolute measurement of Ξ_c decay branching fractions.

This analysis uses $\Upsilon(4S) \rightarrow B\bar{B}$ events in which either a B^+ or a B^0 meson (hereafter denoted B_{reco}) decays into a hadronic final state and is fully reconstructed. We then reconstruct D , D_s , and Λ_c from the recoiling B^- ($\bar{B}^0$) meson and compare the flavor of the charm hadron with that of the B_{reco} , thus allowing separate measurements of the B^- ($\bar{B}^0$) $\rightarrow D^0X$, D^+X , D_s^+X , Λ_c^+X and B^- ($\bar{B}^0$) $\rightarrow$

$\bar{D}^0X$, D^-X , D_s^-X , $\bar{\Lambda}_c^-X$ branching fractions. We extract $\mathcal{B}(B^- \rightarrow \Lambda_c^+\bar{\Lambda}_c^-K^-)$ from the missing-mass spectra of the $\Lambda_c^+K^-$ or $\bar{\Lambda}_c^-K^-$ systems recoiling against the B_{reco} . We can then evaluate indirectly $\mathcal{B}(B^- \rightarrow \Xi_c^-X) = \mathcal{B}(B^- \rightarrow \bar{\Lambda}_c^-X) - \mathcal{B}(B^- \rightarrow \Lambda_c^+\bar{\Lambda}_c^-K^-)$ and compute the average number of charm (anticharm) particles per B^- decay, N_c^- ($N_{\bar{c}}^-$):

$$N_c^- = \sum_{X_c} \mathcal{B}(B^- \rightarrow X_cX), \quad (1)$$

$$N_{\bar{c}}^- = \sum_{\bar{X}_c} \mathcal{B}(B^- \rightarrow \bar{X}_cX), \quad (2)$$

where the sum is performed over $X_c = D^+, D^0, D_s^+, \Lambda_c^+, \Xi_c, (c\bar{c})$ or $\bar{X}_c = D^-, \bar{D}^0, D_s^-, \bar{\Lambda}_c^-, (c\bar{c})$, and $(c\bar{c})$ refers to all charmonium states collectively. We neglect Ξ_c production, as it requires both a $\bar{c}s$ and an $s\bar{s}$ pair in the decay to give $\Xi_c\Omega_c$. We can sum N_c^- and $N_{\bar{c}}^-$ to obtain the average number of charm plus anticharm quarks per B^- decay, $n_c^- = N_c^- + N_{\bar{c}}^-$ (and similarly for $\bar{B}^0$ decays). In addition to the theoretical interest [8–11], the fact that anticorrelated charmed particles are a background for many studies also motivates a more precise measurement of their production rates in B decays.

The measurements presented here are based on a sample of 88.9×10^6 $B\bar{B}$ pairs (81.9 fb^{-1}) recorded at the $\Upsilon(4S)$ resonance with the *BABAR* detector at the PEP-II asymmetric-energy B -meson factory at the Stanford Linear Accelerator Center (SLAC). The *BABAR* detector is described in detail elsewhere [12]. Charged-particle trajectories are measured by a 5-layer double-sided silicon vertex tracker and a 40-layer drift chamber, both operating in a 1.5-T solenoidal magnetic field. Charged-particle identification is provided by the average energy loss (dE/dx) in the tracking devices and by an internally reflecting ring-imaging Cherenkov detector. Photons are detected by a CsI(Tl) electromagnetic calorimeter. We use Monte Carlo simulations of the *BABAR* detector based on GEANT4 [13] to optimize selection criteria and determine selection efficiencies.

We reconstruct B^+ and B^0 decays (B_{reco}) in the modes $B^+ \rightarrow \bar{D}^{(*)0}\pi^+$, $\bar{D}^{(*)0}\rho^+$, $\bar{D}^{(*)0}a_1^+$ and $B^0 \rightarrow D^{(*)-}\pi^+$,




FIG. 1 (color online). m_{ES} spectra of reconstructed (a) B^+ and (b) B^0 candidates. The solid vertical line shows the upper limit of the background control region (hatched), the dotted vertical line the lower limit of the B signal region. The crossed area shows the background under the B signal. The solid curve is the sum of the fitted signal and background; the dashed curve is the background component only.

$D^{(*)-} \rho^+$, $D^{(*)-} a_1^+$, $\bar{D}^0$ candidates are reconstructed in the $K^+ \pi^-$, $K^+ \pi^- \pi^0$, $K^+ \pi^- \pi^+ \pi^-$, and $K_S^0 \pi^+ \pi^-$ ($K_S^0 \rightarrow \pi^+ \pi^-$) decay channels, while D^- are reconstructed in the $K^+ \pi^- \pi^-$ and $K_S^0 \pi^- \pi^-$ modes. D^* candidates are reconstructed in the $D^{*-} \rightarrow \bar{D}^0 \pi^-$ and $\bar{D}^{*0} \rightarrow \bar{D}^0 \pi^0$, $\bar{D}^0 \gamma$ decay modes. The first kinematic variable used to identify fully reconstructed B decays is the beam-energy substituted mass, $m_{ES} = \sqrt{(s/2 + \mathbf{p}_i \cdot \mathbf{p}_B)^2/E_i^2 - \mathbf{p}_B^2}$, where $\mathbf{p}_B$ is the B_{reco} momentum and $(E_i, \mathbf{p}_i)$ is the four-momentum

of the initial e^+e^- system, both measured in the laboratory frame. The invariant mass of the initial e^+e^- system is $\sqrt{s}$. The second variable is $\Delta E = E_B^* - \sqrt{s}/2$, where E_B^* is the B_{reco} candidate energy in the center-of-mass frame. We require $|\Delta E| < n\sigma_{\Delta E}$ with $n = 2$ or 3 , depending on the decay mode, and using the measured resolution $\sigma_{\Delta E}$ for each decay mode.

In the m_{ES} spectra (Fig. 1), we define a signal region with $5.274 < m_{ES} < 5.290$ GeV/c^2 and a background control region with $5.220 < m_{ES} < 5.260$ GeV/c^2 . For each of the B -decay modes, the combinatorial background in the signal region is derived from a fit to the m_{ES} distribution that uses an empirical phase-space threshold function [14] for the background, together with a signal function [15] peaked at the B meson mass. The numbers of reconstructed B^+ and B^0 candidates, $N_{B^+} = 85840 \pm 1910(\text{syst})$ and $N_{B^0} = 48322 \pm 590(\text{syst})$, are then obtained by subtracting this background from the total number of events found in the signal region. These measured B meson yields provide the normalization of all branching-fraction measurements reported below. The systematic uncertainties quoted above are computed by varying the boundaries of the signal and background regions and by comparing the shapes of the threshold function [14] in the data and in the simulation.

The contamination of B^0 events in the B^+ signal induces a background which peaks near the B mass. From the Monte Carlo simulation, the fraction of B^0 events in the reconstructed B^+ signal sample is found to be $c_0 = 0.034$ and the fraction of B^+ events in the reconstructed B^0 signal sample to be $c_+ = 0.019$. A 100% systematic uncertainty is conservatively assigned to these numbers but they will have a small effect on the final results.

We now turn to the analysis of inclusive D , D_s , and Λ_c production in the decays of the $\bar{B}$'s that recoil against the reconstructed B . Charmed particles X_c (correlated production) are distinguished from anticharmed particles $\bar{X}_c$ (anticorrelated production). They are reconstructed from charged tracks that do not belong to the B_{reco} . The decay modes considered are listed in Table I.

For charged B decays, Fig. 2 shows the D , D_s , and Λ_c mass spectra of correlated and anticorrelated candidates

TABLE I. Charmed-particle signal yields and B branching fractions per decay mode. The first uncertainty is statistical, the second is systematic (but does not include the charm branching-fraction uncertainties).

X_c decay mode	$B^- \rightarrow X_c X$		$B^- \rightarrow \bar{X}_c X$		$\bar{B}^0 \rightarrow X_c X$		$\bar{B}^0 \rightarrow \bar{X}_c X$	
	Yield	$\mathcal{B}(\%)$	Yield	$\mathcal{B}(\%)$	Yield	$\mathcal{B}(\%)$	Yield	$\mathcal{B}(\%)$
D^0 [16] $\rightarrow K^- \pi^+$	1273 ± 42	$79.2 \pm 2.6 \pm 3.9$	160 ± 16	$9.3 \pm 1.0 \pm 0.5$	397 ± 24	$50.3 \pm 3.4 \pm 2.4$	139 ± 14	$7.3 \pm 2.2 \pm 0.5$
$\rightarrow K^- \pi^+ \pi^- \pi^+$	998 ± 65	$80.6 \pm 5.3 \pm 7.5$	173 ± 30	$13.4 \pm 2.4 \pm 1.3$	332 ± 36	$56.2 \pm 6.8 \pm 5.4$	83 ± 23	$1.8 \pm 4.4 \pm 0.5$
D^+ $\rightarrow K^- \pi^+ \pi^+$	262 ± 29	$9.8 \pm 1.2 \pm 1.2$	98 ± 20	$3.8 \pm 0.9 \pm 0.4$	452 ± 31	$39.7 \pm 3.0 \pm 2.8$	125 ± 18	$2.3 \pm 1.8 \pm 0.3$
D_s^+ $\rightarrow \phi \pi^+$	11 ± 5	$2.2 \pm 1.1 \pm 0.3$	82 ± 11	$16.5 \pm 2.3 \pm 1.7$	24 ± 6	$8.3 \pm 2.8 \pm 0.8$	28 ± 6	$9.9 \pm 2.9 \pm 1.0$
$\rightarrow \bar{K}^{*0} K^+$	0 ± 3	$0.0 \pm 1.1 \pm 0.2$	55 ± 11	$18.0 \pm 3.5 \pm 1.7$	3 ± 4	$0.0 \pm 2.8 \pm 0.1$	14 ± 5	$9.9 \pm 4.1 \pm 1.2$
$\rightarrow K_S^0 K^+$	0 ± 3	$0.0 \pm 0.9 \pm 0.2$	31 ± 9	$9.2 \pm 2.7 \pm 0.8$	12 ± 5	$5.0 \pm 3.4 \pm 0.4$	23 ± 6	$13.3 \pm 4.3 \pm 1.0$
Λ_c^+ $\rightarrow p K^- \pi^+$	41 ± 9	$3.5 \pm 0.8 \pm 0.3$	33 ± 9	$2.9 \pm 0.8 \pm 0.3$	28 ± 8	$4.9 \pm 1.7 \pm 0.4$	16 ± 6	$2.0 \pm 1.2 \pm 0.2$




FIG. 2 (color online). Correlated (left) and anticorrelated (right) charmed-particle mass spectra in the recoil of B^+ events, for (a),(b) $D^0 \rightarrow K^- \pi^+$; (c),(d) $D^+ \rightarrow K^- \pi^+ \pi^+$; (e),(f) $D_s^+ \rightarrow \phi \pi^+$; and (g),(h) $\Lambda_c^+ \rightarrow p K^- \pi^+$. The solid curve is the sum of a Gaussian signal and of a linear background plus mode-dependent satellite contributions [17]. The shaded areas show the contribution of well reconstructed D , D_s , or Λ_c in the B^+ combinatorial background.

recoiling against B 's reconstructed in the m_{ES} signal region, for some selected decay modes. These spectra are fitted with the sum of a Gaussian signal and a linear background (including a satellite peak for some channels [17]). The shaded areas correspond to well reconstructed D , D_s , or Λ_c from the combinatorial B_{reco} background. They are obtained from data in the m_{ES} background control region, normalized to the number of combinatorial background events expected under the B_{reco} peak. The background-subtracted reconstructed signal yields are listed in Table I. The reconstruction efficiencies for each charmed (anticharmed) final state $X_c \rightarrow f$ ($\bar{X}_c \rightarrow \bar{f}$) are computed from the simulation as a function of the charmed-particle momentum in the B^- center-of-mass

frame and are applied event by event to obtain the efficiency-corrected charm signal yields $N(X_c \rightarrow f)$ [$N(\bar{X}_c \rightarrow \bar{f})$]. The final branching fractions are computed from these yields, the number of B_{reco} , and the intermediate branching fractions $\mathcal{B}(X_c \rightarrow f)$ taken from Ref. [18]. They are given by

$$\mathcal{B}(B^- \rightarrow X_c X) = \frac{N(X_c \rightarrow f)}{N_{B^+} \times \mathcal{B}(X_c \rightarrow f)} - c_0 \mathcal{B}_0. \quad (3)$$

Here the raw branching fraction for $B^- \rightarrow X_c X$ is modified by a small corrective term $c_0 \mathcal{B}_0$ that accounts for the B^0 contamination in the reconstructed B^+ sample. The factor $\mathcal{B}_0$ depends on the measured $\bar{B}^0 \rightarrow X_c X$ and $B^0 \rightarrow X_c X$ branching fractions and on the $B^0 - \bar{B}^0$ mixing parameter χ_d [18]. It ranges from less than 3% for Λ_c to as much as 50% for correlated D^0 and D^+ . Doubly Cabibbo-suppressed D^0 decays are also taken into account. The branching fractions and their errors are given in Table I. The statistical and systematic uncertainties are computed separately for each channel. For example, the 3.9% absolute systematic uncertainty on $\mathcal{B}[B^- \rightarrow D^0(K^- \pi^+)X]$ reflects the quadratic sum of 1.8% attributed to N_{B^+} , 1.3% to the error on the rate of true D 's in the B combinatorial background, 0.8% to the Monte Carlo statistics, 1.2% to the track-finding efficiency, 2.5% to the particle identification, 1.2% to c_0 , and 0.1% to $\mathcal{B}_0$. We combine the results from the different D^0 and D_s decay modes to extract the final branching fractions listed in Table II.

To extract N_c from these numbers, we need to evaluate the contribution of $B^- \rightarrow \Lambda_c^+ \bar{\Lambda}_c^- K^-$. Combining the four-momenta of the recoiling B^- , of a K^- , and of the reconstructed Λ_c^+ or $\bar{\Lambda}_c^-$ candidate, we compute the missing mass: the absence of signal at the Λ_c mass excludes a significant contribution of this process. We therefore take $\mathcal{B}(B^- \rightarrow \Xi_c X) = \mathcal{B}(B^- \rightarrow \bar{\Lambda}_c X)$ in the computation of N_c . Using Eqs. (1) and (2) and taking $\mathcal{B}[B^- \rightarrow (c\bar{c})X] = (2.3 \pm 0.3)\%$ [19,20], one obtains:

$$\begin{aligned} N_c^- &= 0.983 \pm 0.030 \pm 0.046^{+0.028}_{-0.023}, \\ N_{\bar{c}}^- &= 0.330 \pm 0.022 \pm 0.020^{+0.051}_{-0.031}, \\ n_{\bar{c}}^- &= 1.313 \pm 0.037 \pm 0.062^{+0.063}_{-0.042}. \end{aligned}$$

TABLE II. Combined B^- branching fractions. The first uncertainty is statistical, the second is systematic, and the third reflects charm branching-fraction uncertainties [18].

	Correlated	Anticorrelated
X_c	$\mathcal{B}(B^- \rightarrow X_c X)(\%)$	$\mathcal{B}(B^- \rightarrow \bar{X}_c X)(\%)$
D^0	$79.3 \pm 2.5 \pm 4.0^{+2.0}_{-1.9}$	$9.8 \pm 0.9 \pm 0.5^{+0.3}_{-0.3}$
D^+	$9.8 \pm 1.2 \pm 1.2^{+0.8}_{-0.7}$	$3.8 \pm 0.9 \pm 0.4^{+0.3}_{-0.3}$
D_s^+	$0.5 \pm 0.6 \pm 0.2^{+0.2}_{-0.1}$	$14.3 \pm 1.6 \pm 1.5^{+4.9}_{-3.0}$
	<2.2 at 90% C.L.	
Λ_c^+	$3.5 \pm 0.8 \pm 0.3^{+1.3}_{-0.8}$	$2.9 \pm 0.8 \pm 0.3^{+1.1}_{-0.6}$

TABLE III. Combined $\bar{B}^0$ branching fractions. The first uncertainty is statistical, the second is systematic, and the third reflects charm branching-fraction uncertainties [18].

X_c	Correlated $\mathcal{B}(\bar{B}^0 \rightarrow X_c X)(\%)$	Anticorrelated $\mathcal{B}(\bar{B}^0 \rightarrow \bar{X}_c X)(\%)$
D^0	$51.1 \pm 3.1 \pm 2.5^{+1.3}_{-1.3}$	$6.3 \pm 1.9 \pm 0.5^{+0.2}_{-0.2}$
D^+	$39.7 \pm 3.0 \pm 2.8^{+2.8}_{-2.5}$	$2.3 \pm 1.8 \pm 0.3^{+0.2}_{-0.2}$
		<5.1 at 90% C.L.
D_s^+	$3.9 \pm 1.7 \pm 0.4^{+1.3}_{-0.8}$	$10.9 \pm 2.1 \pm 0.8^{+3.8}_{-2.3}$
	<8.7 at 90% C.L.	
Λ_c^+	$4.9 \pm 1.7 \pm 0.4^{+1.8}_{-1.0}$	$2.0 \pm 1.2 \pm 0.2^{+0.7}_{-0.4}$
		<3.8 at 90% C.L.

The reconstruction of D , D_s , and Λ_c from $\bar{B}^0$ decays is performed in the same way as that in the B^- analysis. The corresponding yields are listed in Table I. We then compute for each decay channel $X_c \rightarrow f$ the efficiency-corrected signal yields $N(X_c \rightarrow f)$ [$N(\bar{X}_c \rightarrow \bar{f})$] and define the raw branching fractions $\mathcal{B}_c$ and $\bar{\mathcal{B}}_c$ as

$$\mathcal{B}_c = N(X_c \rightarrow f) / [N_{B^0} \times \mathcal{B}(X_c \rightarrow f)], \quad (4)$$

$$\bar{\mathcal{B}}_c = N(\bar{X}_c \rightarrow \bar{f}) / [N_{B^0} \times \mathcal{B}(X_c \rightarrow f)]. \quad (5)$$

After correcting these numbers for $B^0\bar{B}^0$ mixing, we obtain the final branching fraction for $\bar{B}^0 \rightarrow X_c X$:

$$\mathcal{B}(\bar{B}^0 \rightarrow X_c X) = \frac{\mathcal{B}_c - \chi_d(\mathcal{B}_c + \bar{\mathcal{B}}_c) - c_+ \mathcal{B}_+}{1 - 2\chi_d}, \quad (6)$$

where $\chi_d = 0.181 \pm 0.004$ is the $B^0 - \bar{B}^0$ mixing parameter [18]. The correcting factor $\mathcal{B}_+$ accounts for B^+ contamination in the B^0 sample and depends on $\mathcal{B}(B^- \rightarrow X_c X)$ and $\mathcal{B}(B^+ \rightarrow X_c X)$. The results are given in Table I. Combining the different D^0 or D_s modes, we obtain the final branching fractions listed in Table III.

To compute N_c , we neglect $\bar{B}^0 \rightarrow \Lambda_c^+ \bar{\Lambda}_c^- K^0$ production and assume that $\mathcal{B}(\bar{B}^0 \rightarrow \Xi_c^- X) = \mathcal{B}(\bar{B}^0 \rightarrow \bar{\Lambda}_c^- X)$. Substituting $\bar{B}^0$ for B^- in Eqs. (1) and (2) and taking $\mathcal{B}[B^0 \rightarrow (c\bar{c})X] = (2.3 \pm 0.3)\%$ [19,20], we obtain:

$$\begin{aligned} N_c^0 &= 1.039 \pm 0.051 \pm 0.049^{+0.039}_{-0.031}, \\ N_c^- &= 0.237 \pm 0.036 \pm 0.012^{+0.039}_{-0.024}, \\ n_c^0 &= 1.276 \pm 0.062 \pm 0.058^{+0.066}_{-0.046}. \end{aligned}$$

TABLE IV. Fraction w of anticorrelated charm.

Mode	B^- decays	$\bar{B}^0$ decays
$\bar{D}^0 X$	$0.110 \pm 0.010 \pm 0.003$	$0.110 \pm 0.031 \pm 0.008$
$D^- X$	$0.278 \pm 0.052 \pm 0.009$	$0.055 \pm 0.040 \pm 0.006$
$D_s^- X$	$0.966 \pm 0.039 \pm 0.012$	$0.733 \pm 0.092 \pm 0.010$
$\bar{\Lambda}_c^- X$	$0.452 \pm 0.090 \pm 0.003$	$0.286 \pm 0.142 \pm 0.007$

We also compute the fraction of anticorrelated charm production in B decays, $w(\bar{X}_c) = \mathcal{B}(\bar{B} \rightarrow \bar{X}_c X) / [\mathcal{B}(\bar{B} \rightarrow X_c X) + \mathcal{B}(\bar{B} \rightarrow \bar{X}_c X)]$. Here, many systematic uncertainties cancel (tracking, K identification, D branching fractions, B counting). The results are given in Table IV. We obtain an upper limit on the correlated D_s^+ fraction in B^- decays: $\mathcal{B}(B^- \rightarrow D_s^+ X) / \mathcal{B}(B^- \rightarrow D_s^\pm X) < 0.126$ at 90% C.L.

In conclusion, we have measured for the first time the branching fractions for inclusive decays of B mesons to flavor-tagged D , D_s , and Λ_c , separately for B^- and $\bar{B}^0$. We observe significant production of anticorrelated D^0 and D^+ mesons in B decays (Table IV), with the branching fractions detailed in Tables II and III. The correlated D_s production in B^- decays is measured to be small.

As expected, the sum of all correlated charm branching fractions N_c is compatible with 1, for charged as well as for neutral B 's. The numbers of charmed particles per B^- decay ($n_c^- = 1.313 \pm 0.037 \pm 0.062^{+0.063}_{-0.042}$) and per $\bar{B}^0$ decay ($n_c^0 = 1.276 \pm 0.062 \pm 0.058^{+0.066}_{-0.046}$) are consistent with previous measurements [2,19,21] and with theoretical expectations [8–10].

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from CONACyT (Mexico), A.P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

- [1] Throughout this paper, the named reaction refers also to its complex conjugate.
- [2] CLEO Collaboration, T. E. Coan *et al.*, Phys. Rev. Lett. **80**, 1150 (1998).
- [3] DELPHI Collaboration, J. Abdallah *et al.*, Phys. Lett. B **561**, 26 (2003).

- [4] CLEO Collaboration, R. Ammar *et al.*, Phys. Rev. D **55**, 13 (1997).
- [5] ALEPH Collaboration, R. Barate *et al.*, Eur. Phys. J. C **4**, 387 (1998).
- [6] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. D **68**, 092001 (2003).

B. AUBERT *et al.*PHYSICAL REVIEW D **70** 091106

- [7] CLEO Collaboration, B. Barish *et al.*, Phys. Rev. Lett. **79**, 3599 (1997).
- [8] E. Bagan *et al.*, Phys. Lett. B **351**, 546 (1995).
- [9] G. Buchalla *et al.*, Phys. Lett. B **364**, 188 (1995).
- [10] M. Neubert, in *Proceedings of the 17th International Symposium on Lepton-Photon Interactions, Beijing, China* (World Scientific, Singapore, 1995), p. 298.
- [11] A. L. Kagan, in *Proceedings of the 7th International Symposium on Heavy Flavour Physics, Santa Barbara, CA* (World Scientific, Singapore, 1997).
- [12] BABAR Collaboration, B. Aubert *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [13] GEANT4 Collaboration, S. Agostinelli *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **506**, 250 (2003).
- [14] ARGUS Collaboration, H. Albrecht *et al.*, Z. Phys. C **48**, 543 (1990).
- [15] CRYSTAL BALL Collaboration, T. Skwarnicki *et al.*, Report No. DESY F31-86-02.
- [16] Only $D^0 \rightarrow K^- \pi^+$ and $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$ are used for the charm counting because $D^0 \rightarrow K^- \pi^+ \pi^0$ has a lower significance and $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ is not self-tagging (one cannot distinguish between D^0 and $\bar{D}^0$).
- [17] Satellite contributions include a reflection from $D^0 \rightarrow K^- K^+$ in the $D^0 \rightarrow K^- \pi^+$ mass spectrum and a signal at the D^+ mass (from $D^+ \rightarrow \phi \pi^+$ decays) in the $D_s^+ \rightarrow \phi \pi^+$ mass spectrum.
- [18] Particle Data Group, S. Eidelman *et al.*, Phys. Lett. B **592**, 1 (2004).
- [19] ALEPH Collaboration, CDF Collaboration, DELPHI Collaboration, L3 Collaboration, OPAL Collaboration, and SLD Collaboration, hep-ex/0112028.
- [20] M. Beneke *et al.*, Phys. Rev. D **59**, 054003 (1999).
- [21] CLEO Collaboration, L. Gibbons *et al.*, Phys. Rev. D **56**, 3783 (1997).