

## Energy diffusion in a chaotic adiabatic billiard gas

C. Jarzynski

*Department of Physics and Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720*

(Received 9 March 1993)

A diffusion equation is derived for the energy distribution of a gas of noninteracting point particles following chaotic trajectories inside a slowly-time-dependent container. We discuss the relevance of this problem to results concerning ergodic adiabatic Hamiltonian systems, as well as to one-body dissipation in nuclear dynamics.

PACS number(s): 05.45.+b, 24.10.-i

### INTRODUCTION

This paper considers the problem of a *chaotic adiabatic billiard gas*, a gas of noninteracting point particles bouncing around chaotically inside a container whose shape changes slowly with time. (See Fig. 1.) Unlike an ordinary gas, where particle-particle collisions dominate, producing a Maxwell-Boltzmann distribution of energies, here the evolution of a particle's energy is determined solely by its collisions with the slowly moving walls of the container. Let  $\eta(E, t) dE$  denote the number of particles with energy in a small interval  $dE$  around  $E$ , at time  $t$ . The main result of this paper is a diffusion equation governing the time evolution of  $\eta$ , the distribution of particle energies. We obtain such an equation for both the two- and three-dimensional versions of this problem.

Section I of this paper specifies the problem precisely, and introduces notation. In Sec. II we argue that the distribution of particle energies of a chaotic adiabatic billiard gas evolves diffusively; this suggests a Fokker-Planck equation for the evolution of  $\eta(E, t)$ . In Sec. III we derive explicit expressions for the drift and diffusion coefficients which determine this equation. These are given in terms of the dynamics of particles bouncing around inside *time-independent* containers, obtained by "freezing" the slowly-changing shape of the container at different instants in time. We show in Sec. IV that, under a certain approximation, our results may be further simplified so that the evolution of  $\eta$  is given entirely in terms of the changing shape of the container, without any reference to particle dynamics.

Our interest in this problem is twofold. First, as discussed in Sec. V, our gas can be treated as an *ergodic adiabatic ensemble*, an ensemble of noninteracting systems evolving chaotically under a common, slowly time-varying Hamiltonian. Ott, Brown, and Grebogi [1-3] have used multiple time-scale analysis to study such systems. Their focus has been the goodness of the *ergodic adiabatic invariant*, i.e., the extent to which a certain quantity, shown by Ott [1] to be conserved in the limit of an infinitely slowly evolving Hamiltonian, remains conserved when the Hamiltonian evolves at a slow but finite rate. Recently [4], we have used an alternative approach to this problem to derive an evolution equation for the distribution of energies of an ergodic adiabatic ensemble.

(See also Ref. [5], in which an apparent discrepancy between the multiple time-scale approach of Refs. [1-3] and the diffusion-equation approach of Ref. [4] is resolved.) Such an equation was previously derived by Wilkinson [6]; regrettably, we were unaware of his work, and failed to give due credit. The present paper represents an application of the general approach of Ref. [4] to a specific class of ergodic adiabatic ensembles.

Our other motivation for studying this problem comes from the independent-particle model of nuclear dynamics, in which a nucleus undergoing some dynamical process (e.g., fission or collision with another nucleus) is imagined as a time-dependent container filled with a gas of independent point particles. This simple model provides a mechanism, one-body dissipation for friction in dynamical nuclear processes. A principal result of this approach to nuclear dynamics has been the wall formula [7,8], an expression for the rate at which one-body dissipation transfers energy from the collective degrees of freedom of the idealized nucleus to the individual nucleons. The results of the present paper, as discussed in Sec. VI, extend our understanding of one-body dissipation.

### I. PRELIMINARIES

We take the time-dependent shape of the container to be an externally imposed, rather than a dynamical, quantity: the shape evolves in a predetermined way, independently of the gas of particles. Each bounce of a particle off the moving walls of the container is taken to be specular (the angle of reflection is equal to the angle of incidence) in the instantaneous rest frame of the local piece of wall at which the collision occurs. Effectively, these bounces constitute elastic collisions in which the inertia of the wall is infinitely greater than that of the particle.

We are interested in observing our gas of noninteracting particles as the shape of the container changes slowly. To express "slow" shape evolution mathematically, we make the shape a function of  $\epsilon t$ , where  $t$  is time and  $\epsilon$  is a slowness parameter, formally taken to be small. Thus, let  $\mathcal{S}(\epsilon t)$  denote the shape of the container at time  $t$ . We will be interested in observing our gas for times of order  $\epsilon^{-1}$ , over which the container changes by order

unity. As the extreme limit of slow evolution, we will take the *adiabatic limit* to mean that in which we let  $\epsilon$  go to zero, holding  $\epsilon t_i$  and  $\epsilon t_f$  fixed,  $t_i$  and  $t_f$  being the initial and final times over which we observe the system. In this limit, the container evolves infinitely slowly from the initial shape  $\mathfrak{S}(\epsilon t_i)$  to the final shape  $\mathfrak{S}(\epsilon t_f)$ .

We will frequently refer to the motion of particles inside a *frozen* container, by which we mean the time-independent container obtained by arresting (“freezing”) the slowly evolving shape  $\mathfrak{S}(\epsilon t)$  at some instant in time. Whenever discussing the dynamics of particles inside a frozen container, as opposed to the slowly changing one, we will emphasize the distinction by using  $\mathfrak{S}_\alpha$ , with  $\alpha = \epsilon t$ , to denote the shape of the container frozen at  $\epsilon t$ . When discussing motion inside the time-dependent container, we will retain the notation  $\mathfrak{S}(\epsilon t)$ . The slow evolution of the container from  $\mathfrak{S}(\epsilon t_i)$  to  $\mathfrak{S}(\epsilon t_f)$  defines a continuous sequence of frozen shapes  $\mathfrak{S}_\alpha$ , with  $\alpha$  ranging from  $\epsilon t_i$  to  $\epsilon t_f$ .

The motion of a particle bouncing around inside a frozen container is represented in phase space by a trajectory  $(\mathbf{q}(t), \mathbf{p}(t))$  whose evolution is restricted to an *energy shell*, a surface of constant energy. We make the crucial assumption that, for any of the frozen shapes  $\mathfrak{S}_\alpha$ , an arbitrary nonperiodic trajectory will chaotically and ergodically explore the entire energy shell on which it is found [9]. A consequence of this assumption is that the motion of particles in any of the frozen containers exhibits *chaotic mixing* over the energy shell: any distribution of initial particle positions and velocities will evolve into a uniform distribution of particles throughout the container, with an isotropic distribution of velocities. The time scale over which this mixing occurs is given by the Lyapunov time  $t_L = 1/\lambda$ , where  $\lambda$  is the Lyapunov exponent associated with the chaotic evolution of the trajectories.

We now discuss the relevance of chaotic mixing to a gas of particles in a slowly time-dependent container. First, consider the motion of two particles sharing identical initial conditions at time  $t_0$ , one subsequently evolving inside the time-dependent container  $\mathfrak{S}(\epsilon t)$ , the other inside the frozen container  $\mathfrak{S}_\alpha$ , with  $\alpha = \epsilon t_0$ . Let  $T$  be the length of time over which the paths followed by these two particles remain very close; after this time, they will diverge rapidly.  $T$  can be made arbitrarily large by choosing  $\epsilon$  arbitrarily small, although, due to the assumed chaoticity, a value of  $T$  much larger than the Lyapunov time  $t_L$  would require an extremely small  $\epsilon$ . [By treating motion inside the evolving container as a perturbed version of motion inside the frozen one, with the perturbations, proportional to  $\epsilon$ , introduced at collisions with the wall,  $T$  can be shown to scale like  $t_L \ln(1/\epsilon)$ , for small  $\epsilon$ .] We will henceforth assume  $\epsilon$  to be small enough that

$$T \gtrsim t_L. \quad (1.1)$$

Thus, motion inside the time-dependent container closely mimics that inside the frozen one over times on the order of the Lyapunov time. In this case, chaotic mixing occurs before the particles are affected by the time dependence of the shape; as the container slowly evolves, the continual process of chaotic mixing tends to main-

tain a uniform distribution of particles throughout the container, and an isotropic distribution of velocities.

One more assumption needs to be made in order for the central result of this paper to be valid. Since this assumption involves a correlation sum to be defined below, we postpone its explicit statement to Sec. III, where it is italicized.

We will use the term *chaotic adiabatic billiard gas* to describe a gas of noninteracting particles inside a container whose slowly evolving shape satisfies the assumptions discussed above. Our goal is an evolution equation for the distribution of particle energies,  $\eta(E, t)$ .

## II. DIFFUSION OF ENERGIES

The energy of a given particle changes in small, discrete amounts as the particle collides with the slowly moving walls of the container. We can think of this process in terms of the particle performing a “walk” along the energy axis, with steps determined by the underlying motion of the particle bouncing off the container’s walls. Since this underlying motion is chaotic, correlations between these steps along the energy axis will exist only over a finite time, on the order of the Lyapunov time  $t_L$ . This consideration suggests [4,10] that the distribution of energies of a gas of such particles will, on a time scale much longer than  $t_L$ , evolve by a process of diffusion. We therefore postulate the following Fokker-Planck equation for the time-dependent distribution of energies,  $\eta(E, t)$ :

$$\frac{\partial \eta}{\partial t} = -\frac{\partial}{\partial E}(g_1 \eta) + \frac{1}{2} \frac{\partial^2}{\partial E^2}(g_2 \eta). \quad (2.1)$$

This is a generalized diffusion equation, in which the drift and diffusion coefficients  $g_1$  and  $g_2$  are functions of both energy and time. Specifically, we write  $g_1(E, \epsilon t)$  and  $g_2(E, \epsilon t)$ ; we use explicit functions of  $\epsilon t$  rather than simply  $t$  because we expect the evolution of  $\eta$  at a given time  $t$  to be determined by the instantaneous shape of the container and the way in which it is changing, which depend explicitly on  $\epsilon t$ .

Since we are interested in slow evolution of the shape of the container, we can expand  $g_1$  and  $g_2$  in powers of  $\epsilon$  (making the assumption that integral powers suffice). As discussed below, retaining only  $O(\epsilon^1)$  terms gives an evolution equation for  $\eta$  that corresponds to the adiabatic limit (infinitely slow shape evolution). We are interested in slow but finite evolution of the shape of the container, and therefore want expressions for  $g_1$  and  $g_2$  valid to  $O(\epsilon^2)$ .

In treating the evolution of  $\eta$  as a process of diffusion, we must keep in mind that this picture is valid only over times much longer than the Lyapunov time  $t_L$ . Thus, for Eq. (2.1) to be applicable to our problem, there must exist a time scale which is long compared to  $t_L$ , but short compared to that over which significant changes in the distribution of energies (as well as the shape of the container) occur. We will use the notation  $\Delta t$  to indicate a time on this scale, and will refer to this time as “short” or “long” depending on the context, i.e., whether we are discussing the evolution of  $\eta(E, t)$ , or the motion of particles in the container.

### III. DERIVATION OF DRIFT AND DIFFUSION COEFFICIENTS

To derive expressions for  $g_1$  and  $g_2$ , note that a distribution of energies described initially by a  $\delta$  function along the energy axis,

$$\eta(E, t_0) = \delta(E - E_0), \quad (3.1)$$

will evolve under Eq. (2.1) so that, a short time  $\Delta t$  later, the average energy will have drifted away from  $E_0$  by an amount  $g_1 \Delta t$ , and the distribution will have acquired a variance  $g_2 \Delta t$ , with  $g_1$  and  $g_2$  evaluated at  $E_0, \epsilon t_0$ . The second moment of this new distribution of energies with respect to  $E_0$  is then

$$\begin{aligned} \mathcal{M}_2(\Delta t) &\equiv \int dE \eta(E, t_0 + \Delta t) (E - E_0)^2 \\ &= (g_1 \Delta t)^2 + g_2 \Delta t. \end{aligned} \quad (3.2)$$

By considering a gas of particles sharing a common energy  $E_0$  at time  $t_0$  [such a gas, with the number of particles normalized to unity, is described by Eq. (3.1)], then by solving, in terms of quantities characterizing the subsequent motion of these particles, for  $\mathcal{M}_2(\Delta t)$ , and finally by comparing the result with Eq. (3.2), we will obtain expressions for  $g_1$  and  $g_2$ . We will solve only for the leading term of  $\mathcal{M}_2(\Delta t)$ , which is  $O(\epsilon^2)$ . From this will immediately follow the leading terms of  $g_1$  and  $g_2$ , which are  $O(\epsilon^1)$  and  $O(\epsilon^2)$ , respectively. To obtain the  $O(\epsilon^2)$  term of  $g_1$ , we will invoke a trick using Liouville's theorem.

We therefore begin by considering, at time  $t_0$ , a gas of particles of energy  $E_0$  distributed uniformly with the container, with an isotropic distribution of velocities. Let us introduce the *wall velocity field*,  $\dot{n}$ , a scalar field defined over the surface of the container: the value of  $\dot{n}$  at a particular point on the surface gives the normal outward velocity of the moving wall at that point (see Fig. 1; a negative  $\dot{n}$  indicates a portion of the wall which is moving into the gas). This field contains all information about how the shape of the container is changing at a given




FIG. 1. Two-dimensional version of a chaotic adiabatic billiard gas. The scalar field  $\dot{n}$  gives the rate at which the wall is moving normally outward, as a function of position  $s$  along the wall. It is assumed that “freezing” the shape at any instant will produce a time-independent billiard in which all particle trajectories are chaotic.

instant in time. Since this field changes with time along with the shape of the container, we will write it as  $\dot{n}(\epsilon t)$  (suppressing the dependence on the position on the surface of the wall). We also introduce a *frozen* field  $\dot{n}_\alpha$  — defined over the surface of the frozen shape  $\mathfrak{S}_\alpha$  — which is simply the normal outward wall velocity at the moment of freezing;  $\dot{n}_\alpha$  gives information on how the shape  $\mathfrak{S}(\epsilon t)$  was changing at the instant in which it was frozen into  $\mathfrak{S}_\alpha$ .

To lowest order in the wall velocity (proportional to  $\epsilon$ ), the change in the energy of a particle as it bounces off the wall is  $-2mv\dot{n} \sin \theta$ , where  $m$  is the particle mass,  $v$  is its speed prior to collision,  $\dot{n}$  is the value of the wall velocity field at the point of collision, and  $\theta$  is the angle between the incoming trajectory of the particle and a surface tangent to the wall. (See Figs. 2 and 3.) Between times  $t_0$  and  $t_0 + \Delta t$  the particle bounces many times off the walls of the container, whose shape changes negligibly during that time. The number of collisions,  $B$ , is approximated as

$$B \cong \Delta t / \tau, \quad (3.3)$$

where  $\tau$  is the average time between bounces for a particle inside the container frozen at  $\alpha = \epsilon t_0$ . The total change in the energy of the particle over this time is, to leading order,

$$E - E_0 = -2mv \sum_{b=1}^B \dot{n}_b \sin \theta_b, \quad (3.4)$$

where the  $\dot{n}_b$ 's are the normal outward wall velocities sampled by the sequence of bounces  $b = 1, 2, \dots, B$ , and the  $\theta_b$ 's are the corresponding angles of collision. We are justified in pulling  $v = (2E_0/m)^{1/2}$  outside this sum by the fact that, to lowest order in  $\epsilon$ , the speed of the particle remains constant over time  $\Delta t$ . To obtain  $\mathcal{M}_2(\Delta t)$ , we square the above sum, then average over all particles, i.e., over an ensemble of trajectories evolving from a uniform distribution of initial conditions on the energy shell  $E_0$  at time  $t_0$ . Angular brackets will denote this average:

$$\mathcal{M}_2(\Delta t) = 4m^2 v^2 \sum_{b=1}^B \sum_{b'=1}^B \langle \dot{n}_b \sin \theta_b \dot{n}_{b'} \sin \theta_{b'} \rangle. \quad (3.5)$$

Now, suppose temporarily that, for any initial condition corresponding to energy  $E_0$  at time  $t_0$ , two trajec-




FIG. 2. Particle bouncing off a small segment of wall in a two-dimensional billiard. The value of  $\theta$  ranges from 0 to  $\pi$ .




FIG. 3. Particle bouncing off a small patch of wall in a three-dimensional billiard. The dashed line represents the normal projection of the trajectory onto the wall. The value of  $\theta$  ranges from 0 to  $\pi/2$ ;  $\phi$  ranges from 0 to  $2\pi$ . The line representing  $\phi = 0$  is arbitrary.

tories evolving from that initial condition, one inside the slowly changing container, the other inside the container frozen at  $\alpha = \epsilon t_0$ , remain very close to one another for the entire length of time from  $t_0$  to  $t_0 + \Delta t$ . (Since  $\Delta t \gg t_L$ , this puts a drastic limit, which we later relax, on the magnitude of  $\epsilon$ .) If this condition holds, then, for purposes of evaluating the right-hand side of Eq. (3.5), we may replace the gas of particles evolving for time  $\Delta t$  inside the time-dependent container, with a gas evolving inside the frozen one. With this replacement, Eq. (3.5) becomes

$$\mathcal{M}_2(\Delta t) = 4m^2v^2 \sum_{b=1}^B \sum_{b'=1}^B \left\langle \dot{n}_{\alpha b} \sin \theta_b \dot{n}_{\alpha b'} \sin \theta_{b'} \right\rangle_{\alpha}, \quad (3.6)$$

where the angular brackets  $\langle \rangle_{\alpha}$  indicate an average over an ensemble of trajectories evolving inside the *frozen*  $\mathfrak{S}_{\alpha}$ , with  $\alpha = \epsilon t_0$  (as before, the ensemble is defined by a uniform distribution over the energy shell  $E_0$ ), and  $\dot{n}_{\alpha b}$  gives the value of the frozen field  $\dot{n}_{\alpha}$  at the  $b$ th bounce of one such trajectory.

We rewrite the quantity being summed in Eq. (3.6) as

$$\begin{aligned} & \left\langle (\dot{n}_{\alpha b} \sin \theta_b - \xi_b) (\dot{n}_{\alpha b'} \sin \theta_{b'} - \xi_{b'}) \right\rangle_{\alpha} + \xi_b \xi_{b'} \\ & \equiv c_{b,b'} + \xi_b \xi_{b'}, \end{aligned} \quad (3.7)$$

where  $\xi_b \equiv \langle \dot{n}_{\alpha b} \sin \theta_b \rangle_{\alpha}$  is the value of  $\dot{n}_{\alpha} \sin \theta$  at the  $b$ th bounce, averaged over the ensemble of trajectories in the frozen container. Since the distribution of this ensemble is invariant with time (by virtue of uniform distribution over the energy shell  $E_0$ ),  $\xi_b$  is in fact independent of  $b$ . We will therefore write it simply as  $\xi$ . Similarly,  $c_{b,b'}$ , which measures correlations in  $\dot{n}_{\alpha} \sin \theta$  between the  $b$ th and  $b'$ th bounces, depends on  $b$  and  $b'$  only through the difference  $\Delta b \equiv b' - b$ , and so will be written as  $c_{\Delta b}$ . The double sum in Eq. (3.6) then becomes

$$\sum_{b=1}^B \sum_{b'=1}^B (c_{\Delta b} + \xi^2) = B^2 \xi^2 + B \sum_{\Delta b=-B}^B \left(1 - \frac{|\Delta b|}{B}\right) c_{\Delta b}. \quad (3.8)$$

By chaotic mixing,  $c_{\Delta b} \cong 0$  for  $|\Delta b| > \nu_L$ , where  $\nu_L \cong t_L/\tau$ . Having assumed  $\Delta t \gg t_L$ , we have  $B \gg \nu_L$ , and may therefore approximate the sum appearing on the right hand side of the above expression as  $\sum_{\Delta b=-\infty}^{+\infty} c_{\Delta b}$ ; this is the discrete version of a standard result from the theory of stationary stochastic processes [see Ref. [11] and Eqs. (4.11) and (4.12) of Ref. [4]]. We assume that this sum converges (see Sec. VII for examples in which this assumption fails). We now have, to  $O(\epsilon^2)$ ,

$$\begin{aligned} \mathcal{M}_2(\Delta t) &= 4m^2v^2 \left( B^2 \xi^2 + B \sum_{\Delta b=-\infty}^{+\infty} c_{\Delta b} \right) \\ &= \left( \frac{2mv}{\tau} \xi \Delta t \right)^2 + \frac{4m^2v^2}{\tau} \Delta t \sum_{\Delta b=-\infty}^{+\infty} c_{\Delta b}. \end{aligned} \quad (3.9)$$

Comparison with Eq. (3.2) yields, to  $O(\epsilon^2)$ ,

$$g_1 = \pm \frac{2mv}{\tau} \xi + O(\epsilon^2), \quad (3.10)$$

$$g_2 = \frac{4m^2v^2}{\tau} D, \quad (3.11)$$

where

$$D \equiv \sum_{\Delta b=-\infty}^{+\infty} c_{\Delta b}. \quad (3.12)$$

We now relax the assumption made immediately after Eq. (3.5), and assert that as long as motion inside the frozen container closely mimics that inside the time-dependent one over times on the order of  $t_L$ , rather than the much longer  $\Delta t$ , the steps leading to Eqs. (3.10) and (3.11) will remain valid. (We have already assumed, in Sec. I, that this more relaxed condition holds.) The justification for this assertion is similar to that presented in Ref. [4] [see the paragraph following Eq. (4.14) therein]; its essence is as follows. Due to chaotic mixing, appreciable correlations exist only between bounces separated by times on the order of, or shorter than, the Lyapunov time  $t_L$ . However, if motion inside the frozen container closely resembles that inside the time-dependent one over times up to  $t_L$ , then the correlations that do exist are nearly identical for the two cases. Since it is these correlations that determine the diffusion of particle energies, we are justified in evaluating the right hand side of Eq. (3.5) using particles inside the frozen rather than the time-dependent container.

We henceforth drop the subscript 0 from  $E_0$  and  $t_0$ .

In Appendixes A and B, we evaluate  $\xi$  and  $\tau$  for both two- and three-dimensional containers. The results reduce Eq. (3.10) to

$$g_1(E, \epsilon t) = -\beta(\epsilon t) E + O(\epsilon^2), \quad (3.13)$$

where the factor  $\beta(\epsilon t)$  depends on the dimensionality:

$$\beta = \begin{cases} \dot{A}/A & (2D \text{ container}), \\ 2\dot{V}/3V & (3D \text{ container}), \end{cases} \quad (3.14a)$$

$$(3.14b)$$

where  $A$  and  $V$  denote the area or volume enclosed by the container, and the dot signifies differentiation with respect to time. ( $A$ ,  $\dot{A}$ ,  $V$ , and  $\dot{V}$  are evaluated at  $ct$ .) The ambiguity in sign appearing in Eq. (3.10) has been removed by physical considerations: since there is a net positive amount of work done by a gas inside a container whose area (in the 2D case) or volume (3D) is increasing, the energy drift  $g_1$  associated with  $\dot{A} > 0$  or  $\dot{V} > 0$  must be negative.

In Eq. (3.11), the quantity  $D$  is determined by the frozen shape  $\mathfrak{S}_\alpha$  and the associated frozen wall velocity field  $\dot{n}_\alpha$ , and hence may be written as a function of the value of  $\alpha$ , in this case  $ct$ ; thus,  $D = D(ct)$ . All dependence of  $g_2$  on  $E$  is in the factor  $4m^2v^2/\tau$ . Using the results for  $\tau$  from Appendix B, we get, to  $O(\epsilon^2)$ ,

$$g_2(E, ct) = \gamma(ct) E^{3/2}, \quad (3.15)$$

with

$$\gamma = \begin{cases} (8l/\pi A) (2m)^{1/2} D & \text{(2D)}, \\ (2S/V) (2m)^{1/2} D & \text{(3D)}, \end{cases} \quad (3.16a)$$

$$(3.16b)$$

where  $l(ct)$  is the perimeter of the 2D container, and  $S(ct)$  is the surface area of the 3D one.

It remains to obtain the  $O(\epsilon^2)$  term of  $g_1$ . The strategy for doing so invokes Liouville's theorem, and is detailed in Sec. IV of Ref. [4]. There we find

$$g_1 = g_{11} + \frac{1}{2\Sigma} \frac{\partial}{\partial E} (g_2 \Sigma), \quad (3.17)$$

where  $g_{11}$  is the  $O(\epsilon^1)$  term of  $g_1$  [given above by Eq. (3.13)], and

$$\Sigma(E, ct) \equiv \frac{\partial}{\partial E} \Omega(E, ct), \quad (3.18)$$

where  $\Omega(E, ct)$  represents the volume of phase space enclosed by the energy shell  $E$  at time  $t$ . For a two-dimensional billiard system, this volume is the product of the area of ordinary space enclosed by the container, with the area in momentum space of a circle of radius  $p = (2mE)^{1/2}$ . Thus,

$$\Omega = 2\pi mA E, \quad \Sigma = 2\pi mA. \quad (3.19)$$

In three dimensions, we get

$$\Omega = \frac{4}{3} \pi (2m)^{3/2} V E^{3/2}, \quad \Sigma = 2\pi (2m)^{3/2} V E^{1/2}. \quad (3.20)$$

Using Eq. (3.17) we rewrite Eq. (2.1) as

$$\frac{\partial \eta}{\partial t} = -\frac{\partial}{\partial E} (g_{11} \eta) + \frac{1}{2} \frac{\partial}{\partial E} \left[ g_2 \Sigma \frac{\partial}{\partial E} \left( \frac{\eta}{\Sigma} \right) \right]. \quad (3.21)$$

Combining our results for  $g_{11}$ ,  $g_2$ , and  $\Sigma$  with Eq. (3.21), we finally write the evolution equation for  $\eta$ , to  $O(\epsilon^2)$ , as

$$\frac{\partial \eta}{\partial t} = \beta \frac{\partial}{\partial E} (E \eta) + \frac{\gamma}{2} \frac{\partial}{\partial E} \left( E^{3/2} \frac{\partial \eta}{\partial E} \right) \quad (2D) \quad (3.22a)$$

or

$$\frac{\partial \eta}{\partial t} = \beta \frac{\partial}{\partial E} (E \eta) + \frac{\gamma}{2} \frac{\partial}{\partial E} \left[ E^2 \frac{\partial}{\partial E} \left( E^{-1/2} \eta \right) \right] \quad (3D). \quad (3.22b)$$

Equation (3.22) represents the central result of this paper. The coefficients  $\beta$  and  $\gamma$  are given [Eqs. (3.14) and (3.16)] in terms of the particle mass  $m$ , quantities associated with the changing shape of the container ( $A$ ,  $\dot{A}$ , and  $l$ ; or  $V$ ,  $\dot{V}$ , and  $S$ ), and the function  $D = \sum_{-\infty}^{+\infty} c_{\Delta b}$ . Only the last of these directly involves the dynamics of particles bouncing around inside a container, and is given in terms of motion inside the *frozen* container  $\mathfrak{S}_\alpha$ ,  $\alpha = ct$ . Thus, the time-dependent problem [a gas of particles inside the slowly changing container  $\mathfrak{S}(ct)$ ] is solved in terms of the solutions of a continuous sequence of time-independent problems (motion inside the frozen shapes  $\mathfrak{S}_\alpha$ ). In the following section we show how, in a certain approximation, the quantity  $D$  may be divested of any reference whatsoever to the dynamics of bouncing particles. In this case the evolution of  $\eta$  is given directly in terms of the changing shape of the container. First, however, we discuss the adiabatic limit.

The adiabatic limit, as defined in Sec. I, involves a time  $t_f - t_i$  which approaches infinity like  $\epsilon^{-1}$ . Over such a time, the term involving  $\beta$  ( $\sim \epsilon$ ) in Eq. (3.22) will make an  $O(\epsilon^0)$ , i.e., finite, contribution to the change in  $\eta$ , while the term involving  $\gamma$  ( $\sim \epsilon^2$ ) will make an  $O(\epsilon^1)$ , i.e., vanishing, contribution. Thus, in the adiabatic limit,

$$\frac{\partial \eta}{\partial t} = \beta \frac{\partial}{\partial E} (E \eta), \quad (3.23)$$

for both the 2D and the 3D case. This equation describes a distribution of particles moving along the energy axis under a "velocity" field  $-\beta E$ . The energy  $\mathcal{E}(t)$  of any one of these particles satisfies

$$\frac{d}{dt} \mathcal{E}(t) = -\beta(ct) \mathcal{E}(t) = \begin{cases} -(\dot{A}/A) \mathcal{E}(t) & \text{(2D)}, \\ -(2\dot{V}/3V) \mathcal{E}(t) & \text{(3D)}. \end{cases} \quad (3.24a)$$

$$(3.24b)$$

From this, we get  $(d/dt) \Omega(\mathcal{E}(t), ct) = 0$  [see Eqs. (3.19) and (3.20)]. Equation (3.22) is therefore consistent with the adiabatic invariance of  $\Omega$ , which was demonstrated by Ott [1] to hold generally for ergodic adiabatic Hamiltonian systems.

#### IV. THE QUASILINEAR APPROXIMATION

It may sometimes be the case that the sum  $D = \sum_{-\infty}^{+\infty} c_{\Delta b}$  which appears in  $\gamma$  is dominated by the term  $c_0$ :

$$D \cong c_0. \quad (4.1)$$

We denote this the *quasilinear approximation*, following standard usage [12]. The validity of this approximation, which implies that correlations between the different bounces of a trajectory play a negligible role in the evolution of  $\eta$ , will depend on the details of the shape

$\mathfrak{S}_\alpha$  and the frozen wall velocity field  $\dot{n}_\alpha$ , and may be difficult to assess *a priori*. Roughly speaking, it demands that the container's shape and its motion be sufficiently irregular. We do not pursue here the question of how to define "sufficiently irregular." Rather, for those systems for which Eq. (4.1) happens to be valid, we derive an evolution equation for  $\eta$  wholly in terms of the evolution of the shape  $\mathfrak{S}(\epsilon t)$ , without explicit mention of particle dynamics.

Take Eq. (4.1) to be valid. In Appendix A we solve for  $c_0$ , obtaining

$$c_0 = \frac{2}{3l} \oint ds \left[ \dot{n}^2 - \frac{3\pi^2}{32} \bar{n}^2 \right] \equiv \frac{2}{3l} I_2(\epsilon t) \quad (2D),$$

$$c_0 = \frac{1}{2S} \oint d\sigma \left[ \dot{n}^2 - \frac{8}{9} \bar{n}^2 \right] \equiv \frac{1}{2S} I_3(\epsilon t) \quad (3D). \quad (4.2)$$

Here,  $\oint ds$  and  $\oint d\sigma$  indicate integrals over the entire wall of the container, and  $\bar{n}$  is the average value of  $\dot{n} = \dot{n}(\epsilon t)$  over the wall. Combining these results with Eqs. (3.14), (3.16), and (3.22), we have the simplified results

$$\frac{\partial \eta}{\partial t} = \frac{\dot{A}}{A} \frac{\partial}{\partial E} (E\eta) + \frac{8\sqrt{2m}}{3\pi A} I_2 \frac{\partial}{\partial E} \left( E^{3/2} \frac{\partial \eta}{\partial E} \right) \quad (2D)$$

$$(4.3a)$$

and

$$\frac{\partial \eta}{\partial t} = \frac{2\dot{V}}{3V} \frac{\partial}{\partial E} (E\eta) + \frac{\sqrt{2m}}{2V} I_3 \frac{\partial}{\partial E} \left[ E^2 \frac{\partial}{\partial E} \left( E^{-1/2} \eta \right) \right] \quad (3D). \quad (4.3b)$$

## V. RELATION TO PREVIOUS RESULTS

In Ref. [4] we considered the general problem of systems evolving in phase space under an *ergodic adiabatic Hamiltonian*, a slowly time-varying Hamiltonian which, if frozen at any instant, gives rise to trajectories that ergodically and chaotically explore their energy shells. This problem has been studied by Ott and co-workers [1–3], using multiple-time-scale analysis; by Koonin and Randrup [13], using linear response theory; and by Wilkinson [6], expanding on Ott's results. Let  $\mathbf{z} = (\mathbf{q}, \mathbf{p})$  represent a point in phase space, let  $H(\mathbf{z}, \epsilon t)$  denote the slowly-changing Hamiltonian, and let  $H_\alpha(\mathbf{z})$ , with  $\alpha = \epsilon t$ , denote the time-independent Hamiltonian obtained by freezing  $H(\mathbf{z}, \epsilon t)$  at time  $t$ . A central result of Ref. [4] is an evolution equation (previously derived by Wilkinson) for  $\eta(E, t)$ , the distribution of energies of an ensemble of such systems, all governed by the same ergodic adiabatic Hamiltonian, differing from one another only by their initial conditions in phase space; we call such an ensemble an *ergodic adiabatic ensemble*. This equation is identical to Eq. (2.1) of the present paper, only with

$$g_1(E, \epsilon t) = \bar{u} + \frac{1}{2\Sigma} \frac{\partial}{\partial E} (\Sigma g_2), \quad (5.1)$$

$$g_2(E, \epsilon t) = \int_{-\infty}^{+\infty} ds C(s), \quad (5.2)$$

where  $\Sigma(E, \epsilon t)$  is defined as per Eq. (3.18), and the quantities  $\bar{u}$  and  $C(s)$ , both functions of  $E$  and  $\epsilon t$ , are discussed below. (To avoid notational conflict with Sec. VI of the present paper, we use  $\bar{u}$  to denote the quantity which in Ref. [4] is denoted by  $\bar{v}$ .)

In this section we show that one can consistently treat a chaotic adiabatic billiard gas as an example of an ergodic adiabatic ensemble, by treating the container as the limiting case of a smooth potential well. Thus, Eqs. (5.1) and (5.2) will be shown to reduce, in this limit, to the corresponding expressions derived in Sec. III for the billiard gas.

The quantities  $\bar{u}$  and  $C(s)$  are defined as follows. Let  $\dot{H}(\mathbf{z}, \epsilon t)$  be the slowly time-dependent function obtained by differentiating  $H(\mathbf{z}, \epsilon t)$  with respect to  $t$ ; define  $\dot{H}_\alpha(\mathbf{z})$ , with  $\alpha = \epsilon t$ , as the time-independent function obtained by "freezing"  $\dot{H}(\mathbf{z}, \epsilon t)$  at time  $t$ . Then

$$\bar{u} \equiv \left\{ \dot{H}_\alpha(\mathbf{z}) \right\}, \quad (5.3)$$

$$C(s) \equiv \left\{ \left[ \dot{H}_\alpha(\mathbf{z}) - \bar{u} \right] O_\alpha(s) \left[ \dot{H}_\alpha(\mathbf{z}) - \bar{u} \right] \right\}, \quad (5.4)$$

where the curly brackets indicate an average over all points  $\mathbf{z}$  on the energy shell  $E$  of  $H_\alpha$ , and  $O_\alpha(s)$  is a time evolution operator which acts to the right, evolving a point  $\mathbf{z}$  for a time  $s$  under the frozen Hamiltonian  $H_\alpha$ .

For a particle moving inside a hard-walled container, it is intuitive to think of the container as a potential well  $V(\mathbf{q})$  whose value is zero for  $\mathbf{q}$  inside the container and infinite outside. This formulation, however, does not immediately lend itself to the calculation of  $\bar{u}$  and  $C(s)$  as defined above. We therefore soften the walls of the container by letting the potential rise smoothly from 0 inside to infinity outside, over a wall skin of thickness  $\delta$ ; we let  $\delta$  be arbitrarily small.

The contours of  $V(\mathbf{q})$  in the vicinity of some point on the surface of the wall will have the appearance shown in Fig. 4. If the wall at this point is moving with normal outward velocity  $\dot{n}$ , then, at a point  $\mathbf{q}$  within the wall




FIG. 4. The trajectory of a particle bouncing off a wall of finite skin depth. The parallel lines represent the contours of the potential  $V(\mathbf{q})$  in the vicinity of the bounce.

skin, we have

$$\dot{H} = -\dot{n} \hat{\mathbf{n}} \cdot \nabla V(\mathbf{q}) = -\dot{n} |\nabla V(\mathbf{q})|, \quad (5.5)$$

where  $\hat{\mathbf{n}}$  is the unit vector pointing normally outward from the wall. The frozen value of  $\dot{H}$  is then

$$\dot{H}_\alpha = -\dot{n}_\alpha \hat{\mathbf{n}} \cdot \nabla V(\mathbf{q}). \quad (5.6)$$

By the assumed ergodicity of motion inside the hard-walled container (and by extension in the soft-walled container, for arbitrarily small  $\delta$ ), the phase-space average of  $\dot{H}_\alpha$  over a particular energy shell is equal to the time average of  $\dot{H}_\alpha(\mathbf{z}(t))$ , where  $\mathbf{z}(t)$  is any nonperiodic trajectory of energy  $E$ :

$$\bar{u} = \left\{ \dot{H}_\alpha \right\} = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T dt \dot{H}_\alpha(\mathbf{z}(t)). \quad (5.7)$$

Contributions to this integral occur only along the short segments of  $\mathbf{z}(t)$  that constitute collisions with the wall. The contribution from one such bounce, occurring between times  $t_1$  and  $t_2$  as shown in Fig. 4, is, by Eq. (5.6),

$$\int_{t_1}^{t_2} dt \dot{H}_\alpha(\mathbf{z}(t)) = -\dot{n}_\alpha \hat{\mathbf{n}} \cdot \int_{t_1}^{t_2} dt \nabla V(\mathbf{q}(t)). \quad (5.8)$$

Since  $-\nabla V$  is the force acting on the particle, its integral gives the total change in momentum:

$$\begin{aligned} \int_{t_1}^{t_2} dt \dot{H}_\alpha(\mathbf{z}(t)) &= \dot{n}_\alpha \hat{\mathbf{n}} \cdot [\mathbf{p}(t_2) - \mathbf{p}(t_1)] \\ &= -2mv \dot{n}_\alpha \sin \theta. \end{aligned} \quad (5.9)$$

Thus, Eq. (5.7) becomes

$$\bar{u} = -\frac{2mv}{\tau} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{b=1}^N \dot{n}_{\alpha b} \sin \theta_b, \quad (5.10)$$

the sum being over the bounces occurring between  $t = 0$  and  $t = T$ . The quantity  $\lim_{N \rightarrow \infty} (1/N) \sum_{b=1}^N \dot{n}_{\alpha b} \sin \theta_b$  is the average value of  $n_\alpha \sin \theta$  sampled by a particle bouncing forever off the walls of the frozen container, which, by ergodicity, is equal to the previously defined  $\xi$ . Thus,

$$\bar{u} = -\frac{2mv}{\tau} \xi = -\beta E. \quad (5.11)$$

To solve for  $\int_{-\infty}^{+\infty} ds C(s)$ , note that  $C(s)$  may be written as

$$\left\langle \left[ \dot{H}_\alpha(\mathbf{z}(t)) - \bar{u} \right] \left[ \dot{H}_\alpha(\mathbf{z}(t+s)) - \bar{u} \right] \right\rangle_\alpha, \quad (5.12)$$

where  $\mathbf{z}(t)$  is a trajectory evolving under  $H_\alpha$ , and the angular brackets denote an average over a uniform distribution of such trajectories over the energy shell  $E$ . [Since such a distribution is unchanged by evolution in time, the above expression for  $C(s)$  is independent of  $t$ .] With some manipulation, this allows us to write

$$\int_{-\infty}^{+\infty} ds C(s) = \lim_{T \rightarrow \infty} \frac{1}{T} \left\langle \left( \int_0^T dt \left[ \dot{H}_\alpha(\mathbf{z}(t)) - \bar{u} \right] \right)^2 \right\rangle. \quad (5.13)$$

Using Eqs. (5.9) and (5.11), we have

$$\int_0^T dt \left[ \dot{H}_\alpha(\mathbf{z}(t)) - \bar{u} \right] = -2mv \sum_{b=1}^N (\dot{n}_{\alpha b} \sin \theta_b - \xi), \quad (5.14)$$

where as before the sum is over the bounces of  $\mathbf{z}(t)$  occurring between  $t = 0$  and  $t = T$ . Thus,

$$\int_{-\infty}^{+\infty} ds C(s) = \lim_{T \rightarrow \infty} \frac{1}{T} \left\langle \left[ -2mv \sum_{b=1}^N (\dot{n}_{\alpha b} \sin \theta_b - \xi) \right]^2 \right\rangle, \quad (5.15)$$

$$= \frac{4m^2 v^2}{\tau} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{b=1}^N \sum_{b'=1}^N c_{b'-b}, \quad (5.16)$$

$$= \frac{4m^2 v^2}{\tau} D = \gamma E^{3/2}, \quad (5.17)$$

where the steps taken are similar to those of Sec. III.

By treating the hard-walled container as a limiting case of a potential well [14], we have shown that, in this limit,

$$\bar{u} \rightarrow -\beta E, \quad (5.18)$$

$$\int_{-\infty}^{+\infty} ds C(s) \rightarrow \gamma E^{3/2}. \quad (5.19)$$

When these expressions are plugged into Eqs. (5.1) and (5.2), they give an evolution equation for  $\eta(E, t)$  identical to that obtained in Sec. III. This shows that a chaotic adiabatic billiard gas can be consistently treated as an

example of an ergodic adiabatic ensemble, and so a numerical test of the results presented in the present paper would also stand as a test of the results of Ref. [4].

## VI. ONE-BODY DISSIPATION IN NUCLEAR DYNAMICS

As discussed in Refs. [7,8], it may be instructive to treat a nucleus undergoing some dynamical process (such as fission or heavy-ion collision) as a container, whose shape (but not volume) is allowed to change with time,

filled with a gas of noninteracting point particles. The container is an idealization of the mean field created by the nucleons; the particles represent the individual nucleons moving within this mean field. (Residual nucleon-nucleon interactions are suppressed by Pauli blocking, and are disregarded in this simple approximation.) The solution of this dynamical problem at the classical level is closely related to the problem considered in the present paper.

In applications of this model, one is typically interested in following the shape of the nucleus through some dynamical process. This involves choosing a few reasonable variables to describe the changing shape, then deriving Euler-Lagrange-Rayleigh equations for the evolution of these variables [7,15]. As pointed out in Ref. [7], the particles behave as a source of friction: as they interact with the changing shape of the container (bouncing elastically off its moving walls), there occurs a net flow of energy from the degrees of freedom of the shape, to the degrees of freedom of the gas of particles. This mechanism is known as *one-body dissipation*, and is an example (to our knowledge, the first) of deterministic friction, in which the energy of a few “slow” degrees of freedom is dissipated by their coupling to “fast” deterministic chaotic motion. To incorporate this friction into the equations of motion for the shape of the container, one needs an expression for the rate of this flow of energy, as a function of the way in which the shape is instantaneously changing. In Ref. [7], the wall formula is derived for this rate:

$$\frac{dE_T}{dt} = \rho \bar{v} \oint d\sigma \dot{n}^2. \quad (6.1)$$

Here,  $E_T$  is the total energy of the gas (the sum of the kinetic energies of the individual particles),  $\rho$  is the total mass density of particles inside the container,  $\bar{v}$  is the average speed of the particles, and  $\oint d\sigma \dot{n}^2$  is the surface integral of the square of the normal wall velocity.

The wall formula is derived by treating each infinitesimal area element on the surface of the container as a tiny piston, moving either into or away from the gas of particles. By calculating the work done on the gas by one such piston, then summing over the entire surface (and taking the volume of the container to stay constant), one obtains Eq. (6.1). We will refer to this derivation as the “piston approach” to one-body dissipation [16].

Two key assumptions that enter the derivation of the wall formula are, first, that the motion of the walls is slow compared to that of the particles, and second, that the gas is always distributed uniformly within the container, with an isotropic distribution of velocities. These assumptions are satisfied by a chaotic adiabatic billiard gas, and so the wall formula should be consistent with the results derived in the present paper. To show that this is the case, we first comment that the piston approach disregards any correlations that may exist between the bounces of a particle moving inside the container. Thus, in comparing the wall formula with our results, we use the quasilinear approximation of Sec. IV. The total energy of the gas may be expressed as  $E_T(t) = \int dE \eta(E, t) E$ , where  $\eta$  is the time-dependent distribution of energies. Differentiating with respect to

time, then applying Eq. (4.3b) (with  $\dot{V} = 0$ ), we have

$$\frac{dE_T}{dt} = \frac{\sqrt{2m}}{2V} I_3 \int dE \frac{\partial}{\partial E} \left[ E^2 \frac{\partial}{\partial E} (E^{-1/2} \eta) \right] E, \quad (6.2)$$

where  $I_3 = \oint \dot{n}^2 d\sigma$ . After twice integrating by parts, this becomes

$$\frac{dE_T}{dt} = \frac{\sqrt{2m}}{V} I_3 \int dE \eta E^{1/2}. \quad (6.3)$$

The average speed of the particles is given by

$$\bar{v} = \frac{1}{\mathcal{N}} \int dE \eta (2E/m)^{1/2}, \quad (6.4)$$

where  $\mathcal{N} = \int dE \eta$  is the total number of particles. This enables us to rewrite Eq. (6.3) as

$$\frac{dE_T}{dt} = \frac{m\mathcal{N}}{V} \bar{v} \oint \dot{n}^2 d\sigma, \quad (6.5)$$

which is the wall formula.

Having demonstrated that the results of the present paper (in the simplified form of Sec. IV) agree with the wall formula, we now consider the factor  $\bar{v}$  which appears in the latter. Differentiating both sides of Eq. (6.4) with respect to time, then applying Eq. (4.3b), then integrating by parts twice, we obtain

$$\frac{d\bar{v}}{dt} = \frac{3}{4V} \oint \dot{n}^2 d\sigma. \quad (6.6)$$

Equations (6.5) and (6.6) constitute a closed set of equations, in the sense that, if we know how the shape of the container evolves with time, then Eq. (6.6) may be integrated to yield  $\bar{v}(t)$ , which may then be inserted into the wall formula, which in turn is integrated to give  $E_T(t)$ . Without Eq. (6.6), some assumption must be made about the evolution of  $\bar{v}$  in order for the wall formula to be integrated over any finite length of time. The added understanding of one-body dissipation which one gains from the second wall formula is discussed in greater detail in Refs. [17,18].

We now consider a generalization of Eqs. (6.5) and (6.6). First, note that Eq. (6.5) may be rewritten as

$$\frac{d}{dt} \overline{v^2} = \frac{2}{V} \bar{v} \oint \dot{n}^2 d\sigma, \quad (6.7)$$

where  $\overline{v^2}$  is the average value of particle speed squared. Let  $\overline{v^n}$  denote the average value of the  $n$ th power of particle speed:

$$\overline{v^n}(t) = \int dE \eta(E, t) (2E/m)^{n/2}. \quad (6.8)$$

Differentiating both sides with respect to time, applying Eq. (4.3b), and integrating twice by parts yields

$$\frac{d}{dt} \overline{v^n} = \frac{n(n+2)}{4V} \overline{v^{n-1}} \oint \dot{n}^2 d\sigma; \quad (6.9)$$

Eqs. (6.6) and (6.7) are specific examples of this general formula.



In Ref. [17], Eq. (6.9) is obtained using a generalization of the piston approach described above. A consequence of Eq. (6.9), as shown in Ref. [17], and supported by numerical simulations [18], is that, asymptotically with time, a chaotic adiabatic billiard gas will achieve a distribution of particle velocities which has a universal form:

$$f(\mathbf{v}) \propto \exp(-v/c), \quad (6.10)$$

where  $f(\mathbf{v}) d^3v$  gives the number of particles with velocity in a small region  $d^3v$  around  $\mathbf{v}$ , and the quantity  $c$  is a velocity scale that grows with time. This exponential distribution of velocities stands in contrast to the Maxwell-Boltzmann distribution that occurs when the particles interact with one another.

We conclude this section by drawing attention to the fact that, elsewhere in this paper, we have treated the changing shape of the container as externally imposed, whereas in the nuclear context it is a dynamical quantity. This calls into question the validity of applying Eq. (4.3b) to the problem considered here; might not the evolution of  $\eta$  be affected significantly by allowing the walls to recoil? To answer briefly, we point out that the inertia associated with the collective degrees of freedom of the nucleus, while not infinite, is still much greater than that of an individual nucleon. Therefore the effects of recoil on the evolution of  $\eta$  should constitute a small correction, and Eq. (4.3b) should remain valid to leading order. For a more careful (and general) treatment, see Ref. [19].

## VII. BRIEF DISCUSSION

We conclude with a brief discussion of several issues.

First, the assumption that all of the frozen shapes  $\mathfrak{S}_\alpha$  exhibit global chaos (trajectories chaotically and ergodically explore the energy shell) is admittedly restrictive. Globally chaotic Hamiltonians on the one hand, and integrable Hamiltonians on the other, represent two extremes in the range of dynamical behavior; a generic Hamiltonian will give rise to a mixture of both regular and chaotic trajectories. Therefore, in some sense, this paper is an attempt to understand one of the two extremes. (For an analysis of integrable adiabatic Hamiltonian systems, see Ref. [20]. In mixed systems, the behavior of trajectories near separatrices will be important; this has been investigated in Ref. [21].)

It is possible to rigorously establish the property of global chaos for certain two-dimensional billiard systems [22]. Two of the best-known examples are the Sinai billiard [23] and the Bunimovich stadium [24]. However, in both of these cases there exists a continuous family of periodic trajectories, which, as argued in Ref. [3] (see also references therein, and Ref. [25]), implies that the sum  $\sum_{-\infty}^{+\infty} c_{\Delta b}$  diverges. These systems therefore violate the added assumption made in Sec. III of this paper. (Such systems are said to display *anomalous diffusion*; the problem of how to describe the evolution of  $\eta$  when the frozen shapes exhibit anomalous diffusion would make for an interesting extension of the theory presented here.)

We now propose the “three-leaf clover,” Fig. 5, as a




FIG. 5. The three-leaf clover (heavy outline), a two-dimensional billiard system whose boundary consists of the arcs of six circles. The common radius of the outer circles is  $R$ ; that of the inner ones is  $r$ . By varying  $\alpha = R/r$ , one has a continuous family of such shapes, all globally chaotic. For  $\alpha \geq 1$ , all periodic orbits are isolated.

family of billiard systems satisfying the conditions of the present paper. Varying the parameter  $\alpha = R/r$  gives a continuous family of shapes  $\mathfrak{S}_\alpha$ . By direct application of Theorem 1 of Ref. [22], one can establish the property of global chaos for any of these shapes. Furthermore, it is fairly straightforward to prove that, for  $\alpha \geq 1$ , all periodic orbits inside  $\mathfrak{S}_\alpha$  are isolated, i.e., no continuous families exist. Thus, by filling such a clover with a gas of noninteracting particles, then allowing  $R$  and  $r$  to change slowly with time, always maintaining  $\alpha \geq 1$ , one has an example of a two-dimensional system satisfying the assumptions of this paper. (For another example, see the modified Sinai billiard in Ref. [3].)

Of course, the theory presented in this paper needs to be tested numerically. We are currently running simulations of particles inside a time-dependent three-leaf clover, and plan to publish the results at a later date. However, some numerical results supporting predictions discussed in Sec. VI already exist in the literature.

First, the wall formula, which was shown to follow from Eq. (4.3b), has been studied extensively for a variety of three-dimensional hard-walled cavities [8]. In these studies, the results of numerical simulations agree well with the wall formula, provided that the dynamics in the frozen billiard is sufficiently dominated by chaotic trajectories.

Another result following from Eq. (4.3b) was the second wall formula [Eq. (6.6)], giving the rate of change of  $\bar{v}$ , the average speed of the particles. Combining the two wall formulas, we have obtained [17] a prediction for the amount of energy dissipated after a time long enough that  $\bar{v}$  has changed significantly (i.e., beyond the validity of the original wall formula alone). This prediction has been tested in Ref. [18] for the same set of shapes as the wall formula, and again there is good agreement with theory.

A final prediction made in Sec. VI was that, in the long-time limit, a three-dimensional chaotic adiabatic billiard gas will achieve an exponential distribution of particle velocities. Once more, this prediction has been tested for the same set of shapes [18], and once more the numerics

support the theory.

This agreement between numerical simulation and Eq. (4.3b) raises two questions. First, the billiard systems involved in the computer experiments do not satisfy the assumption of global chaos: in each case, when the shape is frozen, some small but finite fraction of phase space is filled with nonchaotic trajectories. Perhaps, then, the results of this paper, while strictly valid only for the narrow class of systems satisfying the assumptions of Sec. I, in reality work well for a much wider range of billiards. It is reasonable to speculate that, if phase space is dominated by chaos, with only small isolated islands of regular motion, then the results of this paper may still be applicable.

Second, Eq. (4.3b) follows from the quasilinear approximation, in which all correlations between different bounces are ignored. However, we have no reason to believe *a priori* that this approximation holds for the set of shapes tested, which are all simple deformations of a spherical cavity. Is the validity of this approximation more general than one might at first expect, and if so, how can we understand this?

Both of these issues are currently under investigation.

### ACKNOWLEDGMENTS

I would like to thank Władek Świątecki, Robert Littlejohn, and Jim Morehead for many useful discussions. I would also like to thank Jan Błocki at the Institute for Nuclear Studies, Świerk, Poland, and Janusz Skalski at the University of Warsaw, where parts of this manuscript were written, for their hospitality. This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098; and by the Director, Office of Energy Research, Office of Basic Energy Sciences, of the U.S. Department of Energy under Contract No. DE-AC03-75SF00098; and by the National Science Foundation under Grant No. NSF-PYI-84-51276; and by the Polish-American Maria Skłodowska-Curie Joint Fund II in cooperation with the National Science Foundation (U.S.) and the National Atomic Energy Agency (Poland), under project EMB ID#70.

### APPENDIX A

In this appendix we evaluate the quantities  $\xi$  and  $c_0$ , both functions of  $et$ . These quantities are defined with respect to a gas of noninteracting particles evolving inside the frozen shape  $\mathfrak{S}_\alpha$ , with  $\alpha = et$ . The particles are assumed to share a common energy, and to be distributed uniformly within the container, with an isotropic distribution of velocities. While the definitions of  $\xi$  and  $c_0$  involve the dynamics of these particles, our final expressions will be given solely in terms of quantities characterizing the shape of the container and its instantaneous wall velocity field. We use the following notation.  $\mathcal{N}$  is the number of particles in our gas ( $\mathcal{N} \gg 1$ ), and  $v$  is

their common speed.  $A$  and  $l$ , both functions of  $et$ , refer to the area and perimeter of the two-dimensional container;  $\oint ds$  denotes a line integral over the entire wall. Similarly, in three dimensions,  $V$  and  $S$  refer to the volume and surface area of the container, and  $\oint d\sigma$  denotes a surface integral over the wall. Note that

$$\dot{A} \equiv dA/dt = \oint ds \dot{n} \quad (2D),$$

$$\dot{V} \equiv dV/dt = \oint d\sigma \dot{n} \quad (3D), \quad (A1)$$

where  $\dot{n} = \dot{n}(et)$  is the wall velocity field describing the evolution of  $\mathfrak{S}(et)$ .

The quantity  $\xi$  was defined in Sec. III as

$$\xi = \xi_b = \langle \dot{n}_{\alpha b} \sin \theta_b \rangle, \quad (A2)$$

the average value of  $\dot{n}_\alpha \sin \theta$  over the  $b$ th bounce of all particles in the gas. As mentioned, the invariance with time of the distribution of these particles implies that  $\xi$  is independent of  $b$ , and therefore we may alternatively write it as

$$\xi = \langle \langle \dot{n}_\alpha \sin \theta \rangle \rangle, \quad (A3)$$

where the double angular brackets denote an average over *all* bounces of all the particles of the gas. In this form,  $\xi$  becomes easy to evaluate.

To evaluate  $\xi$  in two dimensions, first consider a small segment  $ds$  of the wall of the container. The rate  $r$  at which this segment is being struck by particles making an angle of collision between  $\theta$  and  $\theta + d\theta$  is given by

$$r = j ds \sin \theta, \quad (A4)$$

where  $j = (d\theta/2\pi)(\mathcal{N}v/A)$  is the current density of particles bombarding  $ds$  from this range of angles, and  $\sin \theta$  is a flux factor. The quantity  $\xi = \langle \langle \dot{n}_\alpha \sin \theta \rangle \rangle$  is then the weighted average  $\int r (\dot{n}_\alpha \sin \theta) / \int r$ , where the integral is over the entire wall, and over  $\theta$  from 0 to  $\pi$ . This yields

$$\xi = \frac{\pi \dot{A}}{4l}. \quad (A5)$$

The quantity  $c_0 = c_{b,b} = \langle \langle (\dot{n}_{\alpha b} \sin \theta_b - \xi_b)^2 \rangle \rangle$  is, like  $\xi$ , independent of  $b$ , and may be written as

$$\begin{aligned} c_0 &= \langle \langle (\dot{n}_\alpha \sin \theta - \xi)^2 \rangle \rangle \\ &= \int r (\dot{n}_\alpha \sin \theta - \xi)^2 / \int r, \end{aligned} \quad (A6)$$

which reduces to

$$c_0 = \frac{2}{3l} \oint ds \left[ \dot{n}^2 - \frac{3\pi^2}{32} \bar{n}^2 \right], \quad (A7)$$

where  $\bar{n} \equiv (1/l) \oint ds \dot{n} = \dot{A}/l$  is the average value of  $\dot{n} = \dot{n}(et)$  over the wall of the container. [Since the final expression for  $c_0$  no longer involves the dynamics of particles in the frozen container, our notation has reverted from  $\dot{n}_\alpha$  to  $\dot{n}(et)$ .]

In three dimensions, consider a small patch  $d\sigma$  on the surface of the container. Let  $r$  be the rate at which this patch is being struck by particles coming from a solid angle  $d\Omega$  around the direction  $(\theta, \phi)$ , as defined by Fig. 3. This rate is given by  $r = j d\sigma \sin\theta$ , where

$$j = \frac{d\Omega}{4\pi} \frac{\mathcal{N}}{V} v = \frac{\cos\theta d\theta d\phi}{4\pi} \frac{\mathcal{N}}{V} v. \quad (\text{A8})$$

$\xi = \langle \langle \dot{n}_\alpha \sin\theta \rangle \rangle$  is again equal to  $\int r (\dot{n}_\alpha \sin\theta) / \int r$ , only now the integral is over the entire surface area of the wall,  $\theta$  from 0 to  $\pi/2$ , and  $\phi$  from 0 to  $2\pi$ . This yields

$$\xi = \frac{2\dot{V}}{3S}. \quad (\text{A9})$$

Similarly,

$$\begin{aligned} c_0 &= \int r (\dot{n}_\alpha \sin\theta - \xi)^2 / \int r \\ &= \frac{1}{2S} \oint d\sigma \left[ \dot{n}^2 - \frac{8}{9} \bar{n}^2 \right], \end{aligned} \quad (\text{A10})$$

where  $\bar{n} = (1/S) \oint d\sigma \dot{n} = \dot{V}/S$  is again the average of  $\dot{n}(et)$  over the wall.

## APPENDIX B

In this appendix we solve for  $\tau$ , the average time between the bounces of a particle of speed  $v$  moving chaotically inside a frozen container. Filling the container with a large number  $\mathcal{N}$  of such particles, the total rate at which the walls of the container are being struck is  $R = \mathcal{N}/\tau$ . Alternatively,  $R = \int r$ , in the notation of Appendix A. Setting these two equal yields

$$\tau = \frac{\pi A}{lv} \quad (\text{2D}),$$

$$\tau = \frac{4V}{Sv} \quad (\text{3D}). \quad (\text{B1})$$

- 
- [1] E. Ott, Phys. Rev. Lett. **42**, 1628 (1979).  
 [2] R. Brown, E. Ott, and C. Grebogi, Phys. Rev. Lett. **59**, 1173 (1987).  
 [3] R. Brown, E. Ott, and C. Grebogi, J. Stat. Phys. **49**, 511 (1987).  
 [4] C. Jarzynski, Phys. Rev. A **46**, 7498 (1992).  
 [5] C. Jarzynski, Phys. Rev. Lett. **71**, 839 (1993).  
 [6] M. Wilkinson, J. Phys. A **23**, 3603 (1990).  
 [7] J. Błocki *et al.*, Ann. Phys. **113**, 338 (1978).  
 [8] J. Błocki, J.-J. Shi, and W.J. Świątecki, Nucl. Phys. **A554**, 387 (1993).  
 [9] Thus, we are assuming that the system is uniformly hyperbolic: neighboring trajectories will, with probability 1, diverge away from one another at a rate which is exponential in the long-time limit; and since each trajectory explores the entire energy shell, the Lyapunov exponent associated with this divergence is the same for all trajectories.  
 [10] R. Kubo, M. Toda, and N. Hashitsume, *Statistical Physics II* (Springer-Verlag, Berlin, 1985), Chap. 1.  
 [11] F. Reif, *Fundamentals of Statistical and Thermal Physics* (McGraw-Hill, New York, 1965), Sec. 15.9.  
 [12] A.J. Lichtenberg and M.A. Lieberman, *Regular and Stochastic Motion* (Springer-Verlag, New York, 1983), p. 294.  
 [13] S.E. Koonin and J. Randrup, Nucl. Phys. **A289**, 476 (1977).  
 [14] In Ref. [3], Brown *et al.* define a correlation function  $C(s)$  for billiard systems. Integrating this function from  $s = -\infty$  to  $s = +\infty$  provides an alternative method of obtaining Eq. (5.19).  
 [15] H. Goldstein, *Classical Mechanics*, 2nd ed. (Addison-Wesley, Reading, 1980), p. 24.  
 [16] D. Gross, Nucl. Phys. **A240**, 472 (1975).  
 [17] C. Jarzynski and W.J. Świątecki, Nucl. Phys. **A552**, 1 (1993).  
 [18] J. Błocki, F. Brut, and W.J. Świątecki, Nucl. Phys. **A554**, 107 (1993).  
 [19] M.V. Berry and J.M. Robbins, Proc. R. Soc. London Ser. A **442**, 659 (1993).  
 [20] M. Kruskal, J. Math. Phys. **3**, 806 (1962).  
 [21] J. Tennyson *et al.*, Phys. Rev. Lett. **56**, 2117 (1986).  
 [22] M. Wojtkowski, Commun. Math. Phys. **105**, 391 (1986).  
 [23] Ya. Sinai, Russian Math. Surveys **25**, 137 (1970).  
 [24] L. Bunimovich, Commun. Math. Phys. **65**, 295 (1979).  
 [25] B. Friedman and R.F. Martin, Jr., Phys. Lett. **105A**, 23 (1984).