

Universality of period doubling in coupled maps

Sang-Yoon Kim

Department of Physics, Kangwon National University, Chunchon, Kangwon-Do 200-701, Korea

(Received 18 August 1993; revised manuscript received 8 November 1993)

We study the critical behavior of period doubling in two coupled one-dimensional maps with a single maximum of order z . In particular, the effect of the maximum-order z on the critical behavior associated with coupling is investigated by a renormalization method. There exist three fixed maps of the period-doubling renormalization operator. For a fixed map associated with the critical behavior at the zero-coupling critical point, relevant eigenvalues associated with coupling perturbations vary, depending on the order z , whereas they are independent of z for the other two fixed maps. The renormalization results for the zero-coupling case are also confirmed by a direct numerical method.

PACS number(s): 05.45.+b, 03.20.+i, 05.70.Jk

Universal scaling behavior of period doubling has been found in one-dimensional (1D) maps with a single maximum of order z ($z > 1$),

$$x_{i+1} = f(x_i) = 1 - A |x_i|^z, \quad z > 1. \quad (1)$$

For all $z > 1$, the 1D map (1) exhibits successive period-doubling bifurcations as the nonlinearity parameter A is increased. The period-doubling bifurcation points $A = A_n(z)$ ($n = 0, 1, 2, \dots$), at which the n th period-doubling bifurcation occurs, converge to the accumulation point $A^*(z)$ on the A axis. The scaling behavior near the critical point A^* depends on the maximum-order z , i.e., the parameter and orbital scaling factors, δ and α , vary depending on z [1-4]. Therefore the order z determines universality classes.

Here we study the critical behavior of period doubling in a map T consisting of two identical 1D maps coupled symmetrically:

$$T : \begin{cases} x_{i+1} = F(x_i, y_i) = f(x_i) + g(x_i, y_i), \\ y_{i+1} = F(y_i, x_i) = f(y_i) + g(y_i, x_i), \end{cases} \quad (2)$$

where $f(x)$ is a 1D map (1) with a single maximum of even-order z ($z = 2, 4, 6, \dots$) at $x = 0$, and $g(x, y)$ is a coupling function. The uncoupled 1D map f satisfies a normalization condition $f(0) = 1$, and the coupling function g obeys a condition $g(x, x) = 0$ for any x .

The quadratic-maximum case ($z = 2$) was previously studied in Refs. [5-9]. In this paper, using the renormalization method developed in Ref. [9], we extend the results for the $z = 2$ case to all even-order cases and investigate the dependence of the critical behavior associated with coupling on the order z .

The period-doubling renormalization transformation \mathcal{N} for a coupled map T consists of squaring (T^2) and rescaling (B) operators:

$$\mathcal{N}(T) \equiv BT^2B^{-1}. \quad (3)$$

Here the rescaling operator B is:

$$B = \begin{pmatrix} \alpha & 0 \\ 0 & \alpha \end{pmatrix}, \quad (4)$$

because we consider only in-phase orbits ($x_i = y_i$ for all i).

Applying the renormalization operator \mathcal{N} to the coupled map (2) n times, we obtain the n -times renormalized map T_n of the form,

$$T_n : \begin{cases} x_{i+1} = F_n(x_i, y_i) = f_n(x) + g_n(x, y), \\ y_{i+1} = F_n(y_i, x_i) = f_n(y) + g_n(y, x). \end{cases} \quad (5)$$

Here f_n and g_n are the uncoupled and coupling parts of the n -times renormalized function F_n , respectively. They satisfy the following recurrence equations [9]:

$$\begin{aligned} f_{n+1}(x) &= \alpha f_n \left(f_n \left(\frac{x}{\alpha} \right) \right), \quad (6) \\ g_{n+1}(x, y) &= \alpha f_n \left(f_n \left(\frac{x}{\alpha} \right) + g_n \left(\frac{x}{\alpha}, \frac{y}{\alpha} \right) \right) \\ &\quad + \alpha g_n \left(f_n \left(\frac{x}{\alpha} \right) + g_n \left(\frac{x}{\alpha}, \frac{y}{\alpha} \right), f_n \left(\frac{y}{\alpha} \right) \right) \\ &\quad + g_n \left(\frac{y}{\alpha}, \frac{x}{\alpha} \right) - \alpha f_n \left(f_n \left(\frac{x}{\alpha} \right) \right), \quad (7) \end{aligned}$$

where the rescaling factor α is chosen to preserve the normalization condition $f_{n+1}(0) = 1$, i.e., $\alpha = 1/f_n(1)$. Equations (6) and (7) define a renormalization operator \mathcal{R} for transforming a pair of functions (f, g) ;

$$\begin{pmatrix} f_{n+1} \\ g_{n+1} \end{pmatrix} = \mathcal{R} \begin{pmatrix} f_n \\ g_n \end{pmatrix}. \quad (8)$$

A critical map T_c with the nonlinearity and coupling parameters set to their critical values is attracted to a fixed map T^* under iterations of the renormalization transformation \mathcal{N} ,

$$T^* : \begin{cases} x_{i+1} = F^*(x_i, y_i) = f^*(x_i) + g^*(x_i, y_i), \\ y_{i+1} = F^*(y_i, x_i). \end{cases} \quad (9)$$

Here (f^*, g^*) is a fixed point of the renormalization operator \mathcal{R} with $\alpha = 1/f^*(1)$, which satisfies $(f^*, g^*) = \mathcal{R}(f^*, g^*)$. Note that the equation for f^* is just the fixed-

point equation in the 1D map case. The 1D fixed function f^* varies depending on the order z [4]. Consequently only the equation for the coupling fixed function g^* is left to be solved.

However it is not easy to directly solve the equation for the coupling fixed function. We therefore introduce a tractable recurrence equation for a “reduced” coupling function of the coupling function $g(x, y)$ [9], defined by

$$G(x) \equiv \left. \frac{\partial g(x, y)}{\partial y} \right|_{y=x}. \tag{10}$$

Differentiating the recurrence equation (7) for g with respect to y and setting $y = x$, we have

$$G_{n+1}(x) = \left[f'_n \left(f_n \left(\frac{x}{\alpha} \right) \right) - 2G_n \left(f_n \left(\frac{x}{\alpha} \right) \right) \right] G_n \left(\frac{x}{\alpha} \right) + G_n \left(f_n \left(\frac{x}{\alpha} \right) \right) f'_n \left(\frac{x}{\alpha} \right). \tag{11}$$

Then Eqs. (6) and (11) define a “reduced” renormalization operator $\tilde{\mathcal{R}}$ for transforming a pair of functions (f, G) :

$$\begin{pmatrix} f_{n+1} \\ G_{n+1} \end{pmatrix} = \tilde{\mathcal{R}} \begin{pmatrix} f_n \\ G_n \end{pmatrix}. \tag{12}$$

We look for a fixed point (f^*, G^*) of $\tilde{\mathcal{R}}$, which satisfies $(f^*, G^*) = \tilde{\mathcal{R}}(f^*, G^*)$. Here G^* is just the reduced coupling fixed function of g^* [i.e., $G^*(x) = \partial g^*(x, y)/\partial y|_{y=x}$]. As in the quadratic-maximum case ($z = 2$) [9], we find three solutions for G^* :

$$G^*(x) = 0, \tag{13}$$

$$G^*(x) = \frac{1}{2} f^{*'}(x), \tag{14}$$

$$G^*(x) = \frac{1}{2} [f^{*'}(x) - 1]. \tag{15}$$

Here the first solution, corresponding to the reduced coupling fixed function of the zero-coupling fixed function $g^*(x, y) = 0$, is associated with the critical behavior at the zero-coupling critical point, whereas the second and third solutions dependent on the order z are associated with the critical behavior at other critical points [9].

Consider an infinitesimal reduced coupling perturbation $(0, \Phi(x))$ to a fixed point (f^*, G^*) of $\tilde{\mathcal{R}}$. We then examine the evolution of a pair of functions, $(f^*(x), G^*(x) + \Phi(x))$ under the reduced renormalization transformation $\tilde{\mathcal{R}}$. In the linear approximation we obtain a reduced linearized operator $\tilde{\mathcal{L}}$ of transforming a reduced coupling perturbation Φ :

$$\begin{aligned} \Phi_{n+1}(x) &= [\tilde{\mathcal{L}}\Phi_n](x) \\ &= \left[f^{*'} \left(f^* \left(\frac{x}{\alpha} \right) \right) - 2G^* \left(f^* \left(\frac{x}{\alpha} \right) \right) \right] \Phi_n \left(\frac{x}{\alpha} \right) \\ &\quad + \left[f^{*'} \left(\frac{x}{\alpha} \right) - 2G^* \left(\frac{x}{\alpha} \right) \right] \Phi_n \left(f^* \left(\frac{x}{\alpha} \right) \right). \end{aligned} \tag{16}$$

Here the prime denotes a derivative. If a reduced coupling perturbation $\Phi^*(x)$ satisfies

$$\nu \Phi^*(x) = [\tilde{\mathcal{L}}\Phi^*](x), \tag{17}$$

then it is called a reduced coupling eigenperturbation with coupling eigenvalue (CE) ν .

We first show that CE’s are independent of the order z for the second and third solutions (14) and (15) of $G^*(x)$. In case of the second solution $G^*(x) = \frac{1}{2} f^{*'}(x)$, the reduced linearized operator $\tilde{\mathcal{L}}$ becomes a null operator, independently of z , because the right-hand side of Eq. (17) becomes zero. Therefore there exist no relevant CE’s. For the third case $G^*(x) = \frac{1}{2} [f^{*'}(x) - 1]$, Eq. (17) becomes

$$\nu \Phi^*(x) = \Phi^* \left(\frac{x}{\alpha} \right) + \Phi^* \left(f^* \left(\frac{x}{\alpha} \right) \right). \tag{18}$$

When $\Phi^*(x)$ is a nonzero constant function, i.e., $\Phi^*(x) = c$ (c : nonzero constant), there exists a relevant CE, $\nu = 2$, independently of z .

In the zero-coupling case $G^*(x) = 0$, the eigenvalue equation (17) becomes

$$\begin{aligned} \nu \Phi^*(x) &= f^{*'} \left(f^* \left(\frac{x}{\alpha} \right) \right) \Phi^* \left(\frac{x}{\alpha} \right) \\ &\quad + f^{*'} \left(\frac{x}{\alpha} \right) \Phi^* \left(f^* \left(\frac{x}{\alpha} \right) \right). \end{aligned} \tag{19}$$

Relevant CE’s of Eq. (19) vary depending on the order z , as will be seen below.

An eigenfunction $\Phi^*(x)$ can be separated into two components, $\Phi^*(x) = \Phi^{*(1)}(x) + \Phi^{*(2)}(x)$ with $\Phi^{*(1)}(x) \equiv a_0^* + a_1^*x + \dots + a_{z-2}^*x^{z-2}$ and $\Phi^{*(2)}(x) \equiv a_{z-1}^*x^{z-1} + a_z^*x^z + \dots$, and the 1D fixed function f^* is a polynomial in x^z , i.e., $f^*(x) = 1 + c_z^*x^z + c_{2z}^*x^{2z} + \dots$. Substituting the functions Φ^* , f^* , and $f^{*'}$ into the eigenvalue equation (19), it has the structure

$$\nu a_k^* = \sum_l M_{kl}(\{c^*\}) a_l^*, \quad k, l = 0, 1, 2, \dots \tag{20}$$

We note that each a_l^* ($l = 0, 1, 2, \dots$) in the first and second terms in the right-hand side of Eq. (19) is involved only in the determination of coefficients of monomials x^k with $k = l + mz$ and $k = (z - 1) + mz$ ($m = 0, 1, 2, \dots$), respectively. Therefore any a_l^* with $l \geq z - 1$ (in the right-hand side) cannot be involved in the determination of coefficients of monomials x^k with $k < z - 1$, which implies that the eigenvalue equation (20) is of the form

$$\nu \begin{pmatrix} \Phi^{*(1)} \\ \Phi^{*(2)} \end{pmatrix} = \begin{pmatrix} M_1 & 0 \\ M_3 & M_2 \end{pmatrix} \begin{pmatrix} \Phi^{*(1)} \\ \Phi^{*(2)} \end{pmatrix}, \tag{21}$$

where M_1 is a $(z - 1) \times (z - 1)$ matrix, $\Phi^{*(1)} \equiv (a_0^*, \dots, a_{z-2}^*)$, and $\Phi^{*(2)} \equiv (a_{z-1}^*, a_z^*, \dots)$. From the reducibility of the matrix M into a semiblock form, it follows that to determine the eigenvalues of M it is sufficient to solve the eigenvalue problems for the two submatrices M_1 and M_2 independently.

We first solve the eigenvalue equation of M_1 ($\nu \Phi^{*(1)} = M_1 \Phi^{*(1)}$), i.e.,

$$\nu a_k^* = \sum_l M_{kl}(\{c^*\}) a_l^*, \quad k, l = 0, \dots, z - 2. \tag{22}$$

Note that this submatrix M_1 is diagonal. Hence its eigen-

values are just the diagonal elements:

$$\nu_k = M_{kk} = \frac{f^{*'}(1)}{\alpha^k} = \alpha^{z-1-k}, \quad k = 0, \dots, z-2. \quad (23)$$

Notice that all ν_k 's are relevant eigenvalues.

Although ν_k is also an eigenvalue of M , $(\Phi_k^{*(1)}, 0)$ cannot be an eigenvector of M , because there exists a third submatrix M_3 in M [see Eq. (21)]. Therefore an eigenfunction $\Phi_k^*(x)$ in Eq. (19) with eigenvalue ν_k is a polynomial with a leading monomial of degree k , i.e., $\Phi_k^*(x) = \Phi_k^{*(1)}(x) + \Phi_k^{*(2)}(x) = a_k^* x^k + a_{z-1}^* x^{z-1} + a_z^* x^z + \dots$, where $a_k^* \neq 0$.

We next solve the eigenvalue equation of M_2 ($\nu \Phi^{*(2)} = M_2 \Phi^{*(2)}$), i.e.,

$$\nu a_k^* = \sum_l M_{kl}(\{c^*\}) a_l^*, \quad k, l = z-1, z, \dots \quad (24)$$

Unlike the case of M_1 , $(0, \Phi^{*(2)})$ can be an eigenvector of M with eigenvalue ν . Then its corresponding function $\Phi^{*(2)}(x)$ is an eigenperturbation with eigenvalue ν , which satisfies Eq. (19). One can easily see that $\Phi^{*(2)}(x) = f^{*'}(x)$ is an eigenfunction with CE $\nu = 2$, which is the z th relevant CE in addition to those in Eq. (23). It is also found that there exist an infinite number of additional (coordinate change) eigenfunctions $\Phi^{*(2)}(x) = f^{*'}(x)[f^{*n}(x) - x^n]$ with irrelevant CE's α^{-n} ($n = 1, 2, \dots$), which are associated with coordinate changes [10]. We conjecture that together with the z (noncoordinate change) relevant CE's, they give the whole spectrum of the reduced linearized operator \tilde{L} of Eq. (16) and the spectrum is complete.

Consider an infinitesimal coupling perturbation $g(x, y)$ [$= \varepsilon \varphi(x, y)$] to a critical map at the zero-coupling critical point, in which case the map T of Eq. (2) is of the form,

$$T : \begin{cases} x_{i+1} = F(x_i, y_i) = f_{A^*}(x_i) + \varepsilon \varphi(x, y), \\ y_{i+1} = F(y_i, x_i), \end{cases} \quad (25)$$

where the subscript A^* of f denotes the critical value of the nonlinearity parameter A and ε is an infinitesimal coupling parameter. The map T for $\varepsilon = 0$ is just the zero-coupling critical map consisting of two uncoupled 1D critical maps f_{A^*} . It is attracted to the zero-coupling fixed map (consisting of two 1D fixed maps f^*) under iterations of the renormalization transformation \mathcal{N} of Eq. (3).

The reduced coupling function $G(x)$ of $g(x, y)$ is given by [see Eq. (10)]

$$G(x) = \varepsilon \Phi(x) \equiv \varepsilon \left. \frac{\partial \varphi(x, y)}{\partial y} \right|_{y=x}. \quad (26)$$

Then $\varepsilon \Phi(x)$ corresponds to an infinitesimal perturbation to the reduced zero-coupling fixed function $G^*(x) = 0$ of Eq. (13). The n th image Φ_n of Φ under the reduced linearized operator \tilde{L} of Eq. (16) has the form,

$$\begin{aligned} \Phi_n(x) &= [\tilde{L}^n \Phi](x) \\ &\simeq \sum_{k=0}^{z-2} \alpha_k \nu_k^n \Phi_k^*(x) + \alpha_{z-1} 2^n f^{*'}(x) \quad \text{for large } n, \end{aligned} \quad (27)$$

since the irrelevant part of Φ_n becomes negligibly small for large n .

The stability multipliers $\lambda_{1,n}$ and $\lambda_{2,n}$ of the 2^n -periodic orbit of the map T of Eq. (25) are the same as those of the fixed point of the n -times renormalized map $\mathcal{N}^n(T)$ [9], which are given by

$$\lambda_{1,n} = f_n'(\hat{x}_n), \quad \lambda_{2,n} = f_n'(\hat{x}_n) - 2G_n(\hat{x}_n). \quad (28)$$

Here (f_n, G_n) is the n th image of (f_{A^*}, G) under the reduced renormalization transformation $\tilde{\mathcal{R}}$ [i.e., $(f_n, G_n) = \tilde{\mathcal{R}}^n(f_{A^*}, G)$], and \hat{x}_n is just the fixed point of $f_n(x)$ [i.e., $\hat{x}_n = f_n(\hat{x}_n)$] and converges to the fixed point \hat{x} of the 1D fixed map $f^*(x)$ as $n \rightarrow \infty$. In the critical case ($\varepsilon = 0$), $\lambda_{2,n}$ is equal to $\lambda_{1,n}$ and they converge to the 1D critical stability multiplier $\lambda^* = f^{*'}(\hat{x})$. Since $G_n(x) \simeq [\tilde{\mathcal{L}}_2^n G](x) = \varepsilon \Phi_n(x)$ for infinitesimally small ε , $\lambda_{2,n}$ has the form

$$\begin{aligned} \lambda_{2,n} &\simeq \lambda_{1,n} - 2\varepsilon \Phi_n \\ &\simeq \lambda^* + \varepsilon \left[\sum_{k=0}^{z-2} e_k \nu_k^n + e_{z-1} 2^n \right] \quad \text{for large } n, \end{aligned} \quad (29)$$

where $e_k = -2\alpha_k \Phi_k^*(\hat{x})$ ($k = 0, \dots, z-2$) and $e_{z-1} = -2\alpha_{z-1} f^{*'}(\hat{x})$. Therefore the slope S_n of $\lambda_{2,n}$ at the zero-coupling point ($\varepsilon = 0$) is

$$S_n \equiv \left. \frac{\partial \lambda_{2,n}}{\partial \varepsilon} \right|_{\varepsilon=0} \simeq \sum_{k=0}^{z-2} e_k \nu_k^n + e_{z-1} 2^n \quad \text{for large } n. \quad (30)$$

Here the coefficients $\{e_k; k = 0, \dots, z-1\}$ depend on the initial reduced function $\Phi(x)$, because the α_k 's are determined only by $\Phi(x)$. Note that the magnitude of slope S_n increases with n unless all e_k 's ($k = 0, \dots, z-1$) are zero.

We choose monomials x^l ($l = 0, 1, 2, \dots$) as initial reduced functions $\Phi(x)$, because any smooth function $\Phi(x)$ can be represented as a linear combination of monomials by a Taylor series. Expressing $\Phi(x) = x^l$ as a linear combination of eigenfunctions of $\tilde{\mathcal{L}}_2$, we have

$$\begin{aligned} \Phi(x) = x^l &= \alpha_l \Phi_l^*(x) + \alpha_{z-1} f^{*'}(x) \\ &\quad + \sum_{n=1}^{\infty} \beta_n f^{*n}(x) [f^{*n}(x) - x^n], \end{aligned} \quad (31)$$

where α_l is nonzero for $l < z-1$ and zero for $l \geq z-1$, and all β_n 's are irrelevant components. Therefore the slope S_n for large n becomes

$$S_n \simeq \begin{cases} e_l \nu_l^n + e_{z-1} 2^n & \text{for } l < z-1, \\ e_{z-1} 2^n & \text{for } l \geq z-1. \end{cases} \quad (32)$$

Note that the growth of S_n for large n is governed by two CE's ν_l and 2 for $l < z-1$ and by one CE $\nu = 2$ for $l \geq z-1$.

We numerically study the quartic-maximum case ($z = 4$) in the two coupled 1D maps (25) and confirm the renormalization results (32). In this case we follow the periodic orbits of period 2^n up to level $n = 15$ and obtain the slopes S_n of Eq. (30) at the zero-coupling critical point $(A^*, 0)$ ($A^* = 1.594\,901\,356\,228\dots$) when the reduced function $\Phi(x)$ is a monomial x^l ($l = 0, 1, \dots$).

The renormalization result implies that the slopes S_n for $l \geq z - 1$ obey a one-term scaling law asymptotically:

$$S_n = d_1 r_1^n. \quad (33)$$

We therefore define the growth rate of the slopes as follows:

$$r_{1,n} \equiv \frac{S_{n+1}}{S_n}. \quad (34)$$

Then it will converge to a constant r_1 as $n \rightarrow \infty$. A sequence of $r_{1,n}$ for $\Phi(x) = x^3$ is shown in the second column of Table I. Note that it converges fast to $r_1 = 2$. We have also studied several other reduced-coupling cases with $\Phi(x) = x^l$ ($l > 3$). In all higher-order cases studied, the sequences of $r_{1,n}$ also converge fast to $r_1 = 2$.

When $l < z - 1$, two relevant CE's govern the growth of the slopes S_n . We therefore extend the simple one-term scaling law (33) to a two-term scaling law:

$$S_n = d_1 r_1^n + d_2 r_2^n, \text{ for large } n, \quad (35)$$

where $|r_1| > |r_2|$. This is a kind of multiple-scaling law [11]. Equation (35) gives

$$S_{n+2} = t_1 S_{n+1} - t_2 S_n, \quad (36)$$

where $t_1 = r_1 + r_2$ and $t_2 = r_1 r_2$. Then r_1 and r_2 are solutions of the following quadratic equation,

$$r^2 - t_1 r + t_2 = 0. \quad (37)$$

To evaluate r_1 and r_2 , we first obtain t_1 and t_2 from the S_n 's using Eq. (36):

$$t_1 = \frac{S_{n+1}S_n - S_{n+2}S_{n-1}}{S_n^2 - S_{n+1}S_{n-1}}, \quad t_2 = \frac{S_{n+1}^2 - S_n S_{n+2}}{S_n^2 - S_{n+1}S_{n-1}}. \quad (38)$$

Note that Eqs. (35)–(38) are valid for large n . In fact,

TABLE I. In the quartic-maximum case ($z = 4$), a sequence $\{r_{1,n}\}$ for a one-term scaling law is shown in the second column when $\Phi(x) = x^3$, and two sequences $\{r_{1,n}\}$ and $\{r_{2,n}\}$ for a two-term scaling law are shown in the third and fourth columns when $\Phi(x) = 1$.

n	$\Phi(x) = x^3$	$\Phi(x) = 1$	
	$r_{1,n}$	$r_{1,n}$	$r_{2,n}$
5	1.999 920 2	-4.829 455 8	1.958
6	2.000 009 3	-4.829 422 6	2.090
7	1.999 994 7	-4.829 409 8	1.973
8	2.000 000 4	-4.829 406 8	2.039
9	1.999 999 6	-4.829 405 8	1.984
10	2.000 000 0	-4.829 405 5	2.018
11	2.000 000 0	-4.829 405 5	1.992
12	2.000 000 0	-4.829 405 4	2.009

the values of t_i 's and r_i 's ($i = 1, 2$) depend on the level n . Thus we denote the values of t_i 's in Eq. (38) explicitly by $t_{i,n-1}$'s, and the values of r_i 's obtained from Eq. (37) are also denoted by $r_{i,n-1}$'s. Then each of them converges to a constant as $n \rightarrow \infty$:

$$\lim_{n \rightarrow \infty} t_{i,n} = t_i, \quad \lim_{n \rightarrow \infty} r_{i,n} = r_i, \quad i = 1, 2. \quad (39)$$

The two-term scaling law (35) is very well obeyed. Sequences $r_{1,n}$ and $r_{2,n}$ for $\Phi(x) = 1$ are shown in the third and fourth columns of Table I. They converge fast to $r_1 = \alpha^3$ ($\alpha = -1.6903\dots$) and $r_2 = 2$, respectively. We have also studied two other reduced-coupling cases with $\Phi(x) = x^l$ ($l = 1, 2$). It is found that the sequences $r_{1,n}$ and $r_{2,n}$ for $l = 1(2)$ converge fast to their limit values $r_1 = \alpha^2(2)$ and $r_2 = 2(\alpha)$, respectively.

This work was supported by Non-Directed Research Fund, Korea Research Foundation, 1993.

- [1] M. J. Feigenbaum, J. Stat. Phys. **19**, 25 (1978); **21**, 669 (1979).
- [2] B. Derrida, A. Gervois, and Y. Pomeau, J. Phys. A **12**, 269 (1979).
- [3] B. Hu and I. I. Satija, Phys. Lett. A **98**, 143 (1982).
- [4] J. P. van der Weele, H. W. Capel, and R. Kluiving, Physica A **145**, 425 (1987).
- [5] S. Kuznetsov, Radiophys. Quantum Electron. **28**, 681

- (1985).
- [6] H. Kook, F. H. Ling, and G. Schmidt, Phys. Rev. A **43**, 2700 (1991).
- [7] S.-Y. Kim and H. Kook, Phys. Rev. A **46**, R4467 (1992).
- [8] S.-Y. Kim and H. Kook, Phys. Lett. A **178**, 258 (1993).
- [9] S.-Y. Kim and H. Kook, Phys. Rev. E **48**, 785 (1993).
- [10] See Appendix A of Ref. [9].
- [11] J.-m. Mao and B. Hu, J. Stat. Phys. **46**, 111 (1987).