

Diffusive transport in a one-dimensional disordered potential involving correlations

Cécile Monthus*

Division de Physique Théorique, Unité de recherche des Universités Paris 11 et Paris 6 associée au CNRS,

Institut de Physique Nucléaire, Bâtiment 100, 91406 Orsay, France

and L.P.T.P.E, Université Paris 6, 4 Place Jussieu, 75252 Paris, France

(Received 31 March 1995)

This paper deals with transport properties of one-dimensional Brownian diffusion under the influence of a correlated quenched random force, distributed as a two-level Poisson process. We find in particular that large time scaling laws of the position of the Brownian particle are analogous to the uncorrelated case. We also discuss the probability distribution of the stationary flux going through a sample between two prescribed concentrations, which differs significantly from the uncorrelated case.

PACS number(s): 05.60.+w, 05.40.+j

Disorder significantly influences diffusive transport phenomena [1–3]. Indeed, strong enough disorder does not induce a simple renormalization for transport coefficients of the corresponding pure system, but usually generates “anomalous diffusion” effects. The case of spatially uncorrelated disordered media is now pretty well understood. The effects of spatial correlations on the dynamics have been discussed for the directed random walk problem [4]. More recently, the response to an external applied force has been studied for the one-dimensional case of a Brownian particle diffusing under the influence of a quenched random force $\{F(x)\}$ [5]. For a sample characterized by some particular realization of the stochastic process $\{F(x)\}$, the diffusion is defined by the following Fokker-Planck equation for the probability density $P(x, t|x_0, 0)$

$$\frac{\partial P}{\partial t} = D_0 \frac{\partial}{\partial x} \left[\frac{\partial P}{\partial x} - \beta F(x) P \right]. \quad (1)$$

In this paper, we analyze in detail transport properties of this model in the case where $\{F(x)\}$ is distributed as a two-level Poisson process.

Let us define more precisely the model of disorder we consider and some of its properties. We assume that the quenched random force $F(x)$ takes alternatively a positive value $\phi_0 > 0$ and a negative value $-\phi_1 < 0$ on intervals whose lengths are independent random variables distributed according to the following probability densities, respectively,

$$\begin{aligned} f_0(l) &= \theta(l) n_0 e^{-n_0 l}, \\ f_1(h) &= \theta(h) n_1 e^{-n_1 h}. \end{aligned} \quad (2)$$

The parameters $1/n_0$ and $1/n_1$ are, respectively, the mean length of intervals $\{F(x) = \phi_0\}$ and the mean length

of intervals $\{F(x) = -\phi_1\}$. This choice of exponential distributions for f_0 and f_1 is, in fact, the only one that makes the process $\{F(x)\}$ Markovian. This property enables us to write differential equations for the probability $p_0(x)$ to have $F(x) = \phi_0$ and the probability $p_1(x) = 1 - p_0(x)$ to have $F(x) = -\phi_1$.

$$\begin{aligned} \frac{\partial p_0}{\partial x} &= -n_0 p_0 + n_1 p_1 = n_1 - (n_0 + n_1) p_0, \\ \frac{\partial p_1}{\partial x} &= -n_1 p_1 + n_0 p_0 = n_0 - (n_0 + n_1) p_1. \end{aligned} \quad (3)$$

The mean value F_0 and the two point correlation function $G(x)$ of the process $\{F(x)\}$ read

$$\begin{aligned} F_0 &\equiv \langle \phi \rangle = \phi_0 \frac{n_1}{n_0 + n_1} - \phi_1 \frac{n_0}{n_0 + n_1}, \\ G(x) &\equiv \langle \phi(x) \phi(0) \rangle - \langle \phi \rangle^2 \\ &= \frac{n_0 n_1}{(n_0 + n_1)^2} (\phi_0 + \phi_1)^2 e^{-(n_0 + n_1)|x|}. \end{aligned} \quad (4)$$

The corresponding random potential $U(x) = -\int^x F(y) dy$ seen by the Brownian particle presents an alternance of positive and negative slopes of random lengths as sketched in Fig. 1.

The fundamental random variable associated with classical diffusion under the action of a quenched random force presenting a strictly positive mean $\langle F(x) \rangle \equiv F_0 > 0$, is the exponential functional of the random potential $U(x)$

$$\tau_\infty = \int_0^\infty dx e^{\beta U(x)} = \int_0^\infty dx e^{-\beta \int_0^x F(y) dy}. \quad (5)$$

Indeed, the probability distribution of the functional τ_∞ determines the large time anomalous behavior of the Brownian particle position [3]. In particular, the velocity defined for each sample as

$$V = \lim_{t \rightarrow \infty} \frac{d}{dt} \int_{-\infty}^{+\infty} dx x P(x, t|x_0, 0) \quad (6)$$

is a self-averaging quantity [6] (for any homogeneous ran-

*Electronic address: monthus@ipncls.in2p3.fr




FIG. 1. Example of the random potential $U(x)$ seen by the Brownian particle when the random force $\{F\}$ is a two-level Poisson process.

dom potential presenting only short-range correlations) inversely proportional to the first moment of τ_∞ [3]

$$V = \frac{D_0}{\langle \tau_\infty \rangle}. \quad (7)$$

When the quenched random force is distributed with the Gaussian measure

$$\mathcal{D}F(x) e^{-\frac{1}{2\sigma} \int dx [F(x) - F_0]^2} \quad (8)$$

the probability distribution $\mathcal{P}_\infty(\tau)$ of the functional τ_∞ reads in terms of $\alpha = \sigma\beta^2/2$ and $\mu = 2F_0/\beta\sigma > 0$ [3]

$$\mathcal{P}_\infty(\tau) = \frac{\alpha}{\Gamma(\mu)} \left[\frac{1}{\alpha\tau} \right]^{1+\mu} e^{-1/\alpha\tau} \propto_{\tau \rightarrow \infty} \frac{1}{\tau^{1+\mu}}. \quad (9)$$

The algebraic decay at large τ explains all the dynamical phase transitions between different anomalous behaviors known for this model [3]. In particular, Eq. (7) implies that the value $\mu = 1$ separates a phase of vanishing velocity $V = 0$ for $0 < \mu < 1$, and a phase of finite velocity $V > 0$ for $\mu > 1$. Another interesting physical quantity is the stationary current J_N which goes through a disordered sample of length N between a fixed concentration P_0 and a trap described by the boundary condition $P_N = 0$ [7,8]. In the limit $N \rightarrow \infty$, the stationary flux J_∞ is simply a random variable inversely proportional to the functional τ_∞

$$J_\infty = \frac{D_0 P_0}{\tau_\infty}. \quad (10)$$

Note that unlike the velocity V , this flux is not a self-averaging quantity, but must be described by its full probability distribution.

It is very convenient to introduce the more general functional

$$\begin{aligned} \tau(x, b) &= \int_x^b dy e^{\beta[U(y) - U(x)]} \\ &= \int_x^b dy e^{-\beta \int_x^y F(u) du} \end{aligned} \quad (11)$$

and to consider the random variable τ_∞ as the limit of this process as $x \rightarrow -\infty$

$$\tau_\infty = \lim_{x \rightarrow -\infty} \tau(x, b). \quad (12)$$

The evolution of the functional $\tau(x, b)$ is governed by the stochastic differential equation

$$\frac{\partial \tau}{\partial x} = \beta F(x) \tau(x, b) - 1. \quad (13)$$

The stochastic term $F(x)$ appears multiplicatively, so that the fluctuations of the random force are coupled to the values taken by the random process $\tau(x, b)$ itself. When the random force $\{F\}$ is distributed as a white noise (8), the multiplicative stochastic process $\tau(x, b)$ can be, in fact, related to Brownian motion on a surface of constant negative curvature [9].

We now compute the probability distribution of the functional τ_∞ when the quenched random force $\{F(x)\}$ is a two-level Poisson process, and compare it with the result (9) for the white noise case. The random variable τ_∞ only exists if the force mean value F_0 is strictly positive

$$(n_0 + n_1)F_0 = \phi_0 n_1 - \phi_1 n_0 > 0 \quad \text{that is} \quad \frac{n_1}{\phi_1} - \frac{n_0}{\phi_0} > 0 \quad (14)$$

that we assume from now on. Note that the random variable $\tau(x, b)$ remains confined in an interval $[\tau_{\min}(x, b), \tau_{\max}(x, b)]$ depending on the length $(b - x)$

$$\begin{aligned} \tau_{\min}(x, b) &= \int_x^b dy e^{-\beta\phi_0(y-x)} = \frac{1 - e^{-\beta\phi_0(b-x)}}{\beta\phi_0}, \\ \tau_{\max}(x, b) &= \int_x^b dy e^{+\beta\phi_1(y-x)} = \frac{e^{\beta\phi_1(b-x)} - 1}{\beta\phi_1}. \end{aligned} \quad (15)$$

This is very different from the white noise case, where the random force is not bounded. Even in the limit where the interval length $(b - x)$ tends to infinity, the support of the random variable τ_∞ is not $[0, +\infty]$, but $[1/\beta\phi_0, +\infty]$.

The two-level Poisson process $\{F\}$ is Markovian; so according to the local evolution equation (13), the coupled process $\{F(x), \tau(x)\}$ is still Markovian. Let us define the joint laws

$$P_0(\tau, x) d\tau = \text{Prob} \{F(x) = \phi_0 \text{ and } \tau(x, b) \in [\tau, \tau + d\tau]\}, \quad (16)$$

$$P_1(\tau, x) d\tau = \text{Prob} \{F(x) = -\phi_1 \text{ and } \tau(x, b) \in [\tau, \tau + d\tau]\}.$$

They evolve according to the two coupled Master equations

$$\begin{aligned} -\frac{\partial P_0}{\partial x} &= -\frac{\partial}{\partial \tau} [(1 - \beta\phi_0\tau)P_0] - n_0 P_0 + n_1 P_1, \\ -\frac{\partial P_1}{\partial x} &= -\frac{\partial}{\partial \tau} [(1 + \beta\phi_1\tau)P_1] + n_0 P_0 - n_1 P_1. \end{aligned} \quad (17)$$

Stationary distributions $P_0(\tau)$ and $P_1(\tau)$ in the limit $x \rightarrow -\infty$ are, therefore, the solutions of the system

$$\begin{aligned} \frac{d}{d\tau} [(\beta\phi_0\tau - 1)P_0] - n_0 P_0 + n_1 P_1 &= 0, \\ \frac{d}{d\tau} [(\beta\phi_1\tau + 1)P_1] - n_0 P_0 + n_1 P_1 &= 0, \end{aligned} \quad (18)$$

respectively, normalized on the interval $[1/\beta\phi_0, +\infty]$ by

$$\int_{1/\beta\phi_0}^{\infty} P_0(\tau) d\tau = \frac{n_1}{n_0+n_1}$$

and

$$\int_{1/\beta\phi_0}^{\infty} P_1(\tau) d\tau = \frac{n_0}{n_0+n_1}. \quad (19)$$

It is convenient to set $\tau_0 = 1/\beta\phi_0$, $\tau_1 = 1/\beta\phi_1$, $\nu_0 = n_0\tau_0$, $\nu_1 = n_1\tau_1$, and

$$\nu = \nu_1 - \nu_0 = \frac{1}{\beta} \left[\frac{n_1}{\phi_1} - \frac{n_0}{\phi_0} \right] > 0 \quad (20)$$

which is strictly positive according to the hypothesis $F_0 > 0$ (14). The solutions of (18) and (19) read, using Heaviside function θ ,

$$P_0(\tau) = A \theta(\tau - \tau_0) \frac{(\tau - \tau_0)^{\nu_0 - 1}}{(\tau + \tau_1)^{\nu_1}}$$

where

$$A = \frac{n_1}{n_0+n_1} \frac{\Gamma(\nu_1)}{\Gamma(\nu_0)\Gamma(\nu_1-\nu_0)} (\tau_0 + \tau_1)^{\nu_1 - \nu_0},$$

and

$$P_1(\tau) = B \theta(\tau - \tau_0) \frac{(\tau - \tau_0)^{\nu_0}}{(\tau + \tau_1)^{\nu_1 + 1}}, \quad (21)$$

where

$$B = \frac{n_0}{n_0+n_1} \frac{\Gamma(\nu_1+1)}{\Gamma(\nu_0+1)\Gamma(\nu_1-\nu_0)} (\tau_0 + \tau_1)^{\nu_1 - \nu_0}.$$

Moments of order k of these distributions diverge as soon as $k \geq \nu = (\nu_1 - \nu_0)$. They read otherwise in polynomial forms as

$$\int_0^{\infty} d\tau \tau^k P_0(\tau) = \frac{n_1}{n_0+n_1} \times \tau_0^k F \left[-k, \nu_0, 1 + \nu_0 - \nu_1; 1 + \frac{\tau_1}{\tau_0} \right],$$

$$\int_0^{\infty} d\tau \tau^k P_1(\tau) = \frac{n_0}{n_0+n_1} \times \tau_0^k F \left[-k, \nu_0 + 1, 1 + \nu_0 - \nu_1; 1 + \frac{\tau_1}{\tau_0} \right], \quad (22)$$

where $F(a, b, c, z)$ denotes the hypergeometric function. The velocity can be directly deduced from the first moment of τ_{∞} (7)

$$\frac{1}{V} = \frac{1}{D_0} \langle \tau_{\infty} \rangle = \frac{1}{D_0} \int_0^{\infty} d\tau \tau [P_0(\tau) + P_1(\tau)]. \quad (23)$$

There exists, therefore, a dynamical phase transition at $\nu = (\nu_1 - \nu_0) = 1$.

$$V = 0 \text{ if } \nu \leq 1,$$

$$V = \frac{D_0(\nu - 1)}{\frac{n_1}{n_0+n_1} [\tau_0(\nu_1 - 1) + \tau_1\nu_0] + \frac{n_0}{n_0+n_1} [\tau_0\nu_1 + \tau_1(\nu_0 + 1)]} \text{ if } \nu \geq 1. \quad (24)$$

More generally, the algebraic decay of the distributions $P_{0,1}(\tau)$ in the limit $\tau \rightarrow \infty$

$$P_{0,1}(\tau) \underset{\tau \rightarrow \infty}{\propto} \frac{1}{\tau^{1+\nu}}$$

with

$$\nu = \nu_1 - \nu_0 = \frac{1}{\beta} \left[\frac{n_1}{\phi_1} - \frac{n_0}{\phi_0} \right] \quad (25)$$

will generate the same succession of anomalous diffusion behaviors for the cumulants of the Brownian particle position as a function of the parameter ν , as the one that appears in the white noise case as a function of the parameter μ [3]. In particular, the thermal average of the position of the particle will grow linearly in time only in the finite velocity phase

$$\overline{x(t)} \equiv \int_{-\infty}^{+\infty} dx x P(x, t | x_0, 0) \underset{t \rightarrow \infty}{\propto} Vt \text{ if } \nu > 1. \quad (26)$$

However, in the vanishing velocity phase, the thermal average of the position will grow slower than linearly, as a power of time with the exponent ν which depends continuously on the parameters of the disorder

$$\overline{x(t)} \equiv \int_{-\infty}^{+\infty} dx x P(x, t | x_0, 0) \underset{t \rightarrow \infty}{\propto} t^{\nu} \text{ if } 0 < \nu < 1. \quad (27)$$

Let us now consider the stationary flux J_{∞} going through a sample between two concentrations $P_0 > 0$ and $P_N = 0$ in the limit $N \rightarrow \infty$. The change of variable (10) gives immediately the two joint laws $\mathcal{P}_0(J)$ and $\mathcal{P}_1(J)$ from result (21), with the notations $J_0 = D_0 P_0 / \tau_0$ and $J_1 = D_0 P_0 / \tau_1$

$$\mathcal{P}_0(J) = \mathcal{A} \theta(J) \theta(J_0 - J) J^{\nu_1 - \nu_0 - 1} \frac{(J_0 - J)^{\nu_0 - 1}}{(J_1 + J)^{\nu_1}},$$

$$\mathcal{P}_1(J) = \mathcal{B} \theta(J) \theta(J_0 - J) J^{\nu_1 - \nu_0 - 1} \frac{(J_0 - J)^{\nu_0}}{(J_1 + J)^{\nu_1 + 1}}, \quad (28)$$




FIG. 2. Examples of flux probability distributions $\mathcal{P}_0(J)$ (straight line) and $\mathcal{P}_1(J)$ (dashed line) in the vanishing velocity phase $0 < \nu < 1$ for (a) $\nu_0 < 1$; (b) $\nu_0 > 1$.




FIG. 3. Examples of flux probability distributions $\mathcal{P}_0(J)$ (straight line) and $\mathcal{P}_1(J)$ (dashed line) in the finite velocity phase $\nu > 1$ for (a) $\nu_0 < 1$; (b) $\nu_0 > 1$.

where \mathcal{A} and \mathcal{B} are two normalization constants. These two distributions for the flux are concentrated on the bounded interval $[0, J_0]$. The transition at $\nu \equiv \nu_1 - \nu_0 = 1$ for the self-averaging velocity (24) corresponds for the flux probability distributions to a transition of the behavior in the limit $J \rightarrow 0$

$$\begin{aligned} \mathcal{P}_{0,1}(J) &\underset{J \rightarrow 0}{\propto} J^{\nu-1} \longrightarrow +\infty \text{ if } \nu < 1, \\ \mathcal{P}_{0,1}(J) &\underset{J \rightarrow 0}{\propto} J^{\nu-1} \longrightarrow 0 \text{ if } \nu > 1. \end{aligned} \quad (29)$$

The vanishing velocity phase $\nu = \nu_1 - \nu_0 < 1$ is characterized by the divergence of \mathcal{P}_0 and \mathcal{P}_1 at the origin $J \rightarrow 0$. On the contrary in the finite velocity phase $\nu > 1$, the probability distributions of the flux vanish at the origin.

Some curves $\mathcal{P}_{0,1}(J)$ are drawn in Figs. 2 and 3 for different values of the parameters. They illustrate, in particular, the transitions at $\nu = 1$ we just mentioned (29) and the transition at $\nu_0 = 1$ for the behavior of $\mathcal{P}_0(J)$ as $J \rightarrow J_0$. We refer to the figures in [8] for comparison with the uncorrelated case.

In this paper, we have analyzed some properties of anomalous diffusion in a random medium described by a quenched random force $\{F\}$ distributed as a two-level Poisson process. In particular, we showed that there exists a dimensionless parameter ν , which governs the

asymptotic behaviors of the probability distributions $\mathcal{P}_{0,1}(\tau)$ in the limit $\tau \rightarrow \infty$. As a result, this parameter ν also controls the dynamical phase transition for the velocity and the behaviors of the probability distributions $\mathcal{P}_{0,1}(J)$ of the flux in the limit $J \rightarrow 0$. All this is qualitatively the same as what is known in the white noise case. However, the probability distributions for the random variable τ_∞ and for the stationary flux J_∞ are very different from the white noise case outside the asymptotic regimes $\tau \rightarrow \infty$ and $J \rightarrow 0$. Indeed, they present restricted supports, as a consequence of the bounded character of the random force, and they are given in terms of only rational functions and no exponential.

More generally, the two-level Poisson process we considered is a technically very convenient disorder model, since it allows analytical studies despite the presence of correlations. It has already been used in various contexts, such as one-dimensional quantum localization [10,11], nonlinear systems coupled to a random environment [12], and noise-induced perturbations on Josephson junctions [13]. The limitation to two-level processes simplifies computations, but the approach can be generalized to any finite number of levels for the random process [14].

I wish to thank Alain Comtet for many helpful discussions and for his remarks on the manuscript.

- [1] F. Solomon, *Ann. Probab.* **3**, 1 (1975); H. Kesten, M. Koslov, and F. Spitzer, *Compos. Math.* **30**, 145 (1975); Y. A. G. Sinai, *Theor. Probab. Appl.* **XXVII**, 256 (1982); B. Derrida and Y. Pomeau, *Phys. Rev. Lett.* **48**, 627 (1982); B. Derrida, *J. Stat. Phys.* **31**, 433 (1983).
- [2] J. W. Haus and K. W. Kehr, *Phys. Rep.* **150**, 263 (1987); S. Havlin and D. Ben Avraham, *Adv. Phys.* **36**, 695 (1987).
- [3] J. P. Bouchaud and A. Georges, *Phys. Rep.* **195**, 127 (1990); J. P. Bouchaud, A. Comtet, A. Georges, and P. Le Doussal, *Ann. Phys.* **201**, 285 (1990); A. Georges, Thèse d'état de l'Université Paris 11, 1988.
- [4] C. Aslangul, N. Potier, P. Chvosta, and D. Saint-James, *Phys. Rev. E* **47**, 1610 (1993).
- [5] P. Le Doussal and V. Vinokur, Report No. cond-mat/9501131; S. Scheidl, Report No. cond-mat/9501121.
- [6] C. Aslangul, N. Potier, and D. Saint-James, *J. Phys. (Paris) Colloq.* **50**, 899 (1989).
- [7] S. F. Burlatsky, G. H. Oshanin, A. V. Mogutov, and M. Moreau, *Phys. Rev. A* **45**, 6955 (1992); G. Oshanin, A. Mogutov, and M. Moreau, *J. Stat. Phys.* **73**, 379 (1993); G. Oshanin, S. F. Burlatsky, M. Moreau, and B. Gaveau, *Chem. Phys.* **177**, 803 (1993).
- [8] C. Monthus and A. Comtet, *J. Phys. I (France)* **4**, 635 (1994).
- [9] A. Comtet and C. Monthus, Report No. cond-mat/9506023.
- [10] M. M. Benderskii and L. A. Pastur, *Zh. Eksp. Teor. Fiz.* **57**, 284 (1969) [*Sov. Phys. JETP* **30**, 158 (1970)]; L. A. Pastur and E. P. Fel'dman, *ibid.* **40**, 241 (1974); I. M. Lifshits, S. A. Gredeskul, and L. A. Pastur, *Introduction to the Theory of Disordered Systems* (Wiley, New York, 1987).
- [11] A. Comtet, J. Desbois, and C. Monthus, *Ann. Phys.* **239**, 312 (1995).
- [12] W. Horsthemke and R. Lefever, *Noise-induced Transitions: Theory and Applications in Physics, Chemistry and Biology* (Springer-Verlag, Berlin, 1984).
- [13] G. H. Weiss and M. Gitterman, *J. Stat. Phys.* **70**, 93 (1993).
- [14] P. Erdős and Z. Domanski, in *Analogies in Optics and Microelectronics*, edited by W. Van Haeringen and D. Lenstra (Kluwer Academic, Dordrecht, 1990), p. 49.