
ERRATA

Erratum: Effects of changes in the chain conformation on the kinetics of order-disorder transitions in block copolymer melts
[Phys. Rev. E 56, 3240 (1997)]

Toshihiro Kawakatsu

[S1063-651X(98)11705-1]

PACS number(s): 82.20.Wt, 64.60.Ht, 83.10.Nn, 99.10.+g

Equation (14) should be replaced by

$$\begin{aligned} \Lambda(\mathbf{q}, \mathbf{q}') = & \int d\mathbf{q}'' \int d\mathbf{q}''' [\bar{P}_{AA}(\mathbf{q}, \mathbf{q}'') \bar{P}_T^{-1}(\mathbf{q}'', \mathbf{q}''') \bar{P}_{BB}(\mathbf{q}''', \mathbf{q}') - \bar{P}_{AB}(\mathbf{q}, \mathbf{q}'') \bar{P}_T^{-1}(\mathbf{q}'', \mathbf{q}''') \bar{P}_{BA}(\mathbf{q}''', \mathbf{q}')] \\ & + \bar{P}_{AA}(\mathbf{q}, \mathbf{q}'') \bar{P}_T^{-1}(\mathbf{q}'', \mathbf{q}''') \bar{P}_{AB}(\mathbf{q}''', \mathbf{q}') - \bar{P}_{AB}(\mathbf{q}, \mathbf{q}'') \bar{P}_T^{-1}(\mathbf{q}'', \mathbf{q}''') \bar{P}_{AA}(\mathbf{q}''', \mathbf{q}')], \end{aligned} \quad (1)$$

where $\bar{P}_T^{-1}(\mathbf{q}, \mathbf{q}')$ is defined by the following relation:

$$\int d\mathbf{q}'' \bar{P}_T^{-1}(\mathbf{q}, \mathbf{q}'') [\bar{P}_{AA}(\mathbf{q}'', \mathbf{q}') + \bar{P}_{AB}(\mathbf{q}'', \mathbf{q}') + \bar{P}_{BA}(\mathbf{q}'', \mathbf{q}') + \bar{P}_{BB}(\mathbf{q}'', \mathbf{q}')] = \delta(\mathbf{q} - \mathbf{q}'). \quad (2)$$

As the actual calculations are based on the above expressions, Eqs. (1) and (2), the final results presented in the paper are not affected by this change.