

Field theoretic approach to the counting problem of Hamiltonian cycles of graphs

Saburo Higuchi*

Department of Pure and Applied Sciences, The University of Tokyo, Komaba, Meguro, Tokyo 153-8902, Japan

(Received 14 November 1997; revised manuscript received 3 February 1998)

A field theoretic representation of the number of Hamiltonian cycles of graphs is studied. By integrating out quadratic fluctuations around the saddle point, one obtains an estimate of a number which reflects characteristics of graphs well. The accuracy of the estimate is verified by applying it to two-dimensional square lattices with various boundary conditions. This is an example of how to extract meaningful information from the quadratic approximation of the field theory representation. [S1063-651X(98)14306-4]

PACS number(s): 05.50.+q, 02.10.Eb, 05.20.-y, 82.35.+t

I. INTRODUCTION

Let $G=(V,E)$ be a graph with a set of vertices $V=\{r_j\}$ and edges $E=\{e_k\}$. A Hamiltonian cycle of a graph is a closed path which visits each of the vertices once and only once. I denote the number of all the Hamiltonian cycles of a graph G by $H(G)$:

$$H(G) = \sum_{\text{Hamiltonian cycle on } G} 1. \tag{1}$$

See Fig. 1 for examples.

Hamiltonian cycles have often been used to model collapsed polymer globules [1]. The quantity $H(G)$ corresponds to the entropy of a polymer system in G in a collapsed but disordered phase. One can model even more realistic polymers by introducing a weight that depends on the shape of cycles in Eq. (1). The polymer melting problem is studied by taking into account the bending energy with a weight which depends on the number of turns on the cycle [2]. In Ref. [3], the protein folding problem is studied by incorporating the van der Waals potential as well as the bending energy in model (1).

For homogeneous graphs (lattices) with N vertices, one expects that $H(G)$ behaves like

$$H(G) \rightarrow C(G)N^{\gamma-1}\omega^N \quad (N \rightarrow \infty), \tag{2}$$

where ω is defined by

$$\ln \omega = \lim_{N \rightarrow \infty} \frac{1}{N} \ln H(G). \tag{3}$$

The quantity ω is supposed to be a universal bulk quantity, whereas $C(G)$ and γ depend on the detail of graphs, e.g., boundary conditions [4].

A field theory representation for Eq. (1) for arbitrary graphs was introduced in Ref. [5], and has been used to study the extended models [2,3]. For homogeneous graphs with a number of vertices N and the coordination number q , the saddle point approximation of the representation yields

$$H(G) \approx \left(\frac{q}{e}\right)^N, \tag{4}$$

or $\omega \approx q/e$. This approximation has been proved to be good in many examples [6]. For a square lattice, $q/e=4/e=1.4715\dots$ is quite near to the exact value $\omega \approx 1.473$ estimated by direct enumeration [7] and other methods [4,8,9].

In the saddle point approximation, however, graphs with identical (q,N) are not distinguished. Indeed there is a variety of graphs which have (q,N) in common. Given a graph with a pair (q,N) , it is often possible to change its boundary conditions to modify the ‘‘topology’’ and ‘‘moduli’’ of the graph, keeping the pair. Figure 1 shows examples. Moreover, there are graphs which have identical (q,N) but have distinct local structures, e.g., the two-dimensional (2D) triangular lattice and the 3D cubic lattice.

In this paper, I go beyond the saddle point approximation. I work out the quadratic approximation and find an estimate for $H(G)$ whose G dependence is not merely through (q,N) but is more sensible. To demonstrate the validity of the approximation, I examine 2D square lattices with a variety of boundary conditions and aspect ratios. It is also examined if the estimate $\omega \approx q/e$ is improved.

II. FIELD-THEORETIC REPRESENTATION

The problem of calculating $H(G)$ is mapped into one in a lattice field theory on G [5]. This is done by introducing an




FIG. 1. Examples of graphs and the Hamiltonian cycles on them. One does not distinguish the base points and the directions of a cycle. (a) $H(G)=1$ for this graph. (b) $P(3,4)$, the 2D square lattice with the periodic boundary condition. A Hamiltonian cycle is drawn in the thick line. For this graph, $(q,N)=(4,12)$. (c) $SP(3,4)$, the 2D square lattice with the skew-periodic boundary in the horizontal direction. It is locally isomorphic to $P(3,4)$, but the boundary condition makes it distinct globally.

*Electronic address: hig@rice.c.u-tokyo.ac.jp

$O(n)$ lattice field $\vec{\phi}(r) = (\phi_1(r), \dots, \phi_n(r))$ “living” on $V = \{r_j\}$, with an action

$$S[\vec{\phi}(r)] = \frac{1}{2} \sum_{r, r' \in V, 1 \leq j \leq n} \phi_j(r) (i\Delta^{-1} + \epsilon)_{rr'} \phi_j(r'). \quad (5)$$

The $N \times N$ matrix Δ is the adjacency matrix [10] of the graph G :

$$\Delta_{rr'} = \begin{cases} 1 & \text{if } r, r' \in V \text{ is connected by an } e \in E \\ 0 & \text{otherwise,} \end{cases} \quad (6)$$

and an infinitesimal parameter $\epsilon > 0$ is introduced for convergence.

The integer $H(G)$ is related to a $2N$ point function by

$$H(G) = \lim_{n \rightarrow 0} \lim_{\epsilon \rightarrow +0} \frac{1}{n} \left| \frac{Z_1}{Z_0} \right|, \quad (7)$$

$$Z_0 = \int D\vec{\phi} e^{-S[\vec{\phi}(r)]}, \quad (8)$$

$$Z_1 = \int D\vec{\phi} e^{-S[\vec{\phi}(r)]} \prod_{r \in V} \frac{\vec{\phi}^2(r)}{2}, \quad (9)$$

where $D\vec{\phi} = \prod_{r \in V, 1 \leq j \leq n} d\phi_j(r)$. Equation (7) holds for arbitrary G with $N > 2$, because each term in the diagrammatic expansion corresponds to a Hamiltonian cycle.

III. APPROXIMATION

I first evaluate Eq. (7) by the saddle point method. I concentrate on Z_1 , since $Z_0 \rightarrow 1$ as $n \rightarrow 0$. When a graph G is homogeneous, there are mean field saddle points which are degenerate on

$$\{\vec{\phi} | \vec{\phi}^2(r) \equiv -2iq\epsilon\} \approx \frac{O(n)}{O(n-1)}, \quad (10)$$

where

$$\frac{1}{q\epsilon} := \frac{1}{q} - i\epsilon. \quad (11)$$

This yields estimate (4).

I then consider fluctuations around the saddle point. It can easily be seen that there are zero modes corresponding to the global $O(n)/O(n-1)$ symmetry and sublattice symmetries [5]. Therefore I am led to introduce a gauge fix condition

$$\sum_{r \in V} \phi_j(r) = 0 \quad (2 \leq j \leq n). \quad (12)$$

By the standard Fadeev-Popov method, I have

$$\begin{aligned} Z_1 &= \int D\phi e^{-\frac{1}{2} \sum_{r, r' \in V, 1 \leq j \leq n} \phi_j(r) (i\Delta^{-1} + \epsilon)_{rr'} \phi_j(r')} \\ &\times \left| \sum_{r \in V} \phi_1(r) \right|^{n-1} \frac{\pi^{n/2}}{\Gamma\left(\frac{n}{2}\right)} \prod_{2 \leq j \leq n} \delta\left(\sum_{r \in V} \phi_j(r)\right) \\ &\times \prod_{r \in V} \frac{\vec{\phi}^2(r)}{2}. \end{aligned} \quad (13)$$

The factor $|\sum_r \phi_1(r)|^{n-1} \pi^{n/2} / \Gamma(n/2)$ is the Fadeev-Popov determinant multiplied by the volume of the gauge orbit. Actually, this is nothing but the Jacobian for the n -dimensional radial coordinate for constant modes.

Expanding the field $\vec{\phi}$ around the saddle point as

$$\phi_j(r) = \sqrt{-2iq\epsilon I_{N,n}} \delta_{j1} + \psi_j(r), \quad (14)$$

where

$$I_{N,n} = 1 + \frac{n-1}{2N}, \quad (15)$$

one obtains

$$\begin{aligned} Z_1 &= \left(\frac{q\epsilon I_{N,n}}{ie} \right)^{NI_{N,n}} 2^{(n-1)/2} N^{n-1} \frac{\pi^{n/2}}{\Gamma\left(\frac{n}{2}\right)} \\ &\times \int D\vec{\psi} e^{-\frac{1}{2} \sum_{r, r' \in V} \psi_1(1) A_{rr'}^L(n) \psi_1(r')} \\ &\times e^{-\frac{1}{2} \sum_{r, r' \in V, 2 \leq j \leq n} \psi_j(r) A_{rr'}^T(n) \psi_j(r')} \\ &\times e^{-V_{\text{int}}(\vec{\psi})} \prod_{2 \leq j \leq n} \delta\left(\sum_{r \in V} \psi_j(r)\right), \end{aligned} \quad (16)$$

where $A^L(n)$ and $A^T(n)$ are the inverse propagators of the longitudinal mode ($j=1$) and the transverse modes ($2 \leq j \leq n$), respectively:

$$\begin{aligned} A_{rr'}^L(n) &= i(\Delta^{-1})_{rr'} + \frac{i}{q} I_{N,n}^{-1} \delta_{rr'} + (1 + I_{N,n}^{-1}) \epsilon \delta_{rr'} \\ &+ \frac{1}{N} \frac{i}{q\epsilon} (1 - I_{N,n}^{-1}), \end{aligned} \quad (17)$$

$$A_{rr'}^T(n) = i(\Delta^{-1})_{rr'} - \frac{i}{q} I_{N,n}^{-1} \delta_{rr'} + (1 - I_{N,n}^{-1}) \epsilon \delta_{rr'}. \quad (18)$$

One neglects the interaction terms $V_{\text{int}}(\vec{\psi})$ and performs the Gaussian integrations. Finally, ϵ is sent to $+0$ to yield

$$\begin{aligned} Z_1 &\approx \left(\frac{q I_{N,n}}{ie} \right)^{NI_{N,n}} (2N)^{(n-1)/2} \frac{\pi^{n/2}}{\Gamma\left(\frac{n}{2}\right)} \\ &\times \left[\frac{(2\pi)^{(N-1)/2}}{\det^{1/2} A^T(n)} \right]^{n-1} \frac{(2\pi)^{N/2}}{\det^{1/2} A^L(n)} \times 2. \end{aligned} \quad (19)$$




FIG. 2. The number of Hamiltonian cycles for $P(L_1, L_2)$ for odd L_j as a function of L_2/L_1 , normalized by the saddle point result $(4/e)^{L_1 L_2}$. Plotted are the quadratic approximation to the field theory (solid square: $3 \leq L_1, L_2 \leq 29$; dashed line: the limit $L_1 \times L_2 \rightarrow \infty$), exact results obtained by enumeration (box, $L_1 \times L_2 \leq 39$), and estimates by biased Monte Carlo simulations (solid box, $L_1 \times L_2 \leq 90$).

The last factor 2 corresponds to the residual symmetry $\Sigma_r \phi_1(r) \leftrightarrow -\Sigma_r \phi_1(r)$. The prime on \det means the omission of the eigenvalue for the constant mode.

One notices that the signature of the real part (the last term) of Eq. (18) changes at $n=1$. Thus one has to assume $n > 1$ to derive Eq. (19). Then in Eq. (19) one takes the limit $n \rightarrow 0$ to obtain the final result

$$H(G) \approx \left(\frac{q}{e}\right)^N e^{1/2(I_{N,0})^{N I_{N,0}}} \sqrt{\frac{\pi \det'^{1/2}(A^T(0)\Delta)}{N \det^{1/2}(A^L(0)\Delta)}}. \quad (20)$$

Note that $\det \Delta \det' A^T(0) = q \det'(A^T(0)\Delta)$. Equation (20) is nothing but the estimate I desired to have. The saddle point result $(q/e)^N$ is corrected by the ratio of determinants which contains information of details of the structure of G .

In Ref. [5], it is claimed that the quadratic correction to Eq. (4) vanishes. In the present analysis, the Fadeev-Popov method is worked out to find $\sqrt{\pi/N}$ missing in Ref. [5]. As shown in Sec. IV, the ratio of determinants is not equal to unity and contributes to γ and $C(G)$ nontrivially. Moreover, inclusion of $i = \sqrt{-1}$ and $\epsilon > 0$ in action (5) in the present analysis enables one to discuss the limit of application of the quadratic approximation in Sec. IV.

IV. SQUARE LATTICES

To see how Eq. (20) works, I study the concrete examples $P(L_1, L_2)$ and $SP(L_1, L_2)$ shown in Fig. 1. Both are two-dimensional square lattices with edge lengths L_1 and L_2 . The difference between P and SP lies in boundary conditions. For $P(L_1, L_2)$, the periodic boundary condition is imposed for both two directions. For $SP(L_1, L_2)$, the boundary condition across the edge L_2 is replaced by the skew-periodic one. They are good examples to test Eq. (20). One can switch the boundary condition or vary the aspect ratio L_2/L_1 to make the graph globally distinct while keeping $(q, N) = (4, L_1 \times L_2)$. The saddle point approximation cannot “see” the difference among them, but Eq. (20) has a chance to distinguish them.




FIG. 3. The log plot of the number of Hamiltonian cycles for $SP(L_1, L_2)$ for odd L_j as a function of $\arctan(L_2/L_1)$. Normalized by the saddle point result $(4/e)^{L_1 L_2}$. The plots are the same as in Fig. 2.

Graphs $P(L_1, L_2)$ and $SP(L_1, L_2)$ can be viewed as discrete tori with different moduli parameters. Thus it is also interesting in its own right to determine the asymptotic behaviors of $H(P(L_1, L_2))$ and $H(SP(L_1, L_2))$ in the limit $L_1, L_2 \rightarrow \infty$.

I analytically evaluate Eq. (20) for $P(L_1, L_2)$ and $SP(L_1, L_2)$. In the momentum representation, the determinants become ($\sigma=0$ for P and $\sigma=1$ for SP)

$$\det'(A^T(n)\Delta) = \prod'_{0 \leq n_j \leq L_j - 1} \left\{ 1 - \frac{1}{2I_{N,n}} \left[\cos\left(k_1 + \sigma \frac{k_2}{L_1}\right) + \cos k_2 \right] \right\}, \quad (21)$$

$$\det(A^L(n)\Delta) = 2 \times \prod'_{0 \leq n_j \leq L_j - 1} \left\{ 1 + \frac{1}{2I_{N,n}} \left[\cos\left(k_1 + \sigma \frac{k_2}{L_1}\right) + \cos k_2 \right] \right\}. \quad (22)$$

The constant mode $k_1 = k_2 = 0$ is excluded in Π' . Hereafter, the indices n_j and k_j should be related by $k_j = 2\pi n_j / L_j$.

The formula

$$\prod_{0 \leq m \leq L-1} \left[x^2 - 2x \cos\left(\theta + \frac{2m\pi}{L}\right) + 1 \right] = x^{2L} - 2x^L \cos(L\theta) + 1 \quad (23)$$

enables one to obtain

$$\frac{\det'(A^T(0)\Delta)}{\det(A^L(0)\Delta)} = -2L_1 L_2 I_{L_1 L_2, 0} \prod_{n_2=0}^{L_2-1} \frac{u_-(k_2)}{u_+(k_2 + \pi)}, \quad (24)$$

where

$$u_{\pm}(k) = \begin{cases} y(k)^{L_1} + y(k)^{-L_1} \pm 2 & \text{for } P \\ y(k)^{L_1} + y(k)^{-L_1} - 2 \cos k & \text{for } SP, \end{cases} \quad (25)$$

$$y(k) = 2I_{L_1 L_2, 0} - \cos k + \sqrt{(2I_{L_1 L_2, 0} - \cos k)^2 - 1}. \quad (26)$$

TABLE I. Exact number of Hamiltonian cycles for $L_1 \times L_2$ 2D square lattice with periodic (P) and skew-periodic (SP) boundary conditions. Determined by the direct enumeration.

L_1	L_2	$H(P(L_1, L_2))$	$H(SP(L_1, L_2))$	$H(SP(L_2, L_1))$
3	3	48	55	55
5	3	390	397	866
5	5	23 580	29 001	29 001
7	3	2982	2 989	13 021
7	5	1 045 940	1 108 006	1 820 582
9	3	23 646	23 653	195 157
11	3	196 086	196 093	2 924 373
13	3	1 682 382	1 682 389	43 820 323

Now I specialize to the case where both L_1 and L_2 are odd. In the limit $N=L_1 \times L_2 \rightarrow \infty$ with $R=L_2/L_1$ fixed, it is allowed to approximate Π_{n_2} by $\exp(\int dk_2 \ln)$ after separating quickly oscillating factors. In this limit, assuming $0 < R \leq 1$ for periodic case, I obtain

$$H(P(L_1, L_2)) \approx \left(\frac{q}{e}\right)^N \sqrt{\frac{\pi}{2}} \frac{\sin\left(\frac{1}{2R}\right)^{1/2} \exp\left(\frac{9\pi^2}{16R^2} - \frac{1}{2R}\right)^{1/2}}{\cosh^2\left(\frac{\pi^2}{4R^2} - \frac{1}{2R}\right)^{1/2}}, \quad (27)$$

$$H(SP(L_1, L_2)) \approx \left(\frac{q}{e}\right)^N \sqrt{2\pi} \frac{\left| \sin\left(\frac{1}{2R}\right)^{1/2} \sinh\left(\frac{9\pi^2}{16R^2} - \frac{1}{2R}\right)^{1/2} \right|}{\left| \sinh^2\left(\frac{\pi^2}{4R^2} - \frac{1}{2R}\right)^{1/2} \right|}. \quad (28)$$

One sees that the quadratic approximation predicts the explicit form of $C(G)$ and the fact that $\gamma=1$ in Eq. (2). It is remarkable that the correction is independent of N in the limit. That is, it implies that the estimates for ω are unchanged.

Figures 2 and 3 show the plot of Eqs. (27) and (28) normalized by $(4/e)^{L_1 L_2}$ as functions of L_2/L_1 (solid line). The finite $L_1 L_2$ results (20) for odd L_j such that $3 \leq L_j \leq 29$ are also shown (solid circle). For the purpose of comparison, I plot the exact numbers of Hamiltonian cycles (box) and estimates of them by simulations (solid box) for L_j odd [11].

The exact numbers are determined by direct enumeration by a program in C . Table I shows the result. This program is useful only for small graphs, because the time needed for calculation increases exponentially with $L_1 \times L_2$. The simulation is based on the biased Monte Carlo method with a code in C . The time needed again grows exponentially with

$L_1 \times L_2$ but with a smaller exponent. I have used direct enumeration for $L_1 \times L_2 \leq 39$ and Monte Carlo simulation for $L_1 \times L_2 \leq 90$. The evaluation of Eq. (20) takes time proportional to $L_1 \times L_2$. Therefore, Eq. (20) has an advantage even if it is approximate.

Figures 2 and 3 suggest that the quadratic approximation to the field theory is reliable. The field theory succeeds in predicting that the correction depends almost entirely on L_2/L_1 , and that there is a qualitative difference between P and SP . The L_2/L_1 dependence of the correction generally agrees with the exact result, though there is a slight deviation in the small L_2/L_1 region.

There is a definite discrepancy at $L_2/L_1 = \pi^2/2$ for the skew-periodic case. The quadratic approximation (28) diverges at $L_2/L_1 = \pi^2/2$, as seen in Fig. 3, while the exact result takes finite values and is simply increasing there. Actually, one can argue that $L_2/L_1 \geq \pi^2/2$ is out of range of application of Eq. (20). As I mentioned, I defined the limit $n \rightarrow 0$ as the continuation from $n > 1$ where the Gaussian integral converges. Let us look at the evolution of the spectrum of $(A^{L_1} \Delta)$ on the way from $n > 0$ down to $n = 0$. The eigenvalue for the $k_j = (1 - 1/L_j)\pi$ mode hits zero at some $0 < n_c < 1$ if

$$2I_{L_1 L_2, 0} < \cos\frac{\pi}{L_1 L_2} + \cos\frac{\pi}{L_2}. \quad (29)$$

This condition is equivalent to $L_2/L_1 > \pi^2/2$ in the limit $L_1 \times L_2 \rightarrow \infty$. Thus one suddenly has a zero mode at $n = n_c$, and no symmetry is responsible for it. This suggests that the quadratic approximation breaks down there and that Eq. (28) is not reliable for such values of L_2/L_1 . A careful analysis shows that the periodic case is free from such spurious zero modes.

V. SUMMARY AND DISCUSSIONS

I have found an approximate formula for the number of Hamiltonian cycles of graphs by the quadratic approximation to the field theoretic representation. It has been tested for a 2D square lattice with a variety of boundary conditions. I have obtained a multiplicative correction which is independent of the size $N = \#V$. That is, a dependence on the boundary conditions is obtained, and is shown to be in good agreement with the true behavior.

A natural extension to the present problem is the counting of the number of closed self-avoiding walks that visit fN vertices of a graph, where $0 \leq f \leq 1$ is fixed. In that case, form (2) can still be assumed but $C(G)$, γ , and ω now depend on f . There is a field theoretic representation for $\omega(f)$ called lattice cluster theory [12]. It is not identical with Eq. (7) even for $f = 1$. It is interesting to compare the quadratic approximation of the lattice cluster theory [13,14] with the present result. In the former theory, the estimate for $\omega(f)$ is improved over the saddle point approximation for $f < 1$. For $f = 1$, the estimate for $\omega = \omega(1)$ is unchanged in accord with the present case.

ACKNOWLEDGMENTS

I thank T. Yamaguchi for collaboration in the initial stage of this work. Many helpful conversations with S. Hikami, M. Itakura, K. Minakuchi, R. Pnini, and J. Suzuki are gratefully

acknowledged. This work was partially supported by Grant-in-Aid's for Scientific Research (Grants Nos. 08740196 and 08454106) from the Ministry of Education, Science and Culture, and by CREST from Japanese Science and Technology Corporation.

-
- [1] J. des Cloizeaux and G. Jannik, *Polymers in Solution: Their Modelling and Structure* (Clarendon, Oxford, 1987).
 [2] J. Bascle, T. Garel, and H. Orland, *J. Phys. A* **25**, L1323 (1992).
 [3] S. Doniach, T. Garel, and H. Orland, *J. Chem. Phys.* **105**, 1601 (1996).
 [4] T. G. Schmalz, G. E. Hite, and D. J. Klein, *J. Phys. A* **17**, 445 (1984).
 [5] H. Orland, C. Itzykson, and C. de Dominicis, *J. Phys. (France)* **46**, L353 (1985).
 [6] J. Suzuki, *J. Phys. Soc. Jpn.* **57**, 687 (1988).
 [7] C. J. Camacho and D. Thirumalai, *Phys. Rev. Lett.* **71**, 2505 (1993).
 [8] M. T. Batchelor and C. M. Yung, *J. Phys. A* **27**, 4055 (1994).
 [9] B. Duplantier and H. Saleur, *Nucl. Phys. B* **290**, 291 (1987).
 [10] In the case $\det \Delta = 0$, action (5) is ill-defined. In that case I can

justify Eq. (7) by replacing Δ with, for example,

$$\Delta(\epsilon_0)_{rr'} = \Delta_{rr'} + \epsilon_0 \delta_{rr'},$$

where ϵ_0 is a small number. Then I have $\det(\Delta(\epsilon_0)) \neq 0$, and the field theory yields

$$\langle \phi_j(r) \phi_k(r') \rangle = \delta_{jk} \Delta(\epsilon_0)_{rr'} \quad (\epsilon_0 \rightarrow 0),$$

as desired. So Eq. (7) is valid with the $\epsilon_0 \rightarrow 0$ limit understood. By this regularization, the estimates (4) and (20), which are free from $\Delta^{-1} = \Delta(0)^{-1}$, remain valid.

- [11] For the case with L_j even, the L_2/L_1 dependence does not seem to be the same. Work on this point is in progress.
 [12] K. F. Freed, *J. Phys. A* **18**, 871 (1985).
 [13] A. M. Nemirovsky and M. D. Coutinho-Filho, *J. Stat. Phys.* **53**, 1139 (1988).
 [14] A. M. Nemirovsky and M. D. Coutinho-Filho, *Phys. Rev. A* **39**, 3120 (1989).