

## Optical gap solitons in nonresonant quadratic media

Takeshi Iizuka<sup>1,2</sup> and Yuri S. Kivshar<sup>2</sup>

<sup>1</sup>*Department of Physics, Faculty of Science, Ehime University, Ehime 790-8577, Japan*

<sup>2</sup>*Optical Sciences Centre, Australian National University, Canberra, Australian Capital Territory 0200, Australia*

(Received 30 November 1998)

We demonstrate an important role of the process of optical rectification in the theory of nonlinear wave propagation in quadratically nonlinear [or  $\chi^{(2)}$ ] periodic optical media. We derive a novel physical model for gap solitons in  $\chi^{(2)}$  nonlinear Bragg gratings. [S1063-651X(99)11105-X]

PACS number(s): 41.20.Jb, 42.65.Ky, 42.65.Jx, 42.65.Tg

As has been recently demonstrated, large optical nonlinearities can be generated in noncentrosymmetric media by means of the so-called *cascading effects*, due to parametric wave mixing under the condition of nearly phase-matched second-harmonic generation and other parametric processes [1]. It has been also shown that cascaded nonlinearities can support spatial optical solitons [2] and also different types of gap solitons in periodic Bragg gratings with a quadratic (or  $\chi^{(2)}$ ) nonlinear response [3].

Importantly, when an input electromagnetic wave  $\mathbf{E}$  at frequency  $\omega$  is launched into a noncentrosymmetric material, it generates also a quasistatic electric field (or *dc field*) at frequency zero. This effect is known as *optical rectification*, and it is usually described by a contribution to the medium nonlinear polarization  $P$  of the form  $P_i^0 = \epsilon_0 \chi_{ijk}(0; \omega, -\omega) E_j(\omega) E_k^*(\omega)$ , where  $\chi_{ijk}(0; \omega, -\omega)$  is the nonlinear optical susceptibility describing optical rectification [4]. Such an induced dc field changes a refractive index via the linear electro-optic effect. As has been recently shown by Bosshard *et al.* [5], both theoretically and experimentally, the combined processes of optical rectification and the linear electro-optic effect lead to an *additional, nonresonant* contribution into an effective nonlinear refractive index of noncentrosymmetric materials due to cascading processes.

The effect of optical rectification is *usually neglected* in the theory of quadratic solitons because the equation for the dc field can be integrated explicitly, leading to a nonresonant contribution into the effective cubic nonlinearity of the nonlinear Schrödinger (NLS) equation derived by means of the asymptotic technique in the approximation of cascaded nonlinearities (see, e.g., Ref. [6]). However, for the propagation of spatio-temporal multidimensional optical pulses in nonresonant quadratic media, such a reduction is no longer possible and, as a result, the multidimensional NLS equation becomes coupled to a dc field [7], similar to the integrable case of the Dawey-Stewartson equation [8].

In this paper we show that the physical situation is qualitatively different for periodic quadratically nonlinear optical media. We demonstrate that coupling between the forward and backward waves in one-dimensional *shallow Bragg gratings* with a quadratic nonlinearity is accompanied by a coupling to the induced dc field that appears within the same approximation and cannot be eliminated by integration. This effect has been overseen previously, but it leads to a novel physical model for gap solitons in quadratic media which we introduce and analyze here.

We consider propagation of an optical pulse in a periodic medium with a quadratic  $\chi^{(2)}$  nonlinear response. To derive the coupled-mode equations for the wave envelope, we start from Maxwell's equation,

$$c^2 \nabla^2 E - \frac{\partial^2}{\partial t^2} [\hat{\epsilon}(z, i\partial_t) + \chi^{(2)} E] E = 0, \quad (1)$$

where  $\nabla^2$  stands for the Laplacian,  $c$  is the speed of light in vacuum,  $E$  is the  $x$  element of the electric field,  $\mathbf{E} = E(z, t) \mathbf{e}_x$ , and the quadratic nonlinearity is represented by a tensor element  $\chi^{(2)} = \chi_{xxx}^{(2)}$ . We assume that  $\hat{\epsilon}(z, \omega)$  is a periodic function of  $z$ , so it can be presented in a general form as a Fourier series,

$$\hat{\epsilon}(z, \omega) = \epsilon(\omega) \left( 1 + \sum_{j=1}^{\infty} \epsilon_j e^{2ikz} + \sum_{j=1}^{\infty} \epsilon_j^* e^{-2ikz} \right), \quad (2)$$

where  $d = \pi/k$  is the period of the Bragg-grating structure. Deriving the couple-mode equations below, we assume the case of a *shallow grating*, i.e., that the condition  $\epsilon_j \ll 1$  holds. Additionally, we may consider a periodic modulation of the nonlinear quadratic susceptibility taking  $\chi^{(2)}(z) = \chi^{(2)}(z + d)$  as a periodic function with the same period  $d$ . However, we have verified that this effect does not modify quantitatively the analysis and results presented below, so that we consider the simplest case when  $\chi^{(2)}$  is constant.

For a periodic structure, the Bragg reflection leads to a strong interaction between the forward and backward waves at the Bragg wave number  $k_B \approx k$ . To derive the coupled-mode equations for the wave envelopes, we consider the asymptotic expansion for the electric field in the form

$$E = (E_+ e^{ikz} + E_- e^{-ikz}) e^{-i\omega t} + \text{c.c.} + E^{(0,0)} + E^{(0,2)} e^{2ikz} + E^{(0,-2)} e^{-2ikz} + (E^{(2,0)} + E^{(2,2)} e^{2ikz} + E^{(2,-2)} e^{-2ikz}) e^{-2i\omega t} + \text{c.c.}, \quad (3)$$

where  $E_{\pm} = E_{\pm}(z, t)$  are slowly varying envelopes of the forward (+) and backward (-) waves. The frequency  $\omega$  satisfies the dispersion relation for linear waves,  $c^2 k^2 = \omega^2 \epsilon(\omega)$ . Due to quadratic nonlinearity, the expansion (3) includes higher-order terms at the frequency  $2\omega$  and the zero-frequency term, so that the slowly varying functions  $E^{(n,m)} = E^{(n,m)}(z, t)$  are defined as nonlinear amplitudes of the  $(n, m)$ -order harmonics  $e^{-in\omega t} e^{imkz}$ .

To derive the equations for the coupled-mode theory, we follow the standard procedure of the asymptotic theory for nonlinear waves [9] and introduce a small parameter  $\varepsilon$  according to the relations (another choice of the scaling is discussed below)  $E_{\pm} \sim O(\varepsilon)$ ,  $\partial E_{\pm} / \partial t \sim \partial E_{\pm} / \partial z \sim O(\varepsilon^3)$ , and  $\varepsilon_j \sim O(\varepsilon^2)$ . Then, substituting the expansion (2) and (3) into Eq. (1), we compare the terms of the same order in front of the coefficients  $e^{-in\omega t} e^{imkz}$ . At the orders  $(2,0)$ ,  $(2,\pm 2)$ , and  $(0,\pm 2)$ , we obtain, respectively,

$$E^{(2,0)} = -\frac{2\chi^{(2)}}{\varepsilon(2\omega)} E_+ E_- \sim O(\varepsilon^2),$$

$$E^{(2,\pm 2)} = -\frac{\omega^2 \chi^{(2)}}{[c^2 k^2 - \omega^2 \varepsilon(2\omega)]} E_{\pm}^2 \sim O(\varepsilon^2),$$

$$E^{(0,\pm 2)} = -\frac{\chi^{(2)}}{2c^2 k^2} \frac{\partial^2}{\partial t^2} (E_{\pm}^* E_{\mp}) \sim O(\varepsilon^6),$$

where we have assumed nonresonant interaction with the second harmonic, i.e.,  $\omega^2 \varepsilon(2\omega) \neq c^2 k^2$ .

At the orders  $(1,\pm 1)$  and  $(0,0)$  we obtain a system of coupled nonlinear equations,

$$i \left( \frac{\partial}{\partial t} + v_g \frac{\partial}{\partial z} \right) E_+ + \kappa E_- + (A|E_+|^2 + B|E_-|^2 + CE^{(0,0)}) E_+ = 0, \quad (4)$$

$$i \left( \frac{\partial}{\partial t} - v_g \frac{\partial}{\partial z} \right) E_- + \kappa^* E_+ + (B|E_+|^2 + A|E_-|^2 + CE^{(0,0)}) E_- = 0, \quad (5)$$

$$\left( \frac{\partial^2}{\partial z^2} - \frac{1}{v_0^2} \frac{\partial^2}{\partial t^2} \right) E^{(0,0)} + D \frac{\partial^2}{\partial t^2} (|E_+|^2 + |E_-|^2) = 0, \quad (6)$$

where  $v_g(\omega) = d\omega/dk$ ,  $v_0 = v_g(0)$ ,  $\kappa = \omega^2 \varepsilon(\omega) \varepsilon_1 f^{-1}(\omega)$ ,  $A = 2(\chi^{(2)})^2 \omega^4 \{f(\omega) [c^2 k^2 - \omega^2 \varepsilon(2\omega)]\}^{-1}$ ,  $B = -4(\chi^{(2)})^2 \omega^2 [f(\omega) \varepsilon(2\omega)]^{-1}$ ,  $C = 2\omega^2 \chi^{(2)} f^{-1}(\omega)$ , and  $D = -2\chi^{(2)}/c^2$  with  $f(\omega) \equiv [\omega^2 \varepsilon(\omega)]'$ . If we keep the transverse coordinates  $(x, y)$ , Eq. (6) should also include the transverse Laplacian in the same order. System (4)–(6) describes the interaction between the forward and backward waves coupled to a dc wave induced via the rectification effect. Including the optical Kerr effect, we obtain the same system of the coupled equations as Eqs. (4) and (5) but with the modified constants  $A$  and  $B$ .

An important issue is a link between our model (4)–(6) and the previous studies of gap solitons in a periodic media with a nonlinear quadratic response. As follows from Eqs. (4)–(6), the induced dc field has the order of  $\varepsilon^2$ , given by the assumption of a shallow grating. In spite of the fact that the dc wave  $E^{(0,0)}$  itself is of a higher order in comparison with the forward and backward scattering waves, it becomes coupled to the fields  $E_+$  and  $E_-$  in the main order. On the other hand, if we assume a much stronger dc field [e.g., of order of  $O(1)$ , as in the case of a deep grating], the system (4)–(6) becomes decoupled and the dc wave satisfies an independent equation. As a result, the model (4)–(6) reduces to

a particular form of the conventional model of  $\chi^{(2)}$  gap solitons [3] valid for a nonresonant limit of a large mismatch.

In the case of a single wave propagating in a homogeneous medium, the induced dc field is explicitly given by the host wave [6]. The similar result is valid for the case of a deep grating described by the modulations of the Bloch waves, but not for a shallow grating we discuss here. If we assume the scaling  $\partial E_{\pm} / \partial z \sim O(\varepsilon^2)$  (as usually done in the analysis of higher-dimensional systems such as the Dawey-Stewartson equation) and  $E^{(0,0)}$  of order  $\varepsilon^4$ , then the coupling between the dc wave and host wave can be neglected. However, for isotropic scaling as presented here, the effect of the dc wave is directly included in Eqs. (4) and (5). Interaction between the dc field and fundamental harmonics has also been discussed in Ref. [6], however in that analysis, the dc field appears as a cascading effect and its velocity is almost the same as the phase velocity. We notice that in our case, the dc field is essentially excited by quadratic nonlinearity and no assumption is required for the velocity.

We are looking for spatially localized solutions of Eqs. (4)–(6) for *bright gap solitons* in the form

$$E_+ = \Delta^{-1/2} f(\zeta) e^{i[\theta_1(\zeta) - \Omega t + g/2]},$$

$$E_- = \Delta^{1/2} f(\zeta) e^{i[\theta_2(\zeta) - \Omega t - g/2]}, \quad (7)$$

where  $\zeta = z - Vt$ ; the functions  $f(\zeta)$  and  $\theta_{1,2}(\zeta)$ , and the parameters  $\Omega$ ,  $V$ ,  $\Delta$  are assumed to be real. The parameter  $g$  is the argument of the coupling parameter  $\kappa$ , i.e.,  $\kappa = |\kappa| e^{ig}$ . Substituting the ansatz (7) into Eq. (6), we obtain

$$E^{(0,0)}(\zeta) = -\frac{v_0^2 V^2 D}{(v_0^2 - V^2)} \left( \Delta + \frac{1}{\Delta} \right) f^2(\zeta),$$

and, therefore, the contribution of the dc field should vanish at  $V=0$ .

From Eqs. (4) and (5), we set the parameter  $\Delta$  as  $\sqrt{(v_g - V)/(v_g + V)}$ , and then obtain a system of coupled equations for  $f$ ,  $\theta_- \equiv \theta_1 - \theta_2$ , and  $\theta_+ \equiv \theta_1 + \theta_2$ ,

$$\frac{df}{d\zeta} = \mu f \sin \theta_-, \quad (8)$$

$$\frac{d\theta_-}{d\zeta} + \nu - 2\mu \cos \theta_- + \delta f^2 = 0, \quad (9)$$

$$\frac{d\theta_+}{d\zeta} + \frac{V\nu}{v_g} + \eta f^2 = 0, \quad (10)$$

where  $\mu = |\kappa|/(v_g^2 - V^2)^{1/2}$ ,  $\nu = -(2v_g \Omega)/(v_g^2 - V^2)$ , and

$$\delta = -2(v_g^2 - V^2)^{-1/2} \left( \frac{(v_g^2 + V^2)}{(v_g^2 - V^2)} \tilde{A} + \tilde{B} \right),$$

$$\eta = -4(v_g^2 - V^2)^{-3/2} v_g \tilde{A},$$

$$\tilde{A} = A - \frac{V^2 v_0^2 CD}{(v_0^2 - V^2)}, \quad \tilde{B} = B - \frac{V^2 v_0^2 CD}{(v_0^2 - V^2)}.$$

We notice that in the problem under consideration, we should assume  $|V| < v_g$  and  $|\nu| < 2\mu$ .

Similar to the analysis presented in Ref. [10], from Eqs. (8)–(10) we obtain a closed differential equation for the function  $\theta_-$  in the form of the double sine-Gordon (DSG) equation. The DSG equation can be integrated, and its localized solutions are two types of *kinks* and *antikinks* [11]. Using the relevant solutions, we can then find

$$f(\xi) = \left\{ \frac{\mp (4\mu/\delta)[1 - (\nu/2\mu)^2]}{\cosh(\xi\sqrt{4\mu^2 - \nu^2}) \mp (\nu/2\mu)} \right\}^{1/2},$$

where the signs  $\pm$  stand for the cases  $\delta > 0$  and  $\delta < 0$ , respectively. Functions  $\theta_1$  and  $\theta_2$  are then obtained as

$$\begin{aligned} \theta_- &= \theta_1 - \theta_2 \\ &= -2 \tan^{-1} \left\{ \sqrt{\frac{2\mu - \nu}{2\mu + \nu}} \tanh^{\mp 1} \left( \frac{\sqrt{4\mu^2 - \nu^2}}{2} \xi \right)^{\mp 1} \right\}, \end{aligned} \quad (11)$$

$$\begin{aligned} \theta_+ &= \theta_1 + \theta_2 = -\frac{\nu V}{v_q} \xi \\ &\pm \frac{4\eta v_g}{\delta} \tan^{-1} \left[ \sqrt{\frac{2\mu \pm \nu}{2\mu \mp \nu}} \tanh \left( \frac{\sqrt{4\mu^2 - \nu^2}}{2} \xi \right) \right] + \mathcal{C}_{\pm}, \end{aligned} \quad (12)$$

where  $\mathcal{C}_{\pm}$  are integration constants. In order to obtain solutions for gap solitons, we restrict the possible angle variable by the domain,  $0 \leq \theta_- \leq 2\pi$ . The solution obtained from Eq. (11) describes a two-parameter family of gap solitons, spatially localized waves in the Bragg gratings, which are similar to the gap solitons of the conventional coupled-mode theory. Actually, by renormalizing the variables as  $A \rightarrow \pm\sigma$ ,  $B \rightarrow \pm 1$ ,  $C \rightarrow 0$ ,  $|\kappa| \rightarrow 1$ ,  $v_g \rightarrow 1$ ,  $V \rightarrow \nu$ ,  $\Omega \rightarrow \pm\sqrt{1 - \nu^2} \cos Q$ ,  $\mathcal{C}_{\pm} \rightarrow (1/2 \pm 1/2)\pi \pm (4\sigma\nu\alpha^2)/(1 - \nu^2)\pi + 2\phi$ , we can demonstrate that the solution is essentially the same as that earlier obtained in Ref. [12]. However, the effect of the dc wave is included in the parameters  $\delta$  and  $\eta$ .

Similar to the case of conventional gap solitons [13], in the case  $|\nu| > 2\mu$ , spatially localized solution of Eqs. (8) and (9) do not exist. Instead, the kinks of the DSG equation for  $\theta_-$  give solutions for *dark gap solitons* (see also [13]), localized waves on nonvanishing backgrounds,

$$f(\xi) = \sqrt{\frac{2\mu}{|\delta|} \left( \frac{|\nu|}{2\mu} - 1 \right)} \frac{\sqrt{|\nu|/2\mu \cosh(\sigma\xi) \pm 1}}{\sqrt{(|\nu|/2\mu) \cosh^2(\sigma\xi) - 1}},$$

where

$$\sigma = 4\mu \sqrt{\frac{|\nu|}{2\mu} - 1}.$$

Upper and lower signs correspond to two types of such solitons, with the maximum intensity *large* or *smaller* than the background intensity.

The effective renormalization of the coefficients due to the induced dc field seems extremely important for the soliton stability. Indeed, when  $\nu_0 < v_g$  the coefficients have a

singularity provided  $V \rightarrow \nu_0$ , changing the character of the dependence of the soliton parameters and the system conserved quantities on  $V$ . The recent stability analysis of the conventional gap solitons [14] revealed the existence of two types of instabilities, *oscillatory* and *translational*. The most important, translational instability appears for large  $V$ , so that the induced dc field is expected to have a strong effect on the soliton stability. In fact, we anticipate that all gap solitons for  $\nu_0 < V < v_g$  may become unstable. However, the detailed discussion of the stability is beyond the scope of the present study. It is also worth mentioning that in the limit  $V \rightarrow \nu_0$  when the coefficients grow, the transverse effect in Eq. (6) becomes important and should be included to compensate the singularity.

Finally, we present the system invariants of the model defined by Eqs. (4)–(6). Similar to some other models, we are not able to present Eqs. (4)–(6) in a Hamiltonian form directly, and therefore we introduce an auxiliary function  $\phi$  through the relation  $\alpha \partial\phi/\partial z = E^{(0,0)} - \nu_0^2 D(|E_+|^2 + |E_-|^2)$ , where  $\alpha^2 = \nu_0^2 D/C$ . Then, we define the second canonical variable as  $\psi = \nu_0^2 \partial\phi/\partial t$  and show that Eqs. (4)–(6) can be written as a Hamiltonian system

$$\frac{\partial\phi}{\partial t} = \frac{\delta H}{\delta\psi}, \quad \frac{\partial\psi}{\partial t} = -\frac{\delta H}{\delta\phi}, \quad \frac{\partial E_{\pm}}{\partial t} = i \frac{\delta H}{\delta E_{\pm}^*},$$

with the following Hamiltonian:

$$\begin{aligned} H &= \int_{-\infty}^{+\infty} dz \left\{ \frac{\nu_0^2}{2} \psi^2 - \phi \frac{\partial^2 \phi}{\partial z^2} + \kappa E_+^* E_- + \kappa^* E_+ E_-^* \right. \\ &+ \frac{iv_g}{2} \left( E_+^* \frac{\partial E_+}{\partial z} - E_+ \frac{\partial E_+^*}{\partial z} - E_-^* \frac{\partial E_-}{\partial z} + E_- \frac{\partial E_-^*}{\partial z} \right) \\ &+ \frac{\bar{A}}{2} (|E_+|^4 + |E_-|^4) + \bar{B} |E_- E_+|^2 \\ &\left. + \alpha C \frac{\partial\phi}{\partial z} (|E_+|^2 + |E_-|^2) \right\}, \end{aligned} \quad (13)$$

where  $\bar{A} = A + \nu_0^2 CD$  and  $\bar{B} = B + \nu_0^2 CD$ . Other integrals of motion of the system (4)–(6) are the field momentum,

$$P = \int_{-\infty}^{+\infty} dz \left\{ \left( E_+ \frac{\partial E_+^*}{\partial z} + E_- \frac{\partial E_-^*}{\partial z} \right) - 2 \frac{\partial\phi}{\partial z} \psi \right\},$$

the total number of the forward and backward waves, and an independently conserved number of the dc waves,

$$N = \int_{-\infty}^{+\infty} dz (|E_+|^2 + |E_-|^2), \quad N_0 = \int_{-\infty}^{+\infty} dz \psi.$$

Therefore, in sharp contrast to the conventional coupled-mode theory of gap solitons, the model (4)–(6) possesses one additional integral of motion, and it has no analogy with other soliton-bearing nonintegrable models where the soliton stability has been investigated so far.

In conclusion, we have demonstrated the importance of the optical rectification effect in the theory of gap solitons propagating in periodic optical media with a quadratic nonlinear response. We have derived, for the first time to our knowledge, a novel model of the coupled-mode theory for optical gap solitons in quadratically nonlinear Bragg gratings that describes a coupling of the forward and backward waves

to an induced dc field, and we have found the analytical solutions for moving bright and dark gap solitons.

Takeshi Iizuka acknowledges the hospitality of the Optical Sciences Center and the support of the Japanese Ministry of Education. Yuri Kivshar is a member of the Australian Photonics Cooperative Research Center.

- 
- [1] For an overview, see G. Stegeman, D. J. Hagan, and L. Torner, *Opt. Quantum Electron.* **28**, 1691 (1996).
- [2] For an overview, see Yu. S. Kivshar, in *Advanced Photonics with Second-Order Optically Nonlinear Processes*, edited by A. D. Boardman *et al.* (Kluwer Academic, Dordrecht 1998), p. 451.
- [3] Yu. S. Kivshar, *Phys. Rev. E* **51**, 1613 (1995); Yu. S. Kivshar *et al.*, *Int. J. Mod. Phys. B* **9**, 2963 (1995); C. Conti, S. Trillo, and G. Assanto, *Phys. Rev. Lett.* **78**, 2341 (1997); *Opt. Lett.* **22**, 445 (1997); T. Peschel, U. Peschel, F. Lederer, and B. Malomed, *Phys. Rev. E* **55**, 4730 (1997); H. He and P. D. Drummond, *Phys. Rev. Lett.* **78**, 4311 (1997); C. Conti, G. Assanto, and S. Trillo, *Opt. Lett.* **22**, 1350 (1997); A. Arraf and C. M. de Sterke, *Phys. Rev. E* **58**, 7951 (1998).
- [4] M. Bass, P. A. Franken, J. F. Ward, and G. Weinreich, *Phys. Rev. Lett.* **9**, 446 (1962).
- [5] C. Bosshard, R. Spreiter, M. Zgonik, and P. Günther, *Phys. Rev. Lett.* **74**, 2816 (1995).
- [6] A. G. Kalocsai and J. W. Haus, *Phys. Rev. A* **49**, 574 (1994).
- [7] M. J. Ablowitz, G. Biondini, and S. Blair, *Phys. Lett. A* **236**, 520 (1997).
- [8] See, e.g., M. Boiti, J. J. Leon, L. Martina, and F. Pempinelli, *Phys. Lett. A* **132**, 432 (1988).
- [9] A. Jeffrey and T. Kawahara, *Asymptotic Methods in Nonlinear Wave Theory* (Pitman, Boston, 1982).
- [10] D. L. Mills and S. E. Trullinger, *Phys. Rev. B* **36**, 947 (1987).
- [11] See, e.g., O. M. Braun and Yu. S. Kivshar, *Phys. Rep.* **306**, 1 (1998).
- [12] A. B. Aceves and S. Wabnitz, *Phys. Lett. A* **141**, 37 (1989); D. N. Christodoulides and R. I. Joseph, *Phys. Rev. Lett.* **62**, 1746 (1989).
- [13] J. Feng and F. K. Kneubühl, *Quantum Electron.* **29**, 590 (1993).
- [14] I. V. Barashenkov, D. E. Pelinovsky, and E. V. Zemlyanaya, *Phys. Rev. Lett.* **80**, 5117 (1998); A. De Rossi, C. Conti, and S. Trillo, *ibid.* **81**, 85 (1998).