

Dynamics of FitzHugh-Nagumo excitable systems with delayed coupling

Nikola Burić* and Dragana Todorović

Department of Physics and Mathematics, Faculty of Pharmacy, University of Beograd, Vojvode Stepe 450 Beograd, Yugoslavia

(Received 21 November 2002; revised manuscript received 3 April 2003; published 30 June 2003)

Small lattices of N nearest-neighbor coupled excitable FitzHugh-Nagumo systems, with time-delayed coupling are studied and compared with systems of FitzHugh-Nagumo oscillators with the same delayed coupling. Bifurcations of equilibria in an $N=2$ case are studied analytically, and it is then numerically confirmed that the same bifurcations are relevant for the dynamics in the case $N>2$. Bifurcations found include inverse and direct Hopf and fold limit cycle bifurcations. Typical dynamics for different small time lags and coupling intensities could be excitable with a single globally stable equilibrium, asymptotic oscillatory with symmetric limit cycle, bistable with stable equilibrium and a symmetric limit cycle, and again coherent oscillatory but nonsymmetric and phase shifted. For an intermediate range of time lags, inverse sub-critical Hopf and fold limit cycle bifurcations lead to the phenomenon of oscillator death. The phenomenon does not occur in the case of FitzHugh-Nagumo oscillators with the same type of coupling.

DOI: 10.1103/PhysRevE.67.066222

PACS number(s): 05.45.Xt, 02.30.Ks

I. INTRODUCTION

Excitability is a common property of many physical and biological systems. Since the work of Hodgkin and Huxley [1], and the development of the basic mathematical model by FitzHugh *et al.*, [2] and Nagumo [3] the reported research on the subject has grown enormously. As for a general review, we cite just the classic Refs. [4,5] and Refs. [6,7] for examples of a recent physical, and Refs. [8,9] for neurobiological applications. For instance, a single neuron displays excitable behavior, in the sense that a small perturbation away from its quiescent state, i.e., a stable stationary value of the cross membrane potential, can result in a large excursion of its potential before returning to quiescent. Such generation of a single spike in the electrical potential across the neuron membrane is a typical example of the excitable behavior. Many other cells, besides neurons, are known to generate potential spikes across their membrane. Such excitable units usually appear as constitutive elements of complex systems, and can transmit excitations between them. The dynamics of the complex system depends on the properties of each of the units and on their interactions. In biological, as well as physical, applications the transmission of excitations is certainly not instantaneous, and the representation by nonlocal and instantaneous interactions should be considered only as a very crude approximation. For example, significant delays of more than 4% of the characteristic period of the 40-Hz-frequency oscillations of the brain neurons occur during the nerve conduction between the neurons less than 1 mm apart [10,11].

This paper is devoted to an analysis of a small lattice of a particular type of excitable systems, with a finite nonzero duration of the transfer of the excitations between the neighboring units. Despite its relevance and a large amount of related research (to be summarized and discussed in the last section) excitable systems with time-delayed coupling have not been sufficiently studied. We shall be particularly inter-

ested in the bifurcations that turn on and turn off the oscillatory behavior as the coupling constant and the small time lag are varied.

In this section, we formulate the model that is to be analyzed, and then briefly preview our results and discuss the context of our work.

As a model of each of the excitable units we shall use the paradigmatic example of the FitzHugh-Nagumo system in the form, and for the parameter range, when the system displays the excitable behavior. The dynamical equations of the single uncoupled excitable unit are [11]

$$\begin{aligned}\dot{x} &= -x^3 + (a+1)x^2 - ax - y, \\ \dot{y} &= bx - \gamma y,\end{aligned}\tag{1}$$

where a , b , and γ are positive parameters. In the original interpretation of Eq. (1), as a model of the neuronal excitability, x represents the transmembrane voltage and the variable y should model the time dependence of several physical quantities related to electrical conductances of the relevant ion currents across the membrane. In the model x behaves as an excitable variable and y is the slow refractory variable.

The particular form (1) of the FitzHugh-Nagumo model does not admit periodic solutions for any values of the parameters. Furthermore, we shall restrict our analysis to the range of the parameter values where the system exhibits excitability, with only one attractor in the form of a stable fixed point at the origin. For this to be the case, b and γ should be of the same order of magnitude and considerably smaller than a (see Sec. II). We refer to system (1) in this range of parameters as the excitable FitzHugh-Nagumo model. On the other hand, the minimal modification of Eq. (1), which renders a system to have a stable limit cycle, is obtained by adding to the first equation an external constant current I of a prescribed intensity. We shall refer to such a system with the stable limit cycle as the FitzHugh-Nagumo oscillator as opposed to the excitable FitzHugh-Nagumo (1).

*Email address: buric@phy.bg.ac.yu

The full system is a one-dimensional lattice of N identical excitable units of form (1), given by the equations of the following type:

$$\begin{aligned}\dot{x}_i &= -x_i^3 + (a+1)x_i^2 - ax_i - y_i + cF(x_{i-1}^\tau, x_i, x_{i+1}^\tau), \\ \dot{y}_i &= bx_i - \gamma y_i, \quad i=1, \dots, N,\end{aligned}\quad (2)$$

where

$$x_i^\tau(t) \equiv x_i(t - \tau),$$

τ is a fixed time lag and c is the coupling constant. A general form of the coupling term will be specified later.

Local stability near the rest state of Eq. (2), and global dynamics such as existence of stable limit cycles, and the properties of the oscillations on such a cycle, do depend on the coupling function. However, we shall see that local properties and even the global dynamics are qualitatively the same for a large class of coupling functions that are dependent only on the voltages of the neighbors, for example,

$$F(x_{i-1}^\tau, x_i, x_{i+1}^\tau) = f(x_{i-1}^\tau) + f(x_{i+1}^\tau), \quad f(x) = \tan^{-1}(x).$$

On the other hand, diffusive coupling, i.e., proportional to $x_i(t) - x_{i-1}(t - \tau)$ implies different properties of the global dynamics. Furthermore, important dynamical phenomena that occur for $N=2$ happen also for $N>2$. In fact, most of our results will be derived by considering first the system with only two coupled units, and then checking the conclusions in the case of medium $N>2$ by numerical computations.

It is a well known, and often used, fact that the time delay could destabilize a stationary point and introduces oscillatory behavior. Also, networks of oscillatory units with delayed coupling have been analyzed before. The studied oscillatory systems could be roughly divided into those where the oscillatory units are general limit cycle oscillators, say near the Hopf bifurcation (for example, Refs. [12–14]), phase-coupled phase oscillators [15–20]), or the relaxation oscillators (for example, Refs. [21,22]) typical in the neurobiological applications [8,23,24]. In the later case the form of the coupling takes, more or less, into account the properties of real synaptic interactions between the neurons [24,25].

In the last section we shall more systematically compare, system (2) and our results with several similar or related models. Here we should like to point out that the major part of our analysis deals with the system of coupled excitable units, and the system of FitzHugh-Nagumo oscillators with the same coupling is mentioned only in order to stress the differences. On the other hand, a sufficiently strong instantaneous coupling (time lag equal to zero) between the excitable (not oscillatory) units can introduce the oscillatory solutions. This phenomenon has been known already to Turing [26] and was studied by Smale [27] and Johnson and co-worker [28]. As we shall see, for such sufficiently strong coupling, a time lag that is small on the scale of the interspikes or refractory period induces drastic qualitative changes in the dynamics. Phenomena such as death of oscillations, bistable excitability, and transitions between symmetric in-phase and

nonsymmetric, phase-shifted asymptotic oscillations, all occur in system (2) as the time delay is varied. On the other hand, dynamics of the coupled FitzHugh-Nagumo oscillators with the same type of coupling is quite different.

The results of our study are presented as follows. Sections II and III are concerned with the system with just two excitable units. Analytic results about the codimension 1 bifurcations of the stationary solutions are given in detail in Sec. II for a specific common type of coupling such that there are differences between coupled excitable and coupled oscillatory units. Other types of coupling are briefly discussed. Numerical analyzes of global dynamics and, in particular, of the periodic solutions and their bifurcations are presented in Sec. III. Here we also point out some of the differences between coupled excitable systems vs the oscillators. In Sec. IV we demonstrate, by direct numerical computations, that the phenomena analyzed in Secs. II and III for $N=2$ occur also in a similar way in the system consisting of $N>2$ identical units. Conclusions, discussion and comparison with related works are given in Sec. V.

II. TWO COUPLED UNITS: LOCAL STABILITY AND BIFURCATIONS

In this section we study stability and bifurcations of the zero stationary point of only two coupled identical FitzHugh-Nagumo excitable systems, given by the following equations:

$$\begin{aligned}\dot{x}_1 &= -x_1^3 + (a+1)x_1^2 - ax_1 - y_1 + cf(x_2^\tau), \\ \dot{y}_1 &= bx_1 - \gamma y_1, \\ \dot{x}_2 &= -x_2^3 + (a+1)x_2^2 - ax_2 - y_2 + cf(x_1^\tau), \\ \dot{y}_2 &= bx_2 - \gamma y_2,\end{aligned}\quad (3)$$

where the coupling function satisfies

$$f_0 = 0, \quad f'_0 = \delta > 0, \quad (4)$$

and the subscript 0 denotes that the function is evaluated at $(x_1, x_2, y_1, y_2) = (0, 0, 0, 0)$. In fact, the first condition is not crucial, and is introduced only for convenience.

A. A single neuron

Consider first one of the units in the case of zero coupling. Point $(x, y) = (0, 0)$ is an intersection of the cubic \dot{x} nullcline and the linear \dot{y} nullcline for any value of the parameters a, b, γ , so that it is always a stationary point. Furthermore, it is always a stable stationary point, which could be a node, if $(a - \gamma) > 2\sqrt{b}$, or a focus, when $(a - \gamma) < 2\sqrt{b}$. There could be one more (non-generic case) and two more stationary points, but we shall restrict our attention to the case when $(0, 0)$ is the only stationary solution. This is the case if $4b/\gamma > (a - 1)^2$. We shall make no further assumptions as to the nature of the stable stationary point $(0, 0)$, but, as we shall see, some of the typical behavior of the delayed

coupled systems will be lost in the singular limit $b \rightarrow 0, \gamma \rightarrow 0$, and is difficult to observe very close to this limit. The particular form of the FitzHugh-Nagumo model with no external current and $(0,0)$ stationary point does not possess periodic solutions for any values of the parameters. However, there are solutions that start in a small neighborhood of $(0,0)$, quite rapidly go relatively far away, and then approach back on to the stationary point [see Fig. 4(a)]. Such solutions represent typical excitable behavior. The excitability that is displayed by the FitzHugh-Nagumo system is of, the so called, type II [8], in the sense that there is no clear-cut threshold in the phase space between the excitable orbits and the orbits that return quickly, and directly, to the rest state. In fact, there are orbits that continuously interpolate between the two types of orbits. However, as parameters b and g are decreased, as compared with a fixed a , the excitable behavior quite rapidly (but continuously) becomes dominant. We shall always use such values of the parameters that the excitable behavior is clearly demonstrated, for example $b > \gamma^2$ and $a \gg b, a \gg \gamma$.

In order to stress typical properties of the excitable, but not oscillatory, systems, we shall also need a convenient system with a stable oscillatory behavior. Such a system is obtained by adding an external (say, constant) current I to the \dot{x} equation (or to the other one) of the FitzHugh-Nagumo model. The constant current shifts the intersection of the two nullclines, and if it is such that the intersection lies on the part of the \dot{x} nullcline with a positive slope, then the stationary point is unstable and there is a stable limit cycle. The limit cycle is born in the supercritical Hopf bifurcation of the stationary solution. The limit cycle is of approximately circular shape only if I is quite close to the critical value I_0 , and then the amplitude is of the order of $\sqrt{I - I_0}$. Otherwise it has the shape typical for relaxation oscillators.

B. Instantaneously coupled identical units

As the next step, let us fix parameters a, b , and γ such that each of the units displays the excitable behavior, and consider the coupled system but with the instantaneous coupling $\tau = 0$. Point $(x_1, y_1, x_2, y_2) = (0, 0, 0, 0)$ represents a stationary solution, and its local stability is determined by analyzing the corresponding characteristic equation

$$\begin{aligned} & [(a + \lambda)(\gamma + \lambda) + b - c\delta(\gamma + \lambda)] \\ & \times [(a + \lambda)(\gamma + \lambda) + b - c\delta(\gamma + \lambda)] \\ & = 0. \end{aligned} \tag{5}$$

The sign of the real parts of the four eigenvalues

$$\begin{aligned} 2\lambda_{1,2} &= -(a + \gamma - c\delta) \pm \sqrt{(a - \gamma - c\delta)^2 - 4b}, \\ 2\lambda_{3,4} &= -(a + \gamma + c\delta) \pm \sqrt{(a - \gamma + c\delta)^2 - 4b}, \end{aligned} \tag{6}$$

determines the stability type of the trivial stationary point. If $(a - \gamma) > 2\sqrt{b}$, the point is stable node-node for $0 < c < (a - \gamma - 2\sqrt{b})/\delta$, and if c is larger the eigenvalue $\lambda_{1,2}$ becomes complex and the point becomes stable focus-node. Other-

wise, if $(a - \gamma) < 2\sqrt{b}$, the point is stable focus-focus for $0 < c < (2\sqrt{b} - a + \gamma)/\delta$ and for larger c the eigenvalue $\lambda_{3,4}$ becomes real and the point is again stable focus-node.

Thus, whatever the stability type of the stationary point in the uncoupled case might be, there is the corresponding value of the coupling constant c such that the point becomes focus-node. Then, the complex pair of eigenvalues $\lambda_{1,2}$ correspond to the eigenspace given by $x_1 = x_2$ and $y_1 = y_2$. In such a situation the damped oscillations of the two units interfere synchronously, and at some still larger c_0 given, in both the cases, by

$$c_0 = \frac{a + \gamma}{\delta}, \tag{7}$$

where

$$\text{sgn} \left(\frac{d \text{Re } \lambda_{1,2}}{dc} \right)_{c=c_0} = \text{sgn}(\delta/2) > 0, \tag{8}$$

the point goes through a direct supercritical Hopf bifurcation. As the result, for small $\epsilon = c - c_0 > 0$, the stationary solution acquires an unstable two-dimensional manifold, with a stable limit cycle in it. The unstable manifold is in fact a plane given by the equations $x_1 = x_2 \equiv x$ and $y_1 = y_2 \equiv y$, independently of the form of the coupling function as in Eq. (3). Oscillations on the limit cycle are synchronous, with the linear frequency $\omega = \sqrt{b - \gamma^2}$, and symmetrical. In this paper, by synchronous we mean coherent in-phase oscillations, and by symmetrical we mean that $x_1(t) = x_2(t)$.

The dynamics on the unstable manifold for small ϵ is given by the normal form of the Hopf bifurcation

$$\begin{aligned} \dot{r} &= d\epsilon r + \alpha r^3 + O(\epsilon^2 r, \epsilon r^3, r^5), \\ \dot{\theta} &= \omega + e\epsilon + \beta r^2 + O(\epsilon^2, \epsilon r^2, r^4), \end{aligned} \tag{9}$$

where $\omega = \sqrt{b - \gamma^2}$, $d = \delta/2$, $e = -\gamma\delta/2\omega$, and r and θ are the polar coordinates

$$x = r \sin \theta, \quad y = r \cos \theta.$$

Parameters α and β depend on the particular form of the coupling function. For example, in the case the coupling function is $f(x) = \tan^{-1}(x)$, then

$$\begin{aligned} \alpha &= \frac{-3 + c_0}{8} + \frac{(a + 1)^2 \gamma}{4\omega^2}, \\ \beta &= \frac{(3 + c_0)\gamma}{8\omega} - \frac{(a + 1)^2(5\gamma^2 + 2\omega^2)}{12\omega^3}. \end{aligned}$$

The limit cycle of Eq. (9) is a good approximation only for ϵ quite small. However numerical analysis shows that the limit cycle remains a global attractor in the full four-dimensional phase space of system (3) with no time delay for a large range of $c > c_0$ values, where the approximation by the Hopf normal form (9) is no more valid. Thus there is a range of values of the coupling parameter c , where system




FIG. 1. The figure illustrates continuous transition of the limit cycles from near Hopf to that of a relaxation oscillator for the coupled system with $\tau=0$. The fixed parameters are $a=0.25$, $b = \gamma=0.02$. The cycle is created at $c=0.27$ and the smallest cycle on the figure is for $c=0.2702$, the next to the largest for $c = 0.27048$, and the largest for 0.3.

(3) with no time delay behaves as a system of two coupled identical limit cycle oscillators. Properties of the oscillations on the limit cycle depend on c . Perturbation analyses for ϵ small, or the numerical analyses for larger c , show that oscillations on the limit cycle are synchronous and symmetrical. In Fig. 1, we illustrate the limit cycles in the coupled excitable systems with no time delay. The figure illustrates oscillatory dynamics of both units since on the limit cycle $x_1(t)=x_2(t)$ and $y_1(t)=y_2(t)$. Although the limit cycles deform continuously with c , the deformation from the small Hopf circle all the way up to the large limit cycle of the shape, as for the relaxation oscillators, happens on a small interval of the values of c , smaller than 3% of the interval (c_0, c_1) .

Further increase of c , still with $\tau=0$, leads to a bifurcation of the stationary solution and of the limit cycle. For $c > (a - \gamma + 2\sqrt{b})/\delta$, there is a pair of real-positive and a pair of real-negative eigenvalues at the trivial solution. The limit cycle disappears at some still larger c_1 when there appear other stable stationary solutions of Eq. (3) (with $\tau=0$). This value of the coupling constant $c=c_1$, when there appears nonzero stable stationary solution, obviously depends on the coupling function.

In conclusion, there are three qualitatively different types of dynamics of the instantaneously coupled excitable systems. For $0 < c < c_0$ the coupled system behaves as a pair of excitable units, while for $c_0 < c < c_1$ the system behaves as a pair of identical limit cycle oscillators. For $c > c_1$ there ap-

pears a nontrivial stable stationary state. However, we shall be interested in the influence of time delay only when the coupling constant is in the range $c \in (0, c_1)$, i.e., when the instantaneously coupled system behaves either as excitable $c < c_0$ or as oscillatory $c > c_0$.

Let us now briefly consider the coupled FitzHugh-Nagumo oscillators, just in order to stress the properties that are relevant for comparison of the influence of the time delay on the dynamics of coupled oscillatory vs excitable FitzHugh-Nagumo systems. Thus the external current $I \neq 0$ and in the range such that each of the noncoupled units is an oscillator, either close to the Hopf bifurcation or of the relaxation type. We consider the coupling of the same type as in the case of the coupled excitable units (3) and (4). For convenience, the zero of the coupling function is shifted to coincide with the unstable stationary point of the noninteracting oscillators. The major effect of such coupling is to increase the amplitude of each of the oscillators. The amplitude monotonically increases with the coupling constant c . Furthermore, for the positive coupling constant smaller than some value the asymptotic dynamics of the oscillators is symmetric. However, the oscillations of the two units on the attractor need not be in-phase for larger values of the coupling constant, contrary to the case with oscillations in the instantaneously coupled excitable systems.

Before we pass onto the analysis of the delayed equations, let us mention that the diffusive coupling, when for in the \dot{x}_1 and \dot{x}_2 equations one has $(x_1 - x_2)$ and $(x_2 - x_1)$, respectively, also leads to destabilization of the stationary point and appearance of the stable limit cycle. However, in this case $x_1(t) \neq x_2(t)$ and the corresponding oscillations are coherent but with a constant phase lag. On the other hand, the trivial stationary point of the system with reversed diffusive coupling is stable for any positive c , even with an arbitrary time lag.

C. Delayed coupling

Let us now consider the dynamics in the neighborhood of the stationary point of the delayed system (3). The point $(x_1, y_1, x_2, y_2) = (0, 0, 0, 0)$ is also the stationary solution of Eq. (3), but its stability depends on τ . Linearization of the system and substitution $x_i(t) = A_i e^{\lambda t}$, $y_i(t) = B_i e^{\lambda t}$, $x_i(t - \tau) = A_i e^{\lambda(t - \tau)}$, results in a system of algebraic equations for constants A_i and B_i . This system has a nontrivial solution if the following is satisfied:

$$\Delta_1(\lambda)\Delta_2(\lambda) = 0, \tag{10}$$

where

$$\Delta_1(\lambda) = [\lambda^2 + (a + \gamma)\lambda + a\gamma + b - c\delta\lambda \exp(-\lambda\tau) - c\delta\gamma \exp(-\lambda\tau)], \tag{11}$$

$$\Delta_2(\lambda) = [\lambda^2 + (a + \gamma)\lambda + a\gamma + b + c\delta\lambda \exp(-\lambda\tau) + c\delta\gamma \exp(-\lambda\tau)]. \tag{12}$$

Equation (10) is the characteristic equation of system (3). Infinite dimensionality of the system is reflected in the transcendental character of Eq. (10). However, the spectrum of the linearization of Eqs. (3) is discrete and can be divided into infinite-dimensional hyperbolic and finite-dimensional nonhyperbolic parts [29]. As in the finite-dimensional case, the stability of the stationary point is typically, i.e., in the hyperbolic case, determined by the signs of the real parts of the roots of (10). Exceptional roots, equal to zero or with zero real part, correspond to the finite-dimensional center manifold where the qualitative features of the dynamics, such as local stability, depend on the nonlinear terms.

Let us first answer the question of local stability of the stationary point for all time lags. We have proved (see the Appendix) that the stationary point remains locally stable for all time lags if the coupling constant is below some value c^τ , which is smaller than c_0 , given by

$$c^\tau = \left(\frac{a^2 \gamma^2 - 2b + 2\sqrt{b(2\gamma^2 + 2a\gamma + b^2)}}{\delta^2} \right)^{1/2} < c_0. \quad (13)$$

The previous expression for c^τ is valid if $b > \gamma^2$, which is always satisfied by initial assumptions about the excitable units. Notice that interval (c^τ, c_0) is quite small for the units that display excitable behavior, and shrinks to zero length as $(b + \gamma)/a \rightarrow 0$.

Thus there could be two qualitatively different types of local dynamics around the stationary solution of the time-delayed coupled excitable ($c < c_0$) units. The stationary solution could be a combination of the stable node or the stable focus for $c < c^\tau$ and any τ , and for $c^\tau < c < c_0$ and sufficiently small time lags, or it could be an unstable focus for $c > c^\tau$ and for some time lags larger than a critical value. The smallest critical time lag will be found by studying the bifurcations conditions. In the following section, we shall see that there is also an important global bifurcation due to sufficiently large τ inside interval $(0, c_0)$ which changes the dynamics of the excitations.

Bifurcations due to a nonzero time lag occur when some of the roots of Eq. (10) cross the imaginary axes. Let us first discuss the nonzero pure imaginary roots. Substitution $\lambda = i\omega$, where ω is real and positive, into the first factor gives

$$c \delta (\omega^2 + \gamma^2) \sin(\omega \tau) = -\omega^3 + (b - \gamma^2)\omega,$$

$$c \delta (\omega^2 + \gamma^2) \cos(\omega \tau) = a\omega^2 + (a\gamma + b)\gamma,$$

or into the second factor gives

$$c \delta (\omega^2 + \gamma^2) \sin(\omega \tau) = \omega^3 - (b - \gamma^2)\omega,$$

$$c \delta (\omega^2 + \gamma^2) \cos(\omega \tau) = -a\omega^2 - (a\gamma + b)\gamma.$$

Squaring and adding the above two pairs of equations results in the same equation

$$\omega^6 + (A + \gamma^2)\omega^4 + (\gamma^2 A + B)\omega^2 + b\gamma^2 = 0, \quad (14)$$

where

$$A = a^2 + \gamma^2 - 2b - c^2 \delta^2 \quad \text{and} \quad B = (a\gamma + b)^2 - c^2 \delta^2 \gamma^2. \quad (15)$$

Since $\omega^2 \neq -\gamma^2$, the term $\omega^2 + \gamma^2$ can be factored out from Eq. (14) to obtain

$$\omega^4 + A\omega^2 + B = 0. \quad (16)$$

Solutions of Eq. (16) give the frequencies ω_\pm of possible nonhyperbolic solutions

$$\omega_\pm = [(-A \pm \sqrt{A^2 - 4B})/2]^{1/2}. \quad (17)$$

The corresponding critical time lag follows from Eqs. (13) and (14). Consider the first set (13). Then, if

$$\sin(\omega \tau) = \frac{-\omega_\pm^3 + (b - \gamma^2)\omega_\pm}{c \delta (\omega_\pm^2 + \gamma^2)} > 0, \quad (18)$$

we have

$$\tau_{1,\pm}^j = \frac{1}{\omega_\pm} \left[2j\pi + \cos^{-1} \left(\frac{a\omega_\pm^2 + (a\gamma + b)\gamma}{c \delta (\omega_\pm^2 + \gamma^2)} \right) \right], \quad j=0,1,2 \dots \quad (19)$$

and if

$$\sin(\omega \tau) = \frac{-\omega_\pm^3 + (b - \gamma^2)\omega_\pm}{c \delta (\omega_\pm^2 + \gamma^2)} < 0, \quad (20)$$

we have

$$\tau_{1,\pm}^j = \frac{1}{\omega_\pm} \left[(2j+2)\pi - \cos^{-1} \left(\frac{a\omega_\pm^2 + (a\gamma + b)\gamma}{c \delta (\omega_\pm^2 + \gamma^2)} \right) \right], \quad j=0,1,2 \dots \quad (21)$$

The analogous critical time lags from the second factor of the characteristic equation are given as follows. If

$$\sin(\omega \tau) = \frac{-\omega_\pm^3 + (b - \gamma^2)\omega_\pm}{c \delta (\omega_\pm^2 + \gamma^2)} < 0, \quad (22)$$

we have

$$\tau_{2,\pm}^j = \frac{1}{\omega_\pm} \left[2j\pi + \cos^{-1} \left(\frac{-a\omega_\pm^2 - (a\gamma + b)\gamma}{c \delta (\omega_\pm^2 + \gamma^2)} \right) \right], \quad j=0,1,2 \dots \quad (23)$$

and if

$$\sin(\omega \tau) = \frac{\omega_\pm^3 - (b - \gamma^2)\omega_\pm}{c \delta (\omega_\pm^2 + \gamma^2)} < 0, \quad (24)$$

we have

$$\tau_{2,\pm}^j = \frac{1}{\omega_{\pm}} \left[(2j+2)\pi - \cos^{-1} \left(\frac{-a\omega_{\pm}^2 - (a\gamma + b)\gamma}{c\delta(\omega_{\pm}^2 + \gamma^2)} \right) \right],$$

$$j = 0, 1, 2 \dots (25)$$

The above formulas give bifurcation curves in plane (τ, c) for fixed values of parameters a, b and γ . We denote any bifurcation value of the time lag by τ_c and add the subscripts and superscripts to specify a particular branch of $\tau_c(c)$. The bifurcations are either subcritical Hopf on curves $\tau_{1,-}^j$ and $\tau_{2,-}^j$ leading to the disappearance of one unstable plane, or supercritical Hopf on $\tau_{1,+}^j$ and $\tau_{2,+}^j$ resulting in the appearance of an unstable plane. The type of the bifurcation can be seen by calculation of the variations of the real parts $\text{Re}\lambda$ as the time lag is changed through the critical values. Again, differentiation of the characteristic equation gives

$$\left(\frac{\partial \Delta_1}{\partial \lambda} \Delta_2 + \Delta_1 \frac{\partial \Delta_2}{\partial \lambda} \right) \frac{d\lambda}{d\tau} = - \left(\frac{\partial E q_1}{\partial \tau} \Delta_2 + \Delta_1 \frac{\partial \Delta_2}{\partial \tau} \right)$$

and

$$\text{sgn} \left(\frac{d \text{Re } \lambda}{d\tau} \right)_{\tau=\tau_c} = \text{sgn} \left\{ \text{Re} \left(\frac{d\lambda}{d\tau} \right)^{-1} \right\}_{\tau=\tau_c}$$

$$= \text{sgn} \left(\frac{2\omega^2 + A}{c^2 \delta^2 (\omega^2 + \gamma^2)} \right).$$

Substitution of $\omega = \omega_{\pm}$ finally gives

$$\left(\frac{d \text{Re } \lambda}{d\tau} \right)_{\tau=\tau_+} > 0, \quad \left(\frac{d \text{Re } \lambda}{d\tau} \right)_{\tau=\tau_-} < 0. \quad (26)$$

Let us now discuss the zero solution of Eq. (10). Such a solution would imply that $c = (a\gamma + b)/\gamma\delta$. For all examples of the coupling functions that we have considered, such as linear, sigmoid, \tan^{-1} or \tanh , this value of c was always larger than c_1 , i.e., there were nonzero stable stationary points of Eq. (3). So we disregard such solutions of the characteristic equation Eq. (10), and concentrate only the Hopf bifurcations $\lambda = \pm i\omega$.

The bifurcation curves $\tau_c(c)$, given by Eqs. (19), (21), (23), and (25), are shown in Figs. 2–4, for the first few $j = 0, 1, 2$, and for parameters a, γ and b fixed to some typical values, and for the coupling function with $f'_0 = \delta = 1$. Bracketed letters indicate the number of stable and unstable planes in the considered area of the (c, τ) parameter space. For example, (u^2, u) means two pairs of unstable eigenvalues of the first factor in Eq. (10) and one pair of the unstable eigenvalues of the second factor. Analogously, (s, s) means that all eigenvalues have negative-real parts, i.e., the stationary solution is stable.

Consider first the coupled excitable units when the coupling is in the range $c \in (c^\tau, c_0)$. In this range, condition (20) applies for $\omega = \omega_+$, and condition (18) for $\omega = \omega_-$. Accordingly $\tau_{1,+}^j$ branches should be calculated with formula (21), and $\tau_{2,+}^j$ using Eq. (23). The $-$ branches should be com-

puted with Eq. (19) for $\tau_{1,-}^j$ and with Eq. (25) for $\tau_{2,-}^j$. This gives the bifurcation curves for $c \in (c^\tau, c_0)$ presented in Fig. 2(a). The first unstable (c, τ) domain is between curves $\tau_{2,+}^0$ and $\tau_{2,-}^0$. The unstable plane is given by $x_1 = -x_2$ and $y_1 = -y_2$. The corresponding bifurcation on $\tau_{2,+}^0$ is the inverse Hopf bifurcation which results in the destabilization of the stationary point and a collapse of an unstable limit cycle. The origin of the latter is in a global bifurcation, which will be discussed in the following section, together with the unique global attractor that exists in this parameter domain. The next unstable region between $\tau_{1,+}^0$ and $\tau_{1,-}^0$ is bordered by a direct supercritical Hopf bifurcation at $\tau_{1,+}^0$ and the subcritical Hopf bifurcation at $\tau_{1,-}^0$. The unstable invariant manifold of the stationary point is given by $x_1 = x_2, y_1 = y_2$. The stable limit cycle in it supports coherent in-phase oscillations of the two units. The unstable domains bordered by the different branches start to overlap for sufficiently large time lags, leading to multidimensional unstable manifolds of the stationary point. The global attractors for large time lags are studied in the following section.

Next we consider the range of coupling $c \in (c_0, c_1)$ (see Figs. 3 and 4). Then, for a sufficiently small $\tau > 0$, there is only one pair of roots of Eq. (11) in the right half plane, and all the other roots of Eqs. (11) and (12) have negative real parts. There is an unstable stationary solution and the stable limit cycle. If $c > c_0$ but is smaller than some c_s , the Hopf bifurcation that happens for the smallest time lag corresponds to $\tau_{1,-}^0$. Value c_s corresponds to the intersection of branches $\tau_{1,-}^0$ and $\tau_{2,+}^0$. If $c > c_s$, the first bifurcation occurs for $\tau_{2,+}^0$. In the parameter area below the two curves $\tau_{1,-}^0$ and $\tau_{2,+}^0$, denoted by $(u; s)$, the stationary solution has qualitatively the same properties as for $\tau = 0$, i.e., it is unstable and has the unstable 2D manifold with the stable limit cycle in it. The bifurcation at $\tau_{1,-}^0$ is inverse subcritical Hopf, which results in the stabilization of the stationary solution and in the creation of an unstable limit cycle.

From the set of frames in Fig. 4, we see that as $(b + \gamma) \rightarrow 0$ the value c_s approaches c_0 and the (s, s) domain beyond c_0 shrinks to nothing. In fact, in this singular limit, there are only positive solutions of Eq. (16), and the stabilization of the stationary point by the time delay cannot happen.

In order to claim that the parameter domain denoted (s, s) , where the stationary point is stable, corresponds to the phenomenon of oscillators death, the local stability of the stationary point is not sufficient. We need to analyze the global dynamics of Eq. (3), and this depends on the full form of the coupling function.

Let us briefly discuss the modifications of the presented analyses that would be implied by the substitution of the coupling function of the form as in Eq. (3) by the diffusive or a more general coupling $f(x_1, x_2)$. The analysis of the linear stability in the delayed case, in particular, the formulas for the critical time lags and eigenvalues, would remain unchanged provided that the parameters a and δ are changed as follows:

$$a \rightarrow \bar{a} = a + \partial_1 f_0, \quad \delta \rightarrow \bar{\delta} = \partial_2 f_0.$$




FIG. 2. The figures illustrate typical dynamics below $c_0=0.27$ for $a=0.25, b=\gamma=0.02$: (a) The first few branches of the bifurcation curves $\tau_c(c)$ given by Eqs. (19), (21), (23), and (24), for parameters $a=0.25, b=0.02, \gamma=0.02$, and $c < c_0$. (b) Examples of quickly relaxing (1) and periodic excitable (2) orbits [projections on (x_1, y_1)]. (c) Projection of the global attractor limit cycle on (x_1, x_2) in the (s, u) domain. (d) Projection of the two limit cycle attractors on (x_1, x_2) in the (u, u) domain.

In particular, for the diffusive coupling, with $f(x_1, x_2^T) = (x_1 - x_2^T)$, there will be a bounded (s, s) region, where the stationary point is stable for some finite, nonzero τ and unstable for $\tau=0$, and the other parameters are unchanged. For example, if $a=0.25, b=\gamma=0.02, c=0.132$ the stationary point is stable for $\tau=0.85$ up to $\tau=24.5$, and unstable for $\tau=0$ up to $\tau=0.85$ and above $\tau=24.5$. However, the unstable manifolds with the limit cycles for small τ would not be given by the $x_1 = x_2, y_1 = y_2$ plane.

Contrary to the case of coupled excitable units, the stationary solution of coupled identical FitzHugh-Nagumo oscillators, with the same type of coupling, remains unstable for any value of the time lag. Thus, there could be no oscillator death in the case of coupled FitzHugh-Nagumo oscillators with the considered type of coupling. On the other hand, it is known ([12,13]) that a different type of coupling, such as reverse diffusive, does lead to the amplitude death at least when the oscillators are near the Hopf bifurcation, i.e., $I > I_0; I \approx I_0$. However, as we have pointed earlier, there is no Hopf bifurcation of the trivial stationary point of the excit-

able systems with the reverse diffusive coupling so that the stationary state is in this case always stable.

III. GLOBAL DYNAMICS OF THE SYSTEM WITH TWO UNITS

We study the global dynamics of Eq. (3) by numerical computations of orbits for different typical values of parameters (c, τ) in each of the domains in the local bifurcation diagram of the stationary point up to moderately large values of τ . Our main interest is to determine whether or not there is one or more than one attractors, and, if there are stable oscillatory solutions, what is the dimensionality and properties of the oscillations. As examples of the coupling we used different functions with the same qualitative conclusions, and for illustration we use $f(x) = \tan^{-1}(x)$. On the other hand, diffusive coupling $f(x_1, x_2^T) = (x_1 - x_2^T)$ implies quite different global dynamics, which we shall indicate at the end of the section.

Before presenting the results let us comment on the initial




FIG. 3. First few branches of the bifurcation curves $\tau_c(c)$ given by equations (19), (21), (23), and (24), for parameters $a=0.25$, $b=0.02$, $\gamma=0.02$, and $c > c_0$.

conditions for system (3) that we used in calculations. In order to uniquely fix a solution of the delay-differential equations (3) one needs to specify the solution on interval $[-\tau, 0]$. In our calculations we always use a physically plausible initial functions on $[-\tau, 0]$ given by solutions of Eqs. (3) with $c=0$. Thus, before period τ has elapsed, each of the units was evolving independently of the other unit.

First we discuss the global dynamics for the coupling constant $c < c^\tau$, i.e., when the trivial stationary solution of the whole system is stable for any time lag. Intuitively, one would expect that if the time lag τ is such that 2τ is smaller than the minimal time needed for an excitable orbit to approach the stable stationary solution, then the coupling would just induce both units to fire one spike each, with some time delay, and relax to the stationary solution. However, if 2τ is larger than the indicated minimal time, then the excitation of one of the units would arrive just in time to kick the state of the other unit from close to the stationary into the excitable domain, even if the coupling constant is rather small. Thus, the excitable orbit of the coupled system becomes periodic. Nevertheless, the equilibrium state remains a stable stationary solution. The system as a whole is bistable excitable with periodically spiking excitations. This picture is confirmed by numerical computations. Two typical orbits are illustrated in Fig. 2(b). The periodic orbits are supported on the stable limit cycle. The latter is created in a global fold limit cycle bifurcation, together with an unstable one. As expected, the motion on the limit cycle is coherent and out-of-phase, with a frequency that increases with the time lag.

Qualitatively different global dynamical pictures can occur for the coupling constant in range (c^τ, c_0) and for various

$\tau > 0$. In the domain of the (c, τ) plane bordered by $\tau_{2,+}^0$ and $\tau_{2,-}^0$, there is only one global attractor given by the large stable limit cycle, since the value $\tau_{2,+}^0$ is above the critical time lag of the global fold limit cycle bifurcation discussed above and the stationary point is unstable [see Fig. 2(c)]. The large cycle is not affected qualitatively by the local bifurcation on $\tau_{2,+}^0$ or $\tau_{2,-}^0$, so the dynamics on it is given by coherent out-of-phase oscillations of the coupled units. If the domain is entered by crossing $\tau_{2,+}^0$, the unstable limit cycle collapses on the stationary solution, and if the domain is left through $\tau_{2,-}^0$, yet another unstable limit cycle is created.

The next unstable domain, with c still in (c^τ, c_0) is bounded by $\tau_{1,+}^0$ from below and by $\tau_{1,-}^0$ from above. The supercritical Hopf bifurcation on $\tau_{1,+}^0$ results in the creation of a two-dimensional (2D) unstable manifold of the stationary point given by $x_1 = x_2$ and $y_1 = y_2$. In it, there is a stable limit cycle supporting in-phase coherent oscillations. However, the large limit cycle with out-of-phase oscillations is not affected by the local bifurcation at $\tau_{1,+}^0$, so that in this domain the dynamics is bistable with two limit cycle attractors. This is illustrated in Fig. 2(d).

Next we consider possible attractors for the coupling constant beyond c_0 , i.e., when instantaneously coupled units behave as two limit cycle oscillators. Qualitatively different dynamics corresponding to different domains in (c, τ) are illustrated in Fig. 5. Figure 5(b) corresponds to a typical values of (c, τ) in the (u, s) domain. We have not found any global bifurcation that would occur as (c, τ) are varied inside the (u, s) region. The dynamics is characterized by the 2D unstable manifold of the stationary solution, given by $x_1 = x_2$, $y_1 = y_2$ [see Fig. 5(a)]. There is a globally stable limit cycle inside this manifold. Oscillations of x_1 and x_2 on this limit cycle are obviously in-phase.

Two frames, 5(c) and 5(d), correspond to the situations in (s, s) with one stable stationary solution but with two globally different dynamics. In Fig. 5(c) the system is bistable. There is the stable stationary solution and the stable large limit cycle in the plane $x_1 = x_2$, $y_1 = y_2$. There is also a small unstable limit cycle, which is created in the inverse subcritical Hopf bifurcation at $\tau = \tau_{1,-}^0$. This cycle acts as a threshold between the subexcited damped oscillations and periodic synchronous spiking of both units. As τ is increased, but still for $(c, \tau) \in (s, s)$, the unstable limit cycle approaches the stable one, and they disappear in a fold limit cycle bifurcation, which occurs in the invariant plane $x_1 = x_2$, $y_1 = y_2$. Thus, there is a parameter region inside (s, s) where $(0, 0, 0)$ is globally stable, and that corresponds to the death of the identical oscillators. However, let us stress again that the global dynamics for the parameters in the domain (s, s) could correspond to either spiking excitability, with subthreshold damped oscillations and supthreshold periodic spiking, or to the death of oscillators. In the latter regime the whole system is excitable with the stationary point as the only attractor.

The global dynamics above curve $\tau_{2,+}^0$ is characterized by one large limit cycle as the global attractor. The oscillations on it are coherent and out-of-phase. The same type of the global attractor occurs above the critical line $\tau_{1,+}^0$ as illus-




FIG. 4. The same as Fig. 3, but for few fixed values of a, b, γ : (a) $a=0.25, b=0.005, \gamma=0.005$; (b) $a=0.25, b=0.003, \gamma=0.003$; (c) $a=0.25, b=0.0015, \gamma=0.0015$; (d) $a=0.25, b=0.00075, \gamma=0.00075$.

trated in Figs. 5(e) and 5(f). The oscillations are further illustrated in Fig. 6(b) by plotting the limit cycle as seen in the coordinates $(x_1 - x_2, y_1 - y_2)$. Convergence to the limit cycle is much slower than in the case of the symmetric oscillators that occurs for smaller τ , illustrated in Fig. 6(a). In fact, in all domains up to a quite large value of the time lag the global attractor is a stable limit cycle (could be imbedded in a multidimensional unstable manifold for larger τ), which supports asymmetric phase-shifted oscillations of x_1 and x_2 . However, there are domains for larger values of the time lag, for example, for $\tau=55$ and any $c \in (c_0, c_1)$, where the global attractor is the symmetric limit cycle, with coherent and in-phase oscillations.

It should be pointed out that, for all larger time lags up to quite large values, equal to several refractory times of the noncoupled units, the attractor is always a limit cycle. On the limit cycle, all variables oscillate with the same frequency, and could be either symmetric or phase shifted. The two

regimes interchange as the time lag is increased. These are the only two possible stable attracting patterns, despite the large dimensionality of the unstable manifold of the stationary point. It should be pointed out that in the case of identical FitzHugh-Nagumo oscillators with the same coupling various types of quasiperiodic attractors occur for moderate values of the time lag. However, also in this case, stronger coupling and larger time lags imply synchronization, either identical or phase shifted, as for the coupled excitable systems. The dynamics for time lags much larger than the refractory time has not been systematically studied.

We now briefly comment on the dynamics in the case of the diffusive coupling. As stated before, at some c and for τ zero or small, the only attractor is the stable limit cycle, with coherent and phase-shifted oscillations of the two units. The time delay can stabilize the trivial stationary point, but the system remains bistable with the limit cycle and the stationary point as the attractors, for all values of (c, τ) in the (s, s)




FIG. 5. Phase portraits in (x_1, y_1) (a)–(e) plane, or (x_1, x_2) plane (f). The initial points, if there are different orbits, are indicated by numbers. The fixed parameters are $a=0.25$, $b=0.02$, $\gamma=0.02$, $c=0.3$, except in (a) where $c=0$. The time lag is (a),(b) $\tau=0$; (c) $\tau=4$; (d) $\tau=6$; (e),(f) $\tau=27$. Dynamics illustrated in (e) and (f) is typical also for other values of (c, τ) above the $\tau_{1,+}^I$ curve.

domain. The phenomenon of oscillator death does not occur in the case of the diffusive coupling.

IV. $N > 2$ LATTICE

The goal of this section is to present numerical evidence that for some common types of lattices with $N > 2$ there are

regions in the parameter plane (c, τ) analogous to $(u, s), (s, s), (u, u), \dots$, in Figs. 2 and 3. We have analyzed examples of systems of identical FitzHugh-Nagumo excitable units arranged in linear or circular lattices, with unidirectional or bidirectional symmetrical coupling by few typical coupling functions. Lattices of size $N=10, 20$ and $N=30$ have been studied systematically.




FIG. 6. (a) The asymptotic state is symmetric in any domain below the $\tau_{1,+}^1$ curve in Fig. 3. In domains (u,s) , (s,u) , or (u,u) the symmetric states are the synchronous oscillations, and in (s,s) the stable stationary point. (b) The asymptotic synchronous oscillations are not symmetric for (c,τ) above the $\tau_{1,+}^1$ curve in Fig. 3, as is illustrated for a pair $(c,\tau) \in (u,u)$, but become symmetric for $\tau \geq 55$ (not illustrated, see the main text).

The conclusions are illustrated using the following model:

$$\dot{x}_i = -x_i^3 + (a+1)x_i^2 - ax_i - y_i + cf(x_{i-1}^\tau) + cf(x_{i+1}^\tau), \quad (27)$$

$$\dot{y}_i = bx_i - \gamma y_i, \quad i=2, \dots, N-1,$$

$$\begin{aligned} \dot{x}_{1,N} &= -x_{1,N}^3 + (a+1)x_{1,N}^2 - ax_{1,N} - y_{1,N} + cf(x_{2,N-1}^\tau), \\ \dot{y}_{1,N} &= bx_{1,N} - \gamma y_{1,N}, \end{aligned} \quad (28)$$

where the coupling is given by $f(x) = \tan^{-1}(x)$, and $N = 20$.

First, in the case of instantaneously coupled units, there is the Hopf bifurcation at some $c = c_0$. As in $N=2$, for the coupling constant below some $c^\tau < c_0$ and any time lag the trivial stationary point is stable. If the time lag is sufficiently large, there is also the stable large limit cycle. On it, all units oscillate coherently. However, the nearest neighbors oscillate exactly antiphase, so that two clusters are formed.

The coupling above threshold c_0 , and for small time lags, leads to the appearance of a globally stable limit cycle representing synchronous oscillations in the plane given by $x_2 = \dots = x_{N-1}$, $y_2 = \dots = y_{N-1}$ and $x_1 = x_N$, $y_2 = y_N$, $[(u,s)$ region]. As expected, the synchronization period could be quite large if the value of the coupling constant is near c_0 , i.e., when each of the units is near the Hopf bifurcation.

Increasing the time lag leads to the inverse Hopf bifurcation. For any c in some interval (c_0, c_s) we have been able to find intervals of time lags (τ_c^-, τ_c^+) that correspond to bi stability or to death of all N oscillators $[(s,s)$ region], illustrated in Fig. 7. The same figures illustrate the attractors in the dynamics of any of the identical neurons. Again, the inverse Hopf and the subsequent fold limit cycle bifurcations, due to increasing time delay, are responsible for the

amplitude death in the systems of form (27). On the contrary, the stationary point of a lattice such as Eq. (27) with the same coupling but with Fitzhugh-Nagumo oscillators is always unstable.

Larger time lags do not change the topological nature of the attractor. It is always a limit cycle, but the synchronization pattern between the coherent oscillations of the units depends on τ . Nonsymmetric oscillations with equal frequencies appear after long transients, as is illustrated in Fig. 8(a). Dynamics in the transient period can be quite complicated. Properties of the asymptotic synchronization patterns could depend on the geometry of the lattice.

Existence of the presented types of dynamics, and the order of their appearance as (c,τ) are varied, was confirmed in all examples that we have studied. We conjecture that qualitative properties of the dynamics of all small 1D lattices with nearest-neighbor delayed coupling of the form as in Eqs. (27) between the identical Fitzhugh-Nagumo excitable




FIG. 7. Bistability (a) and oscillator death in lattice (27) with $a=0.25$, $b=0.02$, $\gamma=0.02$ and (a) $(c,\tau)=(0.16,4)$ or (b) $(c,\tau)=(0.16,6)$.




FIG. 8. Asymptotic states of lattice (27) for $(c, \tau) = (0.16, 15)$ are coherent oscillations but with a fixed time lag, represented by the projection of the limit cycle on the $(x_4 - x_{15}, y_4 - y_{15})$ plane in frame (b). For such a small c the synchronization period is more than ten times larger than the characteristic period, as is illustrated in frame (b) with the time dependence of the time-series $x_4(t)$ and $x_{15}(t)$.

systems are the same, at least for not very large values of the time lag, in the sense that the same types of bifurcations appear and determine the dynamics.

V. SUMMARY AND DISCUSSION

We have studied small lattices of excitable identical units with time-delayed coupling, where each unit is given by the excitable FitzHugh-Nagumo model. The coupling is always between the voltages of the nearest neighbors, but could be of a quite general form. Our primary interest was in the bifurcations and the typical dynamics that occur for time lags that are not very large on the time scale set up by the refractory or the interspike period. A detailed study, in the case of only two units, of the local stability and bifurcations of the stationary solution suggested, but does not uniquely determine, the possible global bifurcations and dynamics. These are studied numerically.

There are only few possible types of dynamics, at least for time lags as large as several refractory times. For small coupling constants and small time lags there is only one attractor in the form of the stable stationary solution. The whole system behaves qualitatively as the simple excitable. Relatively small coupling constants $c < c^\tau$ and sufficiently large time lag result in the limit cycle attractor coexistence with the stable stationary solution. The whole system is bistable with spiking excitability. The oscillations on the limit cycle are coherent and out-of-phase. For the coupling constant above c^τ the sequence of Hopf bifurcations due to the time delay of the stationary solution are possible. For $c \in (c^\tau, c_0)$ and small time lags the stable stationary point is the only attractor, but as timelag is increased the system could be either bistable or there could be only one attractor in the form of the limit cycle. The bistability is manifested either in the form of the stable stationary solution and the stable limit cycle, or could be in the form of two stable limit cycles (one in-phase and

one out-of-phase). Interval $c \in (c^\tau, c_0)$ is rather a small part of the c values for which there is only one stationary solution. For $c \approx c_0$ each of the instantaneously coupled units is near a direct supercritical Hopf bifurcation, but as soon as $c - c_0 > 0$ is bigger than some quite small ϵ_0 , the resulting limit cycle has a quite large radius and the harmonics become influential, unlike in the case of the Hopf limit cycle. Increasing the time lag τ could lead to stabilization of the stationary point, via indirect subcritical Hopf, resulting in a bistable dynamics, with a stable stationary point, small unstable limit cycle as a threshold, and a large stable limit cycle. Further increasing τ leads to a fold limit cycle bifurcation, in which the unstable and the stable limit cycles disappear, and the stationary point remains the only attractor. In this, oscillator death regime, the system again displays the simplest form of excitability, as in the case of the weak coupling $c < c_0$ and zero or small time lags. Still further increase of the time lag τ leads to the supercritical Hopf. The oscillations on the limit cycle are coherent but are phase shifted, and the oscillators need not have the same amplitude. The described sequence of bifurcations happens for time lags that are all small, up to 10%, with respect to the refractory period of the single isolated unit. Further increase of the time lag leads to more dimensional unstable manifold of the stationary solution. However, the global attractor is always a simple limit cycle. The asymptotic dynamics is always coherent, and is either in-phase or phase shifted. Unlike the case of coupled FitzHugh-Nagumo oscillators, nothing more complicated than the limit cycle could be the attractor of the coupled excitable FitzHugh-Nagumo systems.

Our analyses shows that the most common type of excitations of the whole system, in response to an impulse submitted to either of the units, is in the form of coherent out-of-phase oscillations. However, if the transmission is sufficiently strong and for moderately large transmission delays of signals between the units, the compound system would respond by synchronous in-phase oscillations. Furthermore, our results suggest that relatively small but non-zero time delay together with sufficiently strong interaction could result in a simple excitable behavior of the compound system. For such values of the parameters the system would operate as a powerful amplifier of a quite small impulse administered to its single unit. Due to the particular model of the excitable system and to the type of coupling that we have studied in detail, the most relevant possible application of our results is in modeling coupled neurons. In fact, relatively recent experiments and analyses [30] show that the FitzHugh-Nagumo equations, despite the common opinion, might represent a better qualitative model of an excitable neuron than the more detailed Hodgkin-Huxley system. Our results indicate that the fine tuning between the synaptic coupling and delay could lead to the in-phase synchronous operation of a collection of neurons.

Although there is a quite substantial amount of work done on the systems of dynamical units with the delayed coupling, such systems are comparatively much less studied than the corresponding systems with the instantaneous coupling. For the purpose of comparison with our work, we shall try to classify the existing contributions into typical groups.

First, we consider the model and the results presented in Refs. [12,13]. In these papers, a network of $N=2$ and $N > 2$ oscillators described by the equations of the normal form of the Hopf bifurcations with delayed inverse diffusive coupling is studied. At zero coupling, and/or for small time lags, all oscillators have small limit cycles just created by the direct Hopf bifurcation. It is shown that the time delay can lead to the stabilization of the trivial stationary point, even for the identical oscillators, which is interpreted as the amplitude death. Analogously, in our case, the Hopf direct bifurcation is responsible for the appearance of the oscillations when the excitable units are instantaneously coupled, and the time delay leads to stabilization of the stationary point. The death of oscillations in our case appears after the fold bifurcation of the stable and the unstable limit cycles, which are created in the same plane. The oscillator death occurs only in a domain in the (c, τ) parameter space smaller than the domain of the stability of the stationary point.

Next, we compare our model and the results with those that appear in the studies of the delayed coupled relaxation oscillators, for example, in Refs. [21,22]. In these studies, each unit is a relaxation oscillator, and the primary objective of the analysis is the periodic orbits that appear in the delayed coupled system. Singular approximation, or an approximate or numerical construction of the Poincaré map, are used to analyze various types of synchronous or asynchronous oscillations. The phenomenon of the oscillator death was not observed [24]. In our case, the noninteracting units are not oscillators and the oscillations are introduced by coupling, via the Hopf bifurcation. The domain of parameters (c, τ) that implies oscillator death shrinks to nothing in the singular limit $(b + \gamma) \rightarrow 0$. Furthermore, the FitzHugh-Nagumo model is type II excitable, which reflects in the type of bifurcations that might occur in the coupled systems.

Less directly related to our work is the analysis of the influence of the time delay in the systems of coupled phase oscillators. In fact, if the rate of attraction to the limit cycles of two voltage-coupled neural oscillators is sufficiently strong, the dynamics can be described by the coupled phase oscillators. The coupling between the phases mimics the voltage coupling, and is not of the diffusive type. The phenomenon of oscillator death in such instantaneously coupled phase oscillators was studied, for example, in Ref. [24]. The influence of time delay in coupled phase oscillators was studied, for example, in Refs. [31] and [19] (and also in Ref. [24]), where it was shown that the time delay cannot introduce significant changes into the dynamics of a class of such systems [31], unless the time lag is of the order of several oscillation periods [19]. Independently of neuronal models, collective behavior of the phase coupled (phase) oscillators with time-delayed coupling has been studied using the dynamical [15–18]), or statistical [20] methods. In our case, the coupling is between the voltages, could be of a quite general form, and all analyses and the observed phenomena occur already for quite small time lags.

Finally, the influence of time delay has been studied in the Cohen-Grasberger-Hopfield (CGH) type of neural networks, as early as in 1967 [32]. More recent references are, for example, [33,34], and for small networks [35–37] (see also

Ref. [38], and the references therein). In the nondelayed case, the stability of the stationary point in such networks is proven using an energy-Lyapunov function. Using the corresponding Lyapunov functional in the delayed case, it was shown in Ref. [34] that the stationary point remains globally stable for sufficiently small time lags. On the other hand, destabilization of the stationary point occurs via the Hopf bifurcation, as was shown in Refs. [35,36], for the networks with $N=2$ and $N=3$ units, and multiple time-delayed coupling.

As is seen, the model treated here and our results have some features common with few other models. As in the CGH networks, each isolated unit has a globally stable stationary point, and the time-delayed weak coupling does nothing to the dynamics, provided that the time lag is sufficiently small. If the coupling is strong enough, the system behaves either as a collection of near-Hopf oscillators, or as a collection of relaxation oscillators. Death of oscillators due to time delay is observed in both types of dynamics, although the phenomenon happens for a smaller range of time lags if the system behaves as a collection of relaxation oscillators.

Let us finally mention few related questions that we shall study in the future. First, it should be interesting to see if the systems of slightly different units share the same type of dynamics. Second, examples of the type-I excitable (and not oscillatory) systems coupled with time delay should be analyzed, in order to underline the role of the type of excitability. Finally, the external pulse perturbations, for example, in Refs. [39–41], could introduce different transitions from excitability to the oscillatory regime, and the consequent changes in the dynamics due to time delay should be analyzed.

ACKNOWLEDGMENT

This work was supported by the Serbian Ministry of Science Contract No. 1443.

APPENDIX

We start with the characteristic function (10) in the form

$$\phi(z) = [(z + \gamma)(z + a) + b]^2 - c^2 \delta^2 (z + \gamma)^2 \exp(-2z\tau),$$

and consider the following expression:

$$f(z) = \frac{\phi(z)}{P_4(z)} = 1 - \frac{c^2 \delta^2 (z + \gamma)^2}{P_4(z)} \exp(-2z\tau),$$

where $P_4(z) = [(z + \gamma)(z + c) + b]^2$ is actually the characteristic function of the single noncoupled unit.

Consider contour C_R in the complex half plane $\text{Re } z > 0$, formed by the segment $[-iR, iR]$ of the imaginary axis and the semicircle with the radius R centered at the origin. As condition $4(b/\gamma) > (a-1)^2$, equivalent to the existence of a unique stationary solution, is by assumption always satisfied, polynomial $P_4(z)$ has no zeros in the half plane $\text{Re } z > 0$. In that case, the number of poles of $f(z)$ is $P_c = 0$. Using the argument principle we infer the number of zeros N_{C_R} of $f(z)$. If $\lim_{R \rightarrow \infty} N_{C_R} = 0$, then all the roots of the character-

istic function $\phi(z)$ satisfy $\text{Re } z > 0$. Thus, we need to find the conditions on parameters a, b, γ , and c , such that the image of contour C_R when $R \rightarrow \infty$ by the function $f(z)$ does not encircle point $z=0$. Then the variation of the argument is zero, so that $\lim_{R \rightarrow \infty} N_{C_R} = 0$, and, consequently, the zeros of the characteristic function satisfy $\text{Re } z > 0$ for any τ . This is the essence of the amplitude-phase method (see, for example, Ref. [42]).

It is enough to consider the image of segment $[-iR, iR]$ by the function

$$\omega_\tau(z) \equiv \frac{c^2 \delta^2 (z + \gamma)^2}{P_4(z)} \exp(-2z\tau),$$

or, in fact, by just $\omega_0(z)$ since $|\omega_0(iy)| < 1$ if and only if $|\omega_\tau(iy)| < 1$, and the image of the semicircle shrinks to a point as $R \rightarrow \infty$.

Since

$$\begin{aligned} |\omega_0(iy)| &= \left| \frac{c \delta (iy + \gamma)}{(iy + \gamma)(iy + a) + b} \right|^2 \\ &= \frac{c^2 \delta^2 (\gamma^2 + y^2)}{y^4 + (a^2 + \gamma^2 - 2b)y^2 + (a\gamma + b)^2} \end{aligned}$$

we obtain that $|\omega_0(z)| < 1$ is equivalent with

$$y^4 + Ay^2 + B > 0,$$

where A and B are given by the same formula as in Eq. (15), i.e.,

$$A = a^2 + \gamma^2 - 2b - c^2 \delta^2 \quad \text{and} \quad B = (a\gamma + b)^2 - c^2 \delta^2 \gamma^2.$$

For the parameters such that $b > \gamma^2$ and $4(b/\gamma) > (a-1)^2$, the above condition is equivalent to

$$c < \left(\frac{a^2 \gamma^2 - 2b + 2\sqrt{b(2\gamma^2 + 2a\gamma + b^2)}}{\delta^2} \right)^{1/2}.$$

The right side is the critical value that we denoted c^τ in the main text.

-
- [1] A.L. Hodgkin and A.F. Huxley, *J. Physiol. (London)* **117**, 500 (1952).
- [2] R. FitzHugh, *J. Gen. Physiol.* **43**, 867 (1960).
- [3] J. Nagumo, S. Arimoto, and S. Yoshizawa, *Proc. IRE* **50**, 2061 (1962).
- [4] A. Winfree, in *1992 Lectures on Complex Systems*, edited by L. Nadel and D. Stein, Santa Fe Institute Studies in the Sciences of Complexity Vol. 5 (Addison-Wesley, Reading, 1993), p. 207.
- [5] A.T. Winfree, *The Geometry of Biological Time*, 2nd ed. (Springer-Verlag, New York, 2000).
- [6] S.M. Wiczkorek, B. Krauskopf, and D. Lenstra, *Phys. Rev. Lett.* **88**, 063901 (2002).
- [7] I.S. Aranson and L. Kramer, *Rev. Mod. Phys.* **74**, 99 (2002).
- [8] E.M. Izhikevich, *Int. J. Bifurcation Chaos Appl. Sci. Eng.* **10**, 1171 (2000).
- [9] W. Gerstner and W.M. Kistler, *Spiking Neuron Models* (Cambridge University Press, Cambridge, 2002).
- [10] G.M. Shepherd, *Neurobiology* (Oxford University Press, New York, 1983).
- [11] J.D. Murray, *Mathematical Biology* (Springer, New York, 1990).
- [12] D.V. Ramana Reddy, A. Sen, and G.L. Johnston, *Phys. Rev. Lett.* **80**, 5109 (1998).
- [13] D.V. Ramana Reddy, A. Sen, and G.L. Johnston, *Physica D* **129**, 15 (1999).
- [14] S. Wirkus and R. Rand, *Proc. DETC* **99**, (1999).
- [15] H.G. Schuster and P. Wagner, *Prog. Theor. Phys.* **81**, 939 (1989).
- [16] E. Niebur, H.G. Schuster, and D. Kammen, *Phys. Rev. Lett.* **67**, 2753 (1991).
- [17] Y. Nakamura, F. Tominaga, and T. Munakata, *Phys. Rev. E* **49**, 4849 (1994).
- [18] S. Kim, S.H. Park, and C.S. Ryu, *Phys. Rev. Lett.* **79**, 2911 (1997).
- [19] E.M. Izhikevich, *Phys. Rev. E* **58**, 905 (1998).
- [20] M.K.S. Yeung and S.H. Strogatz, *Phys. Rev. Lett.* **82**, 648 (1999).
- [21] J.J. Fox, C. Jayaprakash, D. Wang, and S.R. Campbell, *Neural Comput.* **13**, 1003 (2001).
- [22] S.R. Campbell and D. Wang, The Ohio State University CIS-Technical Report No. 47, 1996 (unpublished).
- [23] E.M. Izhikevich, *SIAM (Soc. Ind. Appl. Math.) J. Appl. Math.* **60**, 1789 (2000).
- [24] G.B. Ermentrout and N. Kopell, *SIAM (Soc. Ind. Appl. Math.) J. Appl. Math.* **50**, 25 (1990).
- [25] G.B. Ermentrout and N. Kopell, *Proc. Natl. Acad. Sci. U.S.A.* **95**, 1259 (1998).
- [26] A.M. Turing, *Philos. Trans. R. Soc. London, Ser. B* **237**, 37 (1952).
- [27] S. Smale, in *A Mathematical Model of Two Cells Via Turings Equation*, edited by S. Levin, *Lectures Math. Life Science*, Vol. 6 (American Mathematical Society, Providence, 1974), p. 17.
- [28] L. Segal and J. Jackson, *J. Theor. Biol.* **37**, 545 (1972).
- [29] J. Hale and S.V. Lunel, *Introduction to Functional Differential Equations* (Springer-Verlag, New York, 1993).
- [30] J.R. Clay and A. Shrier, *J. Theor. Biol.* **197**, 207 (1999).
- [31] F.C. Hoppensteadt and E.M. Izhikevich, *Weakly Connected Neural Networks* (Springer-Verlag, New York, 1997).
- [32] S. Grossberg, *Proc. Natl. Acad. Sci. U.S.A.* **58**, 1329 (1967).
- [33] K. Gopalsamy and I. Leung, *Physica D* **89**, 395 (1996).

- [34] H. Ye, A. Michel, and K. Wang, *Phys. Rev. E* **51**, 2611 (1995).
- [35] J. Olier and J. Belair, *Physica D* **102**, 349 (1997).
- [36] L.P. Shayer and S.A. Campbell, *SIAM (Soc. Ind. Appl. Math.) J. Appl. Math.* **61**, 673 (2000).
- [37] I. Ncube, S.A. Campbell, and J. Wu, *Fields Ins. Commun.* **36**, 1 (2002).
- [38] S.A. Campbell (unpublished).
- [39] D.V. Ramana Reddy, A. Sen, and G.L. Johnston, e-print nlin.CD/0205035.
- [40] H. Kitajima and H. Kawakami, in *Bifurcation in Coupled BVP Neurons with External Impulsive Forces*, Proceedings of ISCAS'2001, Sydney, (ISCAS, Sydney, 2001), Vol. III, pp. 285–288.
- [41] S. Combes and A.H. Osbaldestin (unpublished).
- [42] L.E. El'gol'ts and S.B. Norkin, *Introduction to the Theory of Differential Equations with Deviating Argument* (Nauka, Moskva, 1971) (in Russian).