

Velocity distribution in a viscous granular gas

Alexandre Rosas,^{1,2} Daniel ben-Avraham,³ and Katja Lindenberg^{1,4}

¹*Department of Chemistry and Biochemistry, University of California San Diego, La Jolla, California 92093-0340, USA*

²*Departamento de Física, Universidade Federal da Paraíba, João Pessoa, PB 58.059-970, Brazil*

³*Physics Department, Clarkson University, Potsdam New York 13699-5820, USA*

⁴*Institute for Nonlinear Science, University of California San Diego, La Jolla, California 92093-0340, USA*

(Received 16 April 2004; revised manuscript received 12 November 2004; published 17 March 2005)

We investigate the velocity relaxation of a viscous one-dimensional granular gas in which neither energy nor momentum is conserved in a collision. Of interest is the distribution of velocities in the gas as it cools, and the time dependence of the relaxation behavior. A Boltzmann equation of instantaneous binary collisions leads to a two-peaked distribution, as do numerical simulations of grains on a line. Of particular note is that in the presence of friction there is no inelastic collapse, so there is no need to invoke additional assumptions such as the quasielastic limit.

DOI: 10.1103/PhysRevE.71.032301

PACS number(s): 45.70.-n, 05.20.Dd, 05.70.Ln, 83.10.Rs

Velocity distributions in dilute granular gases are generically away from equilibrium because the collision processes are dissipative. Even in the dilute limit the velocities of different particles may be strongly correlated [1], and therefore the usual description in terms of single particle distribution functions may not be sufficient to determine all the properties of the granular gas. Nevertheless, even the single particle velocity distribution $P(v, t)$ contains important information, and is particularly interesting because in a granular gas it typically deviates from the Maxwell-Boltzmann distribution. These deviations have been a subject of intense interest in recent years [2,3] both in granular gases that achieve a steady state because external forcing balances the dissipative collisions among particles [4,5], and in unforced gases as they cool down [6–9].

Typically, gases equilibrate or achieve a steady state via collision processes defined by the conservation of energy and momentum [10]. In granular media the situation is more complex because *energy is not conserved* [11]. A particularly bothersome behavior induced by energy nonconservation is the so-called “inelastic collapse,” whereby the energy of the gas goes to zero in a finite time [6]. In the “quasielastic limit” the inelastic collapse is avoided, and one obtains nontrivial asymptotic velocity distributions [4,7,8,12] with features similar to the ones obtained in this paper.

Friction induces not only nonconservation of energy but also *nonconservation of momentum*, leading to interesting new relaxation behavior [5,13,14]. Our focus here is the effect of viscosity, and consequently, of momentum nonconservation, on the velocity distribution as a one-dimensional dilute granular gas cools down. The model consists of N grains on a line (or a circle, since we use periodic boundary conditions). The grains move freely except during collisions, governed by the Hertz potential, $V(\delta_{k,k+1}) = (a/n)|\delta_{k,k+1}^n$ when $\delta \leq 0$, and $V(\delta_{k,k+1}) = 0$ when $\delta > 0$. Here $\delta_{k,k+1} \equiv y_{k+1} - y_k$ is the relative displacement of granules k and $k+1$ from the positions in which they just touch each other without compression, and a is determined by Young’s modulus and Poisson’s ratio. The exponent n is $5/2$ for spheres, it is 2 for cylinders, and in general depends on geometry. Here we only consider

cylindrical grains, which leads to considerable simplification while still capturing important general features of the non-Maxwellian distributions.

A collision between real granules always involves some momentum and energy loss [15]. Here we take these losses during the collisions. Hence, *during a collision* of the granules k and $k+1$, their equations of motion are respectively

$$\begin{aligned} \frac{d^2 x_k}{dt^2} &= -\gamma \frac{dx_k}{dt} - (x_k - x_{k+1})^{n-1}, \\ \frac{d^2 x_{k+1}}{dt^2} &= -\gamma \frac{dx_{k+1}}{dt} + (x_k - x_{k+1})^{n-1}, \end{aligned} \quad (1)$$

where $x_k = y_k / C_n$, $t = \tilde{t} v_0 / C_n$ (\tilde{t} is the unscaled time), $\gamma = (\tilde{\gamma} / m v_0) C_n$, $C_n \equiv (m v_0^2 / a)^{1/n}$, and $\tilde{\gamma}$ is the friction coefficient. The arbitrary velocity v_0 sets the energy scale.

The one-sidedness of the potential leads to analytic complexities even in the dissipationless one-dimensional case [13,16], and even greater complexities in the presence of dissipation [13]. We explore the effects of friction at *low densities*, implemented via the assumption that the collisions are always *binary*, that is, that only two granules at a time are members of any collision event, and that at any moment there is at most one collision.

Our analytic starting point is the Boltzmann equation for binary collisions in a spatially uniform gas, which describes the rate of change of $P(v, t)$. If u_1 and u_2 are the initial velocities of a pair of particles just before a collision and u'_1 and u'_2 their velocities just after, then the Boltzmann equation is

$$\begin{aligned} \frac{\partial}{\partial t} P(v, t) &= \iint du_1 du_2 \theta(u_1 - u_2) (u_1 - u_2) P(u_1, t) P(u_2, t) \\ &\times [\delta(v - u'_1) + \delta(v - u'_2) - \delta(v - u_1) - \delta(v - u_2)]. \end{aligned} \quad (2)$$

Since the problem is one-dimensional, one can keep track of the precise conditions under which a collision between two particles of given velocities will or will not occur, and how




FIG. 1. Velocity distribution obtained using the simulation algorithm detailed in the text. From left to right and top to bottom, the panels correspond to time 0, 8000, 16000, 32000, 64000, and 128000 in the adimensional units used in this paper. The initial distribution is a symmetric exponential. Notice the change in the scales as time proceeds.

these events will change the distribution function. The particle on the left is that with initial velocity u_1 , and a collision takes place if and only if $u_1 > u_2$, a restriction enforced by the Heaviside theta function $\theta(y)$.

The duration of a collision is $\tau \equiv 2\pi/\sqrt{8-\gamma^2}$, after which the grains lose contact. For $n=2$ the collision time τ is independent of the initial condition. The velocities at the moment of separation are [13]

$$u'_{1,2} = u_1 \frac{\mu^2 \mp \mu}{2} + u_2 \frac{\mu^2 \pm \mu}{2}, \quad (3)$$

where $\mu \equiv e^{-\gamma\tau/2}$. The distance traveled by the granules during a collision is assumed to be negligible. For small damping γ this distance is $(u_1 + u_2)\pi/\sqrt{8}$. This is to be compared to the mean distance between particles, which can be made arbitrarily large by lowering the density. We also take the collision time as instantaneous. For small damping it is $\tau \approx \pi/\sqrt{2}$, to be compared with the typical mean free time of flight of a particle between collisions. With these approximations, the only role played by the viscosity is dictated by the collision rule (3). These assumptions might conceivably be problematic (but unlikely to be important at sufficiently low densities) for the most energetic particles that may travel a relatively long distance during a collision and have relatively short flight times.

In the long-time asymptotic limit we assume a scaling solution of the form $P(v,t) = F(v/\phi(t))/\phi(t)$. This scaling in Eq. (2) with Eq. (3) leads to $\phi(t) \sim t^{-1}$. Haff obtained this behavior (Haff's Law) in his classic paper on a hydrodynamic inelastic hard sphere model for a granular fluid [3].

We have not found an analytic solution of the Boltzmann equation. We therefore simulate the equation numerically and also implement a further approximation that leads to an analytic solution that we can compare with the numerical results.

We directly simulate the Boltzmann equation using the following algorithm. (1) Start with N grains whose velocities are independently assigned according to an initial distribution $P(v,0)$. (2) Choose one pair of grains with probability

proportional to the modulus of their relative velocity and let them collide, using the collision rule. (3) Increment time by twice the inverse of the modulus of the precollisional relative velocity. The factor of 2 accounts for picked pairs that do not collide, since our algorithm forces a collision at each step. (4) Iterate these three steps many times and for many samples.

In our simulations we took $N=100$ and averaged our results over 1000 samples. In Fig. 1 we show the resulting velocity distribution for $\gamma=0.9$ at different times. The initial symmetric and single peaked distribution develops two distinct peaks as it starts to collapse to the ultimate equilibrium distribution, a δ -function at $v=0$. The validity of the scaling solution is evident in the scaled rendition of the velocity distribution shown in Fig. 2. In the underlying simulation the initial distribution was a double peaked Gaussian, chosen because it converges quickly. We find the same asymptotic behavior for the initially exponential distribution.

A two-peaked velocity distribution has previously been




FIG. 2. Asymptotic behavior of the scaled velocity distribution. Different symbols stand for different times (1000×2^n with $n = 0, 1, \dots, 13$). The constant c in the figure is arbitrary and was chosen to facilitate comparison with Fig. 3.

found in a number of studies (see, e.g., [17]), in particular in the context of one-dimensional momentum-conserving granular gases that exhibit inelastic collapse, which occurs because particles whose inelastic collisions are described by a constant restitution coefficient r can collide infinitely often in a finite time. This drives their energy to zero in a finite time [6] unless N is sufficiently large. To avoid the collapse, it is customary to introduce extraneous assumptions such as the “quasielastic limit” $r \rightarrow 1$ and $N \rightarrow \infty$ in such a way as to always remain below a collapse threshold. In this limit, a double-peaked velocity distribution is also observed [6–8,17]. Another way to avoid the collapse is to impose elasticity (or stickyness) on collisions between particles with relative velocity smaller than a given arbitrary positive threshold [17].

An interesting hydrodynamic analysis based on numerical simulations that also arrives at a two-peaked velocity distribution (and that also deals with the issue of the inelastic collapse) leads to the conclusion that the usual hydrodynamic variables (density, velocity, and granular temperature) are not sufficient, and that an additional variable must also be included [8]. Benedetto *et al.* study the problem analytically (again in the quasielastic limit) and also arrive at the conclusion that the velocity distribution is non-Maxwellian. A very recent study of a two-dimensional collection of disks in a channel finds that the shape of the velocity distribution depends on the coefficient of restitution and on the Knudsen number [18]. They find a bimodal distribution when the Knudsen number is high but a unimodal distribution when it is low. We stress again that in the *presence of friction* there is no inelastic collapse, and the double-peaked distribution occurs regardless of the parameter values as long as there is friction.

While the Boltzmann equation does not appear amenable to analytic solution, we can formulate a simpler model that may incorporate its main features. In this model we have an ensemble of rings of size l each containing only two particles. In each ring, the dynamics is perfectly deterministic: the two balls keep colliding with each other, back and forth, moving toward each other with ever decreasing velocities. For each one ring, after n collisions, the velocities u_n and v_n are obtained by repeated application of the collision rule (3),

$$\begin{aligned} u_n &= \frac{\mu^{2n} + (-\mu)^n}{2} u_0 + \frac{\mu^{2n} - (-\mu)^n}{2} v_0, \\ v_n &= \frac{\mu^{2n} - (-\mu)^n}{2} u_0 + \frac{\mu^{2n} + (-\mu)^n}{2} v_0, \end{aligned} \quad (4)$$

where u_0 and v_0 are the initial velocities. Note that each velocity alternates from positive to negative as the particles move in one direction and then another in the ring. The distance traveled between two collisions is l , and the travel time is $l/|u-v|$, where u and v are their current velocities between collisions. Hence the time that has elapsed by the n th collision is

$$t_n = \frac{1}{|\Delta|} \sum_{k=0}^{n-1} \mu^{-k} = \frac{(1/|\Delta|\mu^n) - 1}{\lambda}, \quad (5)$$

where $\Delta = (u_0 - v_0)/l$ and $\lambda = [(1/\mu) - 1]$.

The sign alternation of the velocities in Eq. (4) is not important in the effort to understand the long time behavior of the distribution. We can approximate the velocities by envelope functions $A_+(t)$ and $A_-(t)$ and write $u_n = A_+(t)u_0 + A_-(t)v_0$, $v_n = A_-(t)u_0 + A_+(t)v_0$. The envelope functions can be found by solving Eq. (5) for μ^n , substituting this into Eq. (4) ignoring the minus signs in the $(-\mu)^n$ factors, and setting $t_n \equiv t$. One finds

$$A_{\pm}(t) = \frac{1}{2}(1 + |\Delta|\lambda t)^{-2} \pm \frac{1}{2}(1 + |\Delta|\lambda t)^{-1}. \quad (6)$$

Hence, if initially the particles had a velocity distribution $P(v, 0)$, then the velocity distribution as a function of time is

$$\begin{aligned} P(v, t) &= \int \int \delta(v - A_+(t)u_0 - A_-(t)v_0) \\ &\quad \times P(u_0, 0)P(v_0, 0)du_0dv_0. \end{aligned} \quad (7)$$

Asymptotically, $A_{\pm}(t) = -A_{\mp}(t) \sim 1/2|\Delta|\lambda t$, so that

$$P(v, t) = \int \int \delta\left(v - \frac{l \operatorname{sgn}(\Delta)}{2\lambda t}\right) P(u_0, 0)P(v_0, 0)du_0dv_0. \quad (8)$$

Therefore, $P(u, t)$ consists of two δ peaks, one at positive and one at negative velocities. If the initial normalized distribution is symmetric about zero velocity, then

$$P(v, t) \sim \frac{1}{2}\delta\left(v - \frac{l}{2\lambda t}\right) + \frac{1}{2}\delta\left(v + \frac{l}{2\lambda t}\right). \quad (9)$$

The peaks thus move toward the final velocity $v=0$ as $1/t$. This time dependence is in agreement with the Boltzmann equation analysis.

The two δ peaks here reflect the fact that the magnitude of the velocity difference between the colliding particles eventually becomes independent of the magnitude of the initial velocity difference. Clearly, in the Boltzmann equation this is not quite the case and the peaks have a finite width as they converge. However, this width decreases in time as $1/t$, approaching the behavior of the two-particle ring model asymptotically.

In order to assess the validity of the Boltzmann equation for the viscous granule problem, and to get a sense of the possible effects of the spatial distribution of granules ignored in that formulation, we have carried out numerical simulations of a full chain of 10000 viscous particles. Our collision rules are as indicated in Eq. (3), and, as before, we assume collisions to be instantaneous, but now we place the granules on a line and keep track of their positions so that spatial inhomogeneities can occur if the system is so inclined. The particle density is 10^{-3} with the particles initially distributed uniformly. The initial velocity distribution is a symmetric exponential, and $\gamma=0.9$. As time proceeds, the initial single peak splits into two peaks which move inward and become




FIG. 3. Asymptotic behavior of the scaled velocity distribution. Different symbols stand for different times (1000×2^n with $n = 7, 8, \dots, 13$). The constant c in the figure is arbitrary.

narrower. Both the inward motion of the peaks and the width of the peaks change as $1/t$, as in the Boltzmann approximation. We tested the scaling hypothesis, and in Fig. 3 we show the results averaged over 100 simulations. Clearly, the scaling works quite well.

Comparing the velocity distribution for the Boltzmann approximation (Fig. 1) and the simulations in a line, we find that the Boltzmann model converges to the scaling distribution much more rapidly. Initially the simulation results exhibit the same behavior as the Boltzmann model, but for longer times the behavior of the two distributions for small

velocities begin to differ. In the Boltzmann case, the probability of finding a slow granule is much greater than in the simulation.

We conclude that the Boltzmann problem in which spatial dependences are disregarded captures many essential features of the velocity relaxation in a viscous granular chain, the most important being the appearance of two peaks in the velocity distribution. The Boltzmann problem and even the simpler two-particle simplification of the problem also capture the time dependence of convergence of the two peaks into a single one. The slow ($1/t$) convergence is due to the fact that the collision rate slows down as the gas cools. The Boltzmann approximation does not correctly capture the late time distribution of the slowest particles. This is probably due to spatial correlations that have been ignored.

As mentioned earlier, one-dimensional momentum-conserving granular gases may exhibit “inelastic collapse” [6–8]. To avoid this, the “quasistatic limit” or other *ad hoc* assumptions are frequently invoked, and in this limit, a double-peaked velocity distribution is also observed. In the presence of friction there is no inelastic collapse, and we always observe a double-peaked distribution. A comparison between those results and ours requires an understanding of spatial correlations [19].

The authors are grateful to Andrés Santos for his thorough and critical reading of this manuscript. This work was supported by the Engineering Research Program of the Office of Basic Energy Sciences at the U.S. Department of Energy under Grant No. DE-FG03-86ER13606 (K.L.) and by NSF Grant No. PHY-0140094 (D.b.A.).

-
- [1] S. J. Moon, M. D. Shattuck, and J. B. Swift, *Phys. Rev. E* **64**, 031303 (2001).
- [2] S. Warr, J. M. Huntley, and George T. H. Jacques, *Phys. Rev. E* **52**, 5583 (1995); J. S. Olafsen and J. S. Urbach, *Phys. Rev. Lett.* **81**, 4369 (1998); W. Losert, D. G. W. Cooper, J. Delour, A. Kudrolli, and J. P. Gollub, *Chaos* **9**, 682 (1999); A. Kudrolli and J. Henry, *Phys. Rev. E* **62**, R1489 (2000); F. Rouyer and N. Menon, *Phys. Rev. Lett.* **85**, 3676 (2000); I. Goldhirsch and G. Zanetti, *ibid.* **70**, 1619 (1993); A. Puglisi *et al.*, *ibid.* **81**, 3848 (1998).
- [3] P. K. Haff, *J. Fluid Mech.* **134**, 401 (1983).
- [4] E. L. Grossman and B. Roman, *Phys. Fluids* **8**, 3218 (1996); A. Barrat *et al.*, *J. Phys. A* **35**, 463 (2002).
- [5] S. J. Moon, J. B. Swift, and H. L. Swinney, *Phys. Rev. E* **69**, 011301 (2004).
- [6] S. McNamara and W. R. Young, *Phys. Fluids A* **4**, 496 (1992).
- [7] S. McNamara and W. R. Young, *Phys. Fluids A* **5**, 34 (1993).
- [8] N. Sela and I. Goldhirsch, *Phys. Fluids* **7**, 507 (1995).
- [9] A. Baldassarri, U. M. B. Marconi, and A. Puglisi, *Europhys. Lett.* **58**, 14 (2002); P. L. Krapivsky and E. Ben-Naim, *J. Phys. A* **35**, L147 (2002); M. H. Ernst and R. Brito, *Europhys. Lett.* **58**, 182 (2002); A. V. Bobylev and C. Cercignani, *J. Stat. Phys.* **106**, 547 (2002).
- [10] J. C. Maxwell, *Philos. Trans. R. Soc. London* **157**, 49 (1867).
- [11] *Granular Gases*, edited by T. Pöschel and S. Luding (Springer, Berlin, 2000).
- [12] D. Benedetto, E. Caglioti, J. A. Carrillo, and M. Pulvirenti, *J. Stat. Phys.* **91**, 979 (1998).
- [13] A. Rosas, J. Buceta, and K. Lindenberg, *Phys. Rev. E* **68**, 021303 (2003); A. Rosas and K. Lindenberg, *ibid.* **68**, 041304 (2003).
- [14] D. ben-Avraham, E. Ben-Naim, K. Lindenberg, and A. Rosas, *Phys. Rev. E* **68**, 050103(R) (2003).
- [15] J. Duran, *Sand, Powders and Grains: An Introduction to The Physics of Granular Materials* (Springer-Verlag, New York, 2000).
- [16] V. F. Nesterenko, *J. Appl. Mech. Tech. Phys.* **5**, 733 (1983); E. J. Hinch and S. Saint-Jean, *Proc. R. Soc. London, Ser. A* **455**, 3201 (1999).
- [17] A. Barrat, T. Biben, Z. Rácz, E. Trizac, and F. van Wijland, *J. Phys. A* **35**, 463 (2002).
- [18] M. Bose and V. Kumaran, *Phys. Rev. E* **69**, 061301 (2004).
- [19] A. Rosas, D. ben-Avraham, and K. Lindenberg (unpublished).