

Driven classical diffusion with strong correlated disorder

Leonid P. Pryadko and Jing-Xian Lin

Department of Physics, University of California, Riverside, California 92521, USA

(Received 25 August 2004; revised manuscript received 18 May 2005; published 19 July 2005)

For driven classical diffusion quenched by a strong potential disorder $V(x)$, we identify a prominent cross-over regime between the regimes of very small and very large driving forces F , where the corresponding mobility values $\mu(F)$ differ exponentially. For disorder with power-law correlations at large distances $\langle V(x)V(y) \rangle \sim |x-y|^{-n}$, $n > 0$, the crossover is characterized by power-law dependence of the logarithm of $\mu(F)$ on the driving force $\ln \mu(F) \sim F^{n/(n+1)}$. For finite-range disorder (formally, $n = \infty$), the corresponding dependence is linear (“logarithmic susceptibility”).

DOI: [10.1103/PhysRevE.72.011108](https://doi.org/10.1103/PhysRevE.72.011108)

PACS number(s): 05.40.Jc, 72.20.Ht, 72.80.Ng

It is well known that disorder can dramatically modify transport properties of materials. The effect is present even for classical diffusion [1,2] in one dimension (1D). Here, the renormalization of the carrier mobility μ can be thought of as the inverse of the average Boltzmann factor between the deepest well and the highest hill of the potential energy profile $V(x)$; it is exponential in the ratio $\langle V^2 \rangle / T^2$ [see Eq. (7) below]. Such a renormalization is suppressed if a sufficiently large driving force is applied, in which case the carriers just slide downhill uniformly.

In this paper we study the crossover between these two regimes as a function of the driving force F . In particular, for a strong disorder potential with power-law correlations at large distances [Eqs. (5) and (11)], there is a wide intermediate range of values of the driving force F where the logarithm of the effective mobility scales as a nontrivial power of F (Fig. 1). With Coulomb-like ($n=1$) or longer-range correlations, the effective mobility is a singular function of F already at $F=0$; the applicability region of the linear transport is essentially absent (Fig. 2). The crossover regime remains unchanged in the presence of weak interaction between the particles, introduced here at the level of the self-consistent Poisson equation which describes a Debye-like screening in the presence of strong disorder. We also show that the effective field-dependent mobility $\mu(F)$ is self-averaging, and estimate the relevant length scale.

Over so many years after Kramers’ pioneering work [1], the problem of driven diffusion in the presence of disorder has received a lot of attention [1,2]. In particular, the driven diffusion with correlated disorder was discussed [3,4] by Vinokur *et al.*, who, however, considered only the cases of short-range disorder, and a correlated disorder with infinite local correlations whose phenomenology resembles glass dynamics.

Single-particle diffusion. In 1D single-particle diffusion is described by the Smoluchovskiy equation

$$\eta_0 \dot{x} + \partial_x U(x) = f(t), \quad (1)$$

where $U(x)$ is the external potential and $f(t)$ is the thermal force with the correlator $f(t)f(t') = 2T\eta_0\delta(t-t')$. The usual assumption is that the bare viscous friction coefficient η_0 is determined by fast scattering events off phonons, short-range

disorder, etc. The potential $U(x)$ in Eq. (1) is thus the part of the overall potential remaining after averaging over some distance scale; its precise value depends on the specific physical system. If we assume the carriers have charge e , we can also define the bare mobility in the absence of disorder, $U(x) = -eEx$:

$$\mu_0 \equiv \bar{\dot{x}}/E = e/\eta_0. \quad (2)$$

Equation (1) can be also rewritten as the transport equation for the average particle density $n \equiv n(x, t)$ and the particle current $j \equiv j(x, t)$,

$$\partial_t n + \partial_x j = 0, \quad j = -D_0 \partial_x n - \eta_0^{-1} n \partial_x U(x); \quad (3)$$

the diffusion constant D_0 is related to the viscous friction coefficient η_0 by the Einstein relation $D_0 = T/\eta_0$.

The usual transport problem corresponds to stationary diffusion in the presence of a random Gaussian potential $V(x)$ and a constant driving force F , with the total potential energy $U(x) = V(x) - Fx$. In the case of a periodic potential $V(x) = V(x+a)$, the stationary solution [1,2] corresponds to a con-




FIG. 1. (Color online) Logarithm of the effective mobility renormalization $\ln(\mu(F)/\mu(0))$ [Eqs. (6) and (7)] with the correlation function $g(x) = (1+x^2/\xi_1^2)^{-n/2}$ for strong disorder ($V_0/T=5$) and large driving forces F . Dotted lines guide the eye with the slope of the intermediate asymptote (14). The thin solid line is the analytic result (15) for $n=1$.




FIG. 2. (Color online) As in Fig. 1 but for small driving forces F . Thin solid lines indicate the nonlinear correction calculated analytically: linear in F for $n > 1$, proportional to F^n for $n < 1$, and Eq. (15) for $n = 1$.

stant average current j with periodic boundary conditions $n(a) = n(0)$. Then, by normalizing the density profile $n(x)$ over the period, we can use the current to define the average drift velocity, $\bar{v} = j/\bar{n} = ja$, as well as the effective viscous friction coefficient $\eta \equiv F/\bar{v} = F/ja$,

$$\eta = \frac{\eta_0 F}{aT} \int_0^a dx \int_x^{x+a} dx' \frac{e^{[V(x') - V(x) + F(x-x')]/T}}{1 - e^{-Fa/T}}. \quad (4)$$

Both integrations extend over the entire period, and, as it often happens in classical transport phenomena, for a sufficiently large a the effect of disorder becomes self-averaging [the required size can be large, see Eq. (20) below]. In such cases we can replace Eq. (4) by its average over disorder. We assume that the disorder distribution is Gaussian with the correlations

$$\langle V(x) \rangle = 0, \quad \langle V(x)V(x') \rangle = V_0^2 g(x-x'), \quad (5)$$

where the local r.m.s. value V_0 is taken as the measure of the disorder strength and the correlation function $g(x)$ is defined so that $g(0) \equiv 1$. Then, in the thermodynamical limit $a \rightarrow \infty$, the effective mobility $\mu(F) \equiv e/\langle \eta \rangle$, and [3,5]

$$\frac{\mu_0}{\mu(F)} = e^{V_0^2/T^2} \frac{F}{T} \int_0^\infty dx e^{-Fx/T} e^{-g(x)V_0^2/T^2}, \quad F > 0. \quad (6)$$

In the small- F limit (but at the same time $aF \gg T$), this gives the usual linear response result [6]

$$\langle \eta \rangle / \eta_0 = \mu_0 / \mu(0) = \exp(V_0^2/T^2), \quad (7)$$

which can be understood as the average of the activation exponent of the difference between the highest maximum and the lowest minimum of the potential. Generically, these would be in different parts of the sample and so the friction renormalization factors onto a product of the two averages $\langle e^{V/T} \rangle \langle e^{-V/T} \rangle = \exp(V_0^2/T^2)$, independent of the form of the correlation function $g(x)$.

Similarly, the stationary limit of the dynamical perturbation theory [7] is restored by expanding Eq. (6) in powers of V_0/T and integrating the result by parts

$$\frac{\mu_0}{\mu(F)} = 1 - \frac{V_0^2}{T^2} \int_0^\infty dx e^{-Fx/T} g'(x) + \mathcal{O}(V_0^4/T^4). \quad (8)$$

The disorder correlation function $g(x)$ is expected to decrease with x , remaining substantially different from zero over the distance of the order of the appropriate correlation length ξ . It is clear from the weak-disorder expression (8) that for such finite-range disorder there is a distinct crossover force $F_\xi \sim T/\xi$: while $F \lesssim F_\xi$ have relatively little effect on the mobility, larger values of F begin to suppress the effect of disorder as large-scale potential valleys and hills gradually disappear.

To analyze an analogous effect for *strong* finite-range disorder, note that for large V_0/T , $\Delta_x \equiv e^{-g(x)V_0^2/T^2}$ is exponentially small for $x \lesssim \xi$. This effectively limits the integration in Eq. (6) to the region $x \gtrsim \xi$, so that

$$\ln(\mu(F)/\mu(0)) \sim F\tilde{\xi}/T, \quad (9)$$

where $\tilde{\xi} \approx \xi$ up to a logarithmic correction. Such a dependence on the applied field is analogous to the logarithmic susceptibility [8] typical for systems with activated transport. Here it can be understood as the diffusion limited by far-spaced maxima of the potential, with the particles concentrated in the intermediate low minima; the applied driving force F effectively reduces the energy gap between the minima and the maxima and therefore has an exponential effect on the mobility.

The expression (9) is valid qualitatively as long as the effect of the disorder remains large, $\mu_0/\mu(F) \gg 1$. The precise value of $\tilde{\xi}$ and the prefactor depends on the details of the disorder correlation function. For example, with the exponential correlation function $g(x) = \exp(-x/\xi)$, the integration (6) can be done exactly [3] in terms of the incomplete gamma function; the asymptotic form for $V_0^2/T^2 \gg \max(1, \chi \equiv F\xi/T)$ is [9]

$$\frac{\mu(F)}{\mu(0)} = \frac{(V_0/T)^{2\chi}}{\Gamma(\chi+1)} \xrightarrow{\chi \gg 1} (2\pi\chi)^{-1/2} e^{F\tilde{\xi}/T}, \quad (10)$$

where $\tilde{\xi} = \xi[1 + \ln(V_0^2/(TF\xi))]$.

The disorder-induced transport nonlinearity becomes even more pronounced for long-range potentials, e.g., those with power-law correlations at large distances. With long-range correlations the far-spaced maxima and minima of the potential are not entirely independent and, therefore, even a weak driving force may have a noticeable effect. Specifically, consider a correlation function with the asymptotic form

$$g(x) = (\xi_1/x)^n, \quad x > x_{\min} \gg \xi, \xi_1. \quad (11)$$

With strong enough disorder [$g(x_{\min})V_0^2/T^2 \gtrsim 1$] the integral (6) will be determined by large x , in which case the expression can be rewritten approximately as

$$\frac{\mu(0)}{\mu(F)} \approx \mathcal{I}_n(\alpha), \quad \mathcal{I}_n(\alpha) \equiv \int_0^\infty dx e^{-x-\alpha x^n}, \quad (12)$$

where $\alpha \equiv (F\xi_1/T)^n V_0^2/T^2$ can be small or large within the strong-disorder domain where Eq. (12) is applicable. For sufficiently large α , the integration can be done using the Gaussian approximation around the maximum at $x_0 = (n\alpha)^{1/(n+1)}$,

$$\mathcal{I}_n(\alpha) = \left(\frac{2\pi x_0}{n+1} \right)^{1/2} e^{-x_0(1+1/n)}, \quad \alpha \gg 1. \quad (13)$$

As a result, logarithm of $\mu(F)$ is proportional to a power

$$\ln(\mu(F)/\mu(0)) \sim (F\tilde{\xi}_1/T)^{n/(n+1)}, \quad (14)$$

with a *large* temperature-dependent length parameter $\tilde{\xi}_1 = C_n \xi_1 (V_0/T)^{2/n}$, $C_n \equiv (n+1)^{1+1/n}/n$ [see Eq. (9)]. For small $\alpha \ll 1$, the integration (12) can be done perturbatively in powers of $\alpha \propto F^n V_0^2$ for $n < 1$ or, using the identity $\mathcal{I}_n(\alpha) = \mathcal{I}_{1/n}(\alpha^{1/n})$, in powers of $\alpha^{1/n} \propto F V_0^{2/n}$ for $n > 1$. For Coulomb disorder, $n=1$, the result is expressed in terms of the Macdonald function

$$\mu(0)/\mu(F)|_{n=1} \approx \mathcal{I}_1(\alpha) = 2\alpha^{1/2} K_1(2\alpha^{1/2}). \quad (15)$$

For very small $\alpha \ll 1$, $\mathcal{I}_1(\alpha) \approx 1 - \alpha \ln(e^{1-2\gamma}/\alpha)$, where $\gamma \approx 0.577$ is the Euler's constant; the correction is linear in F up to a logarithm. Clearly, for strong Coulomb or longer-range disorder $n \leq 1$, the mobility $\mu(F)$ is a singular function of the driving force at $F=0$; the linear-transport regime is essentially absent. These asymptotics are illustrated in Figs. 1 and 2 for a model form of the disorder correlation function $g(x) = (1+x^2/\xi_1^2)^{-n/2}$.

Self-averaging. Our conclusions on the scaling of mobility in strongly disordered diffusive 1D systems are based on the average, Eq. (6). To analyze the sample-to-sample fluctuations, consider the irreducible average $\langle\langle \eta^2 \rangle\rangle \equiv \langle \eta^2 \rangle - \langle \eta \rangle^2$ of the effective friction η [Eq. (4)],

$$\begin{aligned} \frac{\langle\langle \eta^2 \rangle\rangle}{\eta_0^2} &= e^{2V_0^2/T^2} \frac{F^2}{aT^2} \int_0^a dx e^{-Fx/T} \Delta_x \int_0^a dy e^{-Fy/T} \Delta_y \\ &\times \int_0^a dz (\Delta_{z+(x+y)/2} \Delta_{z-(x+y)/2} \Delta_{z+(x-y)/2}^{-1} \Delta_{z-(x-y)/2}^{-1} - 1), \end{aligned} \quad (16)$$

where the correlator $\Delta_z \equiv e^{-g(z)V_0^2/T^2}$ is periodic under $z \rightarrow z+a$. Note that both Δ_z and Δ_z^{-1} enter Eq. (16). Therefore, unlike in the average (6), both short- and long-distance disorder correlations affect the variance of η .

For weak disorder, the expansion of Eq. (16) in powers of $V_0/T \ll 1$ begins with the quartic term

$$\begin{aligned} \frac{\langle\langle \eta^2 \rangle\rangle}{\eta_0^2} &= \frac{V_0^4 F^2}{T^4 a T^2} \int_0^a dx e^{-Fx/T} \int_0^a dy e^{-Fy/T} \int_0^a dz \\ &\times g_z (2g_z + g_{z+x+y} + g_{z+x-y} - 2g_{z+x} - 2g_{z+y}) \\ &= \frac{2V_0^4}{aT^4} \int_0^a dz g_z \int_0^a du e^{-Fu/T} (u g''_{z+u} - g'_{z+u}). \end{aligned} \quad (17)$$

The integration is simplified in the limits of weak and large F ; the combined result is

$$\frac{\langle\langle \eta^2 \rangle\rangle}{\eta_0^2} = \frac{V_0^4}{2aT^4} \min(4\xi_2, TF), \quad \xi_2 \equiv \int_0^\infty dx g^2(x). \quad (18)$$

Here, ξ_2 is yet another correlation length, finite for short-range disorder and for long-range disorder with $n > 1/2$. Clearly, for weak disorder the variation of η is small and it is further reduced with increasing system size a .

The situation is different for strong disorder $V_0 \gg T$, which causes an exponential renormalization of η ; large fluctuations are also expected. In this case Δ_x^{-1} has a prominent maximum at the origin. Consequently, the integral (16) gets an exponentially large contribution from a vicinity of the point $z=0$, $x=y$. Using the steepest descent method, we obtain

$$\frac{\langle\langle \eta^2 \rangle\rangle}{\eta_0^2} = \frac{2\pi F^2 e^{4V_0^2/T^2}}{aV_0^2} \int_0^\infty du \frac{e^{-2Fu/T} e^{-4g(u)V_0^2/T^2}}{g''(u) - g''(0)}. \quad (19)$$

For not exceedingly large F the result is determined by values of u away from the origin. Then, the denominator can be replaced by a constant [10] $-g''(0) \equiv 2/\xi_0^2$, and the integral acquires precisely the form of Eq. (6). Generally,

$$\frac{\delta\mu^2}{\mu^2} \approx \frac{\langle\langle \eta^2 \rangle\rangle}{\langle \eta \rangle^2} = C \frac{\pi FT \xi_0^2}{2aV_0^2} e^{2V_0^2/T^2}; \quad (20)$$

while F in the prefactor can be small, it is assumed to be large on the scale of the system size, $Fa/T \gg 1$. The normalization in Eq. (20) is chosen so that for short-range disorder $C \approx 1$ [see Eq. (9)]. For power-law correlation tail $C = \mathcal{I}_n(2^{n+2}\alpha)/[\mathcal{I}_n(\alpha)]^2$, with $\alpha \equiv (F\xi_1/T)^n V_0^2/T^2$ [see Eqs. (11) and (12)]. Overall, we conclude that the effective mobility is a self-averaging quantity in the thermodynamical limit. Of course, the required system size can be large if the fluctuations are strong.

We verified these conclusions by simulating diffusion in a 1D short-range random potential (not shown). With periodic boundary conditions the viscous friction (4) could be obtained by averaging the time t it takes a particle to travel over one period $\Delta x = a$. As expected, with increasing a , the corresponding disorder average $\langle t \rangle / a$ approached the inverse of the average drift velocity.

Stationary diffusion with weak interaction. The considered problem differs from the canonical Kramers problem [1,2] of over-the-barrier transport, where it is the dynamical equilibrium that establishes the exponentially different particle numbers in the “reservoirs” on the two sides of the barrier. Here, we consider a situation corresponding to a typical resistivity measurement in a macroscopic sample where

the total number of particles does not change with the applied field. Then, the macroscopic current would be determined solely by the average drift velocity. It is important that the quantity is self-averaging, as the explicit disorder averaging would not be necessary for large enough samples.

In finite-size systems such a situation arises naturally, e.g., when diffusing particles are charged and the electroneutrality condition needs to be satisfied. The simplest case corresponds to the Debye mean-field screening, where the potential in Eq. (3) is modified by the self-consistent potential $U(x) \rightarrow U(x) + e\varphi(x)$. Specifically, we consider a 1D Poisson equation

$$\varphi'' = -4\pi e(n - \bar{n}), \quad (21)$$

as would be appropriate for diffusion in a 3D system with 1D modulation (layered disorder), a parallel bunch of identical DNA molecules, or electrostatically coupled identical ionic cell channels.

In the static equilibrium $F=0$, the coupled Eqs. (3) and (21) correspond to the nonlinear screening problem; with weak disorder, $V_0 \ll T$, the Debye screening length is κ^{-1} , $\kappa^2 \equiv 4\pi\bar{n}e^2/T$. The linearized self-consistent screening problem can be also solved with the nonzero driving force $F > 0$; the solution involves two screening parameters $s_{\pm} = (\kappa^2 + f^2/4)^{1/2} \pm f/2$, where $f \equiv F/T \approx j/(D_0\bar{n}) \equiv \lambda$. Clearly, in the weak-interaction limit $f \gg \kappa$, the shorter screening length $s_+^{-1} \approx f^{-1} = T/F$ is determined by the driving force, while the longer one diverges, $s_-^{-1} \approx f/\kappa^2$.

With the driving force and a strong disorder, the problem is forbiddingly complicated. However, if the interaction is weak, the additional potential would be small, and the screening equations can be linearized. To this end, it is convenient to eliminate the density n from Eqs. (3) and (21) and write the self-consistent equation for the scaled gradient of the screening potential $\varepsilon \equiv e\varphi'(x)/T$,

$$e^{-V/T} \left(\frac{d}{dx} - s_+ \right) e^{V/T} (\varepsilon' + s_- \varepsilon) = \tau - \left(\varepsilon' - s_- \frac{V'}{T} \right) \varepsilon,$$

where $\tau \equiv \kappa^2(V'/T + \lambda - f)$. For weak interaction, the last term in the right-hand side is quadratic in small κ^2 and can be ignored. The remaining equation should be solved for ε

with zero boundary conditions at infinity. The relation between the driving force $F \equiv fT$ and the diffusion current $j \equiv \lambda\bar{n}D_0$ is established from the self-consistency condition that the screening does not modify the net driving field. Equivalently, the disorder averaged $\langle \varepsilon \rangle = 0$. Approximating $s_+ \approx f$, after some algebra we again arrive at Eq. (6).

Conclusions. We analyzed the stationary 1D problem of a driven diffusion in the presence of a random disorder potential. For large systems and/or in the presence of an interaction fixing the number of particles transport should be described in terms of the effective mobility μ . Strong disorder significantly reduces the mobility and leads to its nontrivial scaling as a function of the driving field F and the temperature. With finite-range disorder, the dependence $\mu(F)$ can be described in terms of the logarithmic susceptibility (9), a generic form for problems with activated transport. For a strongly driven system with power-law disorder correlation tail at large distances, the logarithm of the mobility scales as a power of the driving force (14). The disorder effect is especially pronounced for Coulomb-like correlations [see Eq. (15)]: the field-dependent correction to mobility is singular already at $F=0$.

The results for field-dependent crossover with strong disorder (low temperature) are generic and apply to other systems/dimensions where the mobility is strongly suppressed by disorder. Indeed, e.g., for the classical diffusion with zero-mean Gaussian disorder potential in two dimensions, the effective small-field mobility can be found from the duality arguments [2(c),11] for the equivalent problem of effective conductance of a 2D system with the local conductivity proportional to equilibrium particle density $n \propto e^{-V/T}$, $\mu_0/\mu^{2D}(0) = \langle e^{V/T} \rangle = e^{V_0^2/2T^2}$. The exponent represents activation to the percolation level at zero energy; it is large for large V_0/T . An applied field lowers the activation energy by $\sim F\xi$, which again results in the logarithmic susceptibility form (9). With the long-range power-law disorder in 2D, a more general crossover dependence similar to Eq. (14) is also expected.

The authors are grateful to Mark Dykman and Chandra Varma for encouragement and illuminating discussions.

-
- [1] H. A. Kramers, *Physica* (Amsterdam) **VII**, 284 (1940).
 [2] (a) H. Risken, *The Fokker-Planck Equation* (Springer, Berlin, 1984); (b) P. Hänggi, P. Talkner, and M. Borkovec, *Rev. Mod. Phys.* **62**, 251 (1990); (c) J.-P. Bouchaud and A. Georges, *Phys. Rep.* **195**, 127 (1990).
 [3] P. Le Doussal and V. M. Vinokur, *Physica C* **254**, 63 (1995).
 [4] A. V. Lopatin and V. M. Vinokur, *Phys. Rev. Lett.* **86**, 1817 (2001).
 [5] B. Derrida, *J. Stat. Phys.* **31**, 433 (1983); J. P. Bouchaud *et al.*, *Ann. Phys. (N.Y.)* **201**, 285 (1990); D. A. Gorokhov and G. Blatter, *Phys. Rev. B* **58**, 213 (1998).
 [6] R. Festa and E. Galleani d'Agliano, *Physica A* **90A**, 229 (1978); S. Alexander *et al.*, *Rev. Mod. Phys.* **53**, 175 (1981); K. Golden, S. Goldstein, and J. L. Lebowitz, *Phys. Rev. Lett.* **55**, 2629 (1985); T. Schneider, A. Politi, and M. P. Sørensen, *Phys. Rev. A* **37**, 948 (1988).
 [7] A. I. Larkin and Y. N. Ovchinnikov, *Zh. Eksp. Teor. Fiz.* **65**, 704 (1973) [*Sov. Phys. JETP* **38**, 854 (1974)]; V. M. Vinokur *et al.*, *Phys. Rev. Lett.* **65**, 259 (1990).
 [8] V. N. Smelyanskiy *et al.*, *Phys. Rev. Lett.* **79**, 3113 (1997).
 [9] The full expression in Ref. [3] is misprinted.
 [10] Here ξ_0 is a correlation length describing the short-distance properties of the disorder potential; it may be quite different from the length ξ relevant at large distances.
 [11] A. M. Dykhne, *Zh. Eksp. Teor. Fiz.* **59**, 110 (1970) [*Sov.-Phys. JETP* **32**, 63 (1971)]; K. S. Mendelson, *J. Appl. Phys.* **46**, 917 (1975).