

# Pfaffian solution of a dimer-monomer problem: Single monomer on the boundary

F. Y. Wu

*Department of Physics, Northeastern University, Boston, Massachusetts 02115, USA*

(Received 23 June 2006; published 18 August 2006)

We consider the dimer-monomer problem for the rectangular lattice. By mapping the problem into one of close-packed dimers on an extended lattice, we rederive the Tzeng-Wu solution for a single monomer on the boundary by evaluating a Pfaffian. We also clarify the mathematical content of the Tzeng-Wu solution by identifying it as the product of the nonzero eigenvalues of the Kasteleyn matrix.

DOI: [10.1103/PhysRevE.74.020104](https://doi.org/10.1103/PhysRevE.74.020104)

PACS number(s): 05.50.+q, 04.20.Jb, 02.10.Ox

## I. INTRODUCTION

An outstanding unsolved problem in lattice statistics is the dimer-monomer problem. While it is known [1] that the dimer-monomer system does not exhibit a phase transition, there have been only limited closed-form results. The case of close-packed dimers on planar lattices has been solved by Kasteleyn [2] and by Temperley and Fisher [3,4], and the solution has been extended to nonorientable surfaces [5,6]. But the general dimer-monomer problem has proven to be intractable [7].

In 1974 Temperley [8] pointed out a bijection between configurations of a single monomer on the boundary of a planar lattice and spanning trees on a related lattice. The bijection was used in [8] to explain why enumerations of close-packed dimers and spanning trees on square lattices yield the same Catalan constant. More recently Tzeng and Wu [9] made further use of the Temperley bijection to obtain the closed-form generating function for a single monomer on the boundary. The derivation is, however, indirect since it makes use of the Temperley bijection which obscures the underlining mathematics of the closed-form solution.

Motivated by the Tzeng-Wu result, there has been renewed interest in the general dimer-monomer problem. In a series of papers Kong [10–12] has studied numerical enumerations of such configurations on  $m \times n$  rectangular lattices for varying  $m, n$ , and extracted finite-size correction terms for the single-monomer [10] and general monomer-dimer [11,12] problems. Of particular interest is the finding [10] that in the case of a single monomer the enumeration exhibits a regular pattern similar to that found in the Kasteleyn solution of close-packed dimers. This suggests that the general single-monomer problem might be soluble.

As a first step toward finding that solution it is necessary to have an alternate and direct derivation of the Tzeng-Wu solution without recourse to the Temperley bijection. Here we present such a derivation. Our approach points the way to a possible extension toward the general single-monomer problem. It also shows that, apart from an overall constant, the Tzeng-Wu solution is given by the square root of the product of the nonzero eigenvalues of the Kasteleyn matrix, and thus clarifies its underlining mathematics.

## II. THE SINGLE-MONOMER PROBLEM

Consider a rectangular lattice  $\mathcal{L}$  consisting of an array of  $M$  rows and  $N$  columns, where both  $M$  and  $N$  are odd. The

lattice consists of two sublattices  $A$  and  $B$ . Since the total number of sites  $MN$  is odd, the four corner sites belong to the same sublattice, say,  $A$  and there is one more  $A$  than  $B$  sites. The lattice can therefore be completely covered by dimers if one  $A$  site is left open. The open  $A$  site can be regarded as a monomer.

Assign non-negative weights  $x$  and  $y$ , respectively, to horizontal and vertical dimers. When the monomer is on the boundary, Tzeng and Wu [9] obtained the following closed-form expression for the generating function:

$$G(x, y) = x^{(M-1)/2} y^{(N-1)/2} \times \prod_{m=1}^{(M-1)/2} \prod_{n=1}^{(N-1)/2} \left( 4x^2 \cos^2 \frac{m\pi}{M+1} + 4y^2 \cos^2 \frac{n\pi}{N+1} \right). \quad (1)$$

This result is independent of the location of the monomer provided that it is an  $A$  site on the boundary.

We rederive the result (1) using a formulation that is applicable to any dimer-monomer problem. We first expand  $\mathcal{L}$  into an extended lattice  $\mathcal{L}'$  constructed by connecting each site occupied by a monomer to a new added site, and then consider close-packed dimers on  $\mathcal{L}'$ . Since the newly added sites are all of degree 1, all edges originating from the new sites must be covered by dimers. Consequently, the dimer-monomer problem on  $\mathcal{L}$  (with fixed monomer sites) is mapped to a close-packed dimer problem on  $\mathcal{L}'$ , which can be treated by standard means.

We use the Kasteleyn method [2] to treat the latter problem. Returning to the single-monomer problem let the boundary monomer be at site  $s_0 = (1, n)$  as demonstrated in Fig. 1(a). The site  $s_0$  is connected to a new site  $s'$  by an edge with weight 1 as shown in Fig. 1(b). To enumerate close-packed dimers on  $\mathcal{L}'$  using the Kasteleyn approach, we need to orient, and associate phase factors with, edges so that all terms in the resulting Pfaffian yield the same sign.

A convenient choice of orientation and assignment of phase factors is the one suggested by Wu [13]. While Wu considered the case of  $MN$  even, the consideration can be extended to the present case. Orient all horizontal (vertical) edges in the same direction and the new edge from  $s'$  to  $s_0$ , and introduce a phase factor  $i$  to all horizontal edges as shown in Fig. 1(b). Then all terms in the Pfaffian assume the same sign. To prove this assertion it suffices to show that a typical term in the Pfaffian associated with a dimer configu-




FIG. 1. (a) A dimer-monomer configuration on a  $5 \times 5$  lattice  $\mathcal{L}$  with a single monomer at  $s_0=(1,3)$ . (b) The extended lattice  $\mathcal{L}'$  with edge orientation and a phase factor  $i$  to horizontal edges. (c) The reference dimer configuration  $C_0$ .

ration  $C$  has the same sign as the term associated with a reference configuration  $C_0$ . For  $C_0$  we choose the configuration shown in Fig. 1(c), in which horizontal dimers are placed in the first row with vertical dimers covering the rest of the lattice. Then  $C$  and  $C_0$  assume the same sign.

The simplest way to verify the last statement is to start from a configuration in which every heavy edge in  $C_0$  shown in Fig. 1(c) is occupied by two dimers, and view each of the doubly occupied dimers as a polygon of two edges. Then the “transposition polygon” (cf. [2]) formed by superimposing any  $C$  and  $C_0$  can always be generated by deforming some of the doubly occupied edges into bigger polygons, a process that does not alter the overall sign. It follows that  $C$  and  $C_0$  have the same sign for any  $C$ . This completes the proof.

Here we have implicitly made use of the fact that the monomer is on the boundary. If the monomer resides in the interior of  $\mathcal{L}$ , then there exist transposition polygons encircling the monomer site which may not necessarily carry the correct sign. The Pfaffian, while it can still be evaluated, does not yield the dimer-monomer generating function. We shall consider this general single-monomer problem subsequently [14].

With the edge orientation and phase factors in place, the dimer generating function  $G$  is obtained by evaluating the Pfaffian

$$G(x,y) = \text{Pf}(A') = \sqrt{\text{Det } A'} \quad (2)$$

where  $A'$  is the antisymmetric Kasteleyn matrix of dimension  $(MN+1) \times (MN+1)$ . Explicitly, it reads

$$A' = \begin{pmatrix} 0 & 0 & \cdots & 0 & 1 & 0 & \cdots & 0 \\ 0 & & & & & & & \\ \vdots & & & & & & & \\ 0 & & & & & & & \\ -1 & & & & & A & & \\ 0 & & & & & & & \\ \vdots & & & & & & & \\ 0 & & & & & & & \end{pmatrix}. \quad (3)$$

Here,  $A$  is the Kasteleyn matrix of dimension  $MN$  for  $\mathcal{L}$  given by

$$A = ixT_M \otimes I_N + yI_M \otimes T_N, \quad (4)$$

with  $I_N$  the  $N \times N$  identity matrix and  $T_N$  the  $N \times N$  matrix,

$$T_N = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ -1 & 0 & 1 & \cdots & 0 & 0 \\ 0 & -1 & 0 & \cdots & 0 & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & 0 & \cdots & -1 & 0 \end{pmatrix}. \quad (5)$$

Note that elements of  $A$  are labeled by  $\{(m,n);(m',n')\}$ , where  $(m,n)$  is a site index, and the element 1 in the first row of  $A'$  is at position  $(1,n)$  of  $A$ ,  $n=\text{odd}$ .

Expanding (3) along the first row and column, we obtain

$$\text{Det } A' = C(A; \{(1,n);(1,n)\}) \quad (6)$$

where  $C(A; \{(1,n);(1,n)\})$  is the cofactor of the  $\{(1,n);(1,n)\}$ th element of  $\text{Det } A$ .

The cofactor  $C(\alpha,\beta)$  of the  $(\alpha,\beta)$ th element of any non-singular  $A$  can be computed using the identity

$$C(A; \alpha,\beta) = A^{-1}(\beta,\alpha) \text{Det } A, \quad (7)$$

where  $A^{-1}(\beta,\alpha)$  is the  $(\beta,\alpha)$ th element of  $A$ . However, the formula is not directly useful in the present case since the matrix  $A$  is singular. We shall return to its evaluation in Sec. IV.

### III. EIGENVALUES OF THE DETERMINANT $A$

In this section we enumerate the eigenvalues of  $A$ .

The matrix  $T_N$  is diagonalized by the similarity transformation

$$U_N^{-1} T_N U_N = \Lambda_N$$

where  $U_N$  and  $U_N^{-1}$  are  $N \times N$  matrices with elements

$$U_N(n_1, n_2) = \sqrt{\frac{2}{N+1}} i^{n_1} \sin \frac{n_1 n_2 \pi}{n+1},$$

$$U_N^{-1}(n_1, n_2) = \sqrt{\frac{2}{N+1}} (-i)^{n_2} \sin \frac{n_1 n_2 \pi}{N+1}, \quad (8)$$

and  $\Lambda_N$  is an  $N \times N$  diagonal matrix whose diagonal elements are the eigenvalues of  $T_N$ ,

$$\lambda_m = 2i \cos \frac{m\pi}{N+1}, \quad m = 1, 2, \dots, N. \quad (9)$$

Similarly the  $MN \times MN$  matrix  $A$  is diagonalized by the similarity transformation generated by  $U_{MN} = U_M \otimes U_N$ ; namely,

$$U_{MN}^{-1} A U_{MN} = \Lambda_{MN}, \quad (10)$$

where  $\Lambda_{MN}$  is a diagonal matrix with eigenvalues

$$\lambda_{mn} = 2i \left( ix \cos \frac{m\pi}{M+1} + y \cos \frac{n\pi}{N+1} \right),$$

$$m = 1, 2, \dots, M, \quad n = 1, 2, \dots, N, \quad (11)$$

on the diagonal, and elements of  $U_{MN}$  and  $U_{MN}^{-1}$  are

$$U_{MN}(m_1, n_1; m_2, n_2) = U_M(m_1, m_2)U_N(n_1, n_2),$$

$$U_{MN}^{-1}(m_1, n_1; m_2, n_2) = U_M^{-1}(m_1, m_2)U_N^{-1}(n_1, n_2).$$

Then we have

$$\text{Det } A = \prod_{m=1}^M \prod_{n=1}^N \lambda_{mn}. \quad (12)$$

As in (2), close-packed dimers on  $\mathcal{L}$  are enumerated by evaluating  $\sqrt{\text{Det } A}$ . For  $MN$  even, this procedure gives precisely the Kasteleyn solution [2]. For  $MN$  odd, the case we are considering, the eigenvalue  $\lambda_{mn}=0$  for  $m=(M+1)/2, n=(N+1)/2$ , and hence  $\text{Det } A=0$ , indicating correctly that there is no dimer covering of  $\mathcal{L}$ . However, it is useful for later purposes to consider the product of the nonzero eigenvalues of  $A$ ,

$$P \equiv \prod_{m=1}^M \prod'_{n=1}^N \lambda_{mn}, \quad (13)$$

where the prime over the product denotes the restriction  $(m, n) \neq ((M+1)/2, (N+1)/2)$ .

Using the identity

$$\cos\left(\frac{m}{M+1}\right)\pi = -\cos\left(\frac{M-m+1}{M+1}\right)\pi,$$

one can rearrange factors in the product to arrive at

$$P = Q \prod_{m=1}^{(M-1)/2} \prod_{n=1}^{(N-1)/2} \left(4x^2 \cos^2 \frac{m\pi}{M+1} + 4y^2 \cos^2 \frac{n\pi}{N+1}\right)^2 \quad (14)$$

where the factor  $Q$  is the product of factors with either  $m=(M+1)/2$  or  $n=(N+1)/2$ . That is,

$$Q = \left(\prod_{m=1}^{(M-1)/2} 4x^2 \cos^2 \frac{m\pi}{M+1}\right) \left(\prod_{n=1}^{(N-1)/2} 4y^2 \cos^2 \frac{n\pi}{N+1}\right) \\ = \left(\frac{(M+1)(N+1)}{4}\right) x^{M-1} y^{N-1}, \quad (15)$$

where we have made use of the identity

$$\prod_{n=1}^{(N-1)/2} \left(4 \cos^2 \frac{n\pi}{N+1}\right) = \frac{N+1}{2}, \quad N \text{ odd.}$$

The expression (14) for  $P$  will be used in the next section.

#### IV. EVALUATION OF THE COFACTOR

We now return to the evaluation of the cofactor  $C(A; \{(1, n); (1, n)\})$ . We shall, however, evaluate the cofactor  $C(A; \{(m, n); (m', n')\})$  for general  $m, m', n, n'$ , although only the result of  $m=m'=1, n=n'$  is needed here.

To circumvent the problem of using (7) caused by the vanishing of  $\text{Det } A=0$ , we replace  $A$  by the matrix

$$A(\epsilon) = A + \epsilon I_{MN}, \quad \epsilon \neq 0,$$

whose inverse exists, and take the  $\epsilon \rightarrow 0$  limit to rewrite (7) as

$$C(A; \{(m, n); (m', n')\}) = \lim_{\epsilon \rightarrow 0} \{[A^{-1}(\epsilon)](m', n'; m, n) \text{Det } A(\epsilon)\}. \quad (16)$$

Quantities on the right-hand side of (16) are now well defined and the cofactor can be evaluated accordingly. Consideration of the inverse of a singular matrix along this line is known in mathematics literature as finding the pseudoinverse [15,16]. The method of taking the small- $\epsilon$  limit used here has previously been used successfully in analyses of resistance [17] and impedance [18] networks.

The eigenvalues of  $A(\epsilon)$  are  $\lambda_{mn}(\epsilon) = \lambda_{mn} + \epsilon$  and hence we have

$$\text{Det } A(\epsilon) = \prod_{m=1}^M \prod_{n=1}^N (\lambda_{mn} + \epsilon) = \epsilon P + O(\epsilon^2), \quad (17)$$

where  $P$  is the product of nonzero eigenvalues given by (14).

We next evaluate  $A^{-1}(\epsilon)(m', n'; m, n)$  and retain only terms of the order of  $1/\epsilon$ . Taking the inverse of (10) with  $A(\epsilon)$  in place of  $A$ , we obtain

$$A^{-1}(\epsilon) = U_{MN} \Lambda_{MN}^{-1}(\epsilon) U_{MN}^{-1}.$$

Writing out its matrix elements explicitly, we have

$$A^{-1}(\epsilon)(m', n'; m, n) = \sum_{m''=1}^M \sum_{n''=1}^N \\ \times \frac{U_{MN}(m', n'; m'', n'') U_{MN}^{-1}(m'', n''; m, n)}{\lambda_{m'', n''} + \epsilon}. \quad (18)$$

For  $\epsilon$  small the leading term comes from  $(m'', n'') = ((M+1)/2, (N+1)/2)$  for which  $\lambda_{m'', n''}=0$ . Using  $U_{MN}^{-1}(m, n; m', n') = U_M^{-1}(m, m') U_N^{-1}(n, n')$  and (8), this leads to the expression

$$A^{-1}(\epsilon)(m', n'; m, n) = \left(\frac{1}{\epsilon}\right) \left(\frac{4i^{m'+n'}(-i)^{m+n}}{(M+1)(N+1)}\right) \sin \frac{m'\pi}{2} \\ \times \sin \frac{n'\pi}{2} \sin \frac{m\pi}{2} \sin \frac{n\pi}{2} + O(1).$$

Thus, after making use of (16) and (17) we obtain

$$C(A; \{(m, n); (m', n')\}) = \sin \frac{m\pi}{2} \sin \frac{n\pi}{2} \sin \frac{m'\pi}{2} \sin \frac{n'\pi}{2} \\ \times \left(\frac{4i^{m'+n'}(-i)^{m+n} P}{(M+1)(N+1)}\right). \quad (19)$$

Finally, specializing to  $m=m'=1$ ,  $n=n'$  and combining (2), (6), and (19), we obtain

$$G(x,y) = \sqrt{C(A;\{(1,n);(1,n)\})}$$

$$= \begin{cases} \sqrt{\frac{4P}{(M+1)(N+1)}} & \text{for } n \text{ odd (A site),} \\ 0 & \text{for } n \text{ even (B site).} \end{cases} \quad (20)$$

This gives the result (1) after introducing (14) for  $P$ . It also says that there is no dimer covering if the monomer is on a  $B$  site.

The expression (20) clarifies the underlining mathematical content of the Tzeng-Wu solution (1) by identifying it as the product of the *nonzero* eigenvalues of the Kasteleyn matrix. This is compared to the Kasteleyn result [2] that for  $MN$  even the dimer generating function is given by the product of *all* eigenvalues.

## V. DISCUSSIONS

We have used a direct approach to derive the closed-form expression of the dimer-monomer generating function for the rectangular lattice with a single monomer on the boundary. Our approach is to first convert the problem into one of close-packed dimers without monomers, and consider the latter problem using established means. This approach suggests a possible route toward analyzing the general dimer-monomer problem.

We have also established that the Tzeng-Wu solution (1) is given by the product of the nonzero eigenvalues of the Kasteleyn matrix of the lattice. This is reminiscent of the well-known result in algebraic graph theory [19] that spanning trees on a graph are enumerated by evaluating the product of the nonzero eigenvalues of its tree matrix. The method of evaluating cofactors of a singular matrix as indicated by (16), when applied to the tree matrix of spanning trees, details of which can be easily worked out, offers a simple and direct proof of the fact that all cofactors of a tree matrix are equal and equal to the product of its nonzero eigenvalues. The intriguing similarity of the results suggests that there might be something deeper lurking behind our analysis.

- 
- [1] E. H. Lieb and O. J. Heilmann, Phys. Rev. Lett. **24**, 1412 (1970).  
 [2] P. W. Kasteleyn, Physica (Amsterdam) **27**, 1209 (1961).  
 [3] H. N. V. Temperley and M. E. Fisher, Philos. Mag. **6**, 1061 (1961).  
 [4] M. E. Fisher, Phys. Rev. **124**, 1664 (1961).  
 [5] W. T. Lu and F. Y. Wu, Phys. Lett. A **259**, 108 (1999).  
 [6] G. Tesler, J. Comb. Theory, Ser. B **78**, 198 (2000).  
 [7] M. Jerrum, J. Stat. Phys. **48**, 121 (1987); **59**, 1087 (1990).  
 [8] H. N. V. Temperley, in *Combinatorics: Proceedings of the British Combinatorial Conference*, London Mathematical Society Lecture Notes Series Vol. 13, (Cambridge University Press, Cambridge, U.K., 1974), p. 202.  
 [9] W.-J. Tzeng and F. Y. Wu, J. Stat. Phys. **10**, 671 (2003).  
 [10] Y. Kong, Phys. Rev. E **73**, 016106 (2006).  
 [11] Y. Kong, Phys. Rev. E **74**, 011102 (2006).  
 [12] Y. Kong (private communication).  
 [13] T. T. Wu, J. Math. Phys. **3**, 1265 (1962).  
 [14] W. T. Lu and F. Y. Wu (unpublished).  
 [15] A. Ben-Israel and T. N. E. Greville, *Generalized Inverses: Theory and Applications*, 2nd ed. (Springer-Verlag, New York, 2003).  
 [16] S. H. Friedberg, A. J. Insel, and L. E. Spence, *Linear Algebra*, 4th ed. (Prentice-Hall, New York, 2002), Sec. 6.7.  
 [17] F. Y. Wu, J. Phys. A **37**, 6653 (2004).  
 [18] W.-J. Tzeng and F. Y. Wu, J. Phys. A **39**, 8579 (2006).  
 [19] See, for example, N. L. Bigg, *Algebraic Graph Theory*, 2nd ed. (Cambridge University Press, Cambridge, U.K., 1993).