

## Reply to “Comment on ‘Self-organized criticality and absorbing states: Lessons from the Ising model’ ”

Gunnar Pruessner\*

Department of Mathematics, Imperial College London, 180 Queen’s Gate, London SW7 2AZ, United Kingdom

Ole Peters†

Department of Atmospheric Sciences, University of California, Los Angeles, 405 Hilgard Avenue, Los Angeles, California 90095-1565, USA

(Received 5 January 2008; revised manuscript received 25 March 2008; published 23 April 2008)

In R. Dickman, M. A. Muñoz, A. Vespignani, and S. Zapperi [Braz. J. Phys. **30**, 27 (2000)], Dickman *et al.* suggested that self-organized criticality can be produced by coupling the activity of an absorbing state model to a dissipation mechanism and adding an external drive. We analyzed the proposed mechanism in G. Pruessner and O. Peters [Phys. Rev. E **73**, 025106(R) (2006)] and found that if this mechanism is at work, the finite-size scaling found in self-organized criticality will depend on the details of the implementation of dissipation and driving. In the preceding paper [M. J. Alava, L. Laurson, A. Vespignani, and S. Zapperi, Phys. Rev. E **77**, 048101 (2008)], Alava *et al.* show that one avalanche exponent in the absorbing state approach becomes independent of dissipation and driving. In our reply we clarify their findings and put them in the context of the original paper.

DOI: [10.1103/PhysRevE.77.048102](https://doi.org/10.1103/PhysRevE.77.048102)

PACS number(s): 05.65.+b, 05.50.+q, 05.70.Jk, 64.60.Ht

In [1] we discussed the implications of the absorbing state (AS) mechanism [2] if it were to underlie self-organized criticality (SOC). One of the key ingredients of the AS mechanism is that dissipation and driving are made to vanish in the thermodynamic limit. We showed that criticality would be reached in the thermodynamic limit for almost any choice of the scaling with system size of dissipation and driving (with the effective temperature vanishing, contrary to what is stated in [3]). While this choice is thus not important to answer the question *if* the critical point will be reached, it is important when addressing the question of *how* it will be reached. In particular, we showed that the critical point would be reached in the limit of slow drive  $\omega - \kappa > \beta/\nu$  (“too fast” in [1]), but that the observed finite-size scaling exponents would depend on  $\omega - \kappa$ , and the relative correlation length  $\xi/L$  would vanish asymptotically.

The present discussion can be phrased in terms of two statements. (1) SOC is universal. (2) The AS mechanism is solely responsible for SOC. In [1] we showed that these are mutually exclusive. If the AS mechanism is the reason for SOC, then SOC cannot be universal. This would weaken SOC considerably since studying models as simple as sandpiles is only sensible if these systems show universal behavior. The other possibility is that SOC is universal, which would weaken the AS mechanism, because it does not predict universality. Our analysis was restricted to the finite-size scaling of AS observables, such as the order parameter, the correlation length, and the susceptibility, whereas the preceding paper [3] by Alava *et al.* refers to avalanche characteristics. In the following we assume that what we found out about the universality of AS observables also applies to ava-

lanches. We emphasize that we do not know whether this is true; this assumption is made in order to be able to reply to the comment, which makes the same assumption.

In the following we distinguish between the AS approach intended to explain SOC, and SOC itself. Alava *et al.* do not make this distinction explicit. In the AS approach driving and dissipation rates are tuned (bulk dissipation as  $L^{-\kappa}$  and driving as  $L^{-\omega}$ ), whereas in SOC (boundary dissipation, driving on a separate time scale) they are set implicitly by the dynamics of the system. If the AS approach applies to SOC, then SOC behavior is obtained within the AS approach by taking the thermodynamic limit. Both, our original article [1] as well as the preceding paper [3], are concerned solely with the characterization of the AS approach.

It is important to stress that in [1] we did *not* claim that any exponents, neither those describing avalanches nor those characterizing the activity, in standard SOC models are non-universal. This deserves clarification, because the opposite is stated in the abstract of [3]. As discussed in [1], there are instances of SOC exponents being identical across a wide range of models [4–6], while others are not [5,6].

We do claim (with the proviso stated above), however, that avalanche-size exponents would be nonuniversal if the AS mechanism [2] was applicable to SOC [1]. Alava *et al.* challenge this claim by showing that *within the AS approach*  $\tau_s$  is independent of the scaling of driving and dissipation in the slow driving limit. This limit corresponds to the separation of time scales in SOC.

In [1] we explicitly mention avalanche exponents only once: “In the AS approach also the avalanche-size exponents show a clear, immediate dependence on the choice of the two exponents  $\kappa$  and  $\omega$ .” It is correct that the avalanche-size exponents  $\tau_s$  and  $D_L$  (see below) depend on  $\kappa$  and  $\omega$ , but as Alava *et al.* show, for  $\omega$  large enough  $\tau_s$  becomes independent of  $\kappa$  and  $\omega$ . In this case, as we show below,  $D_L$  nonetheless depends on  $\kappa$ . The nonuniversality of  $D_L$  within the

\*URL: <http://www.ma.imperial.ac.uk/~pruess;g.pruessner@imperial.ac.uk>

†URL: <http://www.santafe.edu/~ole/;ole@santafe.edu>

AS approach implies that it does not explain universal SOC. This is the same conclusion we reached by studying AS observables (order parameter, correlation length, susceptibility).

We agree that the derivation in [1] necessarily breaks down in the large  $\omega$  limit. We explicitly assumed finite bounds for both  $\kappa$  and  $\omega$ , and we did not discuss the case of  $\omega$  being greater than the dynamical exponent, which is the regime studied by Alava *et al.* We agree that this regime is the most important one for SOC.

Alava *et al.* have chosen an observable that is independent of the external drive for sufficiently large  $\omega$ , but its finite-size scaling turns out to depend on the scaling of the dissipation. This can be seen by a finite-size scaling analysis of the characteristic avalanche size  $s_c$ . According to Ref. [3], Eq. (3),  $s_c \propto \xi_s^D$ , where  $\xi_s$  is some “cutoff scale.” Usually, the exponent  $D$  is reserved for the finite-size scaling of  $s_c$ , also known as the avalanche dimension [5], which we call  $D_L$  in the following, so that  $s_c \propto L^{D_L}$ . Combining Eqs. (4) and (3) of [3], they find  $D_L = \kappa / (2 - \tau_s)$  with  $\tau_s$  being independent of external drive and dissipation. This is a very surprising result, because regardless of whether or not the AS mechanism applies, the avalanche dimension  $D_L$  is deeply rooted in the model and in general, directly related to the field theory of the corresponding depinning transition [5,7–9]. There are several models [4–6] which display a universal avalanche dimension  $D_L$  and an avalanche-size exponent  $\tau_s = 2 - \gamma_1 / D_L$ , which depends on the details of the driving of the model [5], with the first moment scaling like  $\langle s \rangle \propto L^{\gamma_1}$ . For these conservative models, it is straightforward to devise a method to produce any exponent  $\tau_s$  in the interval [2

$-2/D_L, 2 - 1/D_L]$  by effectively tuning  $\gamma_1$ . This can be achieved by changing the driving mechanism [10], as the driving mechanism leaves  $D_L$  unchanged.

We would have expected that changing  $\gamma_1$  by introducing dissipation would have the same effect, i.e., varying  $\tau_s$  and constant  $D_L$ . However, Alava *et al.* find that  $D_L$  depends on  $\kappa$ , while  $\tau_s$  remains unchanged, a very interesting numerical finding we do not dispute. It has thus been established that, under the appropriate conditions, both  $\tau_s$  and  $D_L$  can be tuned. Importantly,  $D_L$  can be tuned in the SOC regime of the AS approach (large  $\omega$ ) by changing  $\kappa$ .

Inasmuch as avalanche exponents pertain to the discussion the universality of  $\tau_s$  within the AS approach supports the case for universal SOC being generated by the AS mechanism. But, the implicit finding by Alava *et al.* that  $D_L$  depends on  $\kappa$  confirms that, apparently, the AS approach does not produce universal SOC.

In order to address the question whether the AS mechanism is operating in SOC models (universal or not), one needs to probe its presence either directly or test its implications. In [1] we have shown that the AS mechanism would (almost always) imply a vanishing relative correlation length  $\xi/L$  and a finite-size scaling of the AS order parameter, characterized by exponents  $\beta/\mu$  and  $\gamma/\mu$ , that would depend on the scaling of dissipation and drive, parametrized by  $\kappa$  and  $\omega$ . We stated explicitly how  $\beta/\mu$  and  $\gamma/\mu$  depend on  $\kappa$  and  $\omega$ , while further analysis is necessary for the dependence of the avalanche exponents on  $\kappa$  and  $\omega$ . At the present stage, a more promising route than studying avalanches therefore seems to be the study of AS observables in SOC systems.

- 
- [1] G. Pruessner and O. Peters, Phys. Rev. E **73**, 025106(R) (2006).
- [2] R. Dickman, M. A. Muñoz, A. Vespignani, and S. Zapperi, Braz. J. Phys. **30**, 27 (2000).
- [3] M. J. Alava, L. Laurson, A. Vespignani, and S. Zapperi, preceding paper, Phys. Rev. E **77**, 048101 (2008).
- [4] G.-J. Pan, D.-M. Zhang, H.-Z. Sun, and Y.-P. Yin, Commun. Theor. Phys. **44**, 483 (2005).
- [5] K. Christensen, N. R. Moloney, O. Peters, and G. Pruessner, Phys. Rev. E **70**, 067101 (2004).
- [6] A. Malthe-Sørensen, Phys. Rev. E **59**, 4169 (1999).
- [7] M. Paczuski and S. Boettcher, Phys. Rev. Lett. **77**, 111 (1996).
- [8] G. Pruessner, Phys. Rev. E **67**, 030301(R) (2003).
- [9] M. Alava, in *Advances in Condensed Matter and Statistical Physics*, edited by E. Korutcheva and R. Cuerno (Nova Science Publishers, New York, 2004).
- [10] As long as the model is conservative, individual particles effectively perform a random walk and the first moment of the avalanche size is a function of the spatial distribution of the driving. For example, if a one-dimensional model is driven at site  $x_0$  with two open boundaries, then the first moment is  $(L + 1 - x_0)x_0$ , where  $L$  is the system size. By changing  $x_0$  as a power law of  $L$ , the exponent  $\gamma_1$  changes accordingly. For example,  $x_0 = \sqrt{L}$  produces  $\gamma_1 = 3/2$ .