

Slow excitation trapping in quantum transport with long-range interactions

Oliver Mülken,* Volker Pernice, and Alexander Blumen

Theoretische Polymerphysik, Universität Freiburg, Hermann-Herder-Straße 3, 79104 Freiburg, Germany

(Received 25 April 2008; revised manuscript received 9 July 2008; published 15 August 2008)

Long-range interactions (LRIs) slow down the excitation trapping in quantum transport on a one-dimensional chain with traps at both ends when compared to the case with only nearest-neighbor interactions. This is in contrast to the corresponding classical case, in which LRIs lead to faster excitation trapping. The reason for the slowing down is to be found in subtle changes—due to LRIs—in the spectrum of the Hamiltonian. Perturbation theory allows an analytical analysis of this fact.

DOI: [10.1103/PhysRevE.78.021115](https://doi.org/10.1103/PhysRevE.78.021115)

PACS number(s): 05.60.Gg, 05.60.Cd, 71.35.-y

Building a quantum system from scratch has become possible due to recent experimental advances in controlling and manipulating atoms and molecules. It has actually become possible to tailor the theoreticians' favorite one-dimensional systems using, e.g., ultracold atoms in optical lattices; see [1] and references therein. From a dynamical point of view, this allows for these systems to compare the theoretical predictions for the transport of charge, mass, or energy to the experimental results. In turn, the experimental findings might eventually lead to a refinement of the theoretical models.

The tight-binding approximation for the transport of a quantum particle over a regular structure (network) is a simple description which is equivalent to the so-called continuous-time quantum walk (CTQW) with nearest-neighbor interactions (NNIs) [2,3]. Recently, several experiments have been proposed addressing the CTQW, e.g., based on waveguide arrays [4], atoms in optical lattices [5,6], or structured clouds of ultracold Rydberg atoms [7]. In some of these experiments one finds long-range interactions (LRIs), as in Rydberg gases, where also blockade [9] and antiblockade [10] effects have to be considered. Recently, we analyzed the behavior of CTQWs when the probability of jumps to other sites depends on the mutual distance R between the sites, the LRIs being of the form $R^{-\nu}$ [8]. We found that CTQWs belong to the same universality class for $\nu > 2$, while for classical continuous-time random walks (CTRWs) universality holds only for $\nu > 3$.

Coupling a system to an absorbing site, i.e., to a trap, allows the transport to be monitored by observing the decay of the survival probability of the moving entity, say, the excitation. In the long-time limit and for NNIs the decay is practically exponential for both classical systems modeled by CTRWs [11] and quantum systems modeled by CTQWs [7,12]. At intermediate times, which are experimentally relevant, there appear considerable, characteristic differences between the classical and the quantum situations [7].

Here, we study the quantum dynamics of a one-dimensional CTQW with LRIs in the presence of traps and use the similarity to a CTRW for a comparison with the respective classical case. Without traps, we model the quantum dynamics on a network of connected nodes by a tight-binding Hamiltonian \mathbf{H}_0 . For the corresponding classical pro-

cess, we identify the CTRW transfer matrix \mathbf{T}_0 with \mathbf{H}_0 , i.e., $\mathbf{H}_0 = -\mathbf{T}_0$; see, e.g., [2,3] for details. For undirected networks, \mathbf{H}_0 is related to the connectivity matrix \mathbf{A}_0 of the network by $\mathbf{H}_0 = \mathbf{A}_0$. When the interactions between two nodes vary as $R^{-\nu}$, with $R = |k - j| \geq 1$ being the distance between two nodes j and k , the Hamiltonian has the following structure:

$$\mathbf{H}_0(\nu) = \sum_{n=1}^N \left(\sum_{R=1}^{n-1} R^{-\nu} (|n\rangle\langle n| - |n-R\rangle\langle n|) + \sum_{R=1}^{N-n} R^{-\nu} (|n\rangle\langle n| - |n+R\rangle\langle n|) \right). \quad (1)$$

We restrict ourselves to extensive cases ($\nu > 1$), i.e., we explicitly exclude ultralong-range interactions. The corresponding NNI Hamiltonian is obtained for $\nu = \infty$, in which case only the leading terms with $R = 1$ do not vanish.

The states $|j\rangle$ associated with excitations localized at the nodes j ($j = 1, \dots, N$) form a complete, orthonormal basis set of the whole accessible Hilbert space ($\langle k|j\rangle = \delta_{kj}$ and $\sum_k |k\rangle\langle k| = \mathbf{1}$). In general, the transition probabilities from a state $|j\rangle$ at time $t_0 = 0$ to a state $|k\rangle$ at time t read $\pi_{kj}(t) \equiv |\alpha_{kj}(t)|^2 \equiv |\langle k|\exp[-i\mathbf{H}_0(\nu)t]|j\rangle|^2$. In the corresponding classical CTRW case the transition probabilities follow from a master equation as $p_{kj}(t) = \langle k|\exp(\mathbf{T}_0 t)|j\rangle$ [2,3].

Now, let the nodes m ($m \in \mathcal{M}$ and $\mathcal{M} \subset \{1, \dots, N\}$) be traps for the excitation. Within a phenomenological approach, the new Hamiltonian is $\mathbf{H}(\nu) \equiv \mathbf{H}_0(\nu) - i\mathbf{\Gamma}$, with the trapping operator $i\mathbf{\Gamma} \equiv i\mathbf{\Gamma} \sum_{m \in \mathcal{M}} |m\rangle\langle m|$ (see Ref. [7] for details). As a result, \mathbf{H} is non-Hermitian and has N complex eigenvalues $E_l = \epsilon_l - i\gamma_l$ ($l = 1, \dots, N$) with $\gamma_l > 0$, and N left and N right eigenstates, denoted by $|\Psi_l\rangle$ and $\langle\tilde{\Psi}_l|$, respectively. The transition probabilities follow as

$$\pi_{kj}(t) = \left| \sum_l \exp(-\gamma_l t) \exp(-i\epsilon_l t) \langle k|\Psi_l\rangle \langle\tilde{\Psi}_l|j\rangle \right|^2, \quad (2)$$

where the imaginary parts γ_l of E_l determine the temporal decay. Note that traps do not lead to decoherence but to decaying probabilities. For the incoherent classical process the description by CTRWs is quite similar: The new transfer operator reads $\mathbf{T}(\nu) = \mathbf{T}_0(\nu) - \mathbf{\Gamma} = -\mathbf{A}_0(\nu) - \mathbf{\Gamma}$, which is real and symmetric, leading to the eigenvalues $-\lambda_l$ ($\lambda_l > 0$) and corresponding eigenstates $|\Phi_l\rangle$. Note that due to the different incorporation of the trapping operator in $\mathbf{T}(\nu)$ and $\mathbf{H}(\nu)$ the

*muelken@physik.uni-freiburg.de

corresponding eigenvalues and eigenstates will differ. Without trapping we have $\mathbf{T}_0(\nu) = -\mathbf{H}_0(\nu)$ and thus $\lambda_l \equiv E_l$ and $|\Phi_l\rangle \equiv |\Psi_l\rangle$.

In order to make a global statement for the whole network, we calculate the mean survival probability for a total number of M trap nodes,

$$\Pi_M(t) \equiv \frac{1}{N-M} \sum_{j \in \mathcal{M}} \sum_{k \notin \mathcal{M}} \pi_{kj}(t), \quad (3)$$

i.e., the average of $\pi_{kj}(t)$ over all initial nodes j and all final nodes k , neither of them being a trap node. Classically, we will consider $P_M(t) \equiv 1/(N-M) \sum_{j \in \mathcal{M}} \sum_{k \notin \mathcal{M}} p_{kj}(t)$. For intermediate and long times and a small number of trap nodes, $\Pi_M(t)$ is mainly a sum of exponentially decaying terms [7]:

$$\Pi_M(t) \approx \frac{1}{N-M} \sum_{l=1}^N \exp(-2\gamma_l t). \quad (4)$$

If the imaginary parts γ_l obey a power law with an exponent μ ($\gamma_l \sim a l^\mu$), the mean survival probability scales as $\Pi_M(t) \sim t^{-1/\mu}$.

In Ref. [7] an experimental setup was proposed, which is based on a finite linear chain of clouds of ultracold Rydberg atoms with trapping states at both ends ($m=1, N$). There, the dynamics was approximated by a NNI tight-binding model, which—for a ring without traps—has been shown to behave in the same fashion as systems with LRIs of the form $R^{-\nu}$ for which $\nu > 2$ [8]. The Rydberg atoms interact via dipole-dipole forces, i.e., the potential between two atoms decays roughly as R^{-3} .

For the finite chain with $m=1, N$, Fig. 1 shows a comparison of the quantum mechanical $\Pi_M(t)$ and of the classical $P_M(t)$ behaviors for different ν and Γ , which were obtained by numerically diagonalizing the corresponding Hamiltonian $\mathbf{H}(\nu)$ and transfer matrix $\mathbf{T}(\nu)$, respectively. Clearly, for both Γ values the LRIs lead to a slower decay of $\Pi_M(t)$, i.e., to a slower trapping of the excitation, which is remarkable since the opposite effect is observable for classical systems where the decay of $P_M(t)$ becomes faster for decreasing ν . The classical behavior can be simply understood since, at least for short and intermediate times, increasing the probabilities for long-range jumps also increases the probabilities for visiting new nodes and hence to encounter a trap. By increasing the trapping strength Γ , the difference between the quantum and the classical behavior becomes even more pronounced [compare Figs. 1(a) and 1(b)]. Generally, for $\Pi_M(t)$ the change in Γ results mainly in a rescaled time axis, since the imaginary parts γ_l are of the same order of magnitude when rescaled by Γ . For the specific case of the Rydberg atoms ($\nu=3$ and $\Gamma=1$) one observes the largest difference between the $\Pi_M(t)$ and the $P_M(t)$ behaviors. To understand this phenomenon, we continue to analyze $\Pi_M(t)$ within a perturbation theoretical treatment.

When the strength of the trap, Γ , is small compared to the couplings between neighboring nodes, we can evaluate the eigenvalues using perturbation theory; see, for instance, [13]. Let $|\Psi_l^{(0)}\rangle$ be the l th eigenstate and $E_l^{(0)} \in \mathbb{R}$ be the l th eigen-




FIG. 1. (Color online) ν dependence of the quantum mechanical $\Pi_M(t)$ and the classical $P_M(t)$ decay behaviors for a chain of $N=100$ sites; here $\Gamma=(a)$ 0.001 and (b) 1. The inset in (a) shows a closeup picture of the region where $\Pi_M(t)$ and $P_M(t)$ cross. The inset in (b) shows power-law fits (see arrows) to $\Pi_M(t)$ in the intermediate time regime with exponents $1/\mu$, where the μ are taken from Fig. 3(b).

value of the unperturbed system with Hamiltonian $\mathbf{H}_0(\nu)$. Up to first order the eigenvalues of the perturbed system are given by

$$E_l = E_l^{(0)} - i\Gamma \sum_{m \in \mathcal{M}} |\langle m | \Psi_l^{(0)} \rangle|^2. \quad (5)$$

Therefore, the correction term determines the imaginary parts γ_l , while the unperturbed eigenvalues are the real parts $\epsilon_l = E_l^{(0)}$. Having only a few trap nodes, the sum in Eq. (5) contains only few terms. Moreover, from Eq. (5) we also see that the imaginary parts γ_l are essentially determined by the eigenstates $|\Psi_l^{(0)}\rangle$ of the system without traps. A change in these states will also lead to a change in the γ_l . As we proceed to show, this is exactly what happens by going from NNIs to LRIs: With decreasing ν some of the unperturbed eigenstates $|\Psi_l^{(0)}\rangle$ become more concentrated within the system, i.e., they have a lower contribution from the nodes $m=1, N$. This, in turn, leads to the slower decay of the survival probability, because the imaginary parts are related to $|\langle m | \Psi_l^{(0)} \rangle|^2$ by Eq. (5).

Without loss of generality, an eigenstate of a finite chain with NNIs can be written as ($l=1, \dots, N$)

$$|\Psi_l^{(0)}\rangle = \begin{cases} \sqrt{\frac{1}{N}} \sum_{j=1}^N |j\rangle, & \text{for } l=N, \\ \sqrt{\frac{2}{N}} \sum_{j=1}^N \cos[(2j-1)\theta_l/2] |j\rangle & \text{else,} \end{cases} \quad (6)$$

where for convenience we take $\theta_l \equiv \pi(N-l)/N \in [0, \pi]$; the corresponding eigenvalues are $E_l^{(0)} = 2 - 2 \cos \theta_l$ (note that the smallest eigenvalue is $E_N^{(0)} = 0$). Thus, in first-order perturbation theory we obtain from Eqs. (5) and (6) as imaginary parts $\gamma_N = 2\Gamma/N$ and $\gamma_l = (4\Gamma/N) \cos^2(\theta_l/2) = (2\Gamma/N)[1 + \cos \theta_l]$ for $l=1, \dots, N-1$, which for $l \ll N$ yields $\gamma_l \sim l^2$. In this case the mean survival probability will scale in the corresponding time interval as $\Pi_M(t) \sim t^{-1/2}$.

Formally, we can perform the continuum limit $N \rightarrow \infty$ (by taking now $4\Gamma/N \equiv a$ finite). Then the sum in Eq. (4) turns into an integral such that

$$\Pi_M(t) \sim e^{-at} \frac{1}{\pi} \int_0^\pi d\theta e^{-at \cos \theta} = e^{-at} I_0(at), \quad (7)$$

where $I_0(at)$ is the modified Bessel function of the first kind [14]. From this we get for large t that $\Pi_M(t) \sim t^{-1/2}$, which confirms the previous results. Note, however, that for small N the smallest γ_l value is finite and, therefore, the scaling of γ_l holds only in a quite small interval of l values. Hence, also the time interval in which $\Pi_M(t)$ scales with the exponent $-1/2$ is rather small. A lower bound for scaling is given by the behavior of γ_l for $l \approx N/2$ (corresponding to smaller times than for $l \ll N$). Here, γ_l is linear in l , which leads to a lower bound of $\mu \geq 1$ for the scaling exponent. An exponent μ which is valid over a larger l interval will therefore be in the interval [1,2] and, consequently, the exponent for $\Pi_M(t)$ will lie in the interval $[-1, -1/2]$.

In the case of periodic boundary conditions, one finds translation-invariant Bloch eigenstates regardless of the range of the interaction [8]. In the case of Eq. (1), however, the eigenstates for LRI differ from the ones for NNI [Eq. (6)]; in Eq. (1) the finite extension of the chain destroys the translational invariance. As is immediately clear from Eq. (5), this also implies that the imaginary parts of the eigenvalues, evaluated based on first-order perturbation theory, will change.

For large exponents ν we can regard the LRI as a small perturbation to the NNI, i.e., having $\mathbf{H}_0(\nu) = \mathbf{H}_0 + \mathbf{H}_\nu$, where \mathbf{H}_ν contains only the correction terms to the NNI case \mathbf{H}_0 . This allows us to calculate from the unperturbed states $|\Psi_l^{(0)}\rangle$ the perturbed eigenstates $|\Psi_l\rangle$ up to first order. Taking the states $|\Psi_l\rangle$ to be the eigenstates of the LRI system without traps, we readily obtain the imaginary parts γ_l for small trapping strength from Eq. (5) as $\gamma_l = 2\Gamma |\langle 1 | \Psi_l \rangle|^2$, where

$$\langle 1 | \Psi_l \rangle = \langle 1 | \Psi_l^{(0)} \rangle + \sum_{r \neq l} \frac{\langle \Psi_r^{(0)} | \mathbf{H}_\nu | \Psi_l^{(0)} \rangle}{E_l^{(0)} - E_r^{(0)}} \langle 1 | \Psi_r^{(0)} \rangle. \quad (8)$$

It is straightforward, although cumbersome, to calculate the corrections to the imaginary parts γ_l from Eq. (8). For large ν the coupling to the next-next-nearest neighbor is by a




FIG. 2. (Color online) Correction term $\langle 1 | \Psi_l \rangle - \langle 1 | \Psi_l^{(0)} \rangle$ for $N=100$ and for $\nu=(a) 10$ and (b) 5. The direct numerical evaluation (solid black line) is compared to the perturbation theory expression Eq. (8) (dash-dotted green line) and to the approximate expression Eq. (9) (dashed red line).

factor of $(3/2)^\nu$ smaller, for $\nu=10$ this is about one and a half orders of magnitude. Taking, for fixed ν , only nearest- and next-nearest-neighbor couplings into account allows us to obtain simple analytic expressions. The perturbation term \mathbf{H}_ν is now tridiagonal. Its nonzero elements are $\langle j-2 | \mathbf{H}_\nu | j \rangle = \langle j+2 | \mathbf{H}_\nu | j \rangle = -2^{-\nu}$ and its diagonal elements follow from $\langle j | \mathbf{H}_\nu | j \rangle = -\sum_i \langle i | \mathbf{H}_\nu | j \rangle$, thus $\langle j | \mathbf{H}_\nu | j \rangle = 2^{-\nu}$ for $2 < j < N-1$ and $\langle j | \mathbf{H}_\nu | j \rangle = 2^{-\nu+1}$ else. We hence obtain from Eq. (8)

$$\langle 1 | \Psi_l \rangle = \sqrt{\frac{2}{N}} \cos\left(\frac{\theta_l}{2}\right) + 2^{-\nu} \sqrt{\frac{2}{N}} \sin(2\theta_l) \sin\left(\frac{\theta_l}{2}\right). \quad (9)$$

Figure 2 shows the difference $\langle 1 | \Psi_l \rangle - \langle 1 | \Psi_l^{(0)} \rangle$ for $N=100$ and for $\nu=(a) 10$ and (b) 5. The numerical exact value (solid black line) is obtained by computing separately $\langle 1 | \Psi_l \rangle$ and $\langle 1 | \Psi_l^{(0)} \rangle$ and subsequently taking the difference; the result is then confronted with Eq. (8) (dash-dotted green line), determined numerically, and to Eq. (9) (dashed red line). For $\nu=10$, the agreement between all three curves is remarkably good [see Fig. 2(a)], which justifies the assumptions leading to Eq. (9). For smaller ν [$\nu=5$ in Fig. 2(b)] there is still a reasonable agreement between Eq. (8) and the exact result; however, taking only nearest and next-nearest neighbors into account leads to evident deviations, see the dashed red line in Fig. 2(b).

Now, from Eq. (9) we get

$$\gamma_l \approx \gamma_l^{(0)} + 2^{-\nu} \gamma_l^{(1)} + O(2^{-2\nu}), \quad (10)$$

where $\gamma_l^{(0)}$ is the NNI expression given above and $\gamma_l^{(1)} = (8\Gamma/N) \cos(\theta_l/2) \sin(2\theta_l) \sin(\theta_l/2)$ the correction due to the LRIs. Again, the smallest γ_l values are those for which $l \ll N$, which leads to a decrease of the imaginary parts γ_l because $\gamma_l^{(1)} < 0$ for $l \ll N$. Here, one can approximate the imaginary parts by a power law, i.e., $\gamma_l \sim l^\mu$. A rough estimate of the scaling exponent μ , assuming $\nu \gg 1$ can be readily given. For this we note that from Eq. (10) we have $\ln \gamma_{l+1} - \ln \gamma_l \approx \ln \gamma_{l+1}^{(0)} - \ln \gamma_l^{(0)} + 2^{-\nu} (\gamma_{l+1}^{(1)} / \gamma_{l+1}^{(0)} - \gamma_l^{(1)} / \gamma_l^{(0)})$.




FIG. 3. (Color online) Imaginary parts γ_l (symbols) in ascending order for LRI systems with $\nu=3, 4, 5$, and for NNIs for $N=100$ and $\Gamma=(a)$ 0.001 and (b) 1. The inset shows the power-law behavior of the γ_l in the intermediate l region with the appropriate exponents μ .

Moreover, the term $\mu^{(0)} \equiv (\ln \gamma_{l+1}^{(0)} - \ln \gamma_l^{(0)}) / [\ln(l+1) - \ln l]$ gives the exponent for the NNI case and the term $\mu^{(1)} \equiv (\gamma_{l+1}^{(1)} / \gamma_{l+1}^{(0)} - \gamma_l^{(1)} / \gamma_l^{(0)}) / [\ln(l+1) - \ln l]$ is the LRI correction. Thus

$$\mu \approx \frac{\ln \gamma_{l+1} - \ln \gamma_l}{\ln(l+1) - \ln l} \approx \mu^{(0)} + 2^{-\nu} \mu^{(1)}. \quad (11)$$

Since $\mu^{(1)}$ is strictly positive for small l , the inclusion of LRIs leads to a decrease of γ_l when compared to the NNI case. In turn, this results in a slower decay of $\Pi_M(t)$.

Figure 3 shows the imaginary parts γ_l for a chain of $N=100$ nodes with LRIs ($\nu=3, 4, 5$), and with NNIs. For small l and NNIs, the γ_l obey scaling with the exponent $\mu=2$, as discussed above. Introducing LRIs, i.e., decreasing ν , increases the scaling exponent to $\mu > 2$. Consequently, the scaling exponent $1/\mu$ for $\Pi_M(t)$ decreases, leading to a slowing down of the excitation trapping due to LRIs.

In the classical case decreasing ν leads to a faster excitation trapping, which is observable in a quicker decay of

$P_M(t)$. This can also be deduced from a perturbation theoretical treatment. As can be seen from Fig. 1 (see also Fig. 2 of Ref. [7]), the decay of $P_M(t)$ is exponential already at intermediate times and is dominated by the smallest eigenvalue λ_N and the corresponding eigenstate $|\Phi_N\rangle$ of the transfer operator $\mathbf{T}(\nu)$:

$$P_M(t) = \frac{1}{N-M} \sum_{l=1}^N \exp(-\lambda_l t) \left| \sum_{k \in \mathcal{M}} \langle k | \Phi_l \rangle \right|^2 \\ \approx \frac{1}{N-M} \exp(-\lambda_N t) \left| \sum_{k \in \mathcal{M}} \langle k | \Phi_N \rangle \right|^2. \quad (12)$$

Calculating λ_N and the prefactor $|\sum_{k \in \mathcal{M}} \langle k | \Phi_N \rangle|^2$ for large ν and small Γ shows that with decreasing ν the smallest eigenvalue λ_N increases while the prefactor decreases. Together, this confirms our numerical result of a quicker decay for $P_M(t)$, see Fig. 1.

Finally, we comment on the impact of our results on the experiment proposed in Ref. [7]. Here, clouds of laser-cooled ground state atoms are assembled in a chain by optical dipole traps [15], which are then excited into a Rydberg S state (see [7] for details). The Rydberg atoms interact via long-range dipole-dipole forces which is advantageous in many ways. As can be deduced from Fig. 1, the time intervals over which the decay follows the power law are enlarged by the LRIs. For $\nu=3$ the transition to the long-time exponential decay occurs at times which are about two orders of magnitude larger than the ones found for the NNI case. The difference between a purely coherent (CTQW) and a purely incoherent (CTRW) process is enlarged due to the LRIs, allowing for a better discrimination between the two when clarifying the nature of the energy transfer dynamics in ultracold Rydberg gases.

In conclusion, we have considered the quantum dynamics of excitations with LRIs on a network in the presence of absorbing sites (traps). The LRIs lead to a slowing down of the decay of the average survival probability, which is remarkable since for the corresponding classical process one observes a speedup of the decay. Using perturbation theory arguments we were able to identify the reason for this slowing down; it results from changes in the imaginary parts of the spectrum of the Hamiltonian.

Support from the Deutsche Forschungsgemeinschaft (DFG) and the Fonds der Chemischen Industrie is gratefully acknowledged.

[1] I. Bloch, Nat. Phys. **1**, 23 (2005); I. Bloch, J. Dalibard, and W. Zwerger, Rev. Mod. Phys. **80**, 885 (2008).
 [2] E. Farhi and S. Gutmann, Phys. Rev. A **58**, 915 (1998).
 [3] O. Mülken and A. Blumen, Phys. Rev. E **71**, 016101 (2005).
 [4] H. B. Perets, Y. Lahini, F. Pozzi, M. Sorel, R. Morandotti, and Y. Silberberg, Phys. Rev. Lett. **100**, 170506 (2008).
 [5] W. Dür, R. Raussendorf, V. M. Kendon, and H. J. Briegel, Phys. Rev. A **66**, 052319 (2002).

[6] R. Côté, A. Russel, E. E. Eyler, and P. L. Gould, New J. Phys. **8**, 156 (2006).
 [7] O. Mülken, A. Blumen, T. Amthor, C. Giese, M. Reetz-Lamour, and M. Weidemüller, Phys. Rev. Lett. **99**, 090601 (2007).
 [8] O. Mülken, V. Pernice, and A. Blumen, Phys. Rev. E **77**, 021117 (2008).
 [9] M. D. Lukin, M. Fleischhauer, R. Cote, L. M. Duan, D. Jak-

- sch, J. I. Cirac, and P. Zoller, Phys. Rev. Lett. **87**, 037901 (2001); K. Singer, M. Reetz-Lamour, T. Amthor, L. G. Marcassa, and M. Weidemüller, *ibid.* **93**, 163001 (2004).
- [10] C. Ates, T. Pohl, T. Pattard, and J. M. Rost, Phys. Rev. Lett. **98**, 023002 (2007); Phys. Rev. A **76**, 013413 (2007).
- [11] J. Klafter and R. Silbey, J. Chem. Phys. **72**, 849 (1980); P. Grassberger and I. Procaccia, *ibid.* **77**, 6281 (1982).
- [12] R. M. Pearlstein, J. Chem. Phys. **56**, 2431 (1971); D. L. Huber, Phys. Rev. B **22**, 1714 (1980); **45**, 8947 (1992); P. E. Parris, Phys. Rev. Lett. **62**, 1392 (1989); Phys. Rev. B **40**, 4928 (1989); V. A. Malyshev, R. Rodríguez, and F. Domínguez-Adame, J. Lumin. **81**, 127 (1999).
- [13] J. Sakurai, *Modern Quantum Mechanics*, 2nd ed. (Addison-Wesley, Redwood City, CA, 1994).
- [14] *Handbook of Mathematical Functions*, edited by M. Abramowitz and I. A. Stegun (Dover, New York, 1972).
- [15] R. Grimm, M. Weidemüller, and Y. B. Ovchinnikov, Adv. At., Mol., Opt. Phys. **42**, 95 (2000).