

**Erratum: Dynamics of impacting a bubble by another pulsed-laser-induced bubble:
Jetting, fragmentation, and entanglement [Phys. Rev. E **77**, 026304 (2008)]**

Yen-Hung Chen and Lin I

(Received 25 March 2009; published 13 April 2009)

DOI: [10.1103/PhysRevE.79.049901](https://doi.org/10.1103/PhysRevE.79.049901)

PACS number(s): 47.55.dp, 47.20.Ma, 99.10.Cd

In the references, the contents of [22] and [23] were mistakenly placed in reverse order. They should be inverted as follows: [22] G. Layes, G. Jourdan, and L. Houas, *Phys. Rev. Lett.* **91**, 174502 (2003); and [23] C. D. Ohl and R. Ikink, *Phys. Rev. Lett.* **90**, 214502 (2003).

As stated in the third paragraph of the introduction part, the Richtmyer-Meshkov (RM) instability can lead to vortex formation when the gas bubble with different densities inside and outside the bubble is impacted by a shock wave. In the experiment by Ranjan *et al.* [21], the gas density inside the soap bubble is higher than the surrounding gas outside the bubble. However, it was stated in a reverse way in the published paper. The description of the gas densities for the work by Layes *et al.* was properly stated in the published paper.