


This article was downloaded by:

On: 28 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Physics and Chemistry of Liquids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713646857>

### Solubility of *trans*-Stilbene in Binary Alkane 2-Propanol Solvent Mixtures at 298.2 K

Karina M. De Fina<sup>a</sup>; Carmen E. Hernandez<sup>a</sup>; Tina L. Sharp<sup>a</sup>; William E. Acree Jr.<sup>a</sup>

<sup>a</sup> Department of Chemistry, University of North Texas, Denton, Texas, USA

**To cite this Article** De Fina, Karina M. , Hernandez, Carmen E. , Sharp, Tina L. and Acree Jr., William E.(2000) 'Solubility of *trans*-Stilbene in Binary Alkane 2-Propanol Solvent Mixtures at 298.2 K', *Physics and Chemistry of Liquids*, 38: 1, 89 – 94

**To link to this Article:** DOI: 10.1080/00319100008045299

**URL:** <http://dx.doi.org/10.1080/00319100008045299>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## SOLUBILITY OF *trans*-STILBENE IN BINARY ALKANE + 2-PROPANOL SOLVENT MIXTURES AT 298.2 K

KARINA M. DE FINA, CARMEN E. HERNÁNDEZ,  
TINA L. SHARP and WILLIAM E. ACREE JR.\*

*Department of Chemistry, University of North Texas,  
Denton, Texas 76203-5070, USA*

*(Received 21 September 1998)*

Experimental solubilities are reported for *trans*-stilbene dissolved in six binary alkane + 2-propanol solvent mixtures at 25°C. The alkane cosolvents studied were hexane, heptane, octane, cyclohexane, methylcyclohexane and 2,2,4-trimethylpentane. Results of these measurements are used to test two mathematical representations based upon the combined Nearly Ideal Binary Solvent (NIBS)/Redlich-Kister and Modified Wilson equations. For the six systems studied, both equations provided an accurate mathematical representation of the experimental data, with overall average absolute deviations between measured and calculated values being approximately  $\pm 0.5\%$ .

**Keywords:** *trans*-stilbene solubilities; binary solvent mixtures; alkanes

### INTRODUCTION

Solid–liquid equilibrium data of organic nonelectrolyte systems are becoming increasingly important in the petroleum industry, particularly in light of present trends towards heavier feedstocks and known carcinogenicity/mutagenicity of many of the larger polycyclic aromatic compounds. Solubility data for a number of polycyclic aromatic hydrocarbons (*i.e.*, anthracene and pyrene) and hetero-atom polynuclear aromatics (*i.e.*, carbazole, dibenzothiophene and xanthene) have been published in the recent chemical literature (for listing of references see

---

\*Corresponding author. e-mail: acree@unt.edu

Acree [1–3]). Despite efforts by experimentalists and scientific organizations, both in terms of new experimental measurements and critically-evaluated data compilations, there still exist numerous systems for which solubility data are not readily available.

To address this problem, researchers have turned to predictive methods as a means to generate desired quantities. Numerous equations have been suggested for predicting solute solubilities in binary solvent mixtures. For the most part, the predictive methods do provide fairly reasonable estimates for non-complexing systems. There still remains, however, the need to develop better predictive expressions and mixing models to describe the more nonideal complexing systems believed to contain hydrogen bonding solvent components. Continued development of solution models for describing the thermodynamic properties of a solute in binary solvent systems requires that a large data base be available for assessing the applications and limitations of derived expressions. Currently, only a limited data base exists for crystalline nonelectrolyte solubility in binary solvent mixtures. For this reason, *trans*-stilbene solubilities were determined in six binary alkane + 2-propanol solvent mixtures. Results of these measurements are used to further test the descriptive abilities of two previously derived expressions.

## EXPERIMENTAL METHODS

*trans*-Stilbene (Aldrich, 96%) was recrystallized three times from methanol. 2-Propanol (Aldrich, 99.5+%, anhydrous), hexane (Aldrich, 99+%), heptane (Aldrich, HPLC), octane (Aldrich, 99+%, anhydrous), cyclohexane (Aldrich, HPLC), methylcyclohexane (Aldrich, 99+%, anhydrous) and 2,2,4-trimethylpentane (Aldrich, HPLC) were stored over molecular sieves and distilled shortly before use. Gas chromatographic analysis showed solvent purities to be 99.7 mole percent or better. Binary solvent mixtures were prepared by mass so that compositions could be calculated to 0.0001 mole fraction.

Excess solute and solvent were placed in amber glass bottles and allowed to equilibrate in a constant temperature water bath at  $(25.0 \pm 0.1)^\circ\text{C}$  with periodic agitation for at least three days (often longer). Attainment of equilibrium was verified both by repetitive

measurements after a minimum of three additional days and by approaching equilibrium from supersaturation by pre-equilibrating the solutions at a higher temperature. Aliquots of saturated *trans*-stilbene solutions were transferred through a coarse filter into a tared volumetric flask to determine the amount of sample and diluted quantitatively with methanol for spectrophotometric analysis at 294 nm on a Bausch and Lomb Spectronic 2000. Concentrations of the dilute solutions were determined from a Beer-Lambert law absorbance *versus* concentration working curve derived from measured absorbances of standard solutions of known molar concentration ranging from  $1.38 \times 10^{-5}$  Molar to  $4.62 \times 10^{-5}$  Molar. The calculated molar absorptivity of  $\epsilon \approx 28,850$  Liter  $\text{mol}^{-1} \text{cm}^{-1}$  was constant over the concentration range. Identical molar absorptivities were obtained for select *trans*-stilbene standard solutions that contained up to 5 volume percent of the neat alkane and 2-propanol cosolvents. Experimental *trans*-stilbene solubilities in the six binary alkane + 2-propanol solvent mixtures are listed in Table I. Numerical values represent the average of between four and eight independent determinations, with the measured values being reproducible to within  $\pm 1.5\%$ .

## RESULTS AND DISCUSSION

Acree and Zvaigzne [4] suggested possible mathematical representations for isothermal solubility data based upon either a Combined NIBS/Redlich-Kister model

$$\ln x_A^{\text{sat}} = x_B^o \ln(x_A^{\text{sat}})_B + x_C^o \ln(x_A^{\text{sat}})_C + x_B^o x_C^o \sum_{i=0}^N S_i (x_B^o - x_C^o)^i \quad (1)$$

or Modified Wilson equation

$$\ln[a_A(s)/x_A^{\text{sat}}] = 1 - x_B^o \{1 - \ln[a_A(s)/(x_A^{\text{sat}})_B]\} / (x_B^o + x_C^o \Lambda_{BC}^{\text{adj}}) - x_C^o \{1 - \ln[a_A(s)/(x_A^{\text{sat}})_C]\} / (x_B^o \Lambda_{CB}^{\text{adj}} + x_C^o) \quad (2)$$

where the various  $S_i$  and  $\Lambda_{ij}^{\text{adj}}$  "curve-fit" parameters can be evaluated *via* least squares analysis. In Eqs. (1) and (2)  $x_B^o$  and  $x_C^o$  refer to the

TABLE I Experimental mole fraction solubilities of *trans*-stilbene ( $x_A^{sat}$ ) in binary alkane (*B*) + 2-propanol (*C*) solvent mixtures at 25.0°C

$x_c^o$	$x_A^{sat}$
<b>Hexane (<i>B</i>) + 2-Propanol (<i>C</i>)</b>	
0.0000	0.00960
0.1708	0.00960
0.3037	0.00920
0.5379	0.00746
0.6080	0.00667
0.7153	0.00560
0.8690	0.00398
0.9381	0.00333
1.0000	0.00279
<b>Heptane (<i>B</i>) + 2-Propanol (<i>C</i>)</b>	
0.0000	0.01085
0.1858	0.01094
0.3263	0.01021
0.5623	0.00789
0.6553	0.00685
0.7383	0.00595
0.8824	0.00410
0.9409	0.00340
1.0000	0.00279
<b>Octane (<i>B</i>) + 2-Propanol (<i>C</i>)</b>	
0.0000	0.01241
0.2055	0.01179
0.3621	0.01061
0.5839	0.00826
0.6761	0.00709
0.7662	0.00590
0.8960	0.00404
0.9457	0.00344
1.0000	0.00279
<b>Cyclohexane (<i>B</i>) + 2-Propanol (<i>C</i>)</b>	
0.0000	0.01374
0.1575	0.01351
0.2626	0.01267
0.4753	0.00979
0.5867	0.00829
0.6836	0.00676
0.8349	0.00459
0.9173	0.00363
1.0000	0.00279
<b>Methylcyclohexane (<i>B</i>) + 2-Propanol (<i>C</i>)</b>	
0.0000	0.01413
0.1624	0.01368
0.2984	0.01238
0.5146	0.00973
0.6057	0.00839
0.7188	0.00658

TABLE I (Continued)

$x_c^o$	$x_A^{sat}$
0.8625	0.00448
0.9331	0.00355
1.0000	0.00279
2,2,4-Trimethylpentane (B) + 2-Propanol (C)	
0.0000	0.00803
0.2145	0.00800
0.3638	0.00747
0.5861	0.00613
0.6858	0.00539
0.7620	0.00470
0.8852	0.00366
0.9473	0.00318
1.0000	0.00279

TABLE II Mathematical representation of *trans*-stilbene solubilities in several binary alkane (B) + 2-propanol (C) solvent mixtures

Binary solvent system Component (B) + Component (C)	Eq. (1)		Eq. (2)	
	$S_i^a$	% Dev. <sup>b</sup>	$\Lambda_{ij}^{adj,c}$	% Dev.
Hexane + 2-Propanol	1.628	0.3	1.840	0.6
	-0.155		1.695	
	-0.058			
Heptane + 2-Propanol	1.828	0.7	1.927	0.5
	-0.279		1.927	
	0.135			
Octane + 2-Propanol	1.827	0.5	1.434	0.5
	-0.552		2.217	
	0.187			
Cyclohexane + 2-Propanol	1.749	0.5	2.971	0.8
	0.023		1.782	
Methylcyclohexane + 2-Propanol	1.843	0.4	2.362	0.7
	-0.299		2.014	
	0.059			
2,2,4-Trimethylpentane + 2-Propanol	1.428	0.3	1.492	0.5
	-0.160		1.608	
	-0.089			

<sup>a</sup> Combined NIBS/Redlich-Kister curve-fit parameters are ordered as  $S_0$ ,  $S_1$  and  $S_2$ .

<sup>b</sup> Deviation (%) =  $(100/N) \sum |(x_A^{sat})^{calc} - (x_A^{sat})^{exp}|$ .

<sup>c</sup> Adjustable parameters for the Modified Wilson equation are ordered as  $\Lambda_{BC}^{adj}$  and  $\Lambda_{CB}^{adj}$ .

initial mole fraction composition of the binary solvent calculated as if solute (A) were not present,  $a_A(s)$  is the activity of the solid solute,  $N$  is the number of curve-fit parameters used, and  $(x_A^{sat})_i$  is the saturated

mole fraction solubility of the solute in pure solvent  $i$ . The numerical value of  $a_A(s)$  used in the Modified Wilson computations was  $a_A(s) = 0.06227$  [5].

The ability of Eqs. (1) and (2) to mathematically represent the experimental solubility of *trans*-stilbene in the six binary alkane + 2-propanol mixtures is summarized in Table II in the form of "curve-fit" parameters and percent deviations in back-calculated solubilities. Each percent deviation is based upon the measured *trans*-stilbene solubility data at the seven different binary solvent compositions. Careful examination of Table II reveals that both equations provided a fairly accurate mathematical representation for how the solubility of *trans*-stilbene varies with solvent composition. For the six *trans*-stilbene systems studied, Eqs. (1) and (2) were found to describe the experimental data to within an overall average absolute deviation of approximately  $\pm 0.5\%$ , which is less than the experimental uncertainty.

### **Acknowledgements**

Carmen Hernández and Tina Sharp thank the University of North Texas and the U.S. Department of Education for support provided under the Ronald E. McNair Postbaccalaureate Achievement Program.

### **References**

- [1] Acree, W. E. Jr. (1994). Polycyclic Aromatic Hydrocarbons in Pure and Binary Solvents, Vol. 54, In: *IUPAC Solubility Data Series*; Oxford University Press: Oxford, United Kingdom.
- [2] Acree, W. E. Jr. (1995). Polycyclic Aromatic Hydrocarbons: Binary Nonaqueous Systems: Part 1 (Solutes A–E). Vol. 58, In: *IUPAC Solubility Data Series*; Oxford University Press: Oxford, United Kingdom.
- [3] Acree, W. E. Jr. (1995). Polycyclic Aromatic Hydrocarbons: Binary Nonaqueous Systems: Part 2 (Solutes F–Z). Vol. 59, In: *IUPAC Solubility Data Series*; Oxford University Press: Oxford, United Kingdom.
- [4] Acree, W. E. Jr. and Zvaigzne, A. I. (1991). *Thermochim. Acta*, **178**, 151.
- [5] Fletcher, K. S., McHale, M. E. R., Coym, K. S. and Acree, W. E. Jr. (1997). *Can. J. Chem.*, **75**, 258.