

This article was downloaded by:

On: 28 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Physics and Chemistry of Liquids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713646857>

### Solubility of Xanthene in Organic Nonelectrolyte Solvents: Comparison of Observed Versus Predicted Values Based Upon Mobile Order Theory

Cassandra I. MonÁrrez<sup>a</sup>; Dawn M. Stovall<sup>a</sup>; Jee H. Woo<sup>a</sup>; Priscilla Taylor<sup>a</sup>; William E. Acree Jr.<sup>a</sup>

<sup>a</sup> Department of Chemistry, University of North Texas, Denton, TX, USA

Online publication date: 27 October 2010

**To cite this Article** MonÁrrez, Cassandra I. , Stovall, Dawn M. , Woo, Jee H. , Taylor, Priscilla and Acree Jr., William E.(2002) 'Solubility of Xanthene in Organic Nonelectrolyte Solvents: Comparison of Observed Versus Predicted Values Based Upon Mobile Order Theory', *Physics and Chemistry of Liquids*, 40: 6, 703 – 714

**To link to this Article:** DOI: 10.1080/0031910021000018581

**URL:** <http://dx.doi.org/10.1080/0031910021000018581>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

# SOLUBILITY OF XANTHENE IN ORGANIC NONELECTROLYTE SOLVENTS: COMPARISON OF OBSERVED *VERSUS* PREDICTED VALUES BASED UPON MOBILE ORDER THEORY

CASSANDRA I. MONÁRREZ, DAWN M. STOVALL,  
JEE H. WOO, PRISCILLA TAYLOR and  
WILLIAM E. ACREE, JR.\*

*Department of Chemistry, University of North Texas, P.O. Box 305070,  
Denton, TX 76203-5070, USA*

*(Received 1 May 2002)*

Experimental solubilities are reported at 25.0°C for xanthene dissolved in 34 different organic nonelectrolyte solvents containing ether-, chloro-, hydroxy-, cyano and *t*-butyl-functional groups. Results of these measurements are used to test the applications and limitations of expressions derived from Mobile Order theory. For the 27 solvents for which predictions could be made computations show that Mobile order theory does provide fairly reasonable estimates of the saturation mole fraction solubilities. Average absolute deviation between predicted and observed values is 71.4%.

*Keywords:* Xanthene solubilities; Organic nonelectrolyte solvents; Solubility predictions

## INTRODUCTION

Free energy of solvation is an important thermodynamic variable that quantifies the free energy difference between a molecule in the gas phase and the molecule dissolved in a solvent. Free energies of solva-

---

\*Corresponding author.

tion provide valuable information regarding molecular interactions between dissolved solute and surrounding solvent molecules, and can be used to calculate numerical values of partition coefficients that describe the equilibrium distribution of a solute between two immiscible liquid phases. For example, the octanol–water partition coefficient is the free energy of solvation of the solute molecule in 1-octanol minus its free energy of solvation in water. Solvation free energies and partition coefficients are of critical importance in many pharmaceutical, environmental and chemical engineering applications. Solute partitioning between two immiscible phases is the basis for all chromatographic separations. Correlations have been derived for predicting brain–blood partitioning of organic solutes from solutes' measured and/or calculated free energies of solvation [1,2], and for estimating aqueous solubilities, soil adsorptions, bioaccumulations and toxicities of organic compounds from experimental octanol–water partition coefficient data [3–11].

Historically, many of the very early studies focussed exclusively on developing correlational equations based upon octanol–water partition coefficients. Recent studies have shown that the octanol–water partition coefficient may not necessarily be the best indicator of how likely it is for a particular solute molecule to penetrate a lipid bilayer, skin, brain or central nervous system, or to accumulate in different tissues and body organs. Experimental studies have been expanded to include additional organic solvents, as well as aqueous micellar solvent media, and to use solute descriptors calculable from structural considerations and/or easily measured thermodynamic quantities.

In this regard, Abraham and coworkers [12–19] developed expressions for describing the partition of solutes between water and a given solvent

$$\log P = c + r \cdot R_2 + s \cdot \pi_2^H + a \cdot \sum \alpha_2^H + b \cdot \sum \beta_2^H + v \cdot V_x \quad (1)$$

and between the gas phase and a given solvent

$$\log L = c + r \cdot R_2 + s \cdot \pi_2^H + a \cdot \sum \alpha_2^H + b \cdot \sum \beta_2^H + l \cdot \text{Log } L^{16} \quad (2)$$

The dependent variables in Eqs. (1) and (2) are the  $\log P$  (the partition coefficient of solute(s) between water and a given solvent) and  $\log L$  (Ostwald solubility coefficient). The independent variables are the

solute descriptors as follows:  $R_2$  and  $V_x$  refer to the excess molar refraction and McGowan volume of the solute, respectively,  $\sum \alpha_2^H$  and  $\sum \beta_2^H$  are measures of the solute's hydrogen-bond acidity and hydrogen-bond basicity,  $\pi_2^H$  denotes the solute's dipolarity–polarizability descriptor, and  $\text{Log } L^{16}$  is the solute's gas phase dimensionless Ostwald partition coefficient into hexadecane at 298 K. The Ostwald partition coefficient,  $L$ , is the inverse of the Henry's law constant ( $\text{Pa m}^3 \text{ mole}^{-1}$ ). It should be noted that the various  $c, r, s, a, b, v$  and  $l$  coefficients depend on the solvent phase under consideration. The  $r$ -coefficient gives the tendency of the phase to interact with solutes through polarizability-type interactions, mostly via electron pairs. The  $s$ -coefficient is a measure of the solvent phase dipolarity–polarity, while the  $a$ - and  $b$ -coefficients represent the solvent phase hydrogen-bond basicity and hydrogen-bond acidity, respectively. The  $l$ - and  $v$ -coefficients are a combination of the work needed to create a solvent cavity wherein the solute will reside, and the general dispersion interaction energy between the solute and solvent phase. In the case of partition coefficients, where two solvent phases are involved, the  $c, r, s, a, b, v$  and  $l$  coefficients represent differences in the solvent phase properties.

Several earlier studies [20–24] developed the computational methodology for determining the various solute descriptions from measured solubility data for crystalline nonelectrolyte solutes dissolved in organic solvents for which the solvent regression coefficients were known. Solutes studied included *trans*-stilbene, diuron, monuron, buckminsterfullerene, benzil and ferrocene. Diuron and monuron had the larger numerical values of the  $\sum \alpha_2^H$  and  $\sum \beta_2^H$  solute descriptors. Continued development of additional correlation equations requires the establishment of large solubility and activity coefficient databases for each solvent system to be studied. The databases should contain solutes that span as wide of a range of solute descriptors as possible. For this reason, we are in the process of measuring solubility data for several crystalline organic compounds. In the present communication we report xanthene solubilities at 25°C in 34 different organic solvents of varying polarity and hydrogen bonding capability. Results of these measurements are used to further test the applications and limitations of predictive expressions derived from Mobile Order theory. Subsequent papers will consider the mathematical correlations of xanthene solubilities based upon other solution models.

## MATERIALS AND METHODS

Xanthene (Aldrich, 98%) was recrystallized several times from anhydrous methanol before use. *n*-Hexane (Aldrich, 99%), *n*-heptane (Aldrich, HPLC), *n*-octane (Aldrich 99+ %, anhydrous), *n*-nonane (TCI, 99+ %), *n*-decane (TCI, 99+ %), *n*-hexadecane (Aldrich, 99%), cyclohexane (Aldrich, HPLC), methylcyclohexane (Aldrich, 99+ %, anhydrous), cyclooctane (Lancaster Synthesis, 99+ %), 2,2,4-trimethylpentane (Aldrich, HPLC), methanol (Aldrich, 99.9+ %), ethanol (Aaper Alcohol and Chemical Company, absolute), 1-propanol (Aldrich, 99+ %, anhydrous), 2-propanol (Aldrich, 99+ %, anhydrous), 1-butanol (Aldrich HPLC, 99.8+ %), 2-butanol (Aldrich, 99+ %, anhydrous), 1-pentanol (Aldrich, 99+ %), 1-hexanol (Alfa Aesar, 99+ %), 1-heptanol (Alfa Aesar, 99+ %), 1-octanol (Aldrich, 99+ %, anhydrous), 1-decanol (Alfa Aesar, 99+ %), 2-pentanol (Acros 99+ %), 2-methyl-2-butanol (Acros, 99+ %), 2-methyl-1-propanol (Aldrich, 99+ %, anhydrous), 3-methyl-1-butanol (Aldrich, 99+ %, anhydrous), 4-methyl-2-pentanol (Acros, 99+ %), 2-methyl-1-pentanol (Aldrich, 99%), 2-methyl-2-propanol (Arco Chemical Company, 99+ %), cyclopentanol (Aldrich, 99+ %), dibutyl ether (Aldrich, 99%), methyl *tert*-butyl ether (Arco, 99.9+ %), carbon tetrachloride (Aldrich, 99.5+ %, anhydrous), 1,2-dichloroethane (Aldrich, 99.8%, anhydrous) and acetonitrile (Aldrich, 99.8%, anhydrous) were stored over molecular sieves and distilled shortly before use. Gas chromatographic analysis showed solvent purities to be 99.7 mole percent or better.

Excess solute and solvent were placed in amber glass bottles and allowed to equilibrate in a constant temperature water bath at  $25.0 \pm 0.1^\circ\text{C}$  for at least three days (often longer). Attainment of equilibrium was verified both by repetitive measurements after several additional days and by approaching equilibrium from supersaturation by pre-equilibrating the solutions at a higher temperature. Aliquots of saturated xanthene solutions were transferred through a coarse filter into a tared volumetric flask to determine the amount of sample and diluted quantitatively with 2-propanol for spectrophotometric analysis at 280 nm on a Bausch and Lomb Spectronic 2000. Concentrations of the dilute solutions were determined from a Beer-Lambert law absorbance *versus* concentration working curve for nine standard solutions.

The calculated molar absorptivity of xanthene at 280 nm varied slightly with concentration, ranging from a value of  $\epsilon \approx 2240 \text{ L mol}^{-1} \text{ cm}^{-1}$  ( $1.44 \times 10^{-4} \text{ M}$ ) to  $\epsilon \approx 2190 \text{ L mol}^{-1} \text{ cm}^{-1}$  ( $4.82 \times 10^{-4} \text{ M}$ ).

Experimental molar concentrations were converted to (mass/mass) solubility fractions by multiplying by the molar mass of xanthene, volume(s) of volumetric flask(s) used and any dilutions required to place the measured absorbances on the Beer-Lambert law absorbance *versus* concentration working curve, and then dividing by the mass of the saturated solution analyzed. Mole fraction solubilities were computed from (mass/mass) solubility fractions using the molar masses of the solute and solvent. Experiment mole fraction solubilities,  $X_A^{\text{sat}}$ , are listed in Table I for xanthene dissolved in 34 organic solvents. Numerical values represent the average of between four and eight independent determinations, with the measurements were reproducible to within  $\pm 1.5\%$ .

## RESULTS AND DISCUSSION

Solvents listed in Table I include both noncomplexing alkanes and self-associating alcohols. Of the many solution models proposed in recent years, mobile order theory is perhaps the only one that is capable of describing solute behavior in such a wide range of solvent mixtures. The basic model [25–32] assumes that all molecular groups perpetually move, and that neighbors of a given kind of external atom in a molecule constantly change identity. All molecules of a given kind dispose of the same volume, equal to the total volume  $V$  of the liquid divided by the number  $N_A$  of molecules of the same kind, i.e.,  $\text{Dom } A = V/N_A$ . The center of this domain perpetually moves. The highest mobile disorder is achieved whenever groups visit all parts of their domain without preference. Preferential contacts lead to deviations with respect to this “random” visiting. This is especially true in the case of hydrogen-bonding as specific interactions result in a specific orientation of the “donor” molecule with respect to an adjacent “acceptor” molecule.

In the case of an inert crystalline solute dissolved in a self-associating solvent, mobile order theory expresses the volume fraction saturation

TABLE I Comparison between experimental xanthene mole fraction solubilities and predicted values based upon mobile order theory

<i>Organic solvent</i>	$(X_A^{\text{sat}})^{\text{exp.a}}$	$(X_A^{\text{sat}})^{\text{calc}}$	%Dev <sup>b</sup>
<i>n</i> -Hexane	0.02949	0.03217	9.1
<i>n</i> -Heptane	0.03543	0.03390	-4.3
<i>n</i> -Octane	0.03976	0.03839	-3.4
<i>n</i> -Nonane	0.04306	0.04439	3.1
<i>n</i> -Decane	0.04610	0.04642	0.7
<i>n</i> -Hexadecane	0.06835	0.06431	-5.9
Cyclohexane	0.04203	0.04159	-1.0
Methylcyclohexane	0.04275	0.04459	4.3
Cyclooctane	0.05414	0.05743	6.1
2,2,4-Trimethylpentane	0.02451	0.02551	4.1
Dibutyl ether	0.08310	0.1428	71.8
Methyl <i>tert</i> -butyl ether	0.07846		
1,2-Dichloroethane	0.1549	0.2203	42.2
Carbon tetrachloride	0.1237	0.1431	15.7
Acetonitrile	0.01970	0.1987	908.6
Methanol	0.004455	0.01089	144.4
Ethanol	0.006231	0.01693	171.7
1-Propanol	0.01166	0.02121	81.9
2-Propanol	0.008643	0.02458	184.4
1-Butanol	0.01756	0.02679	52.6
2-Butanol	0.01254	0.02112	68.4
2-Methyl-1-propanol	0.01077	0.01686	56.5
2-Methyl-2-propanol	0.01112	0.01420	27.7
1-Pentanol	0.02212	0.02890	30.7
2-Pentanol	0.01766		
3-Methyl-1-butanol	0.01633		
2-Methyl-2-butanol	0.01946		
1-Hexanol	0.02831	0.02749	-2.9
2-Methyl-1-pentanol	0.01969		
4-Methyl-2-pentanol	0.01762		
1-Heptanol	0.03340	0.03115	-6.7
1-Octanol	0.03800	0.03445	-9.3
1-Decanol	0.04528	0.04026	-11.1
Cyclopentanol	0.02886		

<sup>a</sup>Numerical values represent the average of between four and eight independent determinations, with the measurements being reproducible to  $\pm 1.5\%$ ; <sup>b</sup>Deviations (%) =  $100[(X_A^{\text{sat}})^{\text{calc}} - (X_A^{\text{sat}})^{\text{exp}}]/(X_A^{\text{sat}})^{\text{exp}}$ .

solubility,  $\phi_A^{\text{sat}}$ , as

$$\begin{aligned}
 \ln \phi_A^{\text{sat}} = & \ln a_A^{\text{solid}} - 0.5(1 - V_A/V_{\text{solvent}})\phi_{\text{solvent}} \\
 & + 0.5 \ln[\phi_A^{\text{sat}} + \phi_{\text{solvent}}(V_A/V_{\text{solvent}})] \\
 & - \phi_{\text{solvent}}^2 V_A (\delta'_A - \delta'_{\text{solvent}})^2 (RT)^{-1} \\
 & - r_{\text{solvent}}(V_A/V_{\text{solvent}})\phi_{\text{solvent}}
 \end{aligned} \quad (3)$$

where the  $r_{\text{solvent}} (V_A/V_{\text{solvent}}) \phi_{\text{solvent}}$  term represents the contributions resulting from hydrogen-bond formation between the solvent molecules. For most of the published applications,  $r_{\text{solvent}}$  was assumed to be unity for strongly associated solvents with single hydrogen-bonded chains such as monofunctional alcohols, to be two for water or diols, and to equal zero for nonassociated solvents such as saturated hydrocarbons. A more exact value for alcoholic solvents can be calculated based upon

$$r_{\text{solvent}} = (K_{\text{solvent}} \phi_{\text{solvent}} / V_{\text{solvent}}) / (1 + K_{\text{solvent}} \phi_{\text{solvent}} / V_{\text{solvent}}) \quad (4)$$

with a numerical value of  $K_{\text{solvent}} = 5000 \text{ cm}^3 \text{ mol}^{-1}$  assumed for all monofunctional alcohols.

If complexation does occur between the crystalline solute and solvent

$$\begin{aligned} \ln \phi_A^{\text{sat}} = & \ln a_A^{\text{solid}} - 0.5(1 - V_A/V_{\text{solvent}}) \phi_{\text{solvent}} \\ & + 0.5 \ln[\phi_A^{\text{sat}} + \phi_{\text{solvent}}(V_A/V_{\text{solvent}})] \\ & - \phi_{\text{solvent}}^2 V_A (\delta'_A - \delta'_{\text{solvent}})^2 (RT)^{-1} \\ & + \ln[1 + \phi_{\text{solvent}}(K_{A\text{Solvent}}/V_{\text{solvent}})] \end{aligned} \quad (5)$$

then an addition term involving the solute-solvent equilibrium constant,  $K_{A\text{Solvent}}$ , must be introduced to describe the solubility enhancement that arises as a result of specific interactions. A slightly more complex expression applies in the case of solute complexation with a self-associating solvent. The expression's mathematical form depends to a large extent upon the number and type of hydrogen-bond acid and hydrogen-bond base functional groups in the solute molecule being studied. The symbols  $\delta'_A$  and  $\delta'_{\text{solvent}}$  denote the modified solubility parameters of the solute and solvent, respectively,  $V_i$  is the molar volume, and  $a_A^{\text{solid}}$  is the activity of the solid solute. This latter quantity is defined as the ratio of the fugacity of the solid to the fugacity of the pure hypothetical supercooled liquid. The numerical value of  $a_A^{\text{solid}}$  can be computed from

$$\begin{aligned} \ln a_A^{\text{solid}} = & -\Delta H_A^{\text{fus}}(T_{\text{mp}} - T)/(RT T_{\text{mp}}) + (\Delta C_{p,A}/RT)(T_{\text{mp}} - T) \\ & - (\Delta C_{p,A}/T) \ln(T_{\text{mp}}/T) \end{aligned} \quad (6)$$



the solute's molar enthalpy of fusion,  $\Delta H_A^{\text{fus}}$ , and heat capacity difference,  $\Delta C_{p,A}$ , at the normal melting point temperature,  $T_{\text{mp}}$ . Contributions from nonspecified interaction are incorporated into mobile order theory through the  $\phi_{\text{solvent}}^2 V_A (\delta'_A - \delta'_{\text{solvent}})^2 (RT)^{-1}$  term.

It is noted that other research groups have treated nonspecific interactions differently, and have assumed solution models other than the Scatchard–Hildebrand solubility parameter theory. For example, Ruelle [33] in predicting solubilities in systems involving hydrogen-bond formation between a dissolved solute and surrounding solvent molecules, modified the Scatchard–Hildebrand expression by multiplying it by the fraction of time during which the solute is not bound to the solvent [i.e., during which the distribution between the solvent and unbound solute molecules can still be considered to occur at random]. Ruelle's treatment further assumed that nonspecific interactions involving the bound solute were negligible. The theoretical justification for Ruelle's modification was not given, and it is not clear to us what mathematical form the integral  $(\Delta G_{A\text{Solvent}})_{\text{phys}}$  for the binary solution would have to take in order to give

$$(\Delta G_A)_{\text{phys}} = \left\{ 1 / [1.0 + \max(K_{\text{Oi}}, K_{\text{OHi}})(\phi_{\text{solvent}}/V_{\text{solvent}})] \right\} \cdot \phi_{\text{solvent}}^2 V_A (\delta'_A - \delta'_{\text{solvent}})^2 \quad (7)$$

whenever  $(\Delta G_{A\text{Solvent}})_{\text{phys}}$  is differentiated with respect to the number of moles of solute present. The differentiation is required in deriving the solubility equation. Readers are reminded that any modification to the  $(\Delta G_A)_{\text{phys}}$  expression must also show up in the corresponding  $(\Delta G_{\text{solvent}})_{\text{phys}}$  expression, the latter expression being obtained by differentiating  $(\Delta G_{A\text{Solvent}})_{\text{phys}}$  with respect to the number of molecules of solvent. In Eq. (7)  $\max(K_{\text{Oi}}, K_{\text{OHi}})$  stands for the association constant governing the strongest intermolecular H-bond displayed by the molecular groups in solution. We have elected to use the Scatchard–Hildebrand solubility parameter theory, rather than the Ruelle modification, because we have serious reservations about whether nonspecific interactions for bound molecules are truly negligible as assumed by Ruelle in proposing Eq. (7).

Predictive application of Eqs. (3) and (5) is relatively straightforward. First, an average numerical value of  $\delta'_{\text{xanth}} = 20.34 \text{ MPa}^{1/2}$  is

computed by requiring that each equation (with  $r_{\text{solvent}}=0$  and/or  $K_{A\text{Solvent}}=0$ ) perfectly describes xanthene mole fraction solubility data in *n*-hexane ( $\delta'_{\text{xanth}} = 20.45 \text{ MPa}^{1/2}$ ), *n*-heptane ( $\delta'_{\text{xanth}} = 20.28 \text{ MPa}^{1/2}$ ), and *n*-octane ( $\delta'_{\text{xanth}} = 20.29 \text{ MPa}^{1/2}$ ). The numerical value of a  $\alpha_A^{\text{solid}} = 0.2091$  is calculated using Eq. (6) with  $\Delta H_A^{\text{fus}} = 19.20 \text{ kJ mol}^{-1}$  and  $T_{\text{mp}} = 373.65 \text{ K}$  [34]. The two heat capacity terms in Eq. (6) were set equal to zero as we were unable to find heat capacity data for the subcooled liquid solute. A numerical value of  $V_{\text{xanth}} = 150.0 \text{ cm}^3 \text{ mol}^{-1}$  was used for the molar volume of the hypothetical subcooled liquid solute.

Table I summarizes the predictive ability of Mobile Order theory for the various organic solvents for which both xanthene solubility data and modified solubility parameters could be found. Solvent molar volumes and modified solubility parameters are listed in Table II. We were unable to find modified solubility parameters for all of the secondary and branched alcohols. The modified solubility parameters account for only nonspecific interactions, and in the case of the alcoholic solvents the hydrogen-bonding contributions have been removed. Numerical values of  $\delta'_{\text{solvent}}$  were obtained from published compilations [28,29,31,32], and were either deduced by regressing actual solubility data of solid *n*-alkanes in organic solvent in accordance with the configurational entropic model of Huyskens and Haulait-Pirson [35] or estimated using known values for similar organic solvents. Examination of the entries in Table I reveals that Mobile Order theory does provide fairly reasonable (though by no means perfect) estimates of the solubility behavior of xanthene in a wide range of organic solvents. Average absolute deviation between predicted and observed values is 71.4%. We do note that Mobile Order theory grossly overpredicts the solubility of xanthene in acetonitrile. Similar failures were noted previously in the case of anthracene [36], fluoranthene [37], pyrene [37], phenanthrene [38] and diuron [39]. We suspect that the overprediction results from either failure of the Scatchard-Hildebrand solubility parameter theory to describe the nonspecific interactions between acetonitrile and the afore-mentioned solutes, or perhaps from an incorrect value for this solvent's modified solubility parameter. Deviations between observed and predicted values were significantly reduced, to 39.2%, when acetonitrile was excluded from the comparison.

TABLE II Solvent and solute properties used in mobile order predictions

<i>Component (i)</i>	$V_{ij}/(\text{cm}^3 \text{mol}^{-1})$	$\delta'_i/(\text{MPa}^{1/2})^a$
<i>n</i> -Hexane	131.51	14.56
<i>n</i> -Heptane	147.48	14.66
<i>n</i> -Octane	163.46	14.85
<i>n</i> -Nonane	179.87	15.07
<i>n</i> -Decane	195.88	15.14
<i>n</i> -Hexadecane	294.12	15.61
Cyclohexane	108.76	14.82
Methylcyclohexane	128.32	15.00
Cyclooctane	134.9	15.40
2,2,4-Trimethylpentane	166.09	14.30
Dibutyl ether	170.3	17.45
Methanol	40.7	19.25
Ethanol	58.7	17.81
1-Propanol	75.10	17.29
2-Propanol	76.90	17.60
1-Butanol	92.00	17.16
2-Butanol	92.4	16.60
2-Methyl-1-propanol	92.8	16.14
2-Methyl-2-propanol	94.3	15.78
1-Pentanol	108.6	16.85
1-Hexanol	125.2	16.40
1-Heptanol	141.9	16.39
1-Octanol	158.3	16.38
1-Decanol	191.6	16.35
Tetrachloromethane	97.08	17.04
1,2-Dichloroethane	78.8	20.99
Acetonitrile	52.9	23.62
Xanthene <sup>b</sup>	150.0	20.34 <sup>c</sup>

<sup>a</sup>Tabulated values are taken from a compilation given in Ruelle *et al.* [28,29,31,32];

<sup>b</sup>The numerical value of  $a_A^{\text{solid}} = 0.2091$  was calculated from Eq. (8) using  $\Delta H_A^{\text{fus}} = 19.20 \text{ kJ mol}^{-1}$  and  $T_{\text{mp}} = 373.65 \text{ K}$  [34]; <sup>c</sup>Numerical value was calculated using the measured xanthene mole fraction solubilities in *n*-hexane, *n*-heptane and *n*-octane, in accordance with Eqs. (3) and (5); with  $r_{\text{solvent}} = 0$  and/or  $K_{A\text{Solvent}} = 0$ .

Readers are reminded that in evaluating the applicability of Mobile Order theory one must realize that many of these particular systems are highly nonideal, and that the experimental solubility data covers over a 34-fold range in mole fraction. Had an ideal solution been assumed, then the predicted mole fraction solubility would be  $X_A^{\text{sat}} = a_A^{\text{solid}} = 0.2091$  for each solvent. The ideal solution approximation corresponds to a considerably larger average absolute deviation of 976% between predicted and observed values for the nonalcoholic solvents studied.

### Acknowledgment

This work was supported in part by the University of North Texas Research Council.

### References

- [1] F. Lombardo, J.F. Blake and W.J. Curatolo (1996). *J. Med. Chem.*, **39**, 4750.
- [2] G.M. Keseru and L. Molnar (2001). *J. Chem. Inf. Comput. Sci.*, **41**, 120.
- [3] W.J. Lyman, W.F. Reehl and D.H. Rosenblatt (1982). *Handbook of Chemical Property Estimation Methods*, Chapter 1. McGraw-Hill, New York, NY.
- [4] K. Vereschuren (1983). *Handbook of Environmental Data on Organic Chemicals*, 2nd Edn. Van Nostrand-Reinhold, New York, NY.
- [5] A. Finizio, F. Sicbaldi and M. Vighi (1995). *SAR QSAR Environ. Res.*, **3**, 71.
- [6] A. Finizio, M. Vighi and D. Sandroni (1997). *Chemosphere*, **34**, 131.
- [7] D. Domine, J. Devillers and W. Karcher (1996). J. Devillers (Ed.), *Neural Networks in QSAR and Drug Design*, Chapter 2. Academic Press, New York, NY.
- [8] J.R. Baker, J.R. Mihelcic, D.C. Luehrs and J.P. Hickey (1997). *Water Environ. Res.*, **69**, 136.
- [9] C. Hansch (1993). *Acc. Chem. Res.*, **26**, 147.
- [10] X. Lu, S. Tao, H. Hu and R.W. Dawson (2000). *Chemosphere*, **41**, 1675.
- [11] J.R. Baker, J.R. Mihelcic and E. Shea (2000). *Chemosphere*, **41**, 813.
- [12] M.H. Abraham (1993). *Chrm. Soc. Rev.*, **22**, 73, and references therein.
- [13] M.H. Abraham, H.S. Chadha, G.S. Whiting and R.C. Mitchell (1994). *J. Pharm. Sci.*, **83**, 1085.
- [14] M.H. Abraham, K. Takacs-Novak and R.C. Mitchell (1997). *J. Pharm. Sci.*, **86**, 310.
- [15] M.H. Abraham, H.S. Chadha, G.S. Whiting and R.C. Mitchell (1994). *J. Pharm. Sci.*, **83**, 1257.
- [16] J.A. Gratton, M.H. Abraham, M.W. Bradbury and H.S. Chadha (1997). *J. Pharm. Pharmacol.*, **49**, 1211.
- [17] M.H. Abraham, F. Martins and R.C. Mitchell (1997). *J. Pharm. Pharmacol.*, **49**, 919.
- [18] M.H. Abraham and P.K. Weathersby (1994). *J. Pharm. Sci.*, **83**, 1450.
- [19] M.H. Abraham, R. Kumarsingh, J.E. Cometto-Muniz and W.S. Cain (1998). *Arch. Toxicol.*, **72**, 227.
- [20] M.H. Abraham, C.E. Green, W.E. Acree, Jr., C.E. Hernández and L.E. Roy (1998). *J. Chem. Soc., Perkin Trans.*, **2**, 2677.
- [21] M.H. Abraham, C.E. Green and W.E. Acree, Jr. (2000). *J. Chem. Soc., Perkin Trans.*, **2**, 281.
- [22] M.H. Abraham, N. Benjelloun-Dekhama, J.M.R. Gola, W.E. Acree, Jr., W.S. Cain and J.E. Cometto-Muniz (2000). *New J. Chem.*, **24**, 825.
- [23] C.E. Green, M.H. Abraham, W.E. Acree, Jr., K.M. De Fina and T.L. Sharp (2000). *Pest Manag. Sci.*, **56**, 1046.
- [24] W.E. Acree, Jr. and M.H. Abraham (2002). *J. Solution Chem.*, **31**, 293.
- [25] P.L. Huyskens and G.G. Siegel (1988). *Bull. Soc. Chim. Belg.*, **97**, 821.
- [26] G.G. Siegel, P.L. Huyskens and G. Vanderheyden (1990). *Ber. Bunsenges. Phys.*, **94**, 549.
- [27] P.L. Huyskens (1992). *J. Mol. Struct.*, **274**, 223.
- [28] P. Ruelle, C. Rey-Mermet, M. Buchmann, H. Nam-Tran, U.W. Kesselring and P.L. Huyskens (1991). *Pharm. Res.*, **8**, 840.

- [29] P. Ruelle, M. Buchmann, H. Nam-Tran and U.W. Kesselring (1992). *Int. J. Pharm.*, **87**, 47.
- [30] P. Ruelle, M. Buchmann and U.W. Kesselring (1994). *J. Pharm. Sci.*, **83**, 396.
- [31] P. Ruelle, E. Sarraf and U.W. Kesselring (1994). *Int. J. Pharm.*, **104**, 125.
- [32] P. Ruelle, E. Sarraf, L. Van den Berge, K. Seghers, M. Buchmann and U.W. Kesselring (1993). *Pharm. Acta Helv.*, **68**, 49.
- [33] P. Ruelle (1999). *J. Phys. Org. Chem.*, **12**, 769.
- [34] W.E. Acree Jr. (1991). *Thermochim. Acta*, **189**, 37, and references therein.
- [35] P.L. Huyskens and M.C. Haulait-Pirson (1985). *J. Mol. Liq.*, **31**, 135.
- [36] L.E. Roy, C.E. Hernández and W.E. Acree, Jr. (1999). *Polycyclic Aromat. Compd.*, **13**, 105.
- [37] L.E. Roy, C.E. Hernández and W.E. Acree, Jr. (1999). *Polycyclic Aromat. Compd.*, **13**, 205.
- [38] C.E. Hernández, K.M. De Fina, L.E. Roy, T.L. Sharp and W.E. Acree, Jr. (1999). *Can. J. Chem.*, **77**, 1465.
- [39] K.M. De Fina, T.L. Sharp, M.A. Spurgin, I. Chuca, W.E. Acree, Jr., C.E. Creen and M.H. Abraham (2000). *Can. J. Chem.*, **78**, 184.