

This article was downloaded by:

On: 28 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Physics and Chemistry of Liquids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713646857>

Oxidation of L-Phenylalanine by Manganese(VII) in Concentrated Sulfuric Acid Medium

H. Iloukhani^a; S. R. Ekvan^a; A. A. Rafati^a

^a Department of Chemistry, Faculty of Science, Bu-Ali Sina University, Hamadan, Iran

Online publication date: 27 October 2010

To cite this Article Iloukhani, H. , Ekvan, S. R. and Rafati, A. A.(2003) 'Oxidation of L-Phenylalanine by Manganese(VII) in Concentrated Sulfuric Acid Medium', *Physics and Chemistry of Liquids*, 41: 1, 25 – 31

To link to this Article: DOI: 10.1080/0031910021000018572

URL: <http://dx.doi.org/10.1080/0031910021000018572>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

OXIDATION OF L-PHENYLALANINE BY MANGANESE(VII) IN CONCENTRATED SULFURIC ACID MEDIUM

H. ILOUKHANI*, S.R. EKVAN and A.A. RAFATI

Department of Chemistry, Faculty of Science, Bu-Ali Sina University, Hamadan, Iran

(Received 9 June 2001)

Kinetics and mechanism of manganese(VII) oxidation of L-phenylalanine has been studied in aqueous sulfuric acid medium. Various hypotheses for the mechanism of acid catalysis have been tested and it has been found that rate is related to the activity of water in accord with Bunnett's hypothesis. The energy and entropy of activation and frequency factor have been calculated using the Eyring and Arrhenius plots. Mechanisms in agreement with the observations are reported. The experimental rate law for the reaction was found to be

$$\frac{-d[\text{Mn(VII)}]}{dt} = k_0[\text{L-phenylalanine}][\text{Mn(VII)}]$$

The reaction appears to involve an acid catalysis and the data showed that role of water molecules in the rate-determining step facilitating proton transfer which satisfies Bunnett's theory.

Keywords: Oxidation; L-phenylalanine; Manganese(VII); Bunnett's hypothesis

1. INTRODUCTION

The manganese(VII) oxidation of amino acids has received considerable attention in previous literature due to the importance of these compounds in biological systems. The oxidation has been widely studied in strong acidic media [1–8], where no noticeable auto-catalytic effects were found. However, in neutral or weakly alkaline medium, the literature consulted agrees as to the appearance of the cited auto-catalytic effect [9–13]. The purpose of this study is to analyze the manganese(VII) oxidation of L-phenylalanine in sulfuric acid. The applicability of Bunnett's theory was also tested.

2. EXPERIMENTAL PROCEDURES AND TECHNIQUES

Materials: All reagents used were of analytical grade from Merck, Sigma and Fluka. L-phenylalanine was assayed for amino acid content by standard method [14].

*Corresponding author. E-mail: iloukhani@basu.ac.ir

The solutions were prepared in buffered medium with sulfuric acid and potassium sulfate, with triply distilled water which was deionized. The pH and the ionic strength of the medium were adjusted with the required concentrations of sulfuric acid and potassium sulfate to get the desired values and tested by the Vogel method [15]. Sulfuric acid, potassium sulfate, potassium pyrophosphate, and mercuric chloride were used without further purification.


Kinetic measurements: The course of reaction was followed by measuring the absorbance of unreacted permanganate from time to time at 525 nm using a fw-265 spectrophotometer, supplied with a conventional thermostatic bath (± 0.1 K). The reactions were carried out in the temperature range of 303–323 K.

The reaction was usually followed up to 50% completion, and products of the reaction were identified as ammonia, carbon dioxide, and the corresponding aldehyde [16]. Addition of mercuric chloride to the reaction system did not induce the precipitation of mercurous chloride, indicating that no free radicals are produced in the system [17].

2.1. Stoichiometry

The stoichiometry of the reaction was determined by following the reaction for one hour and estimating both permanganate and aldehyde before and after half an hour. Permanganate was estimated by quenching a known amount of the reaction mixture in a known excess of iron(II) and titrating the unreacted iron(II) against standard cerium(IV). The aldehyde was estimated colorimetrically using chromotropic acid.

It was found that one mole of permanganate oxidized 2.5 moles of aldehyde. The stoichiometry of the reaction is represented by the following equation:



3. RESULT AND DISCUSSION

Under the pseudo-first-order condition of $[\text{L-phenylalanine}] \gg [\text{KMnO}_4]$ in 3.2 M sulfuric acid, plots of $\log(\text{absorbance})$ vs time were linear, showing that the reaction is first-order in $[\text{KMnO}_4]$ (Fig. 1). This was further confirmed by varying $[\text{KMnO}_4]$, which did not show any change, in the pseudo-first-order of rate constants, k'_0 (Table I). The reaction also exhibited a first-order dependence in $[\text{L-phenylalanine}]$ as could be seen from Fig. 2 and Table II.

At constant $[\text{KMnO}_4]_0$, $[\text{L-phenylalanine}]_0$, and temperature an increase in $[\text{H}_2\text{SO}_4]$ increased the rate constant k'_0 from $4.49 \times 10^{-4} \text{ s}^{-1}$ ($2.6 \text{ mol dm}^{-3} \text{ H}_2\text{SO}_4$) to $8.32 \times 10^{-4} \text{ s}^{-1}$ ($4.1 \text{ mol dm}^{-3} \text{ H}_2\text{SO}_4$) (Table III). The addition of sulfate and pyrophosphate ions caused no significant effect on the rate (Table IV).

Activation parameters are calculated using the following Eyring equation:

$$k_o = (RT/Nh) \exp((-\Delta H^\ddagger + T\Delta S^\ddagger)/(RT))$$




FIGURE 1 First-order plot for phenylalanine-permanganate reaction.

TABLE I Effect of concentration of permanganate on the rate of oxidation in strong acid medium at 313 K^a

$[KMnO_4]_0$ (mol dm ⁻³)	$k'_0 \times 10^4$ (s ⁻¹)
2.1×10^{-4}	5.718
3.3×10^{-4}	5.909
4.0×10^{-4}	6.101

^a[L-phenylalanine]₀ = 0.026 mol dm⁻³ and [H₂SO₄] = 3.2 mol dm⁻³.




FIGURE 2 Order of phenylalanine in oxidation reaction, plot of $\log k'_0 + 4$ vs $\log[\text{phenylalanine}] + 4$ at 313.15 K.

Plots of $\ln k_0/T$ vs $1/T$ should be linear. From the intercept and slope values of entropy ΔS^\ddagger and enthalpy ΔH^\ddagger of activation were evaluated, respectively, and results are given in Table V. Frequency factor, A and experimental energy, E_{exp} were also estimated by using Arrhenius plots of k_0 vs $1/T$ (Table V). In agreement with the experimental behavior, mechanisms is proposed to interpret the reaction pathways.

TABLE II Effect of concentration of L-phenylalanine on the rate of oxidation in strong acid medium at different temperatures^a

$[L\text{-Phenylalanine}]_0$ (mol dm ⁻³)	$k'_0 \times 10^4$ (s ⁻¹)		
	303 K	313 K	323 K
0.01	2.45	4.57	6.85
0.015	3.47	5.12	9.53
0.02	4.26	9.70	15.63
0.025	5.94	10.06	19.75
0.03	8.62	14.23	20.14

^a[KMnO₄]₀ = 0.01 [L-phenylalanine]₀ and [H₂SO₄] = 3.2 mol dm⁻³.

TABLE III Effect of concentration of sulfuric acid on the rate of oxidation at 313 K^a

$[H_2SO_4]$ (mol dm ⁻³)	$k'_0 \times 10^4$ (s ⁻¹)
2.6	4.49
3.0	5.90
3.7	6.21
4.1	8.32

^a[L-phenylalanine]₀ = 0.02 mol dm⁻³ and [KMnO₄]₀ ≅ 0.01 [L-phenylalanine]₀.

TABLE IV Effect of concentration of sulfate and pyrophosphate ions on the rate of oxidation at 313 K^a

Concentration (mol dm ⁻³)	$\log k'_0$ (s ⁻¹)
$[SO_4^{2-}] = 0.02$	3.30
$[P_2O_7^{4-}] = 0.02$	3.31

^a[L-phenylalanine]₀ = 0.02 mol dm⁻³, [H₂SO₄] = 3.2 mol dm⁻³ and [KMnO₄]₀ = 0.01 [L-phenylalanine]₀.

TABLE V Activation parameters on the oxidation of L-phenylalanine by permanganate in strong acid medium at different temperature^a

Temp. (K)	$k_0 \times 10^2$ (mol ⁻¹ dm ³ s ⁻¹)	ΔS^\ddagger (J mol ⁻¹ K ⁻¹)	ΔH^\ddagger (kJ mol ⁻¹)	E_{expt} (kJ mol ⁻¹)	$\ln A$
303	1.53	-168.44	30.14	42.72	50.73
313	2.39	-167.88	30.23	42.72	50.69
323	3.66	-167.32	30.32	42.72	50.66

^a[KMnO₄]₀ = 0.0001 mol dm⁻³ and [H₂SO₄] = 3.2 mol dm⁻³.

The expected species on permanganate oxidation in acid solution are manganese(III), manganese(IV), and manganese(VII) [18]. If manganese(III) or manganese(IV) were to be the reactive species, the addition of complexing agents like sulfate and pyrophosphate ions should have decreased the rate. In fact there is no effect of these ions on the rate [19], showing that only manganese(VII) is the most probable reactive species.





FIGURE 3 Bunnett's plot of $\log k'_0 - \log[\text{H}_2\text{SO}_4]$ vs $\log a_{\text{H}_2\text{O}}$.




With increasing $[\text{H}^+]$, the rate enhances due to protonation of the oxidant resulting in the formation of a more powerful oxidant, namely, acid permanganate by the following equilibrium:






The same explanation has also been given in the oxidation of alcohols [18] and esters [20] by acid permanganate.

Plots of $\log k'_0$ vs $\log[\text{H}_2\text{SO}_4]$ and $\log k'_0$ vs H'_0 were found to be linear, indicating that the reaction is acid catalyzed, but none of the above plots gave an ideal slope of unity. In view of the departure from the ideal behavior, the applicability of Bunnett's theory [21] was tested. A plot of $\log k'_0 - \log[\text{H}_2\text{SO}_4]$ vs $\log a_{\text{H}_2\text{O}}$ was linear Fig. 3, and the slope was found to be -0.848 . According to Bunnett's theory if the slope of such a plot is greater than -2 , it will indicate the involvement of a water molecule in the rate-determining step as a proton-abstracting base from the substrate. The values of Hammett's acidity function H'_0 and $\log a_{\text{H}_2\text{O}}$ have been taken from the literature [22,23].

The mechanisms consistent with the observed kinetic data is as follows:





The rate law for the above scheme comes out to be

$$\frac{-d[\text{Mn(VII)}]}{dt} = k_0[\text{L-phenylalanine}][\text{Mn(VII)}]$$

where $k_0 = Kk$.

From the bimolecular rate constants at different temperatures 303–323 K, activation parameters for reaction were evaluated (Table V).

Finally we concluded that the kinetic studies on the manganese(VII) oxidation of L-phenylalanine in moderately acid medium could get an insight in to the mechanism. The order in both [L-phenylalanine] and $[\text{KMnO}_4]$ was found to be unity. The effect of complexing agents like sulfate and pyrophosphate ion was nil, from which manganese(VII) was confirmed to be the reactive species.

The slope of the linear plot $\log k'_0 - \log [\text{H}_2\text{SO}_4]$ vs $\log a_{\text{H}_2\text{O}}$ (Bunnett's plot) was found to be -0.848 , indicating that involvement of the water molecule in the rate-determining step is as a proton abstracting base from the substrate.

Acknowledgments

Authors would like to thank the University authorities for providing the necessary facilities to carry out the work and Mr. E. Samei for his kind helpful for preparing the manuscript.

References

- [1] H. Iloukhani and H. Bahrami (1999). *Int. J. Chem. Kinet.*, **31**(2), 95.
- [2] H. Iloukhani, F. Rajali and A. Zeini (2001). *Phys. Chem. Liq.*, **39**(2), 239.
- [3] M. Moghadasi, N. Rashidi and H. Iloukhani (2001). *Phys. Chem. Liq.*, **39**(2), 269.
- [4] C.S. Ameta, P.N. Pande, H.L. Gupta and H.C. Chowdhry (1980). *Acta Phys. Chem.*, **26**, 89.
- [5] R.S. Verma, M.J. Reddy and V.R. Shastry (1976). *J. Chem. Soc. Perkin Trans.*, **II**, 469.
- [6] V.S. Rao, B. Sethuram and T.N. Rao (1979). *Int. J. Chem. Kinet.*, **11**, 165.
- [7] M. Adinara, B. Sethuram and T.H. Navaneeth Rao (1976). *J. Indian Chem. Soc.*, **53**, 877.
- [8] S. Bhargara, B. Sethuram and T.H. Navaneeth Rao (1976). *Indian. J. Chem.*, **14A**, 770.
- [9] J. De Andres, E. Brillas, J.A. Garrido and J.F. Perez Benito (1988). *J. Chem. Soc. Perkin Trans.*, **II**, 107.
- [10] A. Arrizabalaga, F.J. Andres-Ordan, M.Y. Fernandez-Aranguiz and R. Peche (1996). *Int. J. Chem. Kinet.*, **28**(11), 799.
- [11] A. Arrizabalaga, F.J. Andres-Ordan, M.Y. Fernandez-Aranguiz and R. Peche (1997). *Int. J. Chem. Kinet.*, **29**(3).
- [12] J.F. Perez, F. Mata Perez and E. Brillas (1987). *Can. J. Chem.*, **65**(10), 2329.
- [13] J.A. Gardio, J.F. Perez Benito, R.M. Rodriguez, J. De Andres and E. Brillas (1987). *J. Chem. Res.*, **11**, 380.
- [14] A.I. Vogel (1958). *Qualitative Organic Analysis*, p. 708. Longman and Green, London.

- [15] A.I. Vogel (1960). *Química Analítica Cuantitativa*. Kapelusz, Buenos Aires.
- [16] F. Fiegl (1966). *Testes in Organic Analysis*, p. 195. Elsevier, London.
- [17] A.I. Drummond and W.A. Waters (1953). *J. Chem. Soc.*, 3119.
- [18] R. Stewart (1965). *Oxidation in Organic Chemistry*. Academic, London.
- [19] E.A. Alexander and F.C. Tomkins (1939). *Trans. Faraday Soc.*, **35**, 1156.
- [20] P.V. Subba Rao (1971). *Z. Phys. Chem.*, (Leipzig), **246**, 352.
- [21] J.P. Bunnett (1961). *J. Am. Chem. Soc.*, **83**, 4968.
- [22] L. Hammett and A.J. Deyrup (1932). *J. Am. Chem. Soc.*, **54**, 2721.
- [23] J.P. Bunnett (1961). *J. Am. Chem. Soc.*, **83**, 4956.