

This article was downloaded by:

On: 28 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Physics and Chemistry of Liquids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713646857>

Density oscillations in a model of water and other similar liquids

M. Apostol^a; E. Preoteasa^b

^a Department of Theoretical Physics, Institute of Atomic Physics, Magurele-Bucharest, Romania ^b Life and Environmental Physics Department, Institute of Atomic Physics, Magurele-Bucharest, Romania

To cite this Article Apostol, M. and Preoteasa, E.(2008) 'Density oscillations in a model of water and other similar liquids', *Physics and Chemistry of Liquids*, 46: 6, 653 – 668

To link to this Article: DOI: 10.1080/00319100802146096

URL: <http://dx.doi.org/10.1080/00319100802146096>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Density oscillations in a model of water and other similar liquids

M. Apostol^{a*} and E. Preoteasa^b

^aDepartment of Theoretical Physics, Institute of Atomic Physics, Magurele-Bucharest, Romania; ^bLife and Environmental Physics Department, Institute of Atomic Physics, Magurele-Bucharest, Romania

(Received 21 March 2008; final version received 22 April 2008)

It is suggested that the dynamics of liquid water have a component consisting of O^{-2z} (oxygen) anions and H^{+z} (hydrogen) cations, where z is a (small) reduced effective electron charge. Such a model may apply to other similar liquids. The eigenmodes of density oscillations are derived for such a two-species ionic plasma, included the sound waves, and the dielectric function is calculated. The plasmons may contribute to the elementary excitations in a model introduced recently for the thermodynamics of liquids. It is shown that the sound anomaly in water can be understood on the basis of this model. The results are generalised to an asymmetric short-range interaction between the ionic species as well as to a multi-component plasma, and the structure factor is calculated.

Keywords: density oscillations; water; sound waves; ‘two-sounds anomaly’

PACS: 61.20.Qg; 62.60.+v; 78.40.Dw

1. Introduction

As simple as it may appear, water is still a complex liquid involving various interactions, as well as kinematic and dynamic correlations. It is widely agreed that the water molecule in liquid water preserves to some extent its integrity, especially the directionality of the sp^3 -oxygen orbitals, though it may be affected substantially by hydrogen bonds [1,2]. As such, it is conceived that water has a molecular electric moment, an intrinsic polarisability and hindered rotations (librations) which may affect its orientational polarisability. We examine herein another possible component of the dynamics of the liquid water, as resulting from the dissociation of the water molecule.

The water molecule H_2O has two H–O (hydrogen–oxygen) bonds which make an angle of ca. 109° in accordance with the tetragonal symmetry of the four hybridised sp^3 -oxygen orbitals. The ‘spherical’ diameter of water molecule is $\sim 2.75 \text{ \AA}$ and the inter-molecular spacing in liquid water under normal conditions is $a \sim 3 \text{ \AA}$. This suggests that the water molecule in liquid water, while preserving the directionality of the oxygen electronic orbitals, might be dissociated to a great extent. Dissociation models which assume $OH^- - H^+$ or $OH^- - H_3O^+$ pairs are well known for water. This indicates a certain mobility of hydrogens (and oxygens) in water. We analyse herein the hypothesis that water may

*Corresponding author. Email: apoma@theory.nipne.ro

consist of O^{-2z} anions of mass $M = 16$ amu and density n and H^{+z} cations (protons) of mass $m = 1$ amu and density $2n$, where z is a small reduced effective electron charge (the atomic mass unit, $1 \text{ amu} \simeq 1.7 \times 10^{-24} \text{ g}$). We shall see that such a hypothesis adds another dimension to the dynamics of water. Such a model may apply to other similar liquids.

Due to their large mass, the ions have a classical dynamics. Herein, we limit ourselves to considering the ions motion in water under the action of the Coulomb potentials $\varphi_{OO} = 4z^2 e^2/r$, $\varphi_{HH} = z^2 e^2/r$ and $\varphi_{OH} = -2z^2 e^2/r$, where $-e$ ($\simeq -4.8 \times 10^{-10}$ esu) is the electron charge and r denotes the distance between the ions. For stability, it is also necessary to introduce a short-range repulsive (hard-core) potential χ .¹ It is shown that in the limit $z \rightarrow 0$ water may exhibit an anomalous sound-like mode beside both the ordinary (hydrodynamic) one and the non-equilibrium sound-like excitations governed by short-range interactions. We compute the density oscillations for this model, the dielectric function, the structure factor, and extend the model to a multi-component plasma, including an asymmetric short-range interaction between ion species.

2. Plasmons in a jellium model

Let us consider one species of charged particles, with charge $-ze$, continuously distributed with density n in a neutralising rigid continuous background of positive charge. This is the well-known jellium model [7]. The Coulomb interaction reads

$$U = \frac{1}{2} \int d\mathbf{r} d\mathbf{r}' \varphi(\mathbf{r} - \mathbf{r}') \delta n(\mathbf{r}) \delta n(\mathbf{r}'), \quad (1)$$

where $\delta n(\mathbf{r})$ denotes a small disturbance of density (which preserves the global neutrality). We introduce the Fourier representation

$$\delta n(\mathbf{r}) = \frac{1}{\sqrt{N}} \sum_{\mathbf{q}} \delta n(\mathbf{q}) e^{i\mathbf{q}\mathbf{r}}, \quad \delta n(\mathbf{q}) = \frac{n}{\sqrt{N}} \int d\mathbf{r} \delta n(\mathbf{r}) e^{-i\mathbf{q}\mathbf{r}}, \quad (2)$$

where $N = nV$ is the total number of particles in volume V . Similarly,

$$\varphi(\mathbf{r}) = \frac{1}{V} \sum_{\mathbf{q}} \varphi(\mathbf{q}) e^{i\mathbf{q}\mathbf{r}}, \quad \varphi(\mathbf{q}) = \int d\mathbf{r} \varphi(\mathbf{r}) e^{-i\mathbf{q}\mathbf{r}}, \quad (3)$$

where $\varphi(\mathbf{q}) = 4\pi z^2 e^2/q^2$ is the Fourier transform of the Coulomb potential (interaction). The Coulomb interaction given by (1) becomes

$$U = \frac{1}{2n} \sum_{\mathbf{q}} \varphi(q) \delta n(\mathbf{q}) \delta n(-\mathbf{q}) \quad (4)$$

(where the $q=0$ -term is excluded by the positive background).

The small variations $\delta n(\mathbf{r})$ in density can be represented as $\delta n = -n \text{div } \mathbf{u}$, where \mathbf{u} is a displacement vector [8]. We emphasise that such a representation holds for $\mathbf{q}\mathbf{u}(\mathbf{r}) \ll 1$. It follows $\delta n(\mathbf{q}) = -in\mathbf{q}\mathbf{u}(\mathbf{q})$, and one can see that the Coulomb interaction involves only longitudinal components of the displacement vector $\mathbf{u}(\mathbf{q})$ along the wavevector \mathbf{q} . Therefore, we may write $\mathbf{u}(\mathbf{q}) = (\mathbf{q}/q)u(\mathbf{q})$, with $\delta n^*(-\mathbf{q}) = \delta n(\mathbf{q})$, $\mathbf{u}^*(-\mathbf{q}) = \mathbf{u}(\mathbf{q})$ and $u^*(-\mathbf{q}) = -u(\mathbf{q})$. The Coulomb interaction (4) becomes

$$U = -\frac{n}{2} \sum_{\mathbf{q}} q^2 \varphi(q) u(\mathbf{q}) u(-\mathbf{q}). \quad (5)$$

The kinetic energy associated with the coordinates $u(\mathbf{q})$ is given by

$$T = \frac{1}{2} \int d\mathbf{r} n m \dot{\mathbf{u}}^2 = -\frac{1}{2} m \sum_{\mathbf{q}} \dot{u}(\mathbf{q}) \dot{u}(-\mathbf{q}), \tag{6}$$

where m denotes the particle mass. The equations of motion obtained from the Lagrange function $L = T - U$ are

$$m \ddot{u}(\mathbf{q}) + n q^2 \varphi(q) u(\mathbf{q}) = 0, \tag{7}$$

which leads to the well known plasma oscillations with frequency given by $\omega_p^2 = 4\pi n z^2 e^2 / m$.

3. Plasma oscillations with two species of ions

We apply the above model to the two species of ions O^{-2z} and H^{+z} . The change in density is associated with a displacement vector \mathbf{v} in the former and a displacement vector \mathbf{u} in the latter. First, we note that the Fourier transforms of the Coulomb potentials are given by $\varphi_{OO} = 4\varphi(q)$, $\varphi_{HH} = \varphi(q)$ and $\varphi_{OH} = -2\varphi(q)$, where $\varphi(q) = 4\pi z^2 e^2 / q^2$. Therefore, the interactions can be written as

$$\begin{aligned} U_{OO} &= -\frac{n}{2} \sum_{\mathbf{q}} q^2 [4\varphi(q) + \chi(q)] v(\mathbf{q}) v(-\mathbf{q}), \\ U_{HH} &= -2n \sum_{\mathbf{q}} q^2 [\varphi(q) + \chi(q)] u(\mathbf{q}) u(-\mathbf{q}), \\ U_{OH} &= n \sum_{\mathbf{q}} q^2 [2\varphi(q) - \chi] u(\mathbf{q}) v(-\mathbf{q}), \end{aligned} \tag{8}$$

where $n = N/V$ is the density of water molecules and the Fourier transform, χ , of a hard-core potential has been introduced (the same for both species). The kinetic energy is given by

$$T = -\frac{1}{2} M \sum_{\mathbf{q}} \dot{v}(\mathbf{q}) \dot{v}(-\mathbf{q}) - m \sum_{\mathbf{q}} \dot{u}(\mathbf{q}) \dot{u}(-\mathbf{q}), \tag{9}$$

and the equations of motion read

$$\begin{aligned} m \ddot{u} + 2n q^2 (\varphi + \chi) u - n q^2 (2\varphi - \chi) v &= 0 \\ M \ddot{v} + n q^2 (4\varphi + \chi) v - 2n q^2 (2\varphi - \chi) u &= 0, \end{aligned} \tag{10}$$

where we have dropped out the argument \mathbf{q} .

The solutions of these equations can be obtained straightforwardly. In the long wavelength limit $\mathbf{q} \rightarrow 0$, there are two branches of eigenfrequencies, one given by

$$\omega_p^2 = \frac{16\pi n z^2 e^2}{\mu}, \tag{11}$$

corresponding to plasma oscillations and another given by

$$\omega_s^2 = \frac{9n\chi}{M + 2m} q^2 = v_s^2 q^2, \tag{12}$$

corresponding to sound-like waves propagating with velocity v_s given by (12). $\mu = 2mM/(2m + M)$ is the reduced mass. The plasma oscillations are associated with antiphase oscillations of the relative coordinate ($2mu + Mv = 0$), while the sound waves are associated with in-phase oscillations of the centre-of-mass coordinate ($u - v = 0$).

4. Polarisation

An external electric field arising from a potential $\phi(\mathbf{r})$ gives an additional energy:

$$U_i = q_i \int d\mathbf{r} \phi(\mathbf{r}) \delta n_i(\mathbf{r}) = -i(n_i q_i / n) \sum_{\mathbf{q}} q \phi(\mathbf{q}) u_i(-\mathbf{q}), \tag{13}$$

for a species of ions labelled i , with electric charge q_i and density n_i . We apply this formula to the two-species ionic plasma, and get

$$U_H = -2ize \sum_{\mathbf{q}} q \phi(\mathbf{q}) u(-\mathbf{q}), \quad U_O = 2ize \sum_{\mathbf{q}} q \phi(\mathbf{q}) v(-\mathbf{q}). \tag{14}$$

Adding these two terms to the lagrangian, the equations of motion given by (10) become

$$\begin{aligned} m\ddot{u} + 2nq^2(\varphi + \chi)u - nq^2(2\varphi - \chi)v &= -izeq\phi \\ M\ddot{v} + nq^2(4\varphi + \chi)v - 2nq^2(2\varphi - \chi)u &= 2izeq\phi, \end{aligned} \tag{15}$$

where we have dropped out the argument \mathbf{q} . This is a system of coupled harmonic oscillators under the action of an external force. In the limit of long wavelengths, its solutions are given by

$$\begin{aligned} u &= \frac{izeq}{m} \phi \frac{\omega^2 - (2m/3\mu)\omega_s^2}{(\omega^2 - \omega_p^2)(\omega^2 - \omega_s^2)}, \\ v &= -\frac{2izeq}{M} \phi \frac{\omega^2 - (2M/3\mu)\omega_s^2}{(\omega^2 - \omega_p^2)(\omega^2 - \omega_s^2)}. \end{aligned} \tag{16}$$

On the other hand, equation $n_i \text{div} \mathbf{u}_i = -\delta n_i$ is in fact the Maxwell equation $\text{div} \mathbf{E}_i = 4\pi q_i \delta n_i$, where the electric field is given by $\mathbf{E}_i = -4\pi n q_i \mathbf{u}_i$. We have therefore the internal electric fields $E_u = -8\pi n z e u$ and $E_v = 8\pi n z e v$. The polarisation $P = -(E_u + E_v)/4\pi$ is given by

$$P(\mathbf{q}) = 2nze[u(\mathbf{q}) - v(\mathbf{q})] = \frac{iq}{4\pi} \phi(\mathbf{q}) \frac{\omega_p^2}{\omega^2 - \omega_p^2}. \tag{17}$$

The external field is related to the external potential through $D(\mathbf{q}) = -iq\phi(\mathbf{q})$ and the dielectric function ϵ is given by $D = \epsilon E = \epsilon(D + E_{\text{int}})$, where $E_{\text{int}} = E_u + E_v$ is the internal field. We get the dielectric function:²

$$\epsilon = 1 - \omega_p^2/\omega^2, \tag{18}$$

as expected. As it is well known, its zero gives the longitudinal mode of plasma oscillations.

The ω_p in the nominator of Equation (18) also defines the plasma edge: for frequencies lower than ω_p , the electromagnetic waves are absorbed (the refractive index is given

by $n^2 = \epsilon$). It is well-known that water indeed exhibits a strong absorption in the gigahertz-terrahertz region [9–11]. On the other hand, neutron scattering on heavy water [12,13], as well as inelastic X-ray scattering [14], revealed the existence of a dispersionless mode $\simeq 4 - 5 \text{ meV}$ ($\simeq 10^{13} \text{ s}^{-1}$) in the structure factor, which may be taken tentatively as the ω_p -plasmonic mode given by Equation (11). Making use of this equation we get $\omega_p \simeq 3 \times 10^{14} \text{ zs}^{-1}$ ($n = 1/a^3$, $a = 3 \text{ \AA}$), so we may estimate the reduced effective charge $z \simeq 3 \times 10^{-2}$.

5. Dielectric function

The dielectric function given by Equation (18) has a singularity for $\omega = 0$, as arising from the exact cancellation in the static limit of the external field by the internal field. It is plausible to assume that residual polarisation fields are still present in this static limit, such as, for instance, the intrinsic polarisability. In this case, Equation (18) is modified, and the dielectric function is of the type

$$\epsilon = \frac{\omega^2 - \omega_p^2}{\omega^2 + \omega_0^2}, \tag{19}$$

where ω_0 is a plasma frequency associated with the intrinsic, molecular polarisability.³ As such, it is a very high frequency, and Equation (19) gives a small, negative contribution to the dielectric function in the static limit ($\omega \rightarrow 0$).

The dielectric properties of water are still a matter of debate. It is agreed that the permittivity dispersion of water is described to some extent by a Debye model of the form $\epsilon = a + b/(1 - i\omega\tau)$, where a and b are semi-empirical parameters and $\tau \sim \eta a^3/T$ is a relaxation time; η denotes the viscosity and T is the temperature [15,16]. This Debye model assumes mainly an orientational polarisability of electric dipoles, which, due to the preservation of the directional character of the O–H bonds, is compatible with the plasma model suggested here for water. Therefore, the contribution given by Equation (19) should be added to the above Debye formula for the dielectric function, which becomes

$$\epsilon = a + \frac{b}{1 - i\omega\tau} + \frac{\omega^2 - \omega_p^2}{\omega^2 + \omega_0^2}. \tag{20}$$

Parameters a and b in Equation (20) are related to the static permittivity ϵ_0 and high-frequency permittivity ϵ_∞ through

$$\epsilon_0 = a + b - \omega_p^2/\omega_0^2, \quad \epsilon_\infty = a + 1. \tag{21}$$

We may neglect ω_p^2/ω_0^2 here because it is too small, and we may also take $\epsilon_\infty = 1 (a = 0)$. The static permittivity $\epsilon_0 = b$ is given mainly by the electric dipoles. Let \mathbf{p} be such an electric dipole. Its energy in an electric field \mathbf{D} is $-pD \cos \theta$, where θ is the angle between \mathbf{p} and \mathbf{D} . The thermal distribution of such dipoles is $dw \sim \exp(-pD \cos \theta/T) d(\cos \theta)$, where T denotes the temperature. We easily get the thermal average $\langle \cos \theta \rangle = -L(pD/T)$, where $L(x) = \coth(x - 1)/x$ is the well-known Langevin's function.

We take $p = 2ez_e(a/2) = ez_e a$, where $a \sim 3 \text{ \AA}$ and z_e is a delocalised reduced charge associated with the H–O dipole. We estimate the argument pD/T of the Langevin's function. At room temperature, we find $pD/T \simeq 3 \times 10^{-4} Dz_e$. For $pD/T = 1$ this corresponds to an external field $D = (1/3z_e) \times 10^4 \text{ esu}$, or $D = 10^8/z_e \text{ V m}^{-1}$

(1 esu = $3 \times 10^4 \text{ V m}^{-1}$ [17]). This is an extremely high field, so we are justified to take $pD/T \ll 1$, and $L(pD/T) \simeq pD/3T$. We get therefore a polarisation $P = -np(\cos \theta) = np^2D/3T$, an internal field $E_{\text{int}} = -4\pi P = -4\pi np^2D/3T$, and a permittivity

$$\varepsilon_0 = b = \frac{1}{1 - 4\pi np^2/3T} \quad (22)$$

from $D = \varepsilon E = \varepsilon(D + E_{\text{int}})$. This is the well-known Kirkwood formula [15]. For the empirical value $\varepsilon_0 = 80$, we get (at room temperature) a reduced charge $z_e \simeq 10^{-2}$. This is in good agreement with the $\text{H}^{+z} - \text{O}^{-2z}$ plasma charge z estimated above.

6. Cohesion and thermodynamics

Recently, a model of liquid has been introduced [18] based on an excitation spectrum (per particle) of the form $\varepsilon_n = -\varepsilon_0 + \varepsilon_1(n+1)2$, where ε_0 is a cohesion energy and ε_1 is the quanta of energy of a harmonic oscillator with one degree of freedom; n here represents the quantum number. The model also includes the kinematic correlations (spatial restrictions) of the movement of the liquid molecules. This model leads to consistent thermodynamics for liquids, arising from statistics which are equivalent to the statistics of bosons in two dimensions.

For water, the cohesion energy per particle ε_0 can be estimated from the vaporisation heat ($\simeq 40 \text{ kJ mol}^{-1}$). It gives $\varepsilon_0 \sim 10^3 \text{ K}$. On the other hand, it was shown in two previous papers [19,20] that the transition temperature between a gas and a liquid of identical particles is approximately given by

$$T_t = \frac{4}{3} \frac{\varepsilon_0}{\ln(\varepsilon_0/T_0)}, \quad (23)$$

where $T_0 = \hbar^2 n^{2/3}/m$ is a gas characteristic temperature. We can apply this formula to water dissociation, taking n as the density of hydrogen atoms, m as the mass of two hydrogen atoms and $T_t = 383 \text{ K}$ (at normal pressure; ε_0 depends on the inter-particle spacing). We may neglect the oxygen, as it is too heavy in comparison with the hydrogen atoms. We get $T_0 \simeq 2 \text{ K}$ and the above formula gives $\varepsilon_0 \simeq 2000 \text{ K} \simeq 200 \text{ meV}$ for the cohesion energy of water per molecule, which is consistent with the above estimate ($1 \text{ eV} \simeq 11.6 \times 10^3 \text{ K}$; $n \simeq 1/a^3$ with $a = 3 \text{ \AA}$ and $\hbar \simeq 10^{-27} \text{ erg s}^{-1}$; Bohr radius $a_H = \hbar^2/m_e e^2 \simeq 0.53 \text{ \AA}$, $e^2/a_H \simeq 27.22 \text{ eV}$, where m_e is the electron mass).⁴

The plasma oscillations obtained above can be quantised and the energy levels of the plasma read:

$$E_n = \sum_{\mathbf{q}} \hbar \omega_p(n+1/2) = \frac{V}{(2\pi)^3} \frac{4\pi}{3} q_c^3 \cdot \hbar \omega_p(n+1/2), \quad (24)$$

where q_c is a cutoff wavevector. The prefactor in Equation (24) is $Vq_c^3/6\pi^2 \simeq N(aq_c/4)^3$, so the energy levels given above can be written as

$$E_n = N\varepsilon_1(n+1/2), \quad (25)$$

where $\varepsilon_1 = (aq_c/4)\hbar\omega_p$. These energy levels correspond to a harmonic oscillator with one degree of freedom. It follows that the present description of water as a two-species of highly dissociated ionic plasma provides a further support for the liquid model

mentioned above. If we take $q_c \simeq 1/a$ the energy quanta $\varepsilon_1 = (aq_c/4)^3 \hbar \omega_p = \simeq 3z \text{ meV}$ represents the ε_l parameter in the spectrum of the liquid. (The plasma frequency given by Equation (11) is $\omega_p \simeq 200z \text{ meV}$).

7. Debye screening and the correlation energy

As is well known, the plasma excitations described above represent collective oscillations of the density in the long wavelength limit. At the same time, they induce correlations in the ionic movements. For a classical plasma these correlations are associated with a screening length given by the Debye–Huckel theory as [21]

$$\kappa^{-1} = (T/24\pi n z^2 e^2)^{1/2}, \tag{26}$$

for our case ($\kappa^{-1} = (T/4\pi e^2 \sum_i n_i z_i^2)^{-1}$ where i labels the ionic species with density n_i and charge ez_i). The formula is valid for the Coulomb energy $z^2 e^2/a$ much lower than the temperature T . In the present case, we have $z^2 e^2/a \simeq 45 \text{ K}$ (for $z \simeq 3 \times 10^{-2}$), which shows that the above condition is fulfilled. From (26) we get $\kappa^{-1} \sim 1 \text{ \AA}$ (at room temperature), in agreement with the present molecular-dissociation model. The correlation energy per particle is given by

$$\varepsilon_{\text{corr}} = -\frac{e^2}{a} \sqrt{\frac{\pi e^2}{Ta}} (6z^2)^{3/2} \tag{27}$$

$\{\varepsilon_{\text{corr}} = -(e^2/a) \sqrt{\pi e^2/Ta} (\sum_i n_i z_i^2)^{3/2}\}$. The estimation of this energy gives $\varepsilon_{\text{corr}} \sim 10^2 \text{ K}$ (at room temperature). It contributes to the cohesion energy.

8. Sound anomaly

The sound-like branch $\omega_2 \simeq \omega_s = v_s q$, where $v_s = \sqrt{9n\chi/(M+2m)}$ according to Equation (12), is distinct from the ordinary hydrodynamic sound whose velocity is given by the well-known formula $v_0 = 1/\sqrt{(\kappa n m)}$ for a one-component fluid, where κ is the adiabatic compressibility. For the present two-component fluid ($\text{H}^{+z} - \text{O}^{-2z}$ plasma), the velocity of the ordinary sound is given by $v_0 = 1/\sqrt{\kappa n (M+2m)}$. The former represents a non-equilibrium elementary excitation, whose velocity v_s does not depend on temperature, while the latter proceeds by thermodynamic, equilibrium, adiabatic processes, and its velocity v_0 depends on temperature through the adiabatic compressibility κ . In order to distinguish them from the hydrodynamic sound, we propose to call the sound-like excitations derived here density ‘kinetic’ modes or ‘densitons’. The distinction between the two sounds is made by a threshold wavevector q_t in the following manner. Suppose that there is a finite lifetime, τ , for the sound-like excitations ω_s , propagating with a velocity v_s and a corresponding meanfree path $\Lambda = v_s \tau$. If the sound-like wavelength λ is much longer than the meanfree path, $\lambda \gg \Lambda$, then we are in the collision-like regime ($\omega_s \tau \ll 1$), and the collisions may restore the thermodynamic equilibrium. In this case the hydrodynamic sound propagates, and the sound-like excitations do not. This condition defines the threshold wavevector $q_t = 1/v_s \tau$. In the opposite case, $q \gg q_t$ (collision-less regime), it is the sound-like excitations that propagate, and not the hydrodynamic sound. The finite lifetime τ originates in the residual interactions between the collective modes and the underlying motion of the

individual particles. It is easy to estimate this residual interaction [8]. It is given by $\sqrt{\varepsilon T}$, where ε is the mean energy per particle corresponding to the motion of the individual particles. We get, therefore, $\tau \simeq \hbar/\sqrt{\varepsilon T}$ and the threshold wavevector $q_t = \sqrt{\varepsilon T}/\hbar v_s$. It is difficult to have a reliable estimation of the mean energy ε ; for a reasonable value $\varepsilon = 10 \text{ meV}$ we get $q_t \simeq 0.1 \text{ \AA}^{-1}$ at room temperature for $v = 3000 \text{ m s}^{-1}$, which is in good agreement with experimental data.

Indeed, the phenomenon of two-sound anomaly in water is well documented [22,23]. Neutron, X-ray, Brillouin or ultraviolet light scattering on water revealed the existence of a hydrodynamic sound propagating with velocity $v_0 \simeq 1500 \text{ m s}^{-1}$ for smaller wavevectors and an additional sound propagating with velocity $v_s \simeq 3000 \text{ m s}^{-1}$ for larger wavevectors. In addition, though both sound velocities do exhibit an isotopic effect, their ratio does not. According to the above discussion, we assign this additional, faster sound to the sound-like excitations derived here. We can see that both v_0 and v_s given above exhibit a weak isotopic effect, while their ratio $v_s/v_0 = 3n\sqrt{\kappa\chi}$ does not. From $v_s = \sqrt{9n\chi/(M+2m)} = 3000 \text{ m s}^{-1}$ we get the short-range interaction $\chi \simeq 7 \text{ eV} \cdot \text{\AA}^3$. Similar results are obtained for other forms of dissociation of the water molecule, like $\text{OH}^- - \text{H}^+$ or $\text{OH}^- - \text{H}_3\text{O}^+$, so the $\text{H}^{+z} - \text{O}^{-2z}$ plasma model employed here can be viewed as an average, effective model for various plasma components that may exist in water.

9. Another possible anomalous sound

It is worth calculating the spectrum given by equations of motion (10) without neglecting higher-order contributions in q^2 . The result of this calculation is given by

$$\omega_{1,2}^2 = \frac{1}{2}\omega_p^2 \left[1 + Ax^2 \pm \sqrt{1 + 2Bx^2 + A^2x^4} \right], \quad (28)$$

where

$$A = \frac{1}{9\alpha}(2 + 5\alpha + 2\alpha^2), \quad B = \frac{1}{9\alpha}(2 - 13\alpha + 2\alpha^2), \quad \alpha = m/M \quad (29)$$

and $x = v_s q / \omega_p$. It is shown in Figure 1.

Frequency ω_2 in Equation (28) represents the sound-like branch, which goes like $\omega_2 \simeq \omega_s = v_s q$ in the long wavelength limit and approaches the horizontal asymptote $\omega_2 = \omega_p / \sqrt{A} \simeq \omega_p \sqrt{m/2M}$ for shorter wavelengths. Frequency ω_1 in Equation (28) represents the plasmonic branch ($\omega_1 \simeq \omega_p$ for $q \rightarrow 0$). In the long wavelength limit it is

$$\omega_1 \simeq \omega_p + \frac{(M-m)^2}{9mM} v_s^2 q^2 / \omega_p, \quad q \rightarrow 0. \quad (30)$$

Due to the large disparity between the two masses m and M , we can see that the plasma frequency has an abrupt increase towards the short-wavelength oblique asymptote given by

$$\omega_a \simeq \sqrt{A} v_s q \simeq \sqrt{2M/9m + 5/9} v_s q. \quad (31)$$




Figure 1. The spectrum of the density oscillations given by Equation (28) for the $H^{+z}-O^{-2z}$ plasma with the same short-range interaction between ionic species.

For small values of ω_p (vanishing Coulomb coupling, $z \rightarrow 0$), this asymptotic frequency may look like an anomalous sound propagating with velocity

$$v_a \simeq \sqrt{2M/9m + 5/9} v_s. \tag{32}$$

For water, we get $v_a \simeq 2v_s$ from this formula. However, the ratios v_a/v_s or v_a/v_0 exhibit a rather strong isotopic effect, which is not supported by experimental data.

10. Multi-component plasma

The model presented herein might be generalised to a multi-component plasma consisting of several ionic species labelled i , each with number N_i of particles, density n_i , charge $z_i e$ and mass m_i , such that $\sum_i z_i n_i = 0$.

The Lagrangian of the density oscillations is given by

$$L = -\frac{1}{2n} \sum_{i\mathbf{q}} m_i n_i \dot{u}_i(\mathbf{q}) \dot{u}_i(-\mathbf{q}) + \frac{1}{2n} \sum_{ij\mathbf{q}} n_i n_j q^2 [\varphi_{ij}(q) + \chi(q)] u_i(\mathbf{q}) u_j(-\mathbf{q}) + i \frac{e}{n} \sum_{i\mathbf{q}} n_i z_i q \phi(\mathbf{q}) u_i(-\mathbf{q}), \tag{33}$$

where $\varphi_{ij}(q) = 4\pi z_i z_j e^2 / q^2$. The equations of motion are given by

$$m_i \ddot{u}_i + 4\pi e^2 z_i \sum_j z_j n_j u_j + q^2 \chi \sum_j n_j u_j = -iqe z_i \phi. \tag{34}$$

Downloaded At: 07:32 28 January 2011

Making use of the notations

$$S_1 = \sum_i z_i^2 n_i / m_i, \quad S_2 = \sum_i n_i / m_i, \quad S_3 = \sum_i z_i n_i / m_i, \tag{35}$$

the eigenfrequencies $\omega_{1,2}$ of the system of Equations (34) in the long wavelength limit are given by

$$\omega_1^2 \simeq \omega_p^2 = 4\pi e^2 S_1 = \sum_i \frac{4\pi e^2 z_i^2 n_i}{m_i}, \tag{36}$$

which represents the plasma branch of the spectrum, and

$$\omega_2^2 \simeq \omega_s^2 = (S_2 - S_3^2 / S_1) \chi q^2 = v_s^2 q^2, \tag{37}$$

which represents the sound-like excitations.⁵ The plasma branch of the spectrum has an oblique asymptote given by $\omega_1 \simeq \omega_a = \sqrt{\chi S_2} q$, which may be taken as an anomalous sound propagating with velocity $v_a = \sqrt{\chi S_2}$ for small values of ω_p . The ratio of the two sound velocities is given by

$$v_a / v_s = \frac{1}{\sqrt{1 - S_3^2 / S_1 S_2}}, \tag{38}$$

which is always higher than unity. The sound branch of the spectrum has a horizontal asymptote given by $\omega_2 \simeq \sqrt{1 - S_3^2 / S_1 S_2} \omega_p$. For the $H^{+z} - O^{-2z}$ plasma, we can check from (38) that $v_a / v_s \simeq (2M/9m + 5/9)^{1/2} \simeq 2$, and $\omega_2 \simeq 3\sqrt{m/2M} \omega_p$, as obtained above. As we have discussed above, this ratio exhibits a rather strong isotopic effect, which is not in accordance with experimental data. We assign, therefore, the additional sound to sound-like excitations propagating with velocity v_s given by Equation (37). The ordinary, hydrodynamic sound in a multi-component mixture has the velocity $v_0 = 1/\sqrt{\kappa \sum_i n_i m_i}$. It can be shown that $v_s^2 / v_0^2 \geq n^2 \kappa \chi$ for a neutral multi-component mixture.

The internal field is given by

$$E_{int} = -4\pi e \sum_i z_i n_i u_i; \tag{39}$$

we get easily from Equation (34)

$$E_{int} = -iq\phi \frac{\omega_p^2}{\omega^2 - \omega_p^2} \tag{40}$$

and the dielectric function $\epsilon = 1 - \omega_p^2 / \omega^2$, as expected.

11. Structure factor

The structure factor is defined by

$$\begin{aligned} S(q, \omega) &= \frac{1}{2\pi} \int d\mathbf{r} d\mathbf{r}' dt \langle \delta n(\mathbf{r}, t) \delta n(\mathbf{r}', 0) \rangle e^{i\mathbf{q}(\mathbf{r}-\mathbf{r}') - i\omega t} \\ &= \frac{N}{2\pi n^2} \int dt \langle \delta n(\mathbf{q}, t) \delta n(-\mathbf{q}, 0) \rangle e^{-i\omega t}, \end{aligned} \tag{41}$$

where the brackets stand for the thermal average (we leave aside the central peak). Since

$$\delta n(\mathbf{q}, t) = -iq \sum_i n_i u_i(\mathbf{q}, t), \tag{42}$$

it becomes

$$S(q, \omega) = \frac{Nq^2}{2\pi n^2} \int dt \sum_{ij} n_i n_j \langle u_i(t) u_j(0) \rangle e^{-i\omega t}, \tag{43}$$

where we dropped out the argument \mathbf{q} .

In order to calculate the thermal averages, we turn back to the system of Equations (34) without the external electric field. This system can be written as

$$\begin{aligned} (-\omega^2 + aS_1)x + bS_3y &= 0, \\ aS_3x + (-\omega^2 + bS_2)y &= 0, \end{aligned} \tag{44}$$

where $a = 4\pi e^2$, $b = \chi q^2$, $S_{1,2,3}$ are given by Equation (35) and

$$x = \frac{1}{n} \sum_i z_i n_i u_i, \quad y = \frac{1}{n} \sum_i n_i u_i. \tag{45}$$

In addition,

$$u_i = \frac{anz_i}{m_i \omega^2} x + \frac{bn}{m_i \omega^2} y. \tag{46}$$

The system of Equations (44) has two eigenfrequencies $\omega_{1,2}$, as given by Equations (36) and (37). The corresponding eigenvectors are given by

$$x_1 \sim S_1, \quad y_1 \sim S_3; \quad x_2 \sim bS_3, \quad y_2 \sim -aS_1 \tag{47}$$

in the long wavelength limit. According to Equation (46) the coordinates u_i can be written as

$$u_i^{(1,2)} = \frac{anz_i}{m_i \omega_{1,2}^2} x_{1,2} e^{i\omega_{1,2}t} + \frac{bn}{m_i \omega_{1,2}^2} y_{1,2} e^{i\omega_{1,2}t}, \tag{48}$$

and one can see that they are coordinates of linear harmonic oscillators with frequencies $\omega_{1,2}$ and potential energies

$$\frac{m_i \omega_{1,2}^2 [u_i^{(1,2)}]^2}{2}.$$

The thermal distribution of the coordinate u for such an oscillator is given by $dw = \sqrt{m\omega^2/2\pi T} \exp(-m\omega^2 u^2/2T) du$ in the classical limit, where T denotes the temperature ($T \gg \hbar \omega$). It follows

$$\langle u_i^{(1,2)} u_j^{(1,2)} \rangle = \frac{T}{m_i \omega_{1,2}^2} \delta_{ij}. \tag{49}$$

Writing

$$u_i = u_i^{(1)} e^{i\omega_1 t} + u_i^{(2)} e^{i\omega_2 t} \tag{50}$$

and making use of Equation (49), the structure factor given by Equation (43) becomes

$$S(q, \omega) = NTq^2 \left(\sum_i n_i^2 / n^2 m_i \right) \left[\frac{1}{\omega_1^2} \delta(\omega - \omega_1) + \frac{1}{\omega_2^2} \delta(\omega - \omega_2) \right]. \tag{51}$$

We can see from this equation that the relevant sound contributions are given by

$$S(q, \omega) \simeq \frac{NT}{v_{s,a}^2} \left(\sum_i n_i^2 / n^2 m_i \right) \delta(\omega - v_{s,a}q). \tag{52}$$

The relaxation and damping effects can be included in the above expressions of the structure factor. As it is well-known, they amount to representing the δ -functions by lorentzians.

12. Asymmetric short-range interaction

Up to now, the short-range interaction was assumed to be the same for all ionic species. In general, we may introduce a short-range interaction, χ_{ij} , depending on the nature of the ionic species. If this interaction is separable, the solution given above for a multi-component plasma holds with minor modifications. For a non-separable short-range interaction, appreciable changes may appear in the spectrum, which may exhibit multiple branches. Such a spectrum may serve to identify the nature (mass and charge) of various molecular aggregates in a multi-component plasma. It is worth noting that a range of frequencies $10^{10} - 10^{12} \text{ s}^{-1}$ is documented in living cells by microwave, Raman and optical spectroscopies and by cell-biology studies, upon which the theory of coherence domains in living matter is built [24–30].

We consider here again the $\text{H}^{+z}\text{-O}^{-2z}$ plasma with different short-range interaction $\chi_{\text{HH}} = \chi_1$, $\chi_{\text{OO}} = \chi_2$, $\chi_{\text{OH}} = \chi_3$; it still exhibits two branches of frequencies, a plasmonic one (ω_1) and a sound-like one (ω_2) but the spectrum may have certain peculiarities (the dielectric constant is not affected by this modification). Equations of motion (15) become now

$$\begin{aligned} m\ddot{u} + 2nq^2(\varphi + \chi_1)u - nq^2(2\varphi - \chi_3)v &= -izeq\phi \\ M\ddot{v} + nq^2(4\varphi + \chi_2)v - 2nq^2(2\varphi - \chi_3)u &= 2izeq\phi. \end{aligned} \tag{53}$$

We introduce the notations

$$a = 2nq^2\varphi/m = 8\pi ne^2 z^2/m, \quad b_{1,2,3} = n\chi_{1,2,3}/m. \tag{54}$$

The dispersion relations can be computed straightforwardly. In the long wavelength limit ($q \rightarrow 0$), we get the plasmonic branch

$$\omega_1^2 \simeq (1 + 2\alpha)a + \frac{2b_1 + \alpha^2 b_2 - 4\alpha b_3}{1 + 2\alpha} q^2, \tag{55}$$

where $(1 + 2\alpha)a = 16\pi ne^2 z^2/\mu$ is the plasma frequency, and the sound-like branch

$$\omega_2^2 \simeq \frac{\alpha(4b_1 + b_2 + 4b_3)}{1 + 2\alpha} q^2 = v_s^2 q^2; \tag{56}$$

one can see that the sound velocity v_s is always a real quantity.

The sound-like branch exhibits an asymptote in the short-wavelength limit, given by

$$\omega_2^2 \sim \frac{1}{2} \left[2b_1 + \alpha b_2 - \sqrt{(2b_1 - \alpha b_2)^2 + 8\alpha b_3^2} \right] q^2, \tag{57}$$

whose slope may have either sign or vanish. It is easy to see that this slope is positive for $b_3^2 < b_1 b_2$, negative for $b_3^2 > b_1 b_2$ (when the sound-like branch has a maximum value) and it vanishes for $b_3^2 = b_1 b_2$ (when the sound-like branch has an horizontal asymptote). In the case of a negative slope, the sound velocity may exhibit a negative velocity and the sound may suffer a strong absorption for moderate values of the wavevector, which may indicate an anomalous or unphysical situation.

We return now to the plasmon branch given by Equation (55), and write it as

$$\omega_1^2 = \omega_p^2 + b_2 \frac{2x^2 - 4\alpha\lambda x + \alpha^2}{1 + 2\alpha} q^2, \tag{58}$$

where $\lambda^2 = b_3^2/b_1 b_2$ and $x = \sqrt{b_1/b_2}$. It is easy to see that for $\lambda^2 > 1$ the plasmonic spectrum exhibits a dip around a certain value q_0 of the wavevector q for $(\lambda - \sqrt{\lambda^2 - 1/2})\alpha < \sqrt{b_1/b_2} < (\lambda + \sqrt{\lambda^2 - 1/2})\alpha$; it approaches an asymptote with a positive slope for $q \rightarrow \infty$, which may define again an anomalous sound for small values of ω_p .

We illustrate these anomalies for a particular case of short-range interaction $\chi_{1,2} = 0$ and $\chi_3 = \chi$ ($b_3 = n\chi/m$). The dispersion relations of the system of Equations (53) become

$$\omega_{1,2}^2 = \frac{1}{2} \omega_p^2 \left[1 \pm \sqrt{1 - 4v_s^2 q^2 / \omega_p^2 + \frac{(1 + 2\alpha)^2}{2\alpha} v_s^4 q^4 / \omega_p^4} \right]. \tag{59}$$

The plasmonic branch has a minimum value for $q_0 \simeq 2\sqrt{m/M}\omega_p/v_s$, where the sound-like branch has a maximum value ($\simeq \sqrt{2m/M}\omega_p$). The spectrum is shown in Figure 2. Using $\omega_p \simeq 10^{13} \text{s}^{-1}$ estimated above and the sound velocity $v_s \simeq 3000 \text{ m s}^{-1}$ in water, we get $q_0^{-1} \simeq 6 \text{ \AA}$. We may expand ω_1 in series of $(q - q_0)^2$ around its minimum value at q_0 and get $\omega_1 \simeq \omega_p + (M/4m + 1)(v_s^4 q_0^2 / \omega_p^3)(q - q_0)^2 = \omega_p + (1 + 4m/M)v_s^2(q - q_0)^2 / \omega_p$. This is similar to the rotons-like dispersion relation discussed in connection with the coherence domains in water [31]. Although this might be an interesting suggestion, it is inconsequential here, because ω_p is too small in comparison with the temperatures at which water exists and, therefore, this ‘dip’ feature has no effect for water thermodynamics.

13. Conclusion

We summarise the main features of the model suggested here for liquid water. First, we assume, as it is generally accepted, the four directional sp^3 -oxygen electronic orbitals. The electron delocalisation along two such orbitals together with a corresponding delocalisation of the hydrogen electronic charge lead to the water cohesion. It is represented by the cohesion energy ϵ_0 discussed here. Within such a picture, we can still visualise the oxygen and the hydrogen as neutral atoms, moving around almost freely (as a consequence of the uniformity of the environment; this gives a noteworthy support to the ‘hydrogen bonds’ concept).⁶ To this picture, the present model adds another component, arising from a very small charge transfer between hydrogen and oxygen




Figure 2. Excitation spectrum given by Equation (59) for the $H^{+z} - O^{-2z}$ plasma with short-range potentials $\chi_{OO} = \chi_{HH} = 0$ and $\chi_{OH} = \chi \neq 0$.

atoms, leading to a $H^{+z} - O^{-2z}$ plasma, with the reduced charge z . It may originate in the weak asymmetry of the two occupied sp^3 -oxygen electronic orbitals with respect to the other two unoccupied orbitals. Under these circumstances, the hydrogen and oxygen ions interact, both by Coulomb and by short-range potentials. This interaction gives the plasma frequency and the sound-like excitations frequency. The plasmons contribute to the excitations which give rise to consistent thermodynamics for liquids, in a model introduced recently. In addition, the ionic plasma oscillations entail oscillations of the delocalised electronic cloud, with the same eigenfrequency. Subjected to an external field, these electronic oscillations produce an intrinsic polarisability which removes the $\omega=0$ singularity in the plasma dielectric function (the ω_0 frequency). In addition, the magnitude of the electric moment \mathbf{p} which is responsible for the orientational, static dielectric function is in satisfactory agreement with the plasma charge z derived herein.

On the basis of this model we are able to understand, to some extent, both qualitatively and in some places even quantitatively, the sound anomaly, the dielectric function (permittivity dispersion), the structure factor, cohesion and thermodynamics of water. The model is extended to a multi-component classical plasma, including an asymmetric short-range interaction between the components, which might be relevant for more complex structural aggregates like those in biological matter.

Notes

1. See also in this respect [3–6]. As is well-known, a classical plasma with Coulomb interaction only is unstable.
2. We disregard here the intrinsic and orientational polarisabilities.
3. A static field D produces an electric dipole $p = q_e x$, where q_e is the electric charge and x is a small displacement subjected to the equation of motion $m_e \ddot{x} + m_e \omega_p^2 x = q_e D$, where m_e is the mass of

the electronic cloud. According to the plasma model suggested here, we assume that the electronic cloud in the H–O bonds have the same eigenfrequency ω_p as the H–O ensemble. In the static limit $x = q_e D / m_e \omega_p^2$ (polarisability $\alpha = q_e^2 / m_e \omega_p^2$ in $p = \alpha D$), and we get a polarisation $P = p/a_0^3 = q_e^2 D / m_e a_0^3 \omega_p^2$, where a_0 is of the order of the atomic size. We get an internal field $E_{int} = -4\pi P = -(4\pi q_e^2 / m_e a_0^3) D / \omega_p^2 = -(\omega_0^2 / \omega_p^2) D$, where ω_0 is a frequency of the order of atomic frequencies. Consequently, the dielectric function ε in equation $D = \varepsilon E = \varepsilon(D + E_{int})$ is given by $\varepsilon \simeq -\omega_p^2 / \omega_0^2$ ($\omega_p^2 / \omega_0^2 \ll 1$), which is precisely the static dielectric function given by Equation (19).

4. It is worth noting that the mechanism of vaporisation assumed here implies the dissociation of the water molecule.
5. The sound velocity given by (37) is always a real quantity, as a consequence of the Schwarz–Cauchy inequality.
6. The point of view taken in this article is that the hydrogen bonds in water are introduced in order to account for the uniformity of the environment of a water molecule in liquid water. As such, it helps understand the cohesion. However, a consistent upholding of the hydrogen-bonds concept would mean a vanishing dipole momentum of liquid water. Pauling himself [1], who originally introduced this concept, qualifies it by admitting an asymmetry in the four hydrogen bonds around an oxygen ion, arising from the two-out-of-four occupied orbitals. We suggest that the uniformity of the environment makes the hydrogen atoms (ions) moving as independent entities, while the asymmetry induces a small charge z , so the ion motion is subjected to Coulomb (and short-range interactions). The electric moment is ascribed to the directional character of the sp^3 -oxygen electronic orbitals and the charge transfer between oxygen and hydrogen. Thereby, the hydrogen-bond concept is employed here through its two features, directionality and uniformity, with a slight asymmetry, all viewed as independent qualitative ingredients.

References

- [1] L. Pauling, *General Chemistry* (Dover, New York, 1982).
- [2] F. Francks editor, *Water: A Comprehensive Treatise* (Plenum Press, New York, 1972).
- [3] E. Teller, *Revs. Mod. Phys.* **34**, 627 (1962).
- [4] E.H. Lieb and B. Simon, *Phys. Rev. Lett.* **31**, 681 (1973).
- [5] E.H. Lieb and B. Simon, *Adv. Math.* **23**, 22 (1977).
- [6] L. Spruch, *Revs. Mod. Phys.* **63**, 151 (1991).
- [7] D. Pines, *Elementary Excitations in Solids* (Benjamin, New York, 1963).
- [8] M. Apostol, *Electron Liquid* (Apoma, Magurele, Bucharest, 2000).
- [9] K.H. Tsai and T.-M. Wu, *Chem. Phys. Lett.* **417**, 390 (2005).
- [10] A. Padro and J. Marti, *J. Chem. Phys.* **118**, 452 (2003).
- [11] K.N. Woods and H. Wiedemann, *Chem. Phys. Lett.* **393**, 159 (2004).
- [12] F.J. Bermejo, M. Alvarez, S.M. Bennington, and R. Vallauri, *Phys. Rev.* **E51**, 2250 (1995).
- [13] C. Petrillo, F. Sacchetti, B. Dorner, and J.-B. Suck, *Phys. Rev.* **E62**, 3611 (2000).
- [14] F. Sette, G. Ruocco, M. Krisch, C. Masciovecchio, R. Verbeni, and U. Bergmann, *Phys. Rev. Lett.* **77**, 83 (1996).
- [15] H. Frohlich, *Theory of Dielectrics* (Oxford University Press, Oxford, 1958).
- [16] P. Debye, *Polar Molecules* (Dover, New York, 1945).
- [17] J.D. Jackson, *Classical Electrodynamics* (Wiley, New Jersey, 1999).
- [18] M. Apostol, *J. Theor. Phys.* **125**, 163 (2006).
- [19] M. Apostol, *Mod. Phys. Lett.* **B21**, 893 (2007).
- [20] M. Apostol, *J. Theor. Phys.* **123**, 155 (2006).
- [21] L. Landau and E. Lifshitz, *Course of Theoretical Physics, Vol. 5. Statistical Physics* (Elsevier, Oxford, 1980).
- [22] J. Teixeira, M.C. Bellissent-Funel, S.H. Chen, and B.B. Dorner, *Phys. Rev. Lett.* **54**, 2681 (1985).

- [23] S.C. Santucci, D. Fioretto, L. Comez, A. Gessini, and C. Maschiovecchio, *Phys. Rev. Lett.* **97**, 225701 (2006).
- [24] H. Frohlich, *Phys. Lett.* **A26**, 402 (1968).
- [25] H. Frohlich, *Int. J. Quant. Chem.* **2**, 641 (1968).
- [26] S.J. Webb, M.E. Stoneham, and H. Frohlich, *Phys. Lett* **A63**, 407 (1977).
- [27] S. Webb, *Phys. Repts.* **60**, 201 (1980).
- [28] S. Rowlands, L.S. Sewchand, R.E. Lovlin, J.S. Beck, and E.G. Enns, *Phys. Lett.* **A82**, 436 (1981).
- [29] S.C. Roy, *Phys. Lett.* **A83**, 142 (1981).
- [30] E. Del Giudice, S. Doglia, M. Milani, and G. Vitiello, *Nucl. Phys.* **B275**, 185 (1986).
- [31] G. Preparata, *QED Coherence in Matter* (World Science, Singapore, 1995).