

Phytochemistry, 1972, Vol. 11, p. 853. Pergamon Press. Printed in England.

## LAURACEAE

### TERPENES FROM *LINDERA ERYTHROCARPA*

H. KOMAE and N. HAYASHI

Department of Chemistry, Faculty of General Education, Hiroshima University, Hiroshima, Japan

(Received 3 August 1971)

*Plant.* *Lindera erythrocampa* Makino. *Source.* Ehime Prefecture, Japan. *Uses.* Not known. *Previous work.* None.

*Leaves.* The essential oil (3.8 g,  $d_4^{25}$  0.8606,  $n_D^{25}$  1.4793, 0.06% yield) was isolated from the fresh leaves (6.3 kg) by steam distillation. Caryophyllene (19.7%) and geranyl acetate (31.7%) were isolated by preparative GLC (Carbowax 20 M–20%, at 160°) and identified by IR and NMR. The presence of  $\alpha$ -pinene (6.8%), camphene (4.4%),  $\beta$ -pinene (2.1%), limonene (6.4%) and bornyl acetate (5.9%) were confirmed by GLC (PEG-6000–3%, 50-HB-2000–3% at 60–160°). Unknown constituents (23%).

*Key Word Index*—*Lindera erythrocampa*; Lauraceae; terpenes; geranyl acetate; caryophyllene.

---

Phytochemistry, 1972, Vol. 11, p. 853. Pergamon Press. Printed in England.

### TERPENES FROM *LINDERA GLAUCA*

H. KOMAE and N. HAYASHI

Department of Chemistry, Faculty of General Education, Hiroshima University, Hiroshima, Japan

(Received 3 August 1971)

*Plant.* *Lindera glauca* (Sieb. et Zucc) Blume. *Source.* Hiroshima Prefecture, Japan. *Uses.* Not known. *Previous work.* None.

*Leaves.* The essential oil (120 mg,  $n_D^{25}$  1.4972, 0.054% yield) was isolated from the fresh leaves (2.2 kg) by steam distillation. 1,8-Cineole (8.2%), caryophyllene (15.3%), and bornyl acetate (5.4%) were isolated from the oil by preparative GLC (Carbowax 20 M–20% at 150°) and identified by IR and NMR. Camphene (0.9%),  $\beta$ -pinene (1.1%), limonene (0.8%), were confirmed by GLC (Carbowax 20 M–3%, PEG-6000–3%). Unknown constituents (68.1%).

*Key Word Index*—*Lindera glauca*; Lauraceae; terpenes; caryophyllene.

---

Phytochemistry, 1972, Vol. 11, p. 853 to 854. Pergamon Press. Printed in England.

### TERPENES FROM *LITSEA JAPONICA*

H. KOMAE and N. HAYASHI

Department of Chemistry, Faculty of General Education, Hiroshima University, Hiroshima, Japan

(Received 3 August 1971)

*Plant.* *Litsea japonica* (Thunb.) Juss. *Source.* Shimane Prefecture, Japan. *Uses.* Not known. *Previous work.* None.

*Leaves.* The essential oil (483 mg,  $n_D^{25}$  1.4773, 0.006% yield) was isolated from the fresh leaves (8.0 kg) by steam distillation. Bornyl acetate (1.1%), geranyl acetate (0.9%),  $\gamma$ -elemene (1.1%), and caryophyllene (34.5%) were isolated by preparative GLC (Carbowax 20 M–20% at 160°) and identified by IR and NMR.  $\alpha$ -Pinene (2.0%), camphene (1.6%),  $\beta$ -pinene (0.9%), and limonene (0.3%) were confirmed by GLC (PEG-6000-3%, 50-HB-2000-3%). Unknown constituents (47.6%).

*Key Word Index*—*Litsea japonica*; Lauraceae; terpenes; caryophyllene.

---

Phytochemistry, 1972, Vol. 11, p. 854. Pergamon Press. Printed in England.

## MELIANTHACEAE

### MANGIFERIN IN *BERSAMA YANGAMBIENSIS*

M. VANHAELEN

Department of Pharmacognosy, Pharmacy Institute, Free University of Brussels, Brussels, Belgium

(Received 29 July 1971, in revised form 3 September 1971)

IN PURSUING the chemical analysis of *Bersama yangambiensis* Toussaint<sup>1</sup> (Yangambi, Republic Democratic of Congo), a yellow-brown pigment was detected in the crude ethanolic extract of the stem bark. The method of isolation involved removal of the fatty substances by light petroleum and exhaustive extraction by 94% ethanol. The pigment precipitated when  $\text{CH}_2\text{Cl}_2$  was added to the methanolic solution of the crude extract. Further purification was by fully trimethylsilylation of the precipitate followed by complete de-trimethylsilylation.<sup>2</sup> The spectroscopic properties, physical determinations,  $R_f$ s in different solvents, colour reactions, led to its identification as mangiferin (1,3,6,7-tetrahydroxy-2-C- $\beta$ -D-glucosylxanthone). As well direct comparison with authentic sample, the NMR spectrum of the trimethylsilyl derivative and the mass spectrum of the permethylated derivative, obtained by direct conversion of the TMS ether with  $\text{Me}_2\text{SO}_4$ , supported this identification. The occurrence of mangiferin in *Bersama yangambiensis* is the first record of this pigment in the Melianthaceae; it occurs to the extent of 6.5%. The medicinal use of extracts from *Bersama yangambiensis* by the autochtons for treatment of oedema could be related to the presence of mangiferin.<sup>3,4</sup>

NMR spectrum of mangiferin TMS ether: H-4,  $\delta$  = 6.35 ppm; H-5,  $\delta$  = 6.77 ppm; H-8,  $\delta$  = 7.59 ppm; 2-C- $\beta$ -D glucosyl,  $\delta$  = 3.10–4.90 ppm (7H); -Si ( $\text{CH}_3$ )<sub>3</sub>,  $\delta$  = 0.05–0.50 ppm ( $\approx$  70H).

*Acknowledgement*—We thank Dr. J. B. Harborne for authentic sample of mangiferin.

<sup>1</sup> J. TOUSSAINT, *Bull. Jard. Bot. Etat Brux.* **29**, 69 (1959).

<sup>2</sup> T. J. MABRY, K. R. MARKHAM and M. B. THOMAS, *The Systematic Identification of Flavonoids*, p. 257, Springer-Verlag, Berlin (1970).

<sup>3</sup> R. ANDRIANTSIFERANA, *C.R. Acad. Sci. Paris*, **264D**, 1215 (1967).

<sup>4</sup> C. MENTZER and A. R. RATSIMAMANGA, *Br. Pat.* 1,099,764 (CL.A61k), (17 Jan. 1968).

*Key Word Index*—*Bersama yangambiensis*; Melianthaceae; xanthones; mangiferin.