

ISOLATION OF COLUMBIN FROM *MELOTHRIA MADEROSPATANA*

YU P. CHEN and HONG Y. HSÜ

Brian Research Institute of Taiwan, 116 Chun-Ching South Road, Taipei, Taiwan

and

TSUEN I. RUO, KAZUO IGUCHI and HIROSHI KAKISAWA

Department of Chemistry, Tokyo Kyoiku University, Otsuka, Tokyo, Japan

(Received 5 May 1973. Accepted 5 June 1973)

Key Word Index—*Melothria maderospatana*; Cucurbitaceae; columbin.

Plant. *Melothria maderospatana* Cogn. DC. (Cucurbitaceae) collected in Chaochou, Pingtung, Taiwan. Voucher Specimen is deposited in the Herbarium of Botany, College of Science, National Taiwan University. *Previous work.* None. *Uses.* Antidote (flower) and analgesic (radix) in Taiwan.^{1,2} The ethanolic extract of the roots was concentrated to dryness and the residue was treated repeatedly with portions of warm tartaric acid solution until all of the basic substances were removed. The acid-insoluble residue was recrystallized from acetone to give white needles, m.p. 195–196°.

MS peaks at m/e 358·142 ($C_{20}H_{22}O_6$ requires 358·142) and 314·153 ($C_{20}H_{22}O_6-CO_2$ requires 314·152), IR bands at 3500, 3125, 1745, 1705, 1503 cm^{-1} , and NMR peaks at 7·60(2H,*m*), 6·57(1H,*m*), 6·40(1H,*dd*, J 5·0, 8·0 Hz), 6·12(1H,*dd*, J 1·5, 8·0 Hz), 5·47(1H,*dd*, J 4·5, 11 Hz), 5·21 (1H,*dd*, J 1·5, 5·0 Hz), 3·5(1H,*bs*, D_2O exchangeable OH), 2·47(4H,*m*), 2·23(1H,*d*, J 4·5 Hz), 1·97(1H,*d*, J 11 Hz), 1·6(2H,*m*), 1·10(3H,*s*), and 0·85(3H,*s*) indicated that the compound was columbin. The compound was further characterized by facile transformation to decarboxycolumbin, m.p. 141°, on heating over 195°. The IR and chromatographic behaviour were identical to those of an authentic sample. Columbin, a bitter substance, was isolated from *Jateorhiza palmata* Miers (Menispermaceae) with minor congeners jateorin, chasmanthin, and palmarin.^{3,4}

¹ KAN, W. S. (1971) *Pharmaceutical Botany*, pp. 541, National Research Institute of Chinese Medicine, Taipei, Taiwan.

² SASAKI, S. (1924) *Taiwan Minkan-Yakuyoshokubutsu Shi (Folk-Medicinal Plants in Taiwan)*, pp. 104, Kobunkan, Taipei, Taiwan.

³ BARTON, D. H. R. and ELAD, D. (1956) *J. Chem. Soc.* 2090.

⁴ BALASUBRAMANIAN, S. K., BARTON, D. H. R. and JACKMAN, L. M. (1962) *J. Chem. Soc.* 4816.