

ALKALOIDS OF *CANNABIS SATIVA* LEAVES

FAROUK S. EL-FERALY and CARLTON E. TURNER

Research Institute of Pharmaceutical Sciences, School of Pharmacy, University of Mississippi, University, MS 38677, U.S.A.

(Received 4 April 1975)

Key Word Index—*Cannabis sativa*; Cannabinaceae; marihuana leaves; alkaloids; 4-(β -dimethylaminoethyl)-phenol; hordenine

Plant. *Cannabis sativa* L. Cannabinaceae (leaves) **Source.** Research Institute of Pharmaceutical Sciences, University of Mississippi, Lot CQUF-72, QU-A(1)/C-70, S-62, grown in 1972.

Previous work. On leaves: choline and trigonelline [1], muscarine [2], an unidentified betain [3], and cannabamines A-D, all as yet of unknown structure [4].

Present work. Dried ground leaves were extracted by percolation with 95% EtOH at room temp. After removal of solvent *in vacuo* at 40°, the residue (14%) was partitioned between 2% citric acid and CHCl_3 . The aqueous phase was rendered alkaline with conc. NH_3 and extracted with CHCl_3 . The alkaloidal fraction thus obtained was further purified by dissolving in 1N HCl and extraction with CHCl_3 , then alkalized with conc. NH_3 and reextracted with CHCl_3 . This latter fraction was chromatographed on Si gel G.

Elution with MeOH–conc NH_3 (99:1) gave at first a dark brown material that gave a weak test for alkaloids. This was followed by a light yellow fraction which crystallized spontaneously when the solvent was evaporated and was recrystallized from Me_2CO –hexane to give feathery needles mp

118–119°; IR, $\nu_{\text{max}}^{\text{CHCl}_3}$ 3600, 3040, 2960, 2870, 2838, 2795, 1620, 1603, 1520, 1475, 1380, 1255, 1175, 1145, 1103, 1055, 1040, 1009, 870, 845 and 830 cm^{-1} ; UV, $-\lambda_{\text{max}}^{\text{MeOH}}$ 274 nm ($\epsilon = 1261$) and 211 nm ($\epsilon = 11000$) MS showed M^+ at 165 (14%) with the base peak at 58 ($\text{CH}_2 = \text{N}^+\text{Me}_2$). The NMR spectrum taken in CDCl_3 showed an exchangeable one-proton singlet at δ 8.08, an A_2B_2 pair of doublets integrating for 4 protons centered at δ 6.94 and 6.59 (J 9Hz), a distorted 4-proton triplet at δ 2.69 and a 6-proton singlet at δ 2.36. Direct comparison (mp & mmp, IR NMR) with an authentic sample of hordenine confirmed the identity of the isolated material.

Acknowledgements—The authors are grateful to Dr. A. F. Fentiman for assistance in obtaining the mass spectra. This investigation was supported in part by Grant DA-928 from the National Institute on Drug Abuse and the Research Institute of Pharmaceutical Sciences, School of Pharmacy, University of Mississippi.

REFERENCES

1. Merz, K. W. and Berger, K. B. (1940) *Arch. Pharm.* **49**, 278.
2. Kasniewski, V. (1954) *Deut. Apotheker-Zt.* **94**, 1177.
3. Lousberg, R. J. and Salemink, C. A. (1973) *Pharm. Weekblad* **108**, 1.
4. Klein, F. K., Rapoport, H. and Elliott, H. W. (1971) *Nature* **232**, 258.

L-3-CARBOXY-1,2,3,4-TETRAHYDRO- β -CARBOLINE, A NEW AMINO ACID FROM SEEDS OF *ALEURITES FORDII*

TAKUO OKUDA, TAKASHI YOSHIDA, NORIKO SHIOTA and JUNKO NOBUHARA

Faculty of Pharmaceutical Sciences, Okayama University, Tsushima, Okayama, Japan

(Received 21 January 1975)

Key Word Index—*Aleurites fordii*; Euphorbiaceae; non-protein amino acid; L-3-carboxy-1,2,3,4-tetrahydro- β -carboline; other amino acids.

Plant. *Aleurites fordii* Hemsl. **Source.** Okayama University campus and Handayama Botanical Garden. **Uses.** Tung oil has been used in paints,

varnishes, etc. **Previous work.** Isolation of the toxic principle of fruits [1], fractionation of basic and acidic constituents of tung oil [2]. **Present**