

Structures and Spectral Properties of (O-*sec.*-Butyldithiocarbonato-S,S') Bis(triphenylphosphine) Copper(I) and Silver(I) Complexes

by Q.F. Han, X.J. Yang, L.D. Lu and X. Wang*

Materials Chemistry Laboratory, Nanjing University of Science & Technology,
Nanjing, 210094, P.R. China.

(Received June 11th, 2003; revised manuscript August 11th, 2003)

New complexes $(\text{PPh}_3)_2\text{M}(\text{S}_2\text{COR})$ [R = *sec.*-butyl, M = copper (**1**), silver (**2**)] were synthesized by the reaction of excessive triphenylphosphine and copper(II) or silver(II) xanthate, and characterized using IR, EA, ^1H NMR and TG. Their crystal structures have been determined by X-ray diffraction. The light yellow crystal of the complex (**1**) is triclinic of space group $\text{P}\bar{1}$, with parameters $a = 10.300(2)$ Å, $b = 13.120(3)$ Å, $c = 14.570(3)$ Å, $\alpha = 89.53(3)^\circ$, $\beta = 72.81(3)^\circ$, $\gamma = 78.61(3)^\circ$, and $Z = 2$. The yellow crystal of the complex (**2**) is triclinic of space group $\text{P}\bar{1}$, with parameters $a = 10.330(2)$ Å, $b = 13.410(3)$ Å, $c = 14.420(3)$ Å, $\alpha = 88.61(3)^\circ$, $\beta = 73.60(3)^\circ$, $\gamma = 78.93(3)^\circ$, and $Z = 2$. In the two complexes, the central Cu or Ag atom is in a distorted tetrahedral environment and chelated by two phosphorus atoms from the triphenylphosphine groups and two sulfur atoms from the O-alkyldithiocarbonate. IR and ^1H NMR results supported the structures. The thermal analytical data indicate that the complex (**1**) began to decompose at 122.3° and decomposition was complete at 370.9° , leaving Cu_2S , while the complex (**2**) began to decompose at 133.3° and decomposed completely at 290.3° to Ag_2S .

Key words: copper complex, silver complex, triphenylphosphine, xanthate, synthesis, crystal structure

The ability of the O-alkyldithiocarbonate, ROCS_2^- ligand (xanthate) to bind to metal has been known for many years [1,2]. It forms a chelate with virtually all transition elements and has proven to be a versatile chelating agent for the separation and extracting of metals in analytical chemistry and mineral floating [3,4]. A review described in detail salts, complexes, free acids and esters as well as the reaction of xanthate [5]. Most of the bidentate complexes containing dithiolates such as dithiocarbamate and carbonadithiolate according to Singh *et al.* can react with Lewis bases [triphenylphosphine, phenanthroline, pyridine *et al.*] yielding the mono-, bis- and tetrakis-(Lewis base) adducts [6,7]. There have been many reports about the stu-

dies of the xanthate complexes towards several mono-, bi- and polydentate phosphine ligands [8–12].

Complexes of IB group metals of the general formula (Rxant)ML_n (with Rxant = O-alkylxanthate ion, L = PPh₃; n = 2 when M = Cu, Ag) have been synthesized and characterized by many workers [13,14]. The crystal structures of L₂Cu(MeXant) and L₂Cu(EtXant) have been reported [15–17]. However, the structures of L₂Ag(RXant) have not been proved by the X-ray diffraction, and the complexes L₂Cu(RXant) reported were prepared from copper(II) salts with overelaborated procedures. In this paper, (O-*sec.*-butyldithiocarbonato-S,S') bis(triphenylphosphine) copper(II) and silver(II) complexes were conveniently prepared and their crystal structures were presented. The IR, ¹H NMR and TG spectra of the complexes have also been determined.

EXPERIMENTAL

Preparation of the complexes. Synthesis of complex (PPh₃)₂Cu(S₂COR). To a aqueous solution of C₄H₉OCSSNa (0.17 g, 1.0 mmol) was added an aqueous solution of CuSO₄ (0.16 g, 0.5 mmol) with stirring. A yellowish brown precipitate (*sec.*-butylxanthato copper) was collected by filtration, washed with water, and dried under vacuum (yield 0.35 g, 96%). The dried precipitate was dissolved in ethanol and an excess of triphenylphosphine (0.84 g, 3.0 mmol) was added at 343–353 K with stirring. The solution was then refluxed for 1–2 h. Upon cooling, the light-yellow solution obtained was filtered and left to evaporate. After a few days, a light-yellow solid had separated out and was recrystallized from EtOH/CHCl₃. Single crystals suitable for X-ray analysis were obtained by slow evaporation of an EtOH solution at room temperature.

The synthesis procedure of complex (PPh₃)₂Ag(S₂COR) was similar to the complex **(1)**. The C and H contents of four compounds [Cu(S₂COR)₂, (PPh₃)₂Cu(S₂COR), Ag(S₂COR) and (PPh₃)₂Ag(S₂COR)] were determined by elemental analysis. Anal. Calcd. for C₁₀H₁₈CuO₂S₄: C, 33.15; H, 4.97. Found: C, 33.06; H, 5.14. for C₄₁H₃₉CuOP₂S₂: C, 66.73; H, 5.29. Found: C, 66.78; H, 5.28. for C₅H₉AgOS₂: C, 23.35; H, 3.50. Found: C, 23.48; H, 3.42. for C₄₁H₃₉AgOP₂S₂: C, 62.98; H, 5.01. Found: C, 62.89; H, 5.11.

Physical measurement. Elemental analysis was performed on PE-240 elemental analyzer. IR spectra were measured in KBr discs using a Bruker Vector 22 spectrophotometer. ¹H NMR spectra were obtained in CDCl₃ solution at 300 K on Bruker DRX 300 MHz spectrophotometer. Thermal analyses were performed in nitrogen gas with a Shimadzu TGA-50(10°/min).

Crystallographic data collection and solution of structure. A summary of the key crystallographic information is given in Table 1. The selected crystals of the title Table 1 compounds were mounted on a Bruker P4 diffractometer. Reflection data and reflections for the unit cell determination were measured at 293 K using MoK α radiation ($\lambda = 0.071073$ nm) with a graphite monochromator. Intensities were corrected for Lorentz and polarization effects and empirical absorption correction was applied using SHELXL5 [18]. The structures were solved by direct methods using SHELXS97 [19]. Nonhydrogen atoms were refined on F^2 anisotropically by full-matrix least squares method [20]. The hydrogen atoms were located from the difference map and refined isotropically. The final least-square cycle gave $R_1 = 0.0636$ and $wR_2 = 0.1465$ for 6465 reflections with $I > 2\sigma(I)$; the weighing scheme, $w = 1/[\sigma^2(F_o^2) + (0.1000P)^2]$, where $P = (F_o^2 + F_c^2)/3$ for the complex **(1)**, and $R_1 = 0.0468$ and $wR_2 = 0.0854$ for 6593 reflections with $I > 2\sigma(I)$; the weighing scheme, $w = 1/[\sigma^2(F_o^2) + (0.0300P)^2 + 0.3000P]$, where $P = (F_o^2 + 2F_c^2)/3$ for the complex **(2)**. Atomic scattering factors and anomalous dispersion correction were taken from the International Table for X-ray Crystallography [21]. The crystal structures for the title compounds have been deposited at the CCDC (CCDC-210337 for the complex **(1)**, and CCDC-211071 for the complex **(2)**). These data can be obtained free of charge via www.ccdc.cam.ac.uk/conts/retrieving.html (or from the Cambridge Crystallographic Data Centre, 12, Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033; or deposit@ccdc.cam.ac.uk).

Table 1. Crystal data and data collection details for the title compounds.

	Complex (1)	Complex (2)
Empirical formula	C ₄₁ H ₃₉ CuOP ₂ S ₂	C ₄₁ H ₃₉ AgOP ₂ S ₂
Formula weight	737.32	781.65
Wavelength, Å	0.71073	0.71073
Crystal system, space group	Triclinic, P $\bar{1}$	Triclinic, P $\bar{1}$
Unit cell dimensions	a = 10.300(2) Å α = 89.53(3)° b = 13.120(3) Å β = 72.81(3)° c = 14.570(3) Å γ = 78.61(3)°	a = 10.330(2) Å α = 88.61(3)° b = 13.410(3) Å β = 73.60(3)° c = 14.420(3) Å γ = 78.93(3)°
Volume, Å ³	1841.3(6)	1879.7(7)
Z	2	2
Calculated density, Mg m ³	1.330	1.381
Absorption coefficient, mm ⁻¹	0.825	0.763
F(000)	768	804
Crystal size, mm	0.3×0.2×0.1	0.35×0.260.30
Theta range for data collection	1.47 to 24.97°	1.47 to 24.97
Limiting indices	0 ≤ h ≤ 12, -15 ≤ k ≤ 15, -16 ≤ l ≤ 17	
Reflections collected/unique	6861/6465 [R(int) = 0.0517]	6997/6593 [R(int) = 0.0405]
Data/restraints/parameters	6465/0/424	6593/1/424
Goodness-of-fit on F ²	0.907	1.016
Final R indices [I > 2σ(I)]	R1 = 0.0636, wR2 = 0.1465	R1 = 0.0468, wR2 = 0.0854
R indices (all data)	R1 = 0.1991, wR2 = 0.2010	R1 = 0.1516, wR2 = 0.1066
Largest diff. peak and hole, e. Å ⁻³	0.476 and -0.416	0.510 and -0.496

RESULTS AND DISCUSSION

Analytical data. EA results for Cu(S₂COR)₂, (PPh₃)₂Cu(S₂COR), Ag(S₂COR) and (PPh₃)₂Ag(S₂COR) are listed in the experimental section. These data are in good agreement with the calculated ones from the deduced molecular formula.

IR and NMR spectra properties. Table 2 lists three frequencies of four compounds. The IR spectra of four compounds exhibit characteristic strong bands at 1198–1177 cm⁻¹ for ν (C–O–R) and 1101–1090 cm⁻¹ for ν (C–O). The bands at 1037–1015 cm⁻¹ are the characteristic absorption of ν (C=S) of the dithioformate group [12,14]. The IR spectra data show that the values of ν (C–O–R) are lower and ν (C=S) are higher when copper or silver xanthate are coordinated by triphenylphosphine.

The ¹H NMR spectra of two title complexes, triphenylphosphine and *sec.*-butylxanthato silver are listed in Table 2. As found previously from the ¹H NMR signals, there were not significant differences when xanthate coordinated to triphenylphosphine except for a small increase of the ν values [22]. Similar phenyl resonance of two complexes shows that the P–Cu–P angle is close to the P–Ag–P angle. All these findings support the crystal structures.

Table 2. IR and ^1H NMR spectroscopic data for the compounds^a.

Compound	IR (cm^{-1}) data		
	$\nu(\text{C-O-R})$	$\nu(\text{C-O})$	$\nu(\text{C=S})$
$\text{Cu}(\text{S}_2\text{COR})_2$	1198	1101	1015
$\text{L}_2(\text{S}_2\text{COR})\text{Cu}$	1190	1090	1035
$\text{Ag}(\text{S}_2\text{COR})$	1177	1093	1037
$\text{L}_2(\text{S}_2\text{COR})\text{Ag}$	1198	1095	1022

Compound	^1H NMR (δ ppm)				
	L	CH_3^{b}	CH_2^-	CH_3^-	CH^-
$\text{L}_2(\text{S}_2\text{COR})\text{Cu}$	7.2583	0.9158t	1.7386	1.2478d	5.2258
$\text{L}_2(\text{S}_2\text{COR})\text{Ag}$	7.3205	0.9771t	1.8356	1.3433d	5.2662
$\text{Ag}(\text{S}_2\text{COR})$		0.8906t	1.6286	1.1838d	4.1032
L	7.2653				

^aL = PPh_3 ; R = sec.-butyl. ^bend methyl group. d = doublet; t = triplet.

Thermal gravimetric analyses. In order to examine the thermal stability of the title complexes, thermal gravimetric analyses were carried out from 20°C to 600°C in nitrogen gas. The TGA curve of the complex (**1**) displays a weight loss of 87.568% between 122°C to 370°C , which corresponds to the decomposition of the complex, leaving Cu_2S (calculated: 89.145%). For the complex (**2**), between 133.3°C to 290.3°C , there is a weight loss of 83.142% corresponding to the decomposition to Ag_2S (calculated: 84.121%).

Crystal structure description. The crystal structures of the title compounds are illustrated in Fig. 1. Selected bond lengths and angles are given in Table 3. In two complexes, the copper or silver atom is tetrahedrally coordinated by two sulfur atoms from the bidentate xanthate and two phosphorus atoms from the triphenylphosphine ligand. The geometry of two complexes is highly distorted due to the small bite of the xanthate ligands, the bond angle of S1-Cu-S2 reaching the value of $73.96(9)^\circ$ (range of the other angles $101.45(8)$ – $125.99(8)^\circ$), and the angle of S1-Ag-S2 reaching the value of $67.13(6)^\circ$ (range of the other angles $98.90(6)$ – $125.65(6)^\circ$). The distortion is not restricted to the bond angles but also appears in bond distances. In the complex (**1**), while the Cu-P bond distances ($2.253(2)$, $2.269(2)$ Å), similar each to other, are in excellent agreement with those reported, the Cu-S bond distances ($2.369(2)$, $2.484(2)$ Å) are not equivalent [16]. A similar arrangement is also observed in the complex (**2**), the Ag-P1 bond distance ($2.4687(18)$ Å) close to the bond length of Ag-P2 ($2.4327(15)$), and the Ag-S bond distances ($2.7437(19)$, $2.5759(18)$ Å) are different. An unsymmetrical distribution around the metal atom has been proved in other monomeric copper(II) complexes and in dimeric silver(II) complex and tentatively rationalized on the basis of the well-known tendency of copper(II) or silver(II) to be three- or four-coordinated [23,24]. The P-Cu-P angle ($125.99(8)^\circ$) is almost

equal to the P–Ag–P angle ($125.65(6)^\circ$). The reason why the mean Ag–P (2.4507 \AA) and Ag–S (2.6598 \AA) bond lengths are longer than the mean Cu–P (2.261 \AA) and Cu–S (2.426 \AA) bond distances is probably that the Ag^{1+} radius is bigger than Cu^{1+} radius.




Figure 1. ORTEP perspective of the title compounds showing the atom numbering scheme.

Table 3. Selected bond distances (\AA) and angles ($^\circ$) for the title compounds.

	Complex (1)		Complex (2)	
Cu–P(1)	2.269(2)	Ag–P(1)	2.4687(18)	
Cu–P(2)	2.253(2)	Ag–P(2)	2.4327(15)	
Cu–S(1)	2.369(2)	Ag–S(1)	2.7437(19)	
Cu–S(2)	2.484(2)	Ag–S(2)	2.5759(18)	
S(1)–C(5)	1.664(9)		1.670(6)	
S(2)–C(5)	1.677(8)		1.676(6)	
C(5)–O	1.346(9)		1.344(7)	
O–C(4)	1.427(12)		1.467(8)	
P(1)–Cu–P(2)	125.99(8)	P(1)–Ag–P(2)	125.65(6)	
P(1)–Cu–S(1)	110.80(9)	P(1)–Ag–S(1)	98.90(6)	
P(2)–Cu–S(1)	115.08(9)	P(2)–Ag–S(1)	119.60(6)	
P(1)–Cu–S(2)	101.45(8)	P(1)–Ag–S(2)	111.78(6)	
P(2)–Cu–S(2)	117.12(8)	P(2)–Ag–S(2)	117.69(6)	
S(1)–Cu–S(2)	73.96(9)	S(1)–Ag–S(2)	67.13(6)	

In the complex (**1**), Cu, P1, P2 and C35 atoms are planar, with the maximum deviation from the plane being -0.028 \AA for Cu. Ag atom in the complex (**2**) lies on a plane defined by the atoms Ag, S1, S2 and C5, with the maximum deviation from the plane being -0.113 \AA for C5.

The S_2CO group is planar to a very good approximation. The C(5)–O bond distance [$1.346(9) \text{ \AA}$ in the complex (**1**) and, $1.344(7) \text{ \AA}$ in the complex (**2**)] is shorter than C(4)–O [$1.427(12) \text{ \AA}$ in the complex (**1**) and $1.467(8) \text{ \AA}$ in the complex (**2**)] bond length, which is indicative of considerable double-bond character. The C–S bond distances, $1.664(9)$ – $1.677(8) \text{ \AA}$ and $1.670(6)$ – $1.676(6)$ in complex (**1**) and (**2**), respectively, agree well with those in related compounds [12,16,17].

The complex $(\text{PPh}_3)_2\text{Cu}(\text{S}_2\text{COR})$ was easily synthesized here by the reaction of excessive triphenylphosphine and copper(II) xanthate, and yet similar derivatives reported were obtained by copper(II) salts [16,17]. Therefore, triphenylphosphine here acts as reducing agent that reduces Cu^{2+} to Cu^{1+} and it plays the role in stabilizing the low-valent metal as p-acid ligands as reported [25,26]. There have also been reports that $\text{Au}(\text{S}_2\text{CNC}_5\text{H}_{10})_3$ reduced to gold(I) complex or azide reduction with triphenylphosphine [27,28].

CONCLUSIONS

The new complexes $(\text{PPh}_3)_2\text{M}(\text{S}_2\text{COR})$ ($\text{R} = \textit{sec.}$ -butyl, $\text{M} = \text{copper}$ or silver) have been synthesized and their crystal structures have been determined by X-ray diffraction. The results of IR, EA and ^1H NMR spectroscopic measurement for the complexes agree well with the crystal structures. The thermal analytical data showed that the complexes $(\text{PPh}_3)_2\text{Cu}(\text{S}_2\text{COR})$ and $(\text{PPh}_3)_2\text{Ag}(\text{S}_2\text{COR})$ decomposed completely at 370.9°C and 290.3°C , respectively, leaving Cu_2S and Ag_2S .

REFERENCES

1. Formanek J., Holeckova H., Charewicz W., Walkowiak W. and Gendolla T., *Przem. Chem.*, **64**, 98 (1985).
2. Kotlyar D.G. and Tolley W.K., *Min. Eng.*, **11**, 1033 (1995).
3. Gimeno M.C., Jambriña E., Laguna A., Laguna M., Murray H.H. and Terroba R., *Inorg. Chem. Acta*, **249**, 69 (1996).
4. Allen F.H. and Kennard O., *Chem. Des. Autom. News*, **8**, 31 (1993).
5. Gerhard G., *Sulfur Rep.*, **14**, 1 (1993).
6. Singh M.K., Aggarwal R.C. and Singh B., *Synth. React. Inorg. Met-Org. Chem.*, **15**, 459 (1985).
7. Araya M.A., Cotton F.A., Matonic J.H. and Murillo C.A., *Inorg. Chem.*, **34**, 5424 (1995).
8. Perpignan M.F., Ballester L., Gonzalez-Casso M.E. and Santos A., *J. Chem. Soc., Dalton Trans.*, 281 (1987).
9. Ballester L., Gutierrez A., Perpignan M.F., Gutierrez-Puebla E. and Ruiz-Valero C., *Polyhedron*, **9**, 1341 (1990).
10. Ballester L., Gutierrez A., Perez-Serrano S. and Perpignan M.F., *Inorg. Chem.*, **30**, 2994 (1991).
11. Cotton F.A. and Hong B., *Prog. Inorg. Chem.*, **40**, 179 (1992).
12. Ballester L., Gutierrez A. and Perpignan M.F., *Polyhedron*, **15**, 1103 (1996).
13. Little L.H., Poling G.W. and Leja J., *Can. J. Chem.*, **39**, 745 (1961).

14. Cariati F., Ganadu M.L., Naldini L. and Seneci S., *Gazz. Chim. Ital.*, **109**, 181 (1979).
15. Ciani G., Manassero M., Naldini L. and Sansoni M., Proc. of the Yugoslav Center of Cryst., Sup. to Vol. 11, (1976).
16. Bianchini C., Ghilardi C.A., Meli A., Midollini S. and Orlandini A., *Inorg. Chem.*, **24**, 932 (1985).
17. Jin X.L., Tang K.L., Xia T.B. and Tang Y.Q., *Chinese J. Struct. Chem.*, **14**, 224 (1995).
18. Sheldrick G.M., *SHELXTL V5, The Complete Software Package for Single Crystal Structure Determination*, Siemens, AG, Analytical Systems Aut37, D76181 Karlsruhe 21, Germany (1995).
19. Sheldrick G.M., *Acta Cryst.*, **A46**, 467 (1969).
20. Sheldrick G.M., *SHELXTL 93. Program for Crystal Structure Refinement*, University of Göttingen, Germany (1993).
21. Wilson A.J., *International Table for X-ray Crystallography*, Vol. C, 1992. Kluwer Academic Publishers, Dordrecht: Tables 6.1.1.4, pp. 500–502 and 4.2.6.8, pp. 219–222.
22. Kowala C. and Sawan J.M., *Aust. J. Chem.*, **19**, 999 (1966).
23. Bianchini C., Ghilardi C.A., Meli A. and Orlandini A., *Inorg. Chem.*, **22**, 2188 (1983).
24. Bianchini C., Ghilardi C.A., Meli A., Orlandini A. and Scapacci G., *J. Chem. Soc., Dalton Trans.*, 1969 (1983).
25. Misra T.K. and Sinha C., *Trans. Met. Chem.*, **24**, 168 (1999).
26. Jian F.F., Bei F.L., Lu L.D., Yang X.J., Wang X., Razak I.A., Raj S.S.S. and Fun H.K., *Acta Cryst.*, **C56**, e288 (2000).
27. Razak I.A., Raj S.S.S., Fun H.K., Jian F.F., Bei F.L., Yang X.J., Lu L.D. and Wang X., *Acta Cryst.*, **C56**, 666 (2000).
28. Carrington S., Renault J., Tomasi S., Corbel J.C., Uriac P. and Blagbrough I.S., *Chem. Commun.*, 1341 (1999).