

## Erratum

### Carboxylated polysulfone membranes having a chiral recognition site induced by an alternative molecular imprinting technique\*

Masakazu Yoshikawa<sup>1</sup>, Jun-ichiro Izumi<sup>1</sup>, Takashi Ooi<sup>1</sup>, Toshio Kitao<sup>1</sup>, Michael D. Guiver<sup>2</sup>, and Gilles P. Robertson<sup>2</sup>

1 Department of Polymer Science and Engineering, Kyoto Institute of Technology, Matsugasaki, Kyoto 606-8585, Japan

2 Institute for Chemical Process and Environmental Technology, National Research Council of Canada, Ottawa, Ontario K1A 0R6, Canada

Polymer Bulletin, **40**, 517 (1998)

Figures 2 and 3 were switched round. The corrected figures are:




Figure 2 Adsorption isotherms of D-Glu and L-Glu in the Z-D-Glu imprinted polymeric membrane. ((Z-D-Glu) / (PSf) = 0.5;  $k_D = 1.7 \times 10^3$ ; [PSf] =  $7.4 \text{ mol dm}^{-3}$ ,  $n = 5.7 \times 10^{-2}$ ;  $K_C = 7.9 \times 10^3 \text{ mol}^{-1} \text{ dm}^3$ .)




Figure 3 Adsorption isotherms of D-Glu and L-Glu in the Z-L-Glu imprinted polymeric membrane. ((Z-L-Glu) / (PSf) = 0.5;  $k_D = 1.8 \times 10^3$ ; [PSf] =  $7.4 \text{ mol dm}^{-3}$ ,  $n = 5.7 \times 10^{-2}$ ;  $K_C = 7.9 \times 10^3 \text{ mol}^{-1} \text{ dm}^3$ .)