

Erratum

Erratum to “A novel approach to recycling of polyurethanes: chemical degradation of flexible polyurethane foams by triethyl phosphate”
[Polymer, 41 (2000) 7017][☆]





K. Troev^{*}, G. Grancharov, R. Tsevi, A. Tsekova

Institute of Polymers, Bulgarian Academy of Sciences, 1113 Sofia, Bulgaria

Received 4 October 1999; received in revised form 4 January 2000; accepted 12 January 2000

We wish to point out to readers that errors were present in the published version of the above article. Table 5 should appear as below:

Table 5
Phosphorus containing products from the chemical degradation of flexible polyether polyurethane foam by triethyl phosphate

N	Structure	³¹ P NMR, δ (ppm), J (Hz)	Content (%)	
			Upper layer	Lower layer
4		-0.20, quintet 3J(P,H) = 7.42	61.5	74.2
5		-0.55, q 3J(P,H) = 7.3	21.2	2.6
6		0.24, septet 3J(P,H) = 7.8	5.1	3.3
7		0.55, t 3J(P,H) = 8.0	2.6	3.5

[☆] PII of original article: S0032-3861(00)00054-9

^{*} Corresponding author. Tel.: +359-2-978-2203; fax: +359-2-707-523.

E-mail address: ktroev@polymer.bas.bg (K. Troev).

Table 5 (continued)

N	Structure	^{31}P NMR, δ (ppm), J (Hz)	Content (%)	
			Upper layer	Lower layer
8	$\begin{array}{c} \text{O} \quad \text{O} \quad \text{O} \\ \parallel \quad \parallel \quad \parallel \\ \text{CH}_3\text{CH}_2\text{O}-\text{P}-\text{O}-\text{P}-\text{O}-\text{P}-\text{OCH}_2\text{CH}_3 \\ \quad \quad \\ \text{CH}_3\text{CH}_2\text{O} \quad \text{OH} \quad \text{OH} \end{array}$	m, -11.16, -12.28	9.6	16.3