

Erratum

Erratum to “Analysis of deuterium NMR spectra of probe chains diffusing in a stretched polybutadiene network” [Polymer 44 (2003) 5263–5267][☆]

S. Valić^{a,b,*}, P. Judeinstein^c, B. Deloche^a

^aLaboratoire de Physique des Solides (CNRS-UMR 8502), Université Paris-Sud, 91405 Orsay, France

^bRudjer Bošković Institute, P.O. Box 180, Zagreb, HR-10001 Zagreb, Croatia

^cRMN en Milieu Orienté (CNRS ESA 8074), Bât. 410, Université Paris-Sud, 91405 Orsay, France

The publishers regret that when this article was published Figure 2 was published in the old version without calculated NMR spectra. The new version of Fig. 2, containing the experimental and calculated spectra, is shown below.




Fig. 2. ²H NMR spectra of PBD probe chains diffusing through the PB network observed for various extension ratio λ . Left side: experimental spectra; right side: calculated spectra. The doublets are assigned as following: (a) *cis* units—CD₂ component (▽), (b) *cis* units—CD component (▼), (c) *trans* units—CD₂ component (○) and (d) *trans* units—CD component (●).

[☆] doi of original article 10.1016/S0032-3861(03)00475-0

* Corresponding author.

E-mail address: valic@rudjer.irb.hr (S. Valić).