

This article was downloaded by:

On: 29 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Supramolecular Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713649759>

Novel 3-D Network: From the Reaction of Polyamine with Pyromellitic Dianhydride

Xun-Cheng Su^a; Hua-Kuan Lin^a; Shou-Rong Zhu^a; Lin-Hong Weng^a; Xue-Bing Leng^a; Yun-Ti Chen^a

^a Department of Chemistry, Nankai University, Tianjin, PR China

To cite this Article Su, Xun-Cheng , Lin, Hua-Kuan , Zhu, Shou-Rong , Weng, Lin-Hong , Leng, Xue-Bing and Chen, Yun-Ti(2011) 'Novel 3-D Network: From the Reaction of Polyamine with Pyromellitic Dianhydride', *Supramolecular Chemistry*, 13: 5, 593 – 600

To link to this Article: DOI: 10.1080/10610270108039792

URL: <http://dx.doi.org/10.1080/10610270108039792>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Novel 3-D Network: From the Reaction of Polyamine with Pyromellitic Dianhydride

XUN-CHENG SU, HUA-KUAN LIN*, SHOU-RONG ZHU, LIN-HONG WENG,
XUE-BING LENG and YUN-TI CHEN

Department of Chemistry, Nankai University, Tianjin 300071, PR China

(Received 22 January 2001)

Under mild condition the reaction of pyromellitic dianhydride with polyamine in methanol solution results in ammonium carboxylate salts rather than amide polymer. X-ray analysis indicates a manifestation of 3-D network linked by $N-H^+ \cdots O^- \cdots H-N^+$ hydrogen bonding for the three ammonium salts, in which the porous spaces are filled with water molecules.


Keywords: Self-assembly; Pyromellitic dianhydride; Polyamine; 3-D network

INTRODUCTION

It is well documented that the organic or organic-inorganic self-assembly is the spontaneous association of molecular subunits into structurally well-defined aggregates which are organized by feeble interactions such as hydrogen bond, $\pi-\pi$ stacking, salt bridge and so on [1]. Such molecular solids often possess interesting properties: (1) optical behavior, electrical conductivity, and ferromagnetism; (2) potential ability for selective inclusion and transportation of ions and molecules and the catalysis for

specific chemical reactions [2]. As for organic molecular aggregates, current efforts in assembling molecular solids mainly focus on 1-D and 2-D aggregates [3], while reports about the 3-D networks of organic molecular aggregates are relatively fewer except for the diamonded networks [4]. In addition, the reported organic solids mostly from weak bonds are relatively unstable and most of them will change their forms under fairly high temperature. Such disadvantage thus limits the further potential uses in practice. The carboxyl groups have been extensively applied in the molecular solid design [5], but the studies of their ammonium salts are relatively fewer [6]. It is predicated that the hydrogen bonds formed by carboxylate or water with ammonium will be much augmented by the electrostatic interactions in the ammonium salts, and then the stability of these organic salts will be enhanced as well. It is of much interest to investigate the delicate interactions of polyammonium benzenetetracarboxylate in the solid state so as to thread light on the design and synthesis of new self-assembling solids for functional

*Corresponding author.



SCHEME 1 Schematic illustration of the formation of 3-D networks.

materials, since the polyamine separated by flexible C—C bonds has more binding sites to form hydrogen bond. With this in mind we investigate the reaction of pyromellitic dianhydride with polyamine in methanol solution under mild conditions and the results show that the carboxylate salts are preferably produced rather than amide polymer (Scheme 1). The X-ray analysis, IR and TG analysis have been performed in the investigation of the supramolecular structures self-assembled by carboxylate and ammonium. X-ray analysis indicates that the ammonium carboxylate salts recrystallized in water are of 3-D networks connected by the hydrogen bonds of ammonium and carboxylate, and the stability of 3-D networks is further enhanced by Coulombic forces and water molecules.

RESULTS AND DISCUSSION

Pyromellitic dianhydride is widely used in polymer and material science, and its derivatives have been extensively studied since the few past decades. When it reacts with polyamine at higher temperature, the main product is of amide polymer; while the ammonium salts predominate over others in the mild condition. Under room temperature, the reaction of pyromellitic dianhydride with polyamine in methanol solution leads to formation of ammonium salt as white powders. Single crystals are obtained from the slow evaporation of the respective ammonium salt in aqueous solution. X-ray analysis indicates a manifestation of network linked by $N-H^+ \cdots O^- \cdots H-N^+$ hydrogen

bonding for the crystals **1** to **3**. Crystal structure analysis further indicates that the **1**, **2**, and **3** are of 1:1 aggregates. In case of **2** as shown in Figure 2, the linear polyamine are tetra-protonated and each carboxylic acid is deprotonated, and thus the organic salt are neutral. The bond lengths of C—O are similar (1.24 to 1.26 Å), indicating the charge dispersing average along the O—C—O with the bond angles around 124°. As shown in Figure 1, the four carboxylate are not in a plane, so that they can point out to three different directions. Furthermore, the protonated nitrogen atom in the flexible chain of polyamine is in tetrahedral environment and interacts with carboxylic acid through hydrogen bonds. In Figure 2, it is observed that the rigid benzene carboxylic acid and the flexible polyamine arrange in an interlaced sheet to give rise a three dimensional architecture through the hydrogen bonding of $N-H^+ \cdots O^- \cdots H-N^+$ and $N-H^+ \cdots OCOO^-$. Just different to Zaworotko's report [7], there are no water-microchannels in such 3-D networks. The water molecules, which are involved in the hydrogen bonding with ammonium and carboxylates, occupy the porous spaces and further strengthen the 3-D network structure. Selected distance and angle data estimated from the X-ray crystallographic analyses are summarized in Table II (for the sake of clarity, only those of **2** are listed). In the supramolecular architecture, Coulombic forces between the oppositely charged ions, ammonium and carboxylate, and the included guest water molecules in the cavity further enhance the 3-D stability. It is noteworthy to point out that the delicate structures of **2** differ from **1** and **3**, which have centric symmetry and therefore achiral. The self-assembled aggregate **2** is unique D_2 and is therefore chiral. This is a good example from achiral components to chiral products with losing symmetry by the organic synthesis design [8]. In view of these facts we can study a series of polyamine and pick out the suitable chiral crystals with potential applications in non-linear optics.




FIGURE 1 ORTEP drawings of the structural units for 1, 2 and 3. Thermal ellipsoids are scaled to the 30% probability level. Hydrogen atoms are omitted for clarity.

The 3-D crystals are stable in air and lose no moisture. But if allowed to immerse in methanol or ethanol under ambient conditions, the regular aggregates will lose crystallinity and turn into white powders with losing the included water molecules. Re-immersed in water, the powders will restore their beautiful crystallinity. The experiment implies that the included water

plays an important role to sustain the regular structure of the self-assembly aggregates. To investigate the stability of the included water in such aggregates, TGA experiments were performed under N_2 atmosphere. The results show that at around $80^\circ C$ the crystals will lose water and turn into powders. At around $130^\circ C$ the pattern reaches platform (Fig. 3) and the weight





FIGURE 2 ORTEP views of the three-dimensional hydrogen-bonded network present in crystals. In each view, hydrogen atoms are omitted for clearness and hydrogen bonds are represented by dashed line. (a) view from plane (100) of 1; (b) view from plane (100) of 2; (c) view from plane (100) of 3.



(c)

FIGURE 2 (Continued).




FIGURE 3 Overlay of TG traces of the title coordination polymer, for the sake of the clarity the DTG is deleted. The sample was heated at 20°C/min in a nitrogen atmosphere.

loss is consistent with that of water filled in the porous space (1: 14.83, calcd. 15.25; 2: 14.40, calcd. 17.85; 3: 13.69, calcd. 14.38). TGA indicates the white powders can be stable up to around 250°C, which are in the range of pure ammonium salts. It can be concluded that the hydrogen bonds formed by the included water with

ammonium and carboxylate in the aggregates appear to be critically essential in stabilizing the regular structures of the open 3-D polymeric networks.

In summary, the reaction of pyromellitic dianhydride with polyamine represents a new route to generate 3-D network, which is

connected by hydrogen bonding and Coulombic forces. This type of 3-D structure may shed highlight on the design and synthesis of the functional materials. Some important observations can be concluded. Firstly, the chain length of polyamine may be modified and further substituents (including chiral group) can be introduced. Secondly, chirality of the aggregates salts may be produced as suitable ammine is selected. If possible, a new series of NLO (non-linear optics) materials [9] might be produced.

EXPERIMENTAL SECTION

Materials

The chemicals and solvents used in the synthesis of products were of analytical reagent and were purified according to the literature methods [10].

Measurements

The Fourier-transform IR spectra were taken on a Nicolet 560 ESP spectrometer and elemental analyses were made on a Perkin-Elmer 240C analyzer. 200 MHz ^1H NMR spectroscopic measurements were performed on a Bruker AM-200 NMR spectrometer, using TMS (SiMe_4) as an internal reference at 25°C . Thermogravimetric analysis (TGA) and differential thermal analysis (DTA) were performed at a scan rate of $20^\circ\text{C}/\text{min}$ using a Netzsch TG 209 analyzer under N_2 atmosphere.

General Procedures for Synthesis of 1, 2 and 3

Pyromellitic dianhydride (5 mmol) was warmed to dissolve in 150 mL methanol and then cooled. A solution of the appropriate polyamine (5 mmol) in 30 mL methanol was added to such a stirred solution above at room temperature. The reaction mixture was stirred at room temperature for 5 h and filtered. The powder

product was washed twice with methanol and then recrystallized with hot water to give colourless crystals. Yield: 50.5~78.2%,

The suitable single crystals for X-ray analysis were obtained under the slow evaporation of the respective aqueous solution.

1. ^1H NMR (D_2O , 200 MHz): $\delta = 7.57$ (s, 2H), 3.0~3.1 (m, 12H). IR (KBr pellet): 3421.2 (vs), 3020.6 (s), 2697.7 (b), 1655.9 (s), 1569.5 (vs), 1487.0 (s), 1408.4 (s), 1366.6 (s), 1318.1 (s), 865.7 (m), 814.9 (m) cm^{-1} .

2. ^1H NMR (D_2O , 200 MHz): $\delta = 7.50$ (s, 2H), 3.17 (m, 8H), 2.89 (m, 4H), 1.72 (s, 2H). IR (KBr pellet): 3536.1 (vs), 3415.7 (vs), 3028.3 (b), 2796.1 (b), 1636.4 (s), 1598.9 (vs), 1560.7 (vs), 1537.6 (s), 1486.7 (s), 1411.0 (s), 1371.3 (vs), 1320.7 (s), 862.8 (m), 814.0 (m) cm^{-1} .

3. ^1H NMR (D_2O , 200 MHz): δ 7.64 (s, 2H), 3.37 (m, 4H), 2.94~3.13 (m, 8H), 1.91~1.98 (s, 4H). IR (KBr pellet): 3512.6 (vs), 3353.5 (b), 3008.5 (b), 2804.1 (b), 1603.0 (s), 1559.5 (vs), 1481.0 (s), 1422.9 (s), 1367.2 (vs), 1323.2 (s), 864.9 (m), 810.6 (m), 810.6 (m) cm^{-1} .

X-ray Crystallography

Crystals of 1, 2 and 3 suitable for single-crystal X-ray diffraction with suitable sizes were selected. Structural analyses for 1, 2, and 3 was performed on a Siemens SMART CCD diffractometer with graphite-monochromated $\text{Mo K}\alpha$ ($\lambda = 0.71073 \text{ \AA}$) The structure was solved by direct methods using the SHELXTL-97 package [11]. The ORTEP structures and the selected bond distances and angles are shown in Figure 1 and Table I, respectively.

1: Crystal data: $\text{C}_{16}\text{H}_{32}\text{N}_4\text{O}_{12}$, Mr = 472.46, triclinic, space group P-1. $a = 7.4765$ (13), $b = 7.6547$ (13), $c = 9.6846$ (16) \AA , $\alpha = 76.213$ (3), $\beta = 85.473$ (4), $\gamma = 84.286$ (4) $^\circ$, $V = 527.26$ (15) \AA^3 , $Z = 1$, $\rho_{\text{calcd}} = 1.488 \text{ Mg/mm}^3$, $F(000) = 252$, $T = 298$ (2) K, wavelength 0.71073 \AA . A colorless plate with the dimensions $0.05 \times 0.10 \times 0.20 \text{ mm}$ was

TABLE I Selected bond distance (Å) and angles (°) for 1, 2 and 3

Data for 1			
N(1)—C(6)	1.462(6)	N(2)—C(8)	1.481(5)
N(2)—C(7)	1.490(5)	C(1)—O(1)	1.225(5)
C(1)—O(2)	1.267(4)	C(5)—O(4)	1.245(4)
C(5)—O(3)	1.246(4)		
C(8)—N(2)—C(7)	112.0(3)	O(1)—C(1)—O(2)	123.7(4)
O(1)—C(1)—C(3)	120.2(4)	O(2)—C(1)—C(3)	116.1(4)
O(4)—C(5)—O(3)	124.2(4)	O(4)—C(5)—C(4)	119.1(3)
O(3)—C(5)—C(4)	116.7(3)	N(1)—C(6)—C(7)	113.0(4)
Data for 2			
C(5)—C(10)	1.5091(18)	C(7)—O(1)	1.243(2)
C(7)—O(2)	1.2456(19)	C(8)—O(3)	1.2441(19)
N(1)—C(11)	1.478(2)	N(2)—C(13)	1.490(2)
C(15)—N(3)	1.4841(19)	N(3)—C(16)	1.485(2)
O(1)—C(7)—O(2)	124.31(14)	O(2)—C(7)—C(1)	117.90(14)
O(3)—C(8)—O(4)	125.41(13)	O(8)—C(10)—O(7)	125.06(13)
C(13)—N(2)—C(12)	111.99(13)	C(15)—N(3)—C(16)	113.30(13)
Data for 3			
O(1)—C(1)	1.247(2)	O(2)—C(1)	1.241(2)
O(3)—C(2)	1.246(2)	O(4)—C(2)	1.2663(19)
C(8)—N(2)	1.496(2)	N(2)—C(9)	1.486(2)
O(2)—C(1)—O(1)	123.51(15)	O(2)—C(1)—C(3)	118.62(15)
O(3)—C(2)—O(4)	124.67(14)	N(2)—C(8)—C(7)	112.46(14)
C(9)—N(2)—C(8)	111.46(13)		

TABLE II Geometry of the selected hydrogen bonds in aggregates 2

D—H...A	r(D—H)/nm	r(H...A)/nm	r(D...A)/nm	∠(D...A)/°
N1—H1A...O2(a)	0.894	2.150	2.886	139.11
N1—H1B...O4(b)	0.931	1.897	2.814	167.66
N1—H1C...O1	0.992	1.881	2.857	167.27
N2—H2A...O4(c)	0.860	1.877	2.683	155.49
N2—H2B...O9	0.945	1.832	2.777	179.48
N3—H3B...O3(d)	0.926	1.978	2.839	154.06
N3—H3C...O5(e)	0.876	1.954	2.780	156.76
N4—H4A...O7(f)	0.919	1.978	2.888	170.28
N4—H4B...O5(g)	0.944	1.817	2.739	164.66
N4—H4C...O7(h)	0.908	1.972	2.856	163.92

Symmetry elements: (a) $x+1/2, -y+1/2, -z$; (b) $x+1, y, z$; (c) $x+1/2, -y+1/2, -z$; (d) $-x+1/2, -y+1, z-1/2$; (e) $x+1/2, -y+1/2, -z$; (f) $x, y, z-1$; (g) $x+1/2, -y+1/2, -z$; (h) $x-1/2, -y+1/2, -z$.

measured with a BRUKER SMART 1000. θ range for data collection 2.20 to 24.10°. A total of 1762 were measured, 1592 unique ($R_{\text{int}}=0.0336$). Weighting Scheme $w = 1/[S^2(F_o^2) + (0.0909P)^2 + 0.0000P]$, where $P = (F_o^2 + 2F_c^2)/3$. Goodness of fit on F^2 was 0.993. Final R indices [$I > 2\sigma(I)$]

$R1=0.0621, wR2=0.1450$. Largest difference Fourier peak and hole 0.255 and $-0.235 \text{ e}\text{\AA}^{-3}$.

2: Crystal data: $\text{C}_{17}\text{H}_{36}\text{N}_4\text{O}_{13}$, $M_r=504.50$, orthorhombic, space group $P2_12_12_1$ $a=7.7524(7), b=14.8513(15), c=20.532(2) \text{ \AA}$, $\alpha=90,$

$\beta = 90$, $\gamma = 90^\circ$, $V = 2363.9(4) \text{ \AA}^3$, $Z = 4$, $\rho_{\text{calcd}} = 1.418 \text{ Mg/mm}^3$, $F(000) = 1080$, $T = 293(2) \text{ K}$, wavelength 0.71073 \AA . A colorless plate with the dimensions $0.40 \times 0.35 \times 0.30 \text{ mm}$ was measured with BRUKER SMART 1000. θ range for data collection 1.69 to 25.01° . A total of 9805 were measured, 4162 unique ($R_{\text{int}} = 0.0270$). Weighting Scheme $w = 1/[S^2(F_o^2) + (0.0470P)^2 + 0.2988P]$, where $P = (F_o^2 + 2F_c^2)/3$. Goodness of fit on F^2 was 1.046. Final R indices [$I > 2\sigma(I)$] $R1 = 0.0281$, $wR2 = 0.0749$. Largest difference Fourier peak and hole 0.185 and -0.134 e\AA^{-3} .

3: Crystal data: $\text{C}_{18}\text{H}_{36}\text{N}_4\text{O}_{12}$, $\text{Mr} = 500.51$, triclinic, space group $P-1$. $a = 7.1774(10)$, $b = 7.9927(10)$, $c = 11.4286(14) \text{ \AA}$, $\alpha = 70.879(2)$, $\beta = 84.352(2)$, $\gamma = 68.202(2)^\circ$, $V = 574.96(13) \text{ \AA}^3$, $Z = 1$, $\rho_{\text{calcd}} = 1.446 \text{ Mg/mm}^3$, $F(000) = 268$, $T = 293(2) \text{ K}$, wavelength 0.71073 \AA . A colorless plate with the dimensions $0.35 \times 0.30 \times 0.30 \text{ mm}$ was measured with a BRUKER SMART 1000. θ range for data collection 1.89 to 25.02° . A total of 2417 were measured, 2023 unique ($R_{\text{int}} = 0.0144$). Weighting Scheme $w = 1/[\sigma^2(F_o^2) + (0.0777P)^2 + 0.1852P]$, where $P = (F_o^2 + 2F_c^2)/3$. Goodness of fit on F^2 was 1.048. Final R indices [$I > 2\sigma(I)$] $R1 = 0.0425$, $wR2 = 0.1202$. Largest difference Fourier peak and hole 0.265 and -0.238 e\AA^{-3} .

Acknowledgment

We gratefully acknowledge financial support from the National Natural Science Foundation (29771020 and 29971018) of China, the Nature

Science Foundation of Tianjin and the National Key Laboratory of the Coordination Chemistry, Nanjing University.

Supporting Information Available

Crystallographic files in CIF format are available free from the author.

References

- [1] (a) Desiraju, G. R. (1995). *Angew. Chem. Int. Ed. Engl.*, **34**, 2311–2329; (b) Aakeoy, C. B. and Seddon, K. R. (1993). *Chem. Soc. Rev.*, **22**, 397.
- [2] (a) Brunet, P., Simard, M. and Wuest, J. D. (1997). *J. Am. Chem. Soc.*, **119**, 2723; (b) Russel, V. A., Evans, C. C., Li, W. and Ward, M. D. (1997). *Since*, **276**, 575; (c) *Nonlinear Optical Properties of Organic Molecules and Crystals*, Chemia, D. S. and Zyss, J., Eds., Academic Press, Orlando, FL, 1987; **1**; (d) *Supramolecular Chemistry: Concepts and Perspectives*, VCH Publishers: Weinheim, Germany, 1995.
- [3] Gong, B., Zheng, C., Jankun, E. S., Yan, Y. and Zhang, J. (1998). *J. Am. Chem. Soc.*, **120**, 11194 and references therein.
- [4] Zaworotko, M. J. (1994). *Chem. Soc. Rev.*, **23**, 283.
- [5] Etter, M. C., Lipkowska, Z. U., Ebrahimi, M. Z. and Panunyo, W. T. (1990). *J. Am. Chem. Soc.*, **112**, 8415.
- [6] Koshima, H., Honke, S. and Fujita, J. (1999). *J. Org. Chem.*, **64**, 3916.
- [7] Subramanian, S. and Zaworotko, M. J. (1993). *Chem. Commun.*, p. 952.
- [8] Harris, K. D. M. and Hollingsworth, M. D. (1989). *Nature*, **341**, 19.
- [9] (a) Aakeroy, C. B., Hitchcock, P. B. and Seddon, K. R. (1993). *J. Chem. Soc., Chem. Commun.*, p. 152; (b) Andreu, R., Malfant, I., Lacroix, P. G. and Gornitzka, H. (1999). *Chem. Mater.*, **11**, 840.
- [10] Perrin, D. D. and Armarego, W. L. F., *Purification of Laboratory Chemicals*, 4th edn., Pergamon Press, Oxford, England, 1996.
- [11] Sheldrick, M., G. M., *SHELXL-97, Program for Crystal Structure Refinement*, University of Göttingen, Germany, 1997.