



0957-4166(95)00051-8


Enzymatic Resolution of 2-Hydroxy-4-Phenylbutanoic Acid and 2-Hydroxy-4-Phenylbutenoic Acid

Anju Chadha* and M. Manohar
SPIC Pharma R&D Center
Maraimalai Nagar, Tamil Nadu 603 209
India

Abstract: Racemic 2-hydroxy-4-phenylbutanoic acid and 2-hydroxy-4-phenylbutenoic acid have been resolved using a lipase. In each case, the (R)-2-hydroxy and the (S)-2-acetoxy acids were isolated with high enantiomeric excess and yield.

Introduction: α -hydroxycarboxylic acid derivatives are important building blocks for many biologically active compounds¹. (R)-2-hydroxy-4-phenylbutanoic acid (**1a**) is a precursor for the production of Angiotensin Converting Enzyme (ACE) inhibitors e.g. enalapril^{2,3}. The synthesis of (**1a**) has been reported by enantioselective reduction of the corresponding α -keto acid - both, enzymatically and by chemical enantioselective hydrogenation⁴. Enzymatic resolution of the phenylpropionaldehyde cyanohydrin with the subsequent synthesis of ethyl (R)-2-hydroxy-4-phenylbutyrate is known⁵. The enantioselective hydrolysis of ethyl 2-hydroxy-4-phenylbutanoate has also been reported⁶. The resolution of 2-hydroxy-4-phenylbutanoic acid (**1**) using hexanoic anhydride has been published⁷. In this paper we report a simpler and less expensive procedure. Racemic 2-hydroxy-4-phenylbutanoic acid (**1**) has been resolved using a lipase in t-butyl methyl ether with vinyl acetate as the acylating agent. The enantiomeric excess and yield of the resulting enantiomers are higher than those reported earlier.

We also report the lipase catalysed resolution of the related 2-hydroxy-4-phenyl-3-butenic acid (**3**). The case of the 2-hydroxy-3-enoic acids is special. These are allylic alcohols and at the same time unsaturated carboxylic acids. This is the reason for them to be considered useful for chiral synthesis. They have been prepared from the corresponding 2-oxo acids in enantiomerically pure form by the cells of *Proteus vulgaris* under anaerobic conditions⁸. Another procedure uses an enzyme (Lactate dehydrogenase) along with expensive cofactors⁴. To the best of our knowledge, the enzymatic resolution of the racemic 2-hydroxy-4-phenyl butenoic acid has not been reported so far. In this paper, we report a simple, straightforward lipase catalysed resolution of racemic 2-hydroxy-4-phenyl- butenoic acid with high enantiomeric excess and good yield.



Results And Discussion: Racemic 2-hydroxy-4-phenylbutanoic acid (**1**) was stirred with Lipase PS (Amano) and vinyl acetate in t-butyl methyl ether. The reaction was monitored on HPLC and after ~50% conversion (in 30h) the enzyme was filtered off, the products separated as reported earlier⁷ and purified by silica chromatography. The (R)-2-hydroxy-4-phenylbutanoic acid (**1a**) was recovered in ~45% yield and >99% ee, while the acetate (**2**) was recovered in 35% yield. The acetate was hydrolysed to the (S)-hydroxy acid (**1b**), 84% ee. Sugai and Ohta⁷, have obtained the R(-)-2-hydroxy-4-phenylbutanoic acid from the racemic hydroxy acid in the presence of lipase, using hexanoic anhydride as the acylating agent. The acylating agent used by us is the more commonly used, less expensive vinyl acetate. The enantiomeric excess and the yields of the products are high and, in fact, better than that reported earlier⁷. The enzymatic reduction of the α -keto acid, 2-oxo-4-phenylbutanoic acid has been achieved by many workers⁴ but all these procedures need cofactors. In the case of microbial reduction, an ee of 99% is achieved. However, this method, as reported by Simon et al.⁴ requires stringent anaerobic conditions. Our attempts to reduce the 2-oxo acid and ester with baker's yeast did not yield the required reduced product. Resolution of the racemic 2-hydroxy-4-phenylbutanoic acid using a lipase obviates the need for cofactors.

Immobilised enzymes have been known to be more stable with the distinct advantage of reusability. Even though many commercial processes use immobilised enzymes⁹, reports of immobilised lipases are few¹⁰. We have immobilised the lipase on celite. This immobilised lipase displays a decrease in activity initially which shows signs of picking up after four batch cycles. After immobilisation on Eupergit C, a similar trend is seen. Further experiments on immobilisation are in progress and will be reported elsewhere.

In a related study, 2-hydroxy-4-phenylbutanoic acid (**3**) was also resolved enzymatically. Earlier reports of the formation of (R)-2-hydroxy-4-phenylbutanoic acid⁸, indicate the use of whole cells of *Proteus vulgaris*. This organism requires anaerobic conditions under which quantitative reduction of the corresponding 2-oxo acid is carried out, in the presence of electron donors. The (S)-isomer is not formed. In our procedure, transacylation is done in the presence of a lipase to get the (R)-2-hydroxy acid (**3a**) and the (S)-2-acetoxy acid (**4**), which is hydrolysed to give the (S)-2-hydroxy acid (**3b**). The (R)-2-hydroxy-4-phenylbutanoic acid was recovered in ~42% yield and >99% ee, while the acetate was recovered in 34% yield and hydrolysed to give (S)-2-hydroxy-4-phenylbutanoic acid in 94% ee.

Experimental: To racemic 2-hydroxy-4-phenylbutanoic acid (**1g**) in MTBE was added vinyl acetate (5 ml) and lipase PS (500 mg) from Amano. The mixture was stirred at 25°C for 30h. HPLC (ODS column, CH₃CN:H₂O:THF, 20:65:15 @ 1 ml/min analysis showed a conversion of 51%. The reaction mixture was filtered and the recovered lipase was dried and reused. The filtrate and washings were concentrated and treated with cold hexane to obtain crystalline material (**1a**) (0.45g), m.p. 114°, [α]_D²⁵ = -9 (c=1, EtOH). Reported [α]_D²⁵ = -8.75 (c=1.04, EtOH). The IR and NMR were identical to the data reported earlier. The mother liquor was chromatographed to give the 2-acetoxy compound (**2**) (0.44g), [α]_D²⁵ = -11 (c=0.92, EtOH). This compound was hydrolysed in the presence of acid to give (S)(+)-2-hydroxy-4-phenylbutanoic acid (**1b**) (0.28g), [α]_D²⁵ = +7.5, (c=0.5, EtOH), 84% ee. The (R)(-)-2-hydroxy-4-phenylbutanoic acid was converted to its acetate. The [α]_D²⁵ of this (R)-2-acetoxy-4-phenylbutanoic acid was found to be +11.76 (c=0.51, EtOH). IR(neat) cm⁻¹: 2910, 1740, 1370, 1240. NMR (60 MHz, CDCl₃) δ : 6.95 - 7.15 (aromatic 5H), 4.95 (t, 1H), 2.75 (m, 2H), 2.05 (s, m & 3H, 2H). Chiral chromatography on a Chiracel OD column (Hexane:IPA:TFA, 100:2:0.1, 3 ml/min), showed an ee of >99% for each of the two enantiomers, R(-) and S(+)-2-hydroxy-4-phenylbutanoic acids.

Under similar experimental conditions (R) (-)-2-hydroxy, 4-phenylbutanoic acid (**3a**) was obtained from (**3**) in 42% yield and >99% ee, m.p = 104°, [α]_D²⁵ = -91.5; reported [α]_D²⁵ = -90.6 (c=1.9, MeOH). The (S) (+)-2-hydroxy-4-phenylbutanoic acid (**3b**) (94% ee) [α]_D²⁵ = +85.2 (c=0.55, MeOH) was obtained from the (S) 2-acetoxy, 4-phenylbutanoic acid (**4**) [α]_D²⁵ = +108 (c=0.36, EtOH), m.p. 82°. NMR (60 MHz) CDCl₃(δ): 7.2-7.6 (arom, 5H), 6.9 (d, Ph-CH_a=CH_b, J_{ab} = ~16 Hz), 6.3 (dd, Ph-CH_a=CH_b-CH_cOAc, J_{ab}=16 Hz, J_{bc}=6 Hz), 5.65(d, Ph-CH_a=CH_b-CH_cOAcCOOH, J_{bc}=6 Hz), 2.2 (s, -O-CO-CH₃, 3H). IR (KBr) (cm⁻¹): 3400, 2920, 1690, 1720, 1380, 1280, 1250, 1230, 1100.

Since our final product of interest has an ethyl ester group, we tried the enzymatic resolution of 2-hydroxy-4-phenylbutanoic acid ethyl ester. The lipase PS (Amano) was used in the presence of vinyl acetate and methyl t-butyl ether to give ethyl 2-hydroxy-4-phenylbutanoate [α]_D²⁵ = -7.14 (c=1.12, EtOH) Reported [α]_D²⁵ = 7.8 (c=1, EtOH)¹¹ and ethyl 2-acetoxy-4-phenyl butanoate, [α]_D²⁵ = -15.3 (c=1.5, EtOH). The (R)- ethyl -2-hydroxy-4-phenylbutanoate can thus be used directly for further reactions.

Acknowledgements: The authors thank Dr. P. S. Srinivasan and Mr. R. Venkatesan for making available the 2-oxo-4-phenylbutanoic acid and 2-oxo-4-phenyl-3-butenic acid. Discussions with Dr. N. Kalyanam and constant support from Dr. M. D. Nair are gratefully acknowledged.

References

1. Parker, D.; J.Chem. Soc., Perkin Trans., **1983**, 2, 83.
2. Urbach, H.; Henning, R.; Tetrahedron Lett., **1984**, 25, 1143.
3. Lawton, G.; et al, US patent **1988**, 4, 757, 069.
4. Schmidt, E.; Blaser, H. U.; Fauquex, P. F.; Sedelmeier, G.; Spindler, F.; 'Microbial Reagents in Organic Synthesis', Ed. Servi, S.; Ed. **1992**; 377. Kluwer Academic Publishers, The Netherlands.
5. Wang, Y. -F.; Chen, S. -T.; Kevin, K. C. Liu Wong, C. -H, Tetrahedron Lett., **1989**, 30, 1917.
6. Kalaritis, P.; Regency, R. W.; Partridge, J. J.; Coffen, D. L.; Jour. Org. Chem. **1990**, 55, 812.
7. Sugai, T.; Ohta, H.; Agric. Biol. Chem., **1991**, 55, 293.
8. Yu. H.; Simon, H.; Tetrahedron, **1991**, 47, 9035; Schummer, A.; Yu. H; Simon. H; Tetrahedron, **1991**, 47, 9019.
9. Katchalski-Katzir, E.; Trends in Biotechnology, **1993**, 11, 471.
10. Hsu, S. H.; Wu, S. S.; Wang, Y. F.; Wang, C. H.; Tet.Lett., **1990**, 31, 6403; Maleata, F. X.; Hill Jr., C. G.; Amundson, C. H.; Biotech and Bioeng., **1991**, 38, 853; Bosley, J. A.; Clayton, J.C.. Biotech and Bioeng., **1994**, 43, 934.
11. Atwood, M. R., Hassall, C. H., Krohn, A., Lawton, G., Redshaw, S., Jour. Chem. Soc. Perkin, I., **1986**, 1011.