

0957-4166(95)00213-8

## Synthesis of Optically Active Aporphine and Morphinandienone Alkaloids via *p*-Quinol Esters





Hiroshi Hara,\* Satoshi Komoriya, Takanori Miyashita, and Osamu Hoshino\*

Faculty of Pharmaceutical Sciences, Science University of Tokyo  
12 Ichigaya Funagawara-machi, Shijunju-ku, Tokyo 162, Japan

**Abstract:** Lead tetraacetate oxidation of *N*-trifluoroacetylnorcodamine (**5**) in (*S*)-(+)-2-phenylpropionic acid gave a diastereomeric mixture of two *p*-quinol acylates, which were easily separated to enantiomerically pure **6a** and **6b**. Treatment of the chiral quinol acylates (**6a**) and (**6b**) with trifluoroacetic acid in CH<sub>2</sub>Cl<sub>2</sub> at room temperature afforded (1*R*)-(-)-6-trifluoroacetylwilsonirine (**7a**) and (1*S*)-(+)-(**7b**), respectively. Saponification of **7a** and **7b** gave rise to (-)-wilsonirine (**8a**) and its enantiomer (**8b**), respectively. Similarly, (-)-nordomesticine (**12a**) and (+)-(**12b**) were synthesized in enantiomerically pure form from **10a** and **10b**. On the other hand, acid treatment of **6a** and **6b** in CH<sub>3</sub>CN at lower temperature (-30 °C) followed by *N*-deprotection gave the corresponding normorphinandienones (**15a** and **15b**) as major products, which were transformed to enantiomerically pure (-)-**16a** and (+)-sebiferine (**16b**). In a similar sequence of reactions, (+)-amurine (**19a**) and its enantiomer (**19b**) were synthesized.


A number of aporphine alkaloids have been isolated and characterized.<sup>1</sup> Although there are various enantio- or diastereo-selective synthesis of optically active 1-substituted tetrahydroisoquinolines,<sup>2</sup> few procedures for the synthesis of homochiral aporphines have been reported.<sup>3</sup> We have already developed a facile method<sup>4</sup> for synthesis of racemic aporphine (**3**) via the reactive intermediate, *p*-quinol acetate (**2**), which is readily derived from tetrahydroisoquinolin-7-ol (**1**) bearing an activated benzyl group at the 1-position by lead tetraacetate (LTA) oxidation. We were encouraged to prepare optically active *p*-quinol esters which hold great promise as precursors of tetracyclic alkaloids. We describe here a synthesis of enantiomerically pure aporphines and morphinandienones in a new synthetic manner including LTA oxidation in a chiral acid.

First of all, a readily available chiral acid was sought for formation of chiral *p*-quinol acylates which could be separated into two diastereoisomers. After careful exploration,<sup>5</sup> (*S*)-(+)-2-phenylpropionic acid was found to be adequate for the formation of stable *p*-quinol esters. However, when (±)-codamine (**1**) was submitted to LTA oxidation in the acid, formation of four diastereomers of *p*-quinol esters (**4**), consisting of two pairs of 1,4*a*-*cis* and *trans* isomers, was observed on thin layer chromatography. Separation of the mixture is bothersome and it forced us to choose the *N*-acyl analogue (**5**) as a starting compound, because it has a quasi-axially oriented




arylmethyl group and could be attacked by the acid only from a side opposite to the substituent. Actually, oxidation of ( $\pm$ )-*N*-trifluoroacetyl-1,2,3,4-tetrahydro-7-hydroxy-6-methoxy-1-veratrylisoquinoline (**5**)<sup>6</sup> by LTA (1.2 eq.) in (*S*)-(+)-2-phenylpropionic acid at room temperature for 4.5 h gave a mixture of two diastereomers, which were easily separated by medium pressure liquid chromatography to the *p*-quinol esters (**6a**)<sup>7</sup> [35%, IR (CHCl<sub>3</sub>), cm<sup>-1</sup>: 1735 (ester), 1685, 1650, 1625 (NCOCF<sub>3</sub> and dienone)] and (**6b**)<sup>7</sup> [29%, IR (CHCl<sub>3</sub>), cm<sup>-1</sup>: 1735 (ester), 1685, 1650, 1625 (NCOCF<sub>3</sub> and dienone)]. The absolute configuration of both compounds was determined by transformation of **6b** into the natural alkaloid (**8b**) as described below. Treatment of **6a** with trifluoroacetic acid (TFA) in dichloromethane (CH<sub>2</sub>Cl<sub>2</sub>) afforded an optically active aporphine (**7a**) (64%), the enantiomeric excess (ee) of which was estimated to be 84% by HPLC analysis using a chiral column (DAICEL CHIRALCEL OJ). Enantiomerically pure **7a** was prepared in the following manner. The product was dissolved in hot CH<sub>2</sub>Cl<sub>2</sub> and then after cooling to room temperature the resulting precipitate (mp 188-188.5 °C, 5.7%, 7.1% ee) was removed by filtration. Evaporation of the solvent gave homochiral (*R*)-(-)-6-trifluoroacetylwilsonirine (**7a**) as an amorphous mass, mp 88-89 °C, (48%, 100% ee). Similar treatment of the *p*-quinol acylate (**6b**) provided (*S*)-(+)-6-trifluoroacetylwilsonirine (**7b**) (an amorphous mass, mp 87-89°C, 96.4% ee) in 33.2% yield. Each amide (**7a** or **7b**) was refluxed in 4.3% aqueous K<sub>2</sub>CO<sub>3</sub>-MeOH (1:10) for 5-6 h to give the corresponding (*R*)-(-)-wilsonirine (**8a**) (mp 106-108 °C, 23%, 100% ee) or (*S*)-(+)-wilsonirine (**8b**) (mp 107-109 °C, 39%, 95.9% ee) as colorless crystals, respectively. The physical data of the latter product (**8b**) were consistent with those of natural wilsonirine.<sup>8</sup>

Then this methodology was applied to synthesis of homochiral 9,10-methylenedioxyaporphine, (*S*)-(+)-nordomesticine and its enantiomer. Oxidation of racemic 1-piperonyltetrahydroisoquinolin-7-ol (**9**)<sup>6</sup> afforded two separable *p*-quinol acylates, **10a** (43%) and **10b** (29%). Acid treatment of **10a** and **10b** followed by similar purification as described above gave (*R*)-(-)- (**11a**) (mp 243-244 °C, 32%, 95.3% ee) and (*S*)-(+)-6-trifluoroacetylnordomesticine (**11b**) (mp 244-245 °C, 23%, 94.7% ee), respectively. Alkaline hydrolysis of the trifluoroacetamides (**11a** and **11b**) yielded (*R*)-(-)-nordomesticine (**12a**) (oil, 55%, 100% ee) and (*S*)-(+)- (**12b**) (oil, 70%, 99% ee). <sup>1</sup>H-NMR and high resolution mass spectra of products (**12a** and **12b**) and their diacetates (**13a** and **13b**)<sup>9</sup> supported the structure assignment.




Recently, formation of the morphinandienone skeleton from racemic *N*-trifluoroacetyl-*p*-quinol acetate (**20**) was established in our laboratory.<sup>10</sup> Thus, these reaction conditions were adapted to the reaction of the chiral *p*-quinol acylates in hand. Namely, the reaction of the quinol acylate (**6a**) with TFA in CH<sub>3</sub>CN at -30°C gave an

expected normorphinandienone (**14a**) (35%) together with the noraporphine (**7a**) (16%) and the starting compound (**6a**) (28%). Prolongation of the reaction time did not increase the yield of **14a**.



- 14a:** R=Me, X=COCF<sub>3</sub>  
**15a:** R=Me, X=H  
**16a:** R=X=Me  
**17a:** R+R=CH<sub>2</sub>, X=COCF<sub>3</sub>  
**18a:** R+R=CH<sub>2</sub>, X=H  
**19a:** R+R=CH<sub>2</sub>, X=Me



- 14b:** R=Me, X=COCF<sub>3</sub>  
**15b:** R=Me, X=H  
**16b:** R=X=Me  
**17b:** R+R=CH<sub>2</sub>, X=COCF<sub>3</sub>  
**18b:** R+R=CH<sub>2</sub>, X=H  
**19b:** R+R=CH<sub>2</sub>, X=Me

Similarly, enantiomeric normorphinandienone (**14b**) (31%) and noraporphine (**7b**) (16%) were furnished by treatment of **6b** under the conditions similar to those noted above. IR and <sup>1</sup>H-NMR spectral data of both normorphinandienones (**14a** and **14b**) were completely consistent with those of each authentic racemate.<sup>10</sup> The enantiomeric purity of **14a** and **14b** was estimated after recrystallization to be 92 and 98% ee by HPLC analysis, respectively. Hydrolysis of **14b** with K<sub>2</sub>CO<sub>3</sub> in aqueous MeOH yielded (+)-norsebiferine (**15b**), which was converted by *N*-methylation into (+)-sebiferine (**16b**)<sup>11</sup> (an amorphous mass; 53% from **14b**). The specific rotation ([α]<sub>D</sub><sup>24</sup>) of **16b** (97% ee) was measured to be +12 consistent with the value ([α]<sub>D</sub><sup>20</sup> = +13) for the natural alkaloid.<sup>11</sup> Through the same processes as above, the enantiomer {(-)-**16a**; [α]<sub>D</sub><sup>24</sup> = -13} was synthesized from **14a**. The tactics were applied to the methylenedioxy congeners (**10a** and **10b**). Thus, acid treatment of the *p*-quinol acylate (**10a**) gave an *N*-trifluoroacetylnormorphinandienone (**17a**), which was successively hydrolyzed and *N*-methylated to afford (+)-amurine (**19a**) (mp 203-205 °C, 99% ee, [α]<sub>D</sub><sup>27</sup> = +10.7); (lit.<sup>12</sup> [α]<sub>D</sub><sup>27</sup> = +11). Its enantiomer {(-)-**19b**, mp 200-203 °C, [α]<sub>D</sub><sup>27</sup> = -13.5} was also prepared from **10b** through the same sequence of reactions. <sup>1</sup>H-NMR spectra of both normorphinandienones (**19a** and **19b**) were superimposable on each other.

Although the present methodology serves effectively for synthesis of optically active aporphine and morphinandienone, partial racemization can not be prevented because of the following possibility; (1) acid treatment of some kinds of 1-alkyltetrahydroisoquinolines<sup>13</sup> causes racemization at the 1-position and (2) a proton at the 1-position in *N*-acyl-*p*-quinol acylates (**6** and **10**) seems to be enolizable because of the presence of the conjugated enone, resulting in a little bit of racemization.

In conclusion, we have developed a novel methodology *via* optically active *p*-quinol esters for the synthesis of homochiral aporphine and morphinandienone alkaloids. (*S*)-(+)-2-Phenylpropionic acid employed in the reaction was completely recovered without loss of enantiomeric purity. Further application of the procedure to asymmetric synthesis of 1-phenyl- and 1-phenethyl-tetrahydroisoquinoline alkaloids is now in progress.

## EXPERIMENTAL

All melting points were measured on a Büchi melting point apparatus and are uncorrected. IR spectra were taken with a Hitachi model 260-10 spectrometer in  $\text{CHCl}_3$  solution.  $^1\text{H-NMR}$  spectra were recorded on a JEOL model FX-100 spectrometer in  $\text{CDCl}_3$  solution using tetramethylsilane as internal standard. Mass spectra (MS) were obtained on a Hitachi model M-60 spectrometer. Optical rotations were measured on a JASCO DIP-360 polarimeter. *Ee* values were estimated by HPLC analysis with chiral column (DAICEL CHIRALCEL OC, OD, or OJ). Preparative thin layer chromatography (TLC) was performed on precoated silica gel plates (Merck Kieselgel 60 F254).

Preparation of *p*-Quinol Acylates (6 and 10)

Oxidation was carried out according to a procedure similar to that previously reported except using (*S*)-(+)-2-phenylpropionic acid instead of acetic acid.<sup>4</sup> Usual work-up of the reaction mixture gave yellowish brown oil, which was purified by medium pressure liquid chromatography on silica gel.

From ( $\pm$ )-5 : A mixture of ( $\pm$ )-5 (200 mg, 0.47 mmol) and LTA (250 mg, 0.56 mmol) in (*S*)-(+)-2-phenylpropionic acid (3 ml) was stirred for 4 h. Elution of the residue (317 mg) with hexane-AcOEt (55 : 45) gave two *p*-quinol acylates (**6a** and **6b**, polarity; **6a** < **6b**). (IR. 4*aR*, 2'*S*)-1,2,3,4,4*a*,7-Hexahydro-7-oxo-4*a*-(2'-phenylpropionyloxy)-2-trifluoroacetyl-1-veratrylisoquinoline (**6a**) (oil, 94 mg, 35%);  $^1\text{H-NMR}$   $\delta$ : 1.50 [3H, d, *J* = 7 Hz,  $\text{OCOCH}(\text{CH}_3)\text{Ph}$ ], 3.64, 3.83, 3.84 (each 3H, s,  $\text{OCH}_3$ ); IR  $\text{cm}^{-1}$ : 1735 (ester), 1685, 1650, 1625 (dienone and  $\text{NCOCF}_3$ ). (1*S*, 4*aS*, 2'*S*)-1,2,3,4,4*a*,7-Hexahydro-7-oxo-4*a*-(2'-phenylpropionyloxy)-2-trifluoroacetyl-1-veratrylisoquinoline (**6b**) (oil, 78 mg, 29%);  $^1\text{H-NMR}$   $\delta$ : 1.53 [3H, d, *J* = 7 Hz,  $\text{OCOCH}(\text{CH}_3)\text{Ph}$ ], 3.64, 3.76, 3.82 (each 3H, s,  $\text{OCH}_3$ ); IR  $\text{cm}^{-1}$ : 1735 (ester), 1685, 1650, 1625 (dienone and  $\text{NCOCF}_3$ ).

From ( $\pm$ )-9 : A mixture of ( $\pm$ )-9 (600 mg, 1.47 mmol) and LTA (840 mg, 1.88 mmol) in (*S*)-(+)-2-phenylpropionic acid (9 ml) was stirred for 4.5 h. Elution of the residue (938 mg) with hexane-AcOEt (65 : 35) gave two *p*-quinol acylates (**10a** and **10b**, polarity; **10a** < **10b**). (IR. 4*aR*, 2'*S*)-1,2,3,4,4*a*,7-Hexahydro-7-oxo-4*a*-(2'-phenylpropionyloxy)-2-trifluoroacetyl-1-piperonylisoquinoline (**10a**) (oil, 346 mg, 42%);  $^1\text{H-NMR}$   $\delta$ : 1.50 [3H, d, *J* = 7.1 Hz,  $\text{OCOCH}(\text{CH}_3)\text{Ph}$ ], 3.66 (3H, s,  $\text{OCH}_3$ ), 5.91 (2H, s,  $\text{OCH}_2\text{O}$ ); IR  $\text{cm}^{-1}$ : 1730 (ester), 1665, 1650, 1625 (dienone and  $\text{NCOCF}_3$ ). High resolution MS (*m/z*) Calcd for  $\text{C}_{29}\text{H}_{26}\text{F}_3\text{NO}_7$ ; 557.1660. Found; 557.1690. (1*S*, 4*aS*, 2'*S*)-1,2,3,4,4*a*,7-Hexahydro-7-oxo-4*a*-(2'-phenylpropionyloxy)-2-trifluoroacetyl-1-piperonylisoquinoline (**10b**) (oil, 233 mg, 29%);  $^1\text{H-NMR}$   $\delta$ : 1.53, [3H, d, *J* = 7.1 Hz,  $\text{OCOCH}(\text{CH}_3)\text{Ph}$ ], 3.90 (3H, s,  $\text{OCH}_3$ ), 5.90 (2H, s,  $\text{OCHO}$ ); IR  $\text{cm}^{-1}$ : 1740 (ester), 1690, 1655, 1630 (dienone and  $\text{NCOCF}_3$ ). High resolution MS (*m/z*) Calcd for  $\text{C}_{29}\text{H}_{26}\text{F}_3\text{NO}_7$ ; 557.1658. Found; 557.1658.

Treatment of *p*-Quinol Acylates (6 and 10) with TFA in  $\text{CH}_2\text{Cl}_2$  or  $\text{CH}_3\text{CN}$ 

Acid treatment of *p*-quinol acylates (**6** and **10**) was carried out according to the procedures similar to those (in  $\text{CH}_2\text{Cl}_2$ <sup>4</sup> and  $\text{CH}_3\text{CN}$ <sup>10</sup>) previously reported. Usual work-up of the reaction mixture gave an oily residue, which was separated by preparative TLC.

a) In  $\text{CH}_2\text{Cl}_2$ . From **6a** : A mixture of **6a** (280 mg, 0.49 mmol) and TFA (1.5 ml) in  $\text{CH}_2\text{Cl}_2$  (30 ml) was stirred for 3.5 h. Preparative TLC of the yellowish brown oil (209 mg) with  $\text{CHCl}_3$  afforded (*R*)-(-)-6-trifluoroacetylwilsonirine (**7a**) as an amorphous mass (132 mg, 64%, 84% *ee*). The solid was dissolved in boiling  $\text{CH}_2\text{Cl}_2$  and colorless crystals (11 mg, 5.7%, 7.1% *ee*, mp 188-188.5 °C) were precipitated from the

cooled solution. Evaporation of filtrate *in vacuo* gave optically pure (-)-**7a** [100 mg, 48%, 100% ee (OJ; *i*-PrOH : hexane = 9 : 1)] as an amorphous mass; mp 87-89 °C;  $[\alpha]_D^{24}$  -234.3 ( $c = 1.0$ , CHCl<sub>3</sub>); <sup>1</sup>H-NMR δ: 3.89 (3H, s, OCH<sub>3</sub>), 3.93 (6H, s, 2xOCH<sub>3</sub>), 6.55 (1H, s, 3-H), 6.75 (1H, s, 8-H), 8.09 (1H, s, 11-H); IR cm<sup>-1</sup>: 1685 (NCOCF<sub>3</sub>). High resolution MS (*m/z*) Calcd for C<sub>21</sub>H<sub>20</sub>F<sub>3</sub>NO<sub>5</sub>: 423.1291. Found: 423.1284.

From **6b** : A mixture of **6b** (200 mg, 0.35 mmol) and TFA (1 ml) in CH<sub>2</sub>Cl<sub>2</sub> (20 ml) was stirred for 4.5 h. Preparative TLC of the yellowish brown oil (150 mg) with CHCl<sub>3</sub> afforded (*S*)-(+)-6-trifluoroacetylwilsonirine (**7b**) as an amorphous mass (112 mg, 77%, 85% ee). A manner similar to that noted for **7a** gave colorless crystals (3.2 mg, 2.2%, 3.5% ee, mp 190-191 °C) and (+)-**7b** [49 mg, 33%, 96.4% ee (OJ; *i*-PrOH : hexane = 9 : 1)] as an amorphous mass; mp 87-89 °C;  $[\alpha]_D^{24}$  +253.1 ( $c = 0.7$ , CHCl<sub>3</sub>); <sup>1</sup>H-NMR δ: 3.89 (3H, s, OCH<sub>3</sub>) 3.93 (6H, s, 2xOCH<sub>3</sub>), 6.55 (1H, s, 3-H), 6.75 (1H, s, 8-H), 8.09 (1H, s, 11-H); IR cm<sup>-1</sup>: 1685 (NCOCF<sub>3</sub>). High resolution MS (*m/z*) Calcd for C<sub>21</sub>H<sub>20</sub>F<sub>3</sub>NO<sub>5</sub>: 423.1292. Found: 423.1280.

From **10a** : A mixture of **10a** (346 mg, 0.62 mmol) and TFA (1.7 ml) in CH<sub>2</sub>Cl<sub>2</sub> (35 ml) was stirred for 4 h. Preparative TLC of the yellowish brown oil (256 mg) with CHCl<sub>3</sub> gave (*R*)-(-)-6-trifluoroacetylnordomesticine (**11a**) as an amorphous mass (103 mg, 41%, 94.7% ee). A manner similar to that noted for **7a** gave colorless crystals (17 mg, 6.7%, 79.6% ee, mp 256-257 °C) and (-)-**11a** [82 mg, 32%, 95.3% ee (OD; *i*-PrOH : hexane = 9 : 1)] as an amorphous mass; mp 243-244 °C;  $[\alpha]_D^{21}$  -259.4 ( $c = 0.1$ , CHCl<sub>3</sub>); <sup>1</sup>H-NMR δ: 3.91 (3H, s, OCH<sub>3</sub>) 5.96 (2H, s, OCH<sub>2</sub>O), 6.55 (1H, s, 3-H), 6.73 (1H, s, 8-H), 7.99 (1H, s, 11-H); IR cm<sup>-1</sup>: 1685 (NCOCF<sub>3</sub>). High resolution MS (*m/z*) Calcd for C<sub>20</sub>H<sub>16</sub>F<sub>3</sub>NO<sub>5</sub>: 407.0978. Found: 407.0972.

From **10b**: A mixture of **10b** (233 mg, 0.57 mmol) and TFA (1.1 ml) in CH<sub>2</sub>Cl<sub>2</sub> (23 ml) was stirred for 4.5 h. Preparative TLC of the yellowish brown oil (171 mg) with CHCl<sub>3</sub> gave (*S*)-(+)-6-trifluoroacetylnordomesticine (**11b**) as an amorphous mass (79 mg, 47%, 93.1% ee). A manner similar to that noted for **7a** gave colorless crystals (36 mg, 21%, 92.5% ee, mp 244-245 °C) and (+)-**11b** [39 mg, 23%, 94.7% ee<sup>14</sup> (OD; *i*-PrOH : hexane = 9 : 1)] as an amorphous mass; mp 244-245 °C;  $[\alpha]_D^{21}$  +262.3 ( $c = 0.1$ , CHCl<sub>3</sub>); <sup>1</sup>H-NMR δ: 3.92 (3H, s, OCH<sub>3</sub>), 5.96 (2H, s, OCH<sub>2</sub>O), 6.55 (1H, s, 3-H), 6.73 (1H, s, 8-H), 7.99 (1H, s, 11-H); IR cm<sup>-1</sup>: 1685 (NCOCF<sub>3</sub>). High resolution MS (*m/z*) Calcd for C<sub>20</sub>H<sub>16</sub>F<sub>3</sub>NO<sub>5</sub>: 407.0979. Found: 407.0995.

b) In CH<sub>3</sub>CN. From **6a** : A mixture of **6a** (316 mg, 0.55 mmol) and TFA (1.5 ml) in CH<sub>3</sub>CN (30 ml) was stirred at -30 °C for 7.5 h. Preparative TLC of the yellowish brown oil (277 mg) with CHCl<sub>3</sub>-acetone (35 : 1) gave aporphine (**7a**) (37 mg (16%), an amorphous mass, mp 162-168 °C,  $[\alpha]_D^{26}$  -213.3 ( $c = 0.3$ , CHCl<sub>3</sub>) (84% ee)), the starting *p*-quinol acylate (**6a**) (89 mg, 28%), and (*9R*)-(-)-*N*-trifluoroacetylnorsebiferine (**14a**) (81 mg, 35%) (polarity ; **7a** < **6a** < **14a**). The former two compounds were spectroscopically identified against authentic specimens. Crystallization of the latter product (**14a**) from ether afforded colorless crystals, mp 202-203.5 °C.  $[\alpha]_D^{25}$  -59.1 ( $c = 0.4$ , CHCl<sub>3</sub>) [92% ee (OC; EtOH : hexane = 7 : 1)]; <sup>1</sup>H-NMR δ: 3.80, 3.85, 3.89 (each 3H, s, OCH<sub>3</sub>), 6.32 (1H, s, 5-H), 6.34, 6.39 (1H, each s, 8-H), 6.60 (1H, s, 1-H), 6.82 (1H, s, 4-H); IR cm<sup>-1</sup>: 1690, 1680, 1655, 1628 (NCOCF<sub>3</sub> and dienone). High resolution MS (*m/z*) Calcd for C<sub>21</sub>H<sub>20</sub>F<sub>3</sub>NO<sub>5</sub>: 423.1292. Found: 423.1289.

From **6b** : A mixture of **6b** (175 mg, 0.3 mmol) and TFA (0.85 ml) in CH<sub>3</sub>CN (19 ml) was stirred at -30 °C for 7.5 h. Preparative TLC of the yellowish brown oil (141 mg) with CHCl<sub>3</sub>-acetone (35 : 1) gave noraporphine (**7b**) [21 mg (16%), an amorphous mass, mp 156-159 °C,  $[\alpha]_D^{25}$  +214 ( $c = 0.3$ , CHCl<sub>3</sub>) (86.6% ee)], the starting *p*-quinol acylate (**6b**) (57 mg, 31%), and (*9S*)-(+)-*N*-trifluoroacetylnorsebiferine (**14b**) (41 mg, 31%) (polarity ; **7b** < **6b** < **14b**). The former two compounds were spectroscopically identified with authentic specimens, respectively. Crystallization of the latter product (**14b**) from ether afforded colorless crystals, mp

203-204 °C.  $[\alpha]_D^{25} +60.7$  ( $c = 0.4$ ,  $\text{CHCl}_3$ ) [98.1% ee (OC; EtOH : hexane = 7 : 1)];  $^1\text{H-NMR}$   $\delta$ : 3.80, 3.85, 3.89 (each 3H, s,  $\text{OCH}_3$ ), 6.32 (1H, s, 5-H), 6.34, 6.39 (1H, each s, 8-H), 6.60 (1H, s, 1-H), 6.82 (1H, s, 4-H); IR  $\text{cm}^{-1}$ : 1690, 1680, 1655, 1628 ( $\text{NCOCF}_3$  and dienone). High resolution MS ( $m/z$ ) Calcd for  $\text{C}_{21}\text{H}_{20}\text{F}_3\text{NO}_5$ : 423.1292. Found: 423.1295.

From **10a**: A mixture of **10a** (302 mg, 0.54 mmol) and TFA (1.5 ml) in  $\text{CH}_3\text{CN}$  (30 ml) was stirred at -30 °C for 4 h. Preparative TLC of the yellowish brown oil (268 mg) with hexane-AcOEt (1 : 1) gave the noraporphine (**11a**) {13 mg (6%), an amorphous mass, mp 234-237 °C,  $[\alpha]_D^{26} -211.4$  ( $c = 0.3$ ,  $\text{CHCl}_3$ ) (93.5% ee)}, the starting *p*-quinol acylate (**10a**) (109 mg, 36%), and (9*R*)-(-)-*N*-trifluoroacetylnoramurine (**17a**) (59 mg, 27%) (porality ; **11a** < **10a** < **17a**). The former two compounds were spectroscopically identified by comparison with authentic specimens. Crystallization of the latter product (**17a**) from ether afforded a small amount of colorless crystals (0.7 mg, mp 206-207 °C, 9.6% ee). Evaporation of the filtrate under reduced pressure gave an amorphous mass (**17a**) {42 mg (19%), mp 102-105 °C;  $[\alpha]_D^{27} -31.6$  ( $c = 0.4$ ,  $\text{CHCl}_3$ ) [94.9% ee (OJ; EtOH : hexane = 1 : 1)];  $^1\text{H-NMR}$   $\delta$ : 3.79 (3H, s,  $\text{OCH}_3$ ), 5.93 (2H, AB type,  $\text{OCH}_2\text{O}$ ), 6.28 (1H, s, 5-H), 6.34, 6.37 (1H, each s, 8-H), 6.58 (1H, s, 1-H), 6.86 (1H, s, 4-H); IR  $\text{cm}^{-1}$ : 1690, 1680, 1660, 1630 ( $\text{NCOCF}_3$  and dienone). High resolution MS ( $m/z$ ) Calcd for  $\text{C}_{20}\text{H}_{16}\text{F}_3\text{NO}_5$ : 407.0979. Found: 407.0976.

From **10b**: A mixture of **10b** (226 mg, 0.4 mmol) and TFA (1 ml) in  $\text{CH}_3\text{CN}$  (20 ml) was stirred at -30 °C for 5 h. Preparative TLC of the yellowish brown oil (190 mg) with hexane-AcOEt (1 : 1) gave aporphine (**11b**) {28 mg (17%), an amorphous mass, mp 198-206 °C,  $[\alpha]_D^{26} +107.4$  ( $c = 0.2$ ,  $\text{CHCl}_3$ ) (83.1% ee)}, the starting *p*-quinol acylate (**10b**) (104 mg, 46%), and (9*S*)-(+)-*N*-trifluoroacetylnoramurine (**17b**) (36 mg, 22%) (porality ; **11b** < **10b** < **17b**). The former two compounds were spectroscopically identified by comparison with authentic specimens. Crystallization of the latter product (**17b**) from ether afforded a small amount of colorless crystals (1.4 mg, mp 205-206 °C, 6.6% ee). Evaporation of the filtrate under reduced pressure gave an amorphous mass (**17b**) {25 mg (15%), mp 107-117 °C;  $[\alpha]_D^{27} +37.1$  ( $c = 0.5$ ,  $\text{CHCl}_3$ ) [95.2% ee (OJ; EtOH : hexane = 1 : 1)];  $^1\text{H-NMR}$   $\delta$ : 3.79 (3H, s,  $\text{OCH}_3$ ), 5.93 (2H, AB type,  $\text{OCH}_2\text{O}$ ), 6.28 (1H, s, 5-H), 6.34, 6.37 (1H, each s, 8-H), 6.58 (1H, s, 1-H), 6.86 (1H, s, 4-H). IR  $\text{cm}^{-1}$ : 1690, 1680, 1660, 1630 ( $\text{NCOCF}_3$  and dienone). High resolution MS ( $m/z$ ) Calcd for  $\text{C}_{20}\text{H}_{16}\text{F}_3\text{NO}_5$ : 407.0978. Found: 407.0971.

### General Procedure for Hydrolysis of **7**, **12**, **14**, and **17**

A solution of **7**, **12**, **14**, and **17** and 4.3 % aqueous  $\text{K}_2\text{CO}_3$  in MeOH was refluxed. Usual work-up of the reaction mixture gave an oily residue, which was purified by crystallization or preparative TLC.

(*R*)-(-)-Wilsonirine (**8a**): A mixture of (-)-**7a** (108 mg, 0.25 mmol), aqueous  $\text{K}_2\text{CO}_3$  (2 ml) and MeOH (20 ml) was refluxed for 5 h. A yellowish brown oil (83 mg), which was crystallized from acetone to give (-)-**8a** [19 mg, 23%, 100% ee (OD; *i*-PrOH : hexane = 9 : 1)] as colorless crystals; mp 106-108 °C,  $[\alpha]_D^{24} -53.9$  ( $c = 0.2$ , MeOH);  $^1\text{H-NMR}$   $\delta$ : 3.89 (9H, s,  $3\times\text{OCH}_3$ ), 6.53 (1H, s, 3-H), 6.72 (1H, s, 8-H), 8.03 (1H, s, 11-H). High resolution MS ( $m/z$ ) Calcd for  $\text{C}_{19}\text{H}_{21}\text{NO}_4$ : 327.1469. Found: 327.1465.

(*S*)-(+)-Wilsonirine (**8b**): A mixture of (+)-**7b** (76 mg, 0.18 mmol), aqueous  $\text{K}_2\text{CO}_3$  (1.5 ml) and MeOH (15 ml) was refluxed for 6 h. A yellowish brown oil (63 mg), which was crystallized from acetone to give (+)-**8b** [23 mg, 39%, 95.9% ee (OD; *i*-PrOH : hexane = 9 : 1)] as colorless crystals; mp 107-109 °C,  $[\alpha]_D^{24} +54.8$  ( $c = 0.1$ , MeOH) (lit.<sup>8</sup> mp 108-110 °C,  $[\alpha]_D^{24} +47$ );  $^1\text{H-NMR}$   $\delta$ : 3.89 (9H, s,  $3\times\text{OCH}_3$ ), 6.53 (1H, s, 3-H), 6.72 (1H, s, 8-H), 8.03 (1H, s, 11-H). High resolution MS ( $m/z$ ) Calcd for  $\text{C}_{19}\text{H}_{21}\text{NO}_4$ : 327.1468. Found: 327.1456.

**(R)-(-)-Nordomesticine (12a)** : A mixture of (-)-**11a** (46 mg, 0.15 mmol), aqueous K<sub>2</sub>CO<sub>3</sub> (1.5 ml) and MeOH (15 ml) was refluxed for 5 h. Preparative TLC of the yellowish brown oil (48 mg) with CHCl<sub>3</sub>-MeOH (5 : 1) gave (-)-**12a** [20 mg, 55%, 100% ee (OJ; *i*-PrOH : hexane = 9 : 1)] as an oil; <sup>1</sup>H-NMR δ 3.89 (3H, s, OCH<sub>3</sub>), 5.93 (2H, s, OCH<sub>2</sub>O), 6.53 (1H, s, 3-H), 6.70 (1H, s, 8-H), 7.94 (1H, s, 11-H). High resolution MS (*m/z*) Calcd for C<sub>18</sub>H<sub>17</sub>NO<sub>4</sub>: 311.1156. Found: 311.1162. **(R)-(-)-N,O-Diacetylnordomesticine (13a)**, mp 235-238 °C. [α]<sub>D</sub><sup>22</sup> -319.3 (c = 0.4, CHCl<sub>3</sub>) [97% ee (OD; EtOH : hexane = 1 : 5)]; <sup>1</sup>H-NMR δ: 2.20, 2.33 (each 3H, s, OCOCH<sub>3</sub> and NCOCH<sub>3</sub>), 3.83 (3H, s, OCH<sub>3</sub>), 5.95 (2H, s, OCH<sub>2</sub>O), 6.64 (1H, s, 3-H), 6.73 (1H, s, 8-H), 7.36 (1H, s, 11-H). IR cm<sup>-1</sup>: 1770 (OCOCH<sub>3</sub>), 1635 (NCOCH<sub>3</sub>). High resolution MS (*m/z*) Calcd for C<sub>22</sub>H<sub>21</sub>NO<sub>6</sub>: 395.1367. Found: 395.1374.

**(S)-(+)-Nordomesticine (12b)** : A mixture of (+)-**11b** (37 mg, 0.092 mmol), aqueous K<sub>2</sub>CO<sub>3</sub> (1.3 ml) and MeOH (12 ml) was refluxed for 5.5 h. Preparative TLC of the yellowish brown oil (34 mg) with CHCl<sub>3</sub>-MeOH (5 : 1) gave (+)-**12b** [20 mg, 70%, 99% ee (OJ; *i*-PrOH : hexane = 9 : 1)] as an oil; <sup>1</sup>H-NMR δ: 3.89 (3H, s, OCH<sub>3</sub>), 5.93 (2H, s, OCH<sub>2</sub>O), 6.53 (1H, s, 3-H), 6.70 (1H, s, 8-H), 7.94 (1H, s, 11-H). High resolution MS (*m/z*) Calcd for C<sub>18</sub>H<sub>17</sub>NO<sub>4</sub>: 311.1155. Found: 311.1155. **(S)-(+)-N,O-Diacetylnordomesticine (13b)**, mp 235-236 °C (EtOH) (colorless crystals); [α]<sub>D</sub><sup>22</sup> +326.4 (c = 0.4, CHCl<sub>3</sub>) [99.4% ee (OD; EtOH : hexane = 1 : 5)]; <sup>1</sup>H-NMR δ: 2.20, 2.33 (each 3H, s, OCOCH<sub>3</sub> and NCOCH<sub>3</sub>), 3.83 (3H, s, OCH<sub>3</sub>), 5.95 (2H, s, OCH<sub>2</sub>O), 6.64 (1H, s, 3-H), 6.73 (1H, s, 8-H), 7.36 (1H, s, 11-H); IR cm<sup>-1</sup>: 1770 (OCOCH<sub>3</sub>), 1635 (NCOCH<sub>3</sub>). Anal. Calcd for C<sub>22</sub>H<sub>21</sub>NO<sub>6</sub>: C, 66.83; H, 5.35; N, 3.54. Found: C, 66.83; H, 5.61; N, 3.62.

**(9R)-(+)-Norsebiferine (15a)** : A mixture of (-)-**14a** (21 mg, 0.05 mmol), aqueous K<sub>2</sub>CO<sub>3</sub> (1.2 ml) and MeOH (6 ml) was refluxed for 5 h. Preparative TLC of the yellow oil (19 mg) with CHCl<sub>3</sub>-MeOH-NH<sub>4</sub>OH (100 : 5 : 0.1) gave (+)-**15a** [17 mg, 100%, 96.9% ee (OC; EtOH : hexane = 1 : 1)] as an oil; [α]<sub>D</sub><sup>25</sup> +16.4 (c = 0.3, CHCl<sub>3</sub>); <sup>1</sup>H-NMR δ: 3.80, 3.85, 3.87 (each 3H, s, OCH<sub>3</sub>), 6.27 (1H, s, 5-H), 6.32 (1H, s, 8-H), 6.61 (1H, s, 1-H), 6.78 (1H, s, 4-H); IR cm<sup>-1</sup>: 1670, 1645, 1625 (dienone). High resolution MS (*m/z*) Calcd for C<sub>19</sub>H<sub>21</sub>NO<sub>4</sub>: 327.1469. Found: 327.1466.

**(9S)-(-)-Norsebiferine (15b)** : A mixture of (+)-**14b** (20 mg, 0.047 mmol), aqueous K<sub>2</sub>CO<sub>3</sub> (1.2 ml) and MeOH (6 ml) was refluxed for 5 h. Preparative TLC of the yellow oil (20 mg) with CHCl<sub>3</sub>-MeOH-NH<sub>4</sub>OH (100 : 5 : 0.1) gave (-)-**15b** [17 mg, 100%, 99% ee (OC; EtOH : hexane = 1 : 1)] as an oil; [α]<sub>D</sub><sup>25</sup> -14.5 (c = 0.3, CHCl<sub>3</sub>); <sup>1</sup>H-NMR δ: 3.80, 3.85, 3.87 (each 3H, s, OCH<sub>3</sub>), 6.27 (1H, s, 5-H), 6.32 (1H, s, 8-H), 6.61 (1H, s, 1-H), 6.78 (1H, s, 4-H); IR cm<sup>-1</sup>: 1670, 1645, 1625 (dienone). High resolution MS (*m/z*) Calcd for C<sub>19</sub>H<sub>21</sub>NO<sub>4</sub>: 327.1469. Found: 327.1454.

**(9R)-(+)-Noramurine (18a)** : A mixture of (-)-**17a** (46 mg, 0.11 mmol), aqueous K<sub>2</sub>CO<sub>3</sub> (2.6 ml) and MeOH (13 ml) was refluxed for 2.5 h. Preparative TLC of the yellow oil (35 mg) with CHCl<sub>3</sub>-MeOH-NH<sub>4</sub>OH (100 : 5 : 0.1) gave (+)-**18a** [30 mg, 85%, 93.6% ee (OJ; EtOH : hexane = 9 : 1)] as an oil; [α]<sub>D</sub><sup>25</sup> +29.6 (c = 0.3, CHCl<sub>3</sub>); <sup>1</sup>H-NMR δ: 3.79 (3H, s, OCH<sub>3</sub>), 5.91 (2H, AB type, OCH<sub>2</sub>O), 6.26 (2H, s, olefinic H), 6.59 (1H, s, 1-H), 6.80 (1H, s, 4-H); IR cm<sup>-1</sup>: 1670, 1645, 1625 (dienone). High resolution MS (*m/z*) Calcd for C<sub>18</sub>H<sub>17</sub>NO<sub>4</sub>: 311.1156. Found: 311.1146.

**(9S)-(-)-Noramurine (18b)** : A mixture of (+)-**17b** (28 mg, 0.069 mmol), aqueous K<sub>2</sub>CO<sub>3</sub> (1.6 ml) and MeOH (8 ml) was refluxed for 4.5 h. Preparative TLC of the yellow oil (22 mg) with CHCl<sub>3</sub>-MeOH-NH<sub>4</sub>OH (100 : 5 : 0.1) gave (-)-**18b** [14 mg, 65%, 95.8% ee (OJ; EtOH : hexane = 9 : 1)] as an oil; [α]<sub>D</sub><sup>25</sup> -33.3 (c = 0.3, CHCl<sub>3</sub>); <sup>1</sup>H-NMR δ: 3.79 (3H, s, OCH<sub>3</sub>), 5.91 (2H, AB type, OCH<sub>2</sub>O), 6.26 (2H, s, olefinic H), 6.59



(1H, s, 1-H), 6.80 (1H, s, 4-H); IR  $\text{cm}^{-1}$ : 1670, 1645, 1625 (dienone). High resolution MS ( $m/z$ ) Calcd for  $\text{C}_{18}\text{H}_{17}\text{NO}_4$ : 311.1157. Found: 311.1159.

**General Procedure for Synthesis of (9R)-(-)- and (9S)-(+)-Sebiferine (16a and 16b) and (9R)-(+)- and (9S)-(-)-Amurine (19a and 19b)**

A mixture of the amines (**15** and **18**),  $\text{HCO}_2\text{H}$  and formalin was heated on a boiling water bath. The reaction mixture was concentrated under reduced pressure. Then the residue was basified with saturated aqueous  $\text{NaHCO}_3$  and the product was taken up in  $\text{CHCl}_3$ . Usual work-up of the  $\text{CHCl}_3$  extract gave an oily residue, which was purified by preparative TLC with  $\text{CHCl}_3$ -MeOH (10 : 1).

**(9R)-(-)-Sebiferine (16a)** : A mixture of the amine (+)-(**15a**) (15 mg, 0.045 mmol), 85%  $\text{HCO}_2\text{H}$  (5.2 ml), and 35% formalin (2.6 ml) was heated for 10 h. Preparative TLC of the yellowish brown oil (17 mg) gave (-)-**16a** [6.7 mg, 44%, 97.1% ee (OJ; EtOH : hexane = 1 : 3)] as an amorphous mass;  $[\alpha]_{\text{D}}^{24} -13$  ( $c = 0.1$ ,  $\text{CHCl}_3$ );  $^1\text{H-NMR}$   $\delta$ : 2.49 (3H, s,  $\text{NCH}_3$ ), 3.80, 3.85, 3.87 (each 3H, s,  $\text{OCH}_3$ ), 6.31 (1H, s, 5-H), 6.33 (1H, s, 8-H), 6.61 (1H, s, 1-H), 6.79 (1H, s, 4-H); IR  $\text{cm}^{-1}$ : 1675, 1650, 1630 (dienone). High resolution MS ( $m/z$ ) Calcd for  $\text{C}_{20}\text{H}_{23}\text{NO}_4$ : 341.1624. Found: 341.1622.

**(9S)-(+)-Sebiferine (16b)** : A mixture of the amine (-)-(**15b**) (16 mg, 0.048 mmol), 85%  $\text{HCO}_2\text{H}$  (5.6 ml), and 35% formalin (2.8 ml) was heated for 10 h. Preparative TLC of the yellowish brown oil (19 mg) gave (+)-**16b** [8.6 mg, 53%, 97% ee (OJ; EtOH : hexane = 1 : 3)] as an amorphous mass;  $[\alpha]_{\text{D}}^{24} +12$  ( $c = 0.2$ ,  $\text{CHCl}_3$ ) [lit.<sup>11</sup>  $[\alpha]_{\text{D}}^{20} +13$  ( $c = 0.3$ ,  $\text{CHCl}_3$ )];  $^1\text{H-NMR}$   $\delta$ : 2.49 (3H, s,  $\text{NCH}_3$ ), 3.80, 3.85, 3.87 (each 3H, s,  $\text{OCH}_3$ ), 6.31 (1H, s, 5-H), 6.33 (1H, s, 8-H), 6.61 (1H, s, 1-H), 6.79 (1H, s, 4-H); IR  $\text{cm}^{-1}$ : 1675, 1650, 1630 (dienone). High resolution MS ( $m/z$ ) Calcd for  $\text{C}_{20}\text{H}_{23}\text{NO}_4$ : 341.1623. Found: 341.1634.

**(9R)-(+)-Amurine (19a)**: A mixture of the amine (+)-(**18a**) (28 mg, 0.089 mmol), 98%  $\text{HCO}_2\text{H}$  (10 ml), and 35% formalin (5 ml) was heated for 6.5 h. Preparative TLC of the yellowish brown oil (25 mg) gave (+)-**19a** [7.3 mg, 25%, 99% ee (OJ; EtOH : hexane = 9 : 1)] as colorless crystals; mp 205-206 °C (lit.<sup>12</sup> 213-215 °C),  $[\alpha]_{\text{D}}^{27} +10.7$  ( $c = 0.2$ ,  $\text{CHCl}_3$ ) [lit.<sup>12</sup>  $[\alpha]_{\text{D}}^{20} +11$  ( $c = 0.2$ ,  $\text{CHCl}_3$ )];  $^1\text{H-NMR}$   $\delta$ : 2.47 (3H, s,  $\text{NCH}_3$ ), 3.78 (3H, s,  $\text{OCH}_3$ ), 5.91 (2H, AB type,  $\text{OCH}_2\text{O}$ ), 6.26, 6.31 (each 1H, s, 5- and 8-H), 6.59 (1H, s, 1-H), 6.80 (1H, s, 4-H). High resolution MS ( $m/z$ ) Calcd for  $\text{C}_{19}\text{H}_{19}\text{NO}_4$ : 325.1313. Found: 325.1314.

**(9S)-(-)-Amurine (19b)**: A mixture of the amine (-)-(**18b**) (14 mg, 0.045 mmol), 98%  $\text{HCO}_2\text{H}$  (5 ml), and 35% formalin (2.5 ml) was heated for 7.5 h. Preparative TLC of the yellowish brown oil (15 mg) gave **19b** (7.8 mg, 53%) as an amorphous mass. Crystallization from  $\text{CHCl}_3$ -ether gave (-)-**19b** [1.5 mg, 10%, 99% ee (OJ; EtOH : hexane = 9 : 1)] as colorless crystals; mp 204-205 °C,  $[\alpha]_{\text{D}}^{27} -13.5$  ( $c = 0.1$ ,  $\text{CHCl}_3$ );  $^1\text{H-NMR}$   $\delta$ : 2.50 (3H, s,  $\text{NCH}_3$ ), 3.79 (3H, s,  $\text{OCH}_3$ ), 5.91 (2H, AB type,  $\text{OCH}_2\text{O}$ ), 6.26, 6.31 (each 1H, s, 5- and 8-H), 6.59 (1H, s, 1-H), 6.80 (1H, s, 4-H). High resolution MS ( $m/z$ ) Calcd for  $\text{C}_{19}\text{H}_{19}\text{NO}_4$ : 325.1312. Found: 325.1303.

## REFERENCES

1. a) M. Shamma, *The Isoquinoline Alkaloids. Chemistry and Pharmacology*, A. T. Blomquist and H. Wasserman, Ed., Academic Press, New York and London, **1972**; b) M. Shamma and J. L. Moniot, *Isoquinoline Alkaloids Research: 1972-1977*, Plenum Press, New York and London, **1978**.
2. T. K. Highsmith and A. I. Meyers, *Advances in Heterocyclic Natural Product Synthesis*, W. H. Pearson,

- Ed., Vol. 1, JAI PRESS INC., Greenwich, Connecticut, London, England, 1990, pp. 95-135, and references cited therein.
3. a) A. I. Meyers and D. Dickman, *Tetrahedron Lett.*, **1986**, 27, 1465; b) A. I. Meyers, D. Dickman, and M. Boes, *Tetrahedron*, **1987**, 43, 5095; c) H. Scafer and W. Ludwig, *Angew. Chem. Int. Ed. Engl.*, **1986**, 25, 1025.
  4. H. Hara, O. Hoshino, and B. Umezawa, *Chem. Pharm. Bull.*, **1976**, 24, 262.
  5. (*S*)-(+)-Lactic acid, and ( $\pm$ )- $\alpha$ -chloro- and ( $\pm$ )- $\alpha$ -bromopropionic acids were used as chiral solvent for the oxidation, however, no stable *p*-quinol esters were obtained in each case.
  6. O. Hoshino, H. Ogasawara, M. Suzuki, M. Arasawa, and B. Umezawa, *Heterocycles*, **1990**, 30, 385.
  7. Without further purification, the *p*-quinol esters were submitted to acid catalyzed cyclization.
  8. S. R. Johns, J. A. Lambertson, C. S. Li, and A. A. Sioumis, *Aust. J. Chem.*, **1970**, 23, 363.
  9. S. R. Johns, J. A. Lambertson, and A. A. Sioumis, *Aust. J. Chem.*, **1966**, 19, 2331.
  10. O. Hoshino, H. Ogasawara, M. Arasawa, M. Suzuki, and K. Iizima, *Heterocycles*, **1993**, 35, 1005.
  11. J. T. Etse and P. G. Waterman, *Phytochemistry*, **1986**, 25, 1903.
  12. R. Hocquemiller, A. Öztekin, F. Roblot, M. Hutin, and A. Cavé, *J. Nat. Prod.*, **1984**, 47, 342.
  13. J. Knabe and P. Horn, *Arch. Pharm.*, **1967**, 300, 726; B. Umezawa, O. Hoshino, H. Hara, and J. Sakakibara, *Chem. Pharm. Bull.*, **1968**, 16, 566.; G. Grethe, H. L. Lee, M. R. Uskokovic, and A. Brossi, *Helv. Chim. Acta*, **1970**, 53, 874.
  14. The value of ee could not be raised up because of unsuccessful separation of the racemate and (+)-enantiomer by crystallization.

(Received in Japan 10 May 1995)