



Stereochemistry of Nucleophilic Ring-Opening Reactions of Optically Active N-Acetyl-2-Methoxycarbonylaziridine.

Paolo Davoli, Arrigo Forni, Irene Moretti,* Fabio Prati

Dipartimento di Chimica, Università di Modena, via Campi 183, 41100 Modena, Italy

Abstract: The S_N2 -like mechanism of the nucleophilic attack of sodium azide on (S)-(-)-N-acetyl-2-methoxycarbonylaziridine was verified through the chemical correlation of the ring-opening products with (S)-(+)- or (R)-(-)-2,3-diaminopropanoic acid monohydrochloride.

Introduction.

Aziridines, like epoxides and cyclopropanes, have high ring-strain, which makes for easy ring-opening in the presence of nucleophiles. They can therefore be used as useful intermediates for the synthesis of biologically and pharmacologically active compounds. Ring-opening occurs even more readily in aziridines when an electron-withdrawing group, such as an acyl or sulphonyl substituent, is introduced on the ring-nitrogen atom. There are many reports in the literature about the stereochemical and regiochemical control exhibited by aziridine-ring substituents and nucleophiles and about the conditions pertaining to the ring-opening reactions.¹⁻⁴


In this work we perform a detailed study of the mechanism by which sodium azide induces an acid-catalyzed ring-opening reaction of optically active N-acetyl-2-methoxycarbonylaziridine **1**. (S)-(-)-N-acetyl-2-methoxycarbonylaziridine **1** was treated with sodium azide in the presence of boron trifluoride etherate as Lewis acid catalyst. The ring-opening products were correlated with the enantiomerically and configurationally known⁵ 2,3-diaminopropanoic acid monohydrochloride, which is commercially available. This correlation allowed us to define unequivocally the stereochemical course of the nucleophilic attack on the aziridine ring.

Results and discussion.

(S)-(-)-N-acetyl-2-methoxycarbonylaziridine **1**: resolution.-- Resolution of **1** through stereoselective enzymatic hydrolysis, catalyzed by *Candida cylindracea* lipase (CCL), is reported to afford only 50-60% enantiomeric excess (ee).⁶ Enantiomerically pure aziridine **1** was obtained here in good chemical yield from the corresponding N-butyryl-2-methoxycarbonylaziridine **2**, which can be resolved in 90% ee by CCL-enzymatic hydrolysis, following Scheme 1.

Racemic aziridine **2** was synthesized and resolved by CCL-catalysed hydrolysis, following the procedure reported in the literature.⁶ The resolution was performed in phosphate buffer (0.1 mol dm⁻³ and NaCl 0.1 mol dm⁻³; pH 7.5) at 37 °C using an enzyme /aziridine ratio (w/w) of 1:60. After 30 min, at 60% conversion, the reaction mixture revealed the presence of N-H aziridine **3**, as hydrolysis product, together with unchanged aziridine **2**. The mixture was extracted with methylene chloride and directly acetylated with acetyl chloride in the


presence of triethylamine to convert the unstable and volatile aziridine **3** into the corresponding N-acetylaziridine **1**. Aziridine (S)-(-)-**2**, in a nearly enantiomerically pure form (90% ee), and aziridine (R)-(+)-**1** with 50% ee were isolated by chromatography.




Enzymatic deacylation of enantiomerically pure aziridine **2** in hexane, using a CCl₄/aziridine ratio of 1:1 at 37 °C for 45 min in the presence of an equimolar amount of *n*.butanol and water in trace, gave the derivative **3** in quantitative yield. After removal of the enzyme by filtration, the solution was acylated to afford (S)-(-)-**1** in 70 % chemical yield and 90% ee.

The enantiomeric purities of aziridines **1** and **2** were determined by analysis of the ¹H-NMR spectra recorded in CDCl₃ and in the presence of the chiral shift-reagent, Eu(hfc)₃, tris [3-(heptafluoropropyl)-hydroxymethylene-(+)-camphorato]europium-(III). The absolute configurations of **1** and **2** are known from the literature.⁶

Ring-opening of (S)-(-)-1 by sodium azide.— Treating (S)-(-)-**1** with sodium azide in DMF and in the presence of boron trifluoride ethyl etherate, as Lewis acid catalyst, at 37 °C for 3 days, two ring-opening products were obtained in a ratio of 1:1 in 50% total chemical yield. ¹H-NMR and mass data confirm that the nucleophilic attack occurred at both the C₃ and C₂ ring-carbon atoms, thus affording two regioisomers, namely, (+)-2-acetamido-3-azidopropanoic acid methyl ester **4** and (+)-3-acetamido-2-azidopropanoic acid methyl ester **5**, respectively. Moreover, the ¹H-NMR spectra recorded in C₆D₆ and in the presence of the chiral solvating agent (R)-(-)-2,2,2-trifluoro-1-(9-anthryl)ethanol showed that the enantiomeric purities of **4** and **5** were not less than 90%, Scheme 2.




Chemical correlations.— In order to verify the mechanism of the aziridine-ring nucleophilic attack we correlated the unknown compounds **4** and **5** with the known compound (S)-(+)- or (R)-(-)-2,3-diaminopropanoic acid monohydrochloride **6**.⁵ Catalytic hydrogenation of azide (+)-**4** in methanol, with Pd 10% on carbon as catalyst, afforded (-)-2-acetamido-3-aminopropanoic acid methyl ester **7**; the hydrolysis of (-)-**7** in aqueous HCl 10% afforded (S)-(+)-**6** with 90% ee, Scheme 3.



Scheme 3

Similarly, the catalytic hydrogenation of (+)-**5** provided (-)-3-acetamido-2-aminopropanoic acid methyl ester **8**, which gave, by hydrolysis in HCl 10%, (R)-(-)-**6** with 90% ee, Scheme 4.


Since reduction and hydrolysis do not involve the carbon stereogenic centre of azido-derivatives, compounds (+)-**4** and (-)-**7** must have the same S configuration at the stereogenic centre of (+)-**6**. At the same time, compounds (+)-**5** and (-)-**8** must have the same R configuration at the stereogenic centre of (-)-**6**.




Scheme 4

This result unambiguously indicates that, when the nucleophilic attack is at the C₃ ring-carbon atom of aziridine (S)-(-)-**1**, the ring-opening product (+)-**4** has the same configuration as the aziridine **1**. On the other hand, when the nucleophilic attack occurs at the C₂ stereogenic carbon atom of (S)-(-)-**1**, a ring-opening product (+)-**5** of opposite configuration with respect to aziridine **1** is recovered. Clearly, nucleophilic attack on aziridine (S)-(-)-**1** occurs by a S_N2-like mechanism with total inversion.

It is also noteworthy that the same correlation, effected by reduction of the azides (+)-**4** and (+)-**5** with triphenylphosphine⁷ in THF, surprisingly provided only the derivative **8** in both the enantiomeric forms, (+) and (-)-**8**, respectively. Acidic hydrolysis of (+)-**8** and (-)-**8** afforded (S)-(+)-**6** and (R)-(-)-**6**, respectively, thus confirming the above correlation, Scheme 5a,b.




Scheme 5a



Scheme 5b

The recovery of (+)-8 from (+)-4, as reduction product, seems to indicate that reduction proceeds through the formation of an iminophosphorane intermediate which spontaneously cyclizes to the imidazoline 9 by an aza-Wittig intramolecular reaction, as already reported for ω -azido ketones,⁷ Scheme 6.



Scheme 6

Easy imidazoline hydrolysis would produce derivative (+)-8. Since hydrolysis and phosphorane rearrangement do not involve the azide-stereogenic centre, we can deduce that (+)-8 and (+)-4 must again have the same *S* configuration as the correlated compound (+)-6.

Conclusion.

Nucleophilic ring-opening attack of sodium azide on monosubstituted *N*-activated aziridine (*S*)-(-)-1, under acid catalysis, occurs at both the *C*₂ and *C*₃ aziridine carbon atoms and with a *S*_N2-like mechanism with total inversion to the *C*₂ stereogenic centre. This agrees with reports in the literature^{1,3} regarding the parent compounds. Thus, α -aminoacids with the same configuration and β -aminoacids with opposite configuration can be synthesized from aziridine (*S*)-(-)-1 by *C*₃- and *C*₂-nucleophilic attack, respectively, at the aziridine ring.

Experimental

¹H-NMR spectra were recorded in CDCl₃ solution on a Bruker AMX 400 WB spectrometer. Chemical shifts are reported in δ values from TMS as internal standard (s singlet, d doublet, m multiplet, t triplet, br broad signal). Coupling constants (*J*) are given in Hz. Optical rotations were measured at 20 °C on a Perkin-Elmer 241 polarimeter in chloroform solutions and are in 10⁻¹ deg cm² g⁻¹. Enantiomeric purities (ee's) were evaluated

from the $^1\text{H-NMR}$ spectra recorded in CDCl_3 and in the presence of the chiral lanthanide shift-reagent (CLSR) $\text{Eu}(\text{hfc})_3$, tris [3-(heptafluoropropyl-hydroxymethylene)-(+)-camphorato]europium (III), in the CLSR/compound molar ratio 0.5-1, or in C_6D_6 and in the presence of a 5-fold excess of the chiral solvating agent (CSA) (R)-(-)-2,2,2-trifluoro-1-(9-anthryl)ethanol. Accuracy was within $\pm 2\%$. Mass spectra were determined on a Hewlett-Packard 5970 mass selective detector. GLC analyses were performed on a Hewlett-Packard 5890 A gas chromatograph (capillary column DB-1, $5\ \mu\text{m}$, $30\ \text{m} \times 0.53\ \text{mm}$ I.D.). Chromatographic purification of the compounds was performed on silica gel (ϕ 0.05-0.20 mm). $\text{Eu}(\text{hfc})_3$ and (R)-(-)-2,2,2-trifluoro-1-(9-anthryl)ethanol were purchased from Ega-Chemie and used without purification. The enzyme *Candida cylindracea* lipase (type VII) was purchased by Aldrich and used without purification. (S)-(+)-2,3-diaminopropanoic acid monohydrochloride was furnished by Fluka.

Racemic N-acetyl-2-methoxycarbonylaziridine **1** and N-butyryl-2-methoxycarbonylaziridine **2** were synthesized as described in the literature.⁶

(S)-(-)-N-Butyryl-2-methoxycarbonylaziridine 2: Following the procedure described elsewhere,⁶ racemic aziridine **2** (3 g) was added to $0.1\ \text{mol dm}^{-3}$ potassium phosphate buffer [120 mL containing NaCl ($0.1\ \text{mol dm}^{-3}$)], pH 7.5 at $37\ ^\circ\text{C}$ and treated with *Candida cylindracea* lipase (CCL, 50 mg) with vigorous stirring. Hydrolysis was followed by GLC and TLC and stopped after 30 min at 60% conversion. The reaction-mixture, containing the unchanged aziridine **2** and the N-unsubstituted aziridine **3**, was extracted with dichloromethane; the organic extracts were dried over Na_2SO_4 and treated with acetyl chloride and triethylamine at $0\ ^\circ\text{C}$ to convert the volatile aziridine **3** into the aziridine **1**. The mixture was stirred for 1 h, washed with water, dried (Na_2SO_4) and concentrated. The residue was chromatographed on column (diethyl ether-light petroleum 50:50) to afford (S)-(-)-**2** (1.1 g, 37%), $[\alpha]_{\text{D}} -76.1$ (c 0.9), 93% ee; δ_{H} 0.91 (3H, t), 1.70 (2H, m), 2.39 (2H, m), 2.50 (1H, dd, J 1.7, 5.5), 2.57 (1H, dd, J 1.7, 3.0), 3.12 (1H, dd, J 3.0, 5.5), 3.80 (3H, s); MS m/z 171 (M^+). (R)-(+)-**1** (1.2 g, 48%) was obtained with $[\alpha]_{\text{D}} +34.5$ (c 1.1), 50% ee; δ_{H} 2.16 (3H, s), 2.5 (1H, dd, J 5.5, 1.7), 2.58 (1H, dd, J 3.0, 1.7), 3.16 (1H, dd, J 3.0, 5.5), 3.8 (3H, s); MS m/z 144 ($\text{M}+1^+$).

(S)-(-)-N-acetyl-2-methoxycarbonylaziridine 1: *n*-Butanol (1.1 mL, 12 mmol) and CCL (1.76 g) were added to a solution of (S)-(-)-N-butyryl-2-methoxycarbonylaziridine **2**, (1.76 g, 10 mmol), 90% ee, in hexane (25 mL) previously saturated with water, and the suspension was vigorously stirred at $37\ ^\circ\text{C}$. After 45 min, GLC analysis revealed complete conversion of compound **2** into **3**. The enzyme was removed by filtration and the solution treated at $0\ ^\circ\text{C}$ with triethylamine (12 mmol) and acetyl chloride (11 mmol). After 30 min, the mixture was washed with water, dried (Na_2SO_4) and the solvent removed under reduced pressure. Column chromatography (diethyl ether-light petroleum 50:50), afforded compound **1** as a colourless oil (1.05 g, 70%) showing $[\alpha]_{\text{D}} -70.4$ (c 1.0), 90% ee; δ_{H} 2.16 (3H, s), 2.5 (1H, dd, J 5.5, 1.7), 2.58 (1H, dd, J 3.0, 1.7), 3.16 (1H, dd, J 3.0, 5.5), 3.8 (3H, s); MS m/z 144 ($\text{M}+1^+$).

Methyl (+)-2-acetamido-3-azidopropanoate 4 and methyl (+)-3-acetamido-2-azidopropanoate 5: Boron trifluoride ethyl etherate (206 μL , 1.7 mmol) was gradually added to a solution of (S)-(-)-**1**, 90% ee (200 mg, 1.4 mmol) and NaN_3 (272 mg, 4.2 mmol) in anhydrous DMF (5 mL) at $37\ ^\circ\text{C}$ under nitrogen. After 70 h the reaction mixture was poured in water (40 mL), extracted with dichloromethane and dried (Na_2SO_4). After removal of the solvent, the residue was chromatographed (ethyl acetate-hexane 80:20) to isolate compounds **4** and **5**. Compound **4** showed $[\alpha]_{\text{D}} +74.2$ (c 1.4), 90% ee; δ_{H} 2.07 (3H, s), 3.75 (1H, dd, J 12.6, 3.5), 3.77 (1H, dd, J 12.6, 3.5), 3.82 (3H, s), 4.76 (1H, dt, J 7.2, 3.5), 6.34 (1H, br); MS m/z 187

(M+1⁺). Compound **5** had $[\alpha]_D +81.6$ (c 1.6), 90% ee; δ_H 2.00 (3H, s), 3.51 (1H, dt, *J* 14.0, 6.7), 3.68 (1H, ddd, *J* 14.0, 6.1, 5.3), 3.82 (3H, s), 4.20 (1H, dd, *J* 6.7, 5.3), 5.90 (1H, br); MS *m/z* 187 (M+1⁺).

General procedure for catalytic hydrogenation of the azido derivatives (+)-4 and (+)-5: A solution of each azido-derivative in methanol, at room temperature, was stirred with Pd 10% on carbon, as catalyst, under 1 atm of hydrogen until TLC analysis (ethyl acetate-hexane 80:20) showed that no azide remained. The catalyst was removed by filtration and the solution concentrated *in vacuo*. The identity of the reduction products was confirmed by ¹H-NMR and mass spectroscopies.

Methyl (S)-(-)-2-acetamido-3-aminopropanoate 7: From (+)-4 (0.13 g), 90% ee, after 2 h, compound **7** (95%) was obtained with $[\alpha]_D -22.6$ (c 2.3, CH₃OH), 90% ee; δ_H 1.20 (2H, br), 2.06 (3H, s), 3.06 (1H, dd, *J* 13.2, 4.4), 3.11 (1H, dd, *J* 13.2, 4.4), 3.78 (3H, s), 4.61 (1H, dt, *J* 8.8, 4.4), 6.53 (1H, br); MS *m/z* 161 (M+1⁺). The absolute configuration of (-)-7 was assigned by conversion into (S)-(-)-2,3-diaminopropanoic acid monohydrochloride **6**. The hydrolysis of (-)-7 in aqueous HCl 10% at 60 °C for 12 h afforded (S)-(+)-6, m.p. 235-238 °C (dec), with $[\alpha]_D + 21.6$ (c 0.67, HCl 0.5 N), 90% ee.

Methyl (R)-(-)-3-acetamido-2-aminopropanoate 8: From (+)-5, (0.1 g), 90% ee, after 1.30 h, product **8** (93%) was obtained with $[\alpha]_D - 27.3$ (c 0.65, CH₃OH), 90% ee; δ_H 1.89 (2H, br), 2.01 (3H, s), 3.34 (1H, ddd, *J* 13.1, 6.5, 5.0), 3.61 (1H, dd, *J* 6.5, 5.0), 3.66 (1H, ddd, *J* 13.1, 6.5, 5.0), 3.76 (3H, s), 6.09 (1H, br); MS *m/z* 131 (M-OCH₃⁺). Hydrolysis of (-)-8 in aqueous HCl 10% afforded (R)-(-)-6, $[\alpha]_D - 22.0$ (c 0.75, HCl 0.5 N), 90% ee.

General procedure for reduction of (+)-4 and (+)-5 with triphenylphosphine: PPh₃ (1 mmol) was added to a solution of (+)-4 or (+)-5 (1 mmol) in THF at room temperature with vigorous stirring. After 25 h, water (1.5 mmol) was added and the solution refluxed for 38 h until the disappearance of the azide was verified by TLC analysis (ethyl acetate-hexane 80:20). After the removal of the solvent, the residue was chromatographed on column to afford (+)-8 or (-)-8 in no more than 70% chemical purity: further purification attempts failed. (+)-8 was obtained from (+)-4 and (-)-8 from (+)-5. The hydrolysis in HCl 10% of (+)-8 and (-)-8 afforded, respectively, (S)-(+)-6 and (R)-(-)-6 with 90% ee.

Acknowledgements

The authors thank the Ministero dell'Università e della Ricerca Scientifica e Tecnologica, Rome, for financial support.

References

1. Tanner, D. *Angew. Chem. Int. Ed. Engl.*, **1994**, 33, 599 and references cited.
2. Falkenstein, R.; Mall, T.; Speth, D.; Stamm, H. *J. Org. Chem.*, **1993**, 58, 7377.
3. Legters, J.; Thijs, L.; Zwanenburg, B. *Recl. Trav. Chim. Pays-Bas*, **1992**, 111, 16.
4. Dubois, L.; Mehta, A.; Tourette, E.; Dodd, R. H. *J. Org. Chem.*, **1994**, 59, 434.
5. Birnbaum, S. M.; Koegel, R. J.; Fu, S.J.; Greenstein, J. P. *J. Biol. Chem.*, **1952**, 198, 335.
6. Bucciarelli, M.; Forni, A.; Moretti, I.; Prati, F.; Torre, G. *J. Chem. Soc., Perkin Trans 1*, **1993**, 3041.
7. Knouzi, N.; Vaultier, M.; Carrié, R. *Bull. Soc. Chim. Fr.*; **1985**, 5, 815.