



S0957-4166(96)00139-5

First Asymmetric Synthesis of (-)-(2*S*, 3*R*)-Methanoproline

Alain Hercouet*, Bernard Bessières and Maurice Le Corre.

Laboratoire de Synthèse Organique, Associé au CNRS. Université de Rennes I.
Avenue du Général Leclerc, 35042 Rennes Cedex, France.


Abstract: The title amino acid was synthesized in enantiomerically pure form, starting from (*S*)-(-)-butanetriol **1**, by condensation of cyclic sulfate **3** with methyl benzylideneglycinate. Copyright © 1996 Elsevier Science Ltd

The potential value of 2,3-methanoaminoacids as enzyme inhibitors or in the synthesis of conformationally constrained peptidomimetics was recognized some years ago¹. During the course of our studies toward the synthesis of cyclopropane aminoacids^{2,3}, we recently reported the synthesis of (-)-(1*S*, 2*R*)-allonorcoronamic acid using a chiral cyclic sulfate as precursor³.

We now wish to report the first asymmetric synthesis of (-)-(2*S*, 3*R*)-methanoproline **7**, a known ethylene biosynthesis inhibitor which has already been prepared by Stammer⁴ in the racemic form, and then resolved a few years later⁵.

Thus (*S*)-(-)-butanetriol⁶ **1** was converted to chlorosulfite **2**⁷ which was purified by flash chromatography (Scheme 1). The following oxidation, using the Sharpless procedure⁸, gave sulfate **3**.

Scheme 1



Condensation of this sulfate on methyl benzylideneglycinate⁹, at room temperature in DME in the presence of two equivalents of sodium hydride, was achieved following the procedure recently reported³, to give the alkylated imine **4** in quantitative yield. This reaction is diastereospecific, only the Z isomer is obtained.

Hydrolysis of the amino protective group by 1 N HCl gave the hydrochloride **5**¹⁰ in 80% yield, that cyclized to the desired aminoester **6** when treated with K₂CO₃. Hydrolysis of the ester group was realized by refluxing in 6 N HCl. The aminoacid zwitterion **7**¹¹ was then quantitatively obtained using Dowex 50X8. This new synthesis can be performed on a multigram scale with 43% overall yield from **2** and without purification of intermediates.

Spectral data and specific rotation of aminoacid **7** are in accordance with those obtained by Stammer⁵.

Acknowledgement: We are grateful to the *Conseil Régional de Bretagne* for a doctoral fellowship to B. B..

REFERENCES

- For recent reviews see: (a), Stammer C. H., *Tetrahedron*, **1990**, 46, 2231. (b) Alami A., Calmes M., Daunis J. and Jacquier R., *Bull. Soc. Chim. Fr.*, **1993**, 130, 5. (c) Burgess K., Ho K. K. and Moye-Sherman D., *Synlett*, **1994**, 575.
- Le Corre M., Hercouët A. and Bessières B., *Tetrahedron: Asymmetry*, **1995**, 6, 683.
- Hercouët A., Bessières B. and Le Corre M., *Tetrahedron: Asymmetry*, **1996**, 7, 283.
- Switzer F. L., Van Halbeek H., Holt E. M., Stammer C. H. and Salveit M. E., *Tetrahedron*, **1989**, 45, 6091.
- Matsui S., Srivastava V. P., Holt E. M., Taylor E. W. and Stammer C. H., *Int. J. Peptide Protein Res.*, **1991**, 37, 306.
- Hanessian S., Ugolini A., Dubé D. and Glamyan A., *Can. J. Chem.*, **1984**, 62, 2146.
- 4-chloroethyl-2-oxo-1,3,2-dioxathiolane **2**. SOCl₂ (21.3 mL, 300 mmol) is added dropwise to a solution of **1** (10.61 g, 100 mmol) in CCl₄ (100 mL) and the solution is refluxed for 3h. The solvent is evaporated and the resulting oil is chromatographed over SiO₂, eluting with Et₂O to yield the two diastereomers (50:50) of **2** (11.76 g, 69 mmol, 69%). ¹H NMR (300 MHz, CDCl₃) δ 2.06-2.48 (m, 4H); 3.65- 3.76 (m, 4H); 4.05-4.07 (m, 1H), 4.38-4.44 (m, 1H), 4.53-4.58 (m, 1H); 4.76-4.81 (m, 2H) and 5.12-5.21 (m, 1H). ¹³C NMR (75.5 MHz, CDCl₃) δ 35.45 and 36.61; 40.25 and 40.77; 70.43 and 71.26; 77.48 and 80.42.
- Gao Y. and Sharpless K. B., *J. Am. Chem. Soc.*, **1988**, 110, 7538.
- Stork G., Leong A. Y. W. and Touzin A. M., *J. Org. Chem.*, **1976**, 41, 3491.
- NMR of crude **5**: ¹H (300 MHz, CDCl₃) δ 1.62-1.74 (m, 2H), 1.98-2.05 (m, 1H), 2.19-2.26 (m, 1H), 2.41-2.46 (m, 1H), 3.67-3.82 (m, 2H), 3.78 (s, 3H) and 9.14 (br s, 3H). ¹³C (75.5 MHz, CDCl₃) δ 19.26, 24.19, 30.03, 38.40, 44.05, 53.38 and 169.73.
- (-)-(2*S*, 3*R*)-Methanoproline **7**: M.p. 217°C (dec) (Litt.⁵ M.p. 215°C (dec)), [α]_D²⁰ -51.2 (c 1, MeOH) (Litt.⁵ [α]_D²⁰ -48.2 (c 1, MeOH)). ¹H NMR (300 MHz, D₂O) δ 1.27-1.31 (m, 1H), 1.44-1.49 (m, 1H), 2.03-2.21 (m, 3H), 2.86-2.91 (m, 1H) and 3.39-3.46 (m, 1H). ¹³C NMR (75.5 MHz, D₂O) δ 15.57, 27.28, 27.31, 45.40, 51.16 and 176.81.