


The first asymmetric synthesis of the naturally occurring (+)-Kotatin and the assignment of its absolute configuration

Guo-Qiang Lin* and Min Zhong

Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 354 Fenglin Lu, Shanghai, 200032, China

Abstract: The first asymmetric synthesis of the naturally occurring (+)-Kotatin is described. The key steps involve the intramolecular oxidative coupling of the cyanocuprate intermediate and the Fries rearrangement. The absolute configuration of (+)-Kotatin was assigned as aS by CD spectroscopic method. © 1997 Elsevier Science Ltd

In 1971, the fungal metabolites Kotatin **1** and demethylkotatin **2** were isolated as the first enantiomerically pure 8,8'-bicoumarins from *Aspergillus clavatus* by Büchi and coworkers¹. Several years later, Cutler and coworkers² isolated another optically active 8,8'-bicoumarin named as Orlandin **3** from *Aspergillus niger*. Since then other two groups of optically active bicoumarins, *i.e.*, Desertorins³ and Isokotatinins⁴ were isolated from *Emericella desertorum* and *Aspergillus Alliaceus*, respectively.



- 1** Kotatin $R_1=R_2=Me$
2 Demethylkotatin $R_1=H, R_2=Me$
3 Orlandin $R_1=R_2=H$


Wheat coleoptile and chick bioassays indicated that the biological activity of Orlandin **3** in plants, and of Kotatin **1** in chick, is intimately associated with the 7,7'-hydroxyl groups. Orlandin **3** was nontoxic to day-old cockerels but significantly inhibited wheat coleoptile growth. Kotatin **1** was inactive in the coleoptile elongation but toxic to chicks². This result indicated that the bicoumarin structure offers an interesting model to determine functional group activity in plant and animal bioassays.

The optical activity of those bicoumarins was caused by the restricted rotation around the carbon-carbon single bond connecting the two aryl monomers. Although racemic Kotatin was synthesized by Büchi *et al.*^{1a}, the stereochemistry of it still remained unsolved for over twenty years. As a continuation of our efforts in this area, we report here the first asymmetric synthesis of the naturally occurring (+)-Kotatin, in which the asymmetric intramolecular oxidative coupling of the cyanocuprate intermediate of **8** developed by Lipshutz's group⁵ and the Fries rearrangement of **15** were employed as the key steps. The absolute configuration of (+)-Kotatin was assigned as aS based on the comparison of the CD Cotton effects of the intermediate **16** with that of the known compound⁶.

As shown in Scheme 1, 1,4-di-*O*-benzyl-L-threitol **4**⁷ was converted to its monosilyl ether **5**. Mitsunobu reaction of **5** with 2-bromo-3-methoxy-5-methylphenol⁸ gave **6** in 85% yield. Cleavage of

* Corresponding author. Email: lingq@pub.sioc.ac.cn

the silyl ether of **6** with *n*-Bu₄NF in THF gave **7** in 91% yield, which was followed by treatment **7** with 2-bromo-3-methoxy-5-methylphenol again to afford **8**⁹ in 54% yield. Treatment of **8** with *n*-BuLi followed by addition of CuCN-TMEDA (1:3) led to formation *in situ* of a higher order cyanocuprate intermediate, which transformed to **9**¹⁰ upon exposure to dry oxygen at -78°C in 60% yield. Catalytic hydrogenation of **9** gave the threitol **10** in 90% yield. The threitol **10** was converted to the ditosylate **11** in 92% yield, which upon treatment with NaI gave the diiodide **12** in 96% yield. Reduction of **12** by activated zinc powder in ethanol provided the biphenol **13** in 80%. The enantiomeric excess of **13** was determined to be 82% by the examination of the ¹H NMR spectra of its corresponding (*S*)-Mosher's ester **14**. The enantiomerically pure **13**¹¹ was obtained by recrystallization from ethyl acetate and hexane.




Reagents and conditions: a. TBDMSCl, imidazole, DMF, r.t., 24 h, 84%; b. 2-bromo-3-methoxy-5-methylphenol, DEAD, *n*-Bu₃P, THF, r.t., 24 h, 85%; c. *n*-Bu₄NF, THF, 2 h, 91%; d. 2-bromo-3-methoxy-5-methylphenol, DEAD, *n*-Bu₃P, THF, r.t., 42 h, 54%; e. *n*-BuLi, THF, -78°C, 1 h; CuCN-TMEDA(1:3), -78°C → -40°C, 1 h; dry O₂, -78°C, 4 h, 60%; f. 10% Pd/C, H₂, EtOAc, 12 h, 90%; g. TsCl, py., 0 °C, 8 h, 92%; h. NaI, acetone, reflux, 3 h, 96%; i. activated Zn powder, EtOH, reflux, 1 h, 80%; j. (*S*)- α -methoxy- α -(trifluoromethyl)phenylacetyl chloride, 4-DMAP, Et₃N, CH₂Cl₂, r.t., 24 h, 90%.

Scheme 1.

Subsequently, our efforts were made to complete the synthesis of **1** (Scheme 2).

The diacetate **15**, generated from **13** on treatment with acetic anhydride in pyridine, underwent Fries rearrangement promoted by TiCl₄ as Lewis acid to afford **16**¹² in 68% yield. The dicarbonate **17** prepared from **16** and methyl chloroformate in pyridine, was subjected to treatment of *t*-BuOK in *t*-BuOH to afford the desired cyclized product **18**. Then the crude **18** was directly methylated with NaH/HMPA/(CH₃)₂SO₄ to give the naturally occurring (+)-Kotanin **1**¹³ [[α]_D²⁰ +38.4 (c 0.44, CHCl₃); lit.^{1b} [α]_D²³ +40.0 (c 1.65, CHCl₃)] in 42% yield.

The CD Cotton effects of (+)-2,2'-dihydroxy-3,3'-diacetyl-4,4'-dimethyl-6,6'-dimethoxybiphenyl **16** were in accordance with that of (a*S*)-(+)-2,2'-dihydroxy-3,3'-diacetyl-4,4',6,6'-tetramethoxybiphenyl **19**, which exhibits positive first and negative second Cotton effects at the ¹L_a transition region⁶. Accordingly, the absolute configuration of the biaryls **9**, **10**, **11**, **12**, **13**, **15**, **16**, **17**, **18** and the naturally occurring (+)-Kotanin **1** were all assigned as a*S*. This assignment was also



Reagents and conditions: a. $(\text{CH}_3\text{CO})_2\text{O}$, py., 2 h, 93%; b. TiCl_4 , benzene, reflux, 4 h, 68%; c. ClCOOMe , py., 4-DMAP, 50 °C, 8 h, 85%; d. *t*-BuOK, *t*-BuOH, 60 °C, 2 h; e. NaH, HMPA, r.t.; $(\text{CH}_3)_2\text{SO}_4$, HMPA, r.t., 20 min, 42% (17→1).

Scheme 2

in agreement with Lipshutz's conclusion⁵ that the (2*S*,3*S*)-tetraether generally induced the formation of (a*S*)-biaryl in the coupling process (8→9).

The absolute configuration of the naturally occurring (+)-Isokotanin A, which belongs to 6,6'-bicumarin, was determined to be a*R* by us¹⁴, but that of the naturally occurring (+)-Kotanin, which was belong to 8,8'-bicumarin, was established as a*S*. The biosynthetic passway of (+)-Isokotanin A and (+)-Kotanin remains as an interesting issue.

In summary, we have accomplished the first asymmetric synthesis of the naturally occurring (+)-Kotanin and assigned its absolute configuration as a*S* by CD spectra.

References

1. a. Büchi, G.; Klaubert, D. H.; Shank, R. C.; Weinreb, S. M.; Wogan, G. N., *J. Org. Chem.*, **1971**, 36, 1143. b. Büchi, G.; Luk, K. C.; Kobbe, B.; Townsend, J. M., *J. Org. Chem.*, **1977**, 42, 244.
2. Cutler, H. G.; Crumley, F. G.; Cox, R. H.; Hernandez, O.; Cole, R. J.; Dorner, J. W., *J. Agric. Food Chem.*, **1979**, 27, 592.
3. Nozawa, K.; Seya, H.; Nakajima, S.; Udagawa, S.; Kawai, K., *J. Chem. Soc. Perkin Trans. I*, **1987**, 1735.
4. Laakso, J. A.; Narske, E. D.; Gloer, J. B.; Wicklow, D. T.; Dowd, P. F., *J. Natural Product*, **1994**, 57, 128.
5. Lipshutz, B. H.; Kayser, F.; Liu, Z.-P., *Angew. Chem. Int. Ed. Engl.*, **1994**, 33, 1842 and references cited therein.
6. (a*R*)-**19**. $[\alpha]^{20}_{\text{D}} -27.6$ (c 0.56, CHCl_3). CD (EtOH) λ_{ext} 298 nm ($\Delta\epsilon -3.29$), 276 ($\Delta\epsilon +4.60$), 241 ($\Delta\epsilon -1.34$), 235 ($\Delta\epsilon -0.26$), 229 ($\Delta\epsilon -2.63$), 212 ($\Delta\epsilon +8.67$). for (a*S*)-**19**. $[\alpha]^{19}_{\text{D}} +27.2$ (c 0.40, CHCl_3). CD (EtOH) λ_{ext} 297 nm ($\Delta\epsilon +3.38$), 275 ($\Delta\epsilon -2.03$), 240 ($\Delta\epsilon +3.78$), 235 ($\Delta\epsilon +2.43$), 230 ($\Delta\epsilon +3.78$), 211 ($\Delta\epsilon -6.21$).
7. Mash, E. A.; Nelson, K. A.; Densen, E. V.; Hemperly, S. B., *Org. Synth.*, Coll. Vol. VIII; John Wiley and Sons. Inc.: New York, **1993**: p. 155.
8. 2-Bromo-3-methoxy-5-methylphenol was synthesized according to the following lit.: a. Mirrington, R. N.; Feutrill, G. I. *Org. Synth.*, Coll. Vol. IV; John Wiley and Sons. Inc., New York, **1993**: p. 155. b. Cannon, J. R.; Cresp, T. M.; Metcalf, B. W.; Sargent, M. V.; Vinciguerra, G.; Elix, J. A., *J. Chem. Soc. (C)*, **1971**, 3495. c. Feutrill, G. I.; Mirrington, R. N., *Aust. J. Chem.*, **1972**, 25, 1719.
9. **8**, $[\alpha]^{22}_{\text{D}} -0.74$ (c 4.34, CHCl_3). FT-IR (film): 3064, 3030, 3005, 2980, 2863, 1587, 1496, 1457, 1412, 1367, 1326, 1236, 1112, 1040, 910, 813, 738, 699, 626, 573, 531, 462 cm^{-1} . ^1H NMR (300 MHz, C_6D_6) δ 7.12–7.27 (m, 10H), 6.79 (s, 2H), 6.09 (s, 2H), 5.12 (t, $J=3.6\text{Hz}$, 2H), 4.39, 4.35 (AB, $J_{\text{AB}}=12.0\text{Hz}$, 4H), 4.16 (dd, $J=4.1, 10.2\text{Hz}$, 2H), 3.95 (dd, $J=4.8, 10.3\text{Hz}$, 2H), 3.32 (s, 6H), 2.02 (s, 6H) ppm. MS m/z (EI, 70 eV): 700(M^+), 619, 593, 513, 297, 267, 227, 181, 138, 108, 91(100), 77. Calcd. for $\text{C}_{34}\text{H}_{36}\text{O}_6\text{Br}_2$: C, 58.30; H, 5.18. Found: C, 58.49; H, 5.32.

10. (aS)-**9**, $[\alpha]^{22}_D -12.0$ (c 0.68, CHCl₃). FT-IR (film): 3031, 2932, 2860, 1606, 1566, 1497, 1455, 1407, 1367, 1311, 1222, 1099, 948, 880, 738, 699, 583 cm⁻¹. ¹H NMR (300 MHz, CDCl₃) δ 7.30–7.38 (m, 10H), 6.70 (d, *J*=1.0Hz, 2H), 6.60 (d, *J*=1.0Hz, 2H), 4.61, 4.59 (AB, *J*_{AB}=12.1Hz, 4H), 4.10 (t, *J*=2.1Hz, 2H), 3.78 (s, 6H), 3.74 (dd, *J*=0.8, 10.9Hz, 2H), 3.62 (tt, *J*=3.3, 10.9Hz, 2H), 2.37 (s, 6H) ppm. ¹H NMR (300 MHz, C₆D₆) δ 7.30–7.32 (m, 4H), 7.15–7.18 (m, 6H), 6.99 (s, 2H), 6.46 (s, 2H), 4.43 (s, 4H), 4.35 (m, 2H), 3.50–3.62 (m, 4H), 3.42 (s, 6H), 2.23 (s, 6H) ppm. ¹³C NMR (75.5 MHz, CDCl₃) δ 159.393, 157.912, 139.388, 138.173, 128.433, 127.807, 127.699, 115.167, 114.589, 108.320, 85.099, 73.719, 70.744, 55.911, 21.820 ppm. MS *m/z* (EI, 70 ev): 541, 540(M⁺, 10.76), 432, 401, 341, 324, 311, 285, 256, 213, 195, 175, 151, 129, 105, 91(100), 77. HRMS calcd. for C₃₄H₃₆O₆(M⁺): 540.2513, found 540.2475.
11. (aS)-**13**, m.p. 122–123°C (EtOAc/hexane). $[\alpha]^{22}_D -143.2$ (c 0.66, CHCl₃). FT-IR (KBr): 3477, 3426, 2944, 2850, 1615, 1573, 1465, 1334, 1318, 1199, 1164, 1101, 1099, 1003, 938, 836, 812, 589, 562, 510, 470 cm⁻¹. ¹H NMR (300 MHz, C₆D₆) δ 6.74 (s, 2H), 6.15 (s, 2H), 5.18 (s, 2H), 3.20 (s, 6H), 2.15 (s, 6H) ppm. MS *m/z* (EI, 70 ev): 276, 275, 274(M⁺, 100), 259, 243, 228, 213, 199, 188, 177, 162, 151, 141, 128, 115, 105, 91, 77. Calcd. for C₁₆H₁₈O₄: C, 70.06; H, 6.61. Found: C, 69.95; H, 6.65.
12. (aS)-**16**, m.p. 217–218°C (CHCl₃/EtOH). $[\alpha]^{22}_D +109.2$ (c 0.70, CHCl₃). CD (EtOH) λ_{ext} 335nm (Δε +3.71), 310(Δε +2.17), 297(Δε +3.10), 274(Δε -6.82), 227(Δε +22.32), 210(Δε -32.89). FT-IR (KBr): 3503, 3370, 3011, 2978, 2941, 2850, 2739, 2687, 1600, 1495, 1464, 1420, 1400, 1349, 1271, 1214, 1183, 1162, 1110, 1044, 953, 874, 822, 610, 586, 537 cm⁻¹. ¹H NMR (300 MHz, C₆D₆) δ 14.10 (s, 2H), 6.10 (s, 2H), 3.34 (s, 6H), 2.07 (s, 6H), 2.04 (s, 6H) ppm. MS *m/z* (EI, 70 ev): 360, 359, 358(M⁺, 43.38), 343(100), 325, 301, 259, 255, 241, 213, 193, 164, 141, 128, 115, 91, 77. Calcd. for C₂₀H₂₂O₆: C, 67.03; H, 6.18. Found: C, 67.32; H, 6.18.
13. (aS)-**1**, m.p. >300°C (CHCl₃/EtOH). $[\alpha]^{20}_D +38.4$ (c 0.44, CHCl₃). FT-IR (KBr): 2928, 2853, 1711, 1611, 1588, 1457, 1412, 1365, 1330, 1259, 1204, 1136, 1125, 1204, 1136, 1125, 1097, 1054, 1032, 973, 838, 809, 733, 580 cm⁻¹. ¹H NMR (300 MHz, CDCl₃) δ 6.72 (s, 2H), 5.51 (s, 2H), 3.93 (s, 6H), 3.78 (s, 6H), 2.70 (s, 6H) ppm. MS *m/z* (EI, 70ev): 440, 439, 438(M⁺, 24.60), 410, 409, 408, 407(100), 379, 293, 265, 219, 165, 139, 111, 91, 69, 57, 43. HRMS calcd. for C₂₄H₂₂O₈(M⁺): 438.1315, found 438.1342.
14. Lin, G.-Q.; Zhong, M., *Tetrahedron Lett.*, **1996**, 37, 3015.

(Received in Japan 9 December 1996; accepted 4 March 1997)