




B,B-Dihaloterpenylboranes as reagents for the diastereo- and enantioselective synthesis of *syn*-aldols

P. Veeraraghavan Ramachandran, Wei-chu Xu[†] and Herbert C. Brown^{*}

H. C. Brown and R. B. Wetherill Laboratories of Chemistry, Purdue University, West Lafayette, Indiana 47907-1393, USA

Abstract: A series of known and new *B,B*-dihaloterpenylboranes, readily synthesized from the corresponding terpenes, were examined for the diastereo- and enantioselective synthesis of *syn*-aldols. Thus, *lpc*BCl₂, *lcp*BCl₂, *lcp*fBCl₂, *cis*-MyrBCl₂, *2-lcr*BCl₂, *4-lcr*BCl₂, and *2-lcr*BBr₂ in the presence of *i*-Pr₂NEt, convert 3-pentanone to ≥99% *Z*-enolates, converted by aldehydes to pure *syn*-aldols in 7–74% enantiomeric excess (ee). The most efficient of these reagents, *B,B*-dibromo-2-isocaranylborane, was tested for the enolboration of 3-pentanone, followed by aldolization with a series of aldehydes. © 1997 Elsevier Science Ltd

The diastereoselective syntheses of cross aldols via enolboration of ketones have been essentially mastered over the past two decades.¹ Although most of the literature reports on enolborations utilize R₂BX (X=Cl, Br, I, OTf, etc.) reagents,¹ there have been only scanty reports involving the corresponding RBX₂ reagents.² As part of our ongoing projects on enolboration, we recently reported the utilization of *B,B*-dihalomonalkylboranes as reagents for the stereoselective enolization of ethyl ketones, followed by *syn*-selective aldolization with a representative series of aldehydes (Eq. 1).³ All of the reagents tested provided the *syn*-aldols selectively.



Enantioselective aldol reactions have been achieved by way of substrate- and/or reagent-controlled enolborations.⁴ The reagents that have been developed for the latter strategy include several R₂BX derivatives prepared from terpenes, such as α -pinene,⁵ carene,⁶ and menthene.⁷ However, to the best of our knowledge, there has been no report of the utilization of dihaloborane derivatives, with an optically active alkyl group attached, for enantioselective enolboration–aldolization. Encouraged by our earlier success with the RBX₂ reagents (Eq. 1),³ we undertook a study of the enolboration of ethyl ketones with a series of chiral dihaloborane reagents (1–6) prepared from readily available terpenes employing the BCl₃–Me₃SiH procedure developed by Soundararajan and Matteson.⁸ Promising preliminary results are reported herein.

It has been recorded, in the case of *lpc*₂BX reagents, that the isopinocampheyl (*lpc*) moiety usually provides high enantioselectivity for *syn*-selective, and low enantioselectivity for *anti*-selective aldol reactions.⁵ For example, while diisopinocampheylboron triflate generally provides high enantiomeric excess (ee) for *syn*-aldols, diisopinocampheylboron chloride affords poor ee for *anti*-aldols. We were

^{*} Corresponding author. Email: HCBROWN@CHEM.PURDUE.EDU


[†] Postdoctoral Research Associate on a grant from the Office of Naval Research.

Table 1. Enolboration–aldolization of diethyl ketone/benzaldehyde with TerBCl₂ reagents under standard conditions


reagent	R*BCl ₂ R*	enolate ^a		yield isol.	aldol ^b		ee ^d %	config. ^d
		%Z	%E		syn ^c %	anti %		
1	Ipc	≥99	≤1	88	≥99	≤1	15	1 <i>R</i> ,2 <i>R</i>
2	Eap	≥99	≤1	92	≥99	≤1	19	1 <i>R</i> ,2 <i>R</i>
3	Lgf	≥99	≤1	95	≥99	≤1	8	1 <i>S</i> ,2 <i>S</i>
4	Myr	≥99	≤1	90	≥99	≤1	7	1 <i>S</i> ,2 <i>S</i>
5	2-Icr	≥99	≤1	93	≥99	≤1	59	1 <i>R</i> ,2 <i>R</i>
6	4-Icr	≥99	≤1	98	≥99	≤1	28	1 <i>S</i> ,2 <i>S</i>

^aThe enolization was carried out in CH₂Cl₂ at –78 °C for 1 h. The *Z/E* ratio is based on the *syn/anti* ratio of the aldol. ^bThe aldolization was carried out at –78 °C for 2 h. ^cThe *syn/anti* ratio is determined on the basis of ¹H NMR. ^dDetermined by ¹H and ¹⁹F NMR of the corresponding MTPA derivative.

curious to examine this 'super chiral auxiliary',⁹ α-pinene, for *syn*-selective aldol reactions under the changed steric and electronic environments.



i-Pr₂EtN was added to a flask containing isopinocampheylchloroborane (IpcBCl₂, **1**) and 3-pentanone in CH₂Cl₂ at –78 °C. The enolboration was complete in 1 h. The ¹¹B NMR spectrum of an aliquot of the reaction mixture shifted from δ 62 ppm (singlet), corresponding to the reagent, to a singlet at δ 41 ppm, corresponding to an alkylchloroborinate. Following addition of benzaldehyde to the reaction mixture at –78 °C, the aldolate was produced within 2 h (¹¹B NMR δ 31 ppm).³ Upon oxidation, the product aldol was obtained as ≥99% *syn* isomer, as established by the ¹H NMR spectrum. Since all of the dihaloboranes consistently yield *syn*-aldols, the diastereoselectivity achieved with this reagent is not surprising. However, the enantiomeric excess (ee) realized, 15% as determined by the ¹⁹F NMR spectral analysis of the MTPA ester,¹⁰ is very disappointing (eq 2).



In our asymmetric reduction program, we observed that increasing the steric requirements at the 2-position of the apopinene moiety in the reagent containing such a group results in a considerable increase in the ee of the product alcohols produced.¹¹ On this basis, we examined EapBCl₂ (**2**) for the enolboration–aldolization of 3-pentanone–benzaldehyde. The modest improvement in ee achieved (19%) fell short of our expectations. We then investigated reagents **3–6**. The reagents **3** and **4** derived from the terpenes containing terminal olefins, viz. longifolene and β-pinene, yielded poor enantioselectivity (7–8% ee). This may be due to the fact that the nearest chiral center is one carbon removed from the boron atom.

Our experiences in the allyl- and crotylboration program suggested that the reagents derived from the two carenes might prove superior than those derived from α-pinene.¹² Indeed reagents **5** and **6** offered reasonably good ee for the aldol. The results on the enolboration–aldolization of 3-pentanone–benzaldehyde with TerBCl₂ reagents are summarized in Table 1.


Since 2-IcrBCl₂ provided the best results, the corresponding dibromoborane (2-IcrBBr₂, **7**) was prepared from 2-carene and examined for the same reaction (Scheme 1). This reagent provides improved enantioselectivity (74% ee) for the product aldol. We investigated the generality of this

Table 2. Enolboration–aldolization of diethyl ketone/RCHO with 2-*lcr*BBr₂ (7) under standard conditions

RCHO R	enolate ^a		yield isol.	aldol ^b		ee ^d %	config. ^d
	%Z	%E		<i>syn</i> ^c %	<i>anti</i> %		
Ph	≥99	≤1	88	≥99	≤1	74	1 <i>R</i> ,2 <i>R</i>
Et	≥99	≤1	92	≥99	≤1	59	1 <i>S</i> ,2 <i>R</i>
<i>i</i> -Pr	≥99	≤1	93	≥99	≤1	58	1 <i>S</i> ,2 <i>R</i>
<i>t</i> -Bu	≥99	≤1	98	≥99	≤1	14	1 <i>R</i> ,2 <i>R</i>

^{a-d}See footnotes *a-d* in Table 1.

reagent by carrying out the cross aldol reaction with a representative series of aldehydes and obtained 14–74% ee for the products. The aldols from unhindered aliphatic aldehydes were obtained in 58–59% ee. When the aldehyde was hindered, the ee dropped considerably. For example, the reaction of pivalaldehyde with the corresponding enolate from 3-pentanone, affords only 14% ee. The results are summarized in Table 2.



In conclusion, we have synthesized a series of *B,B*-dihaloterpenylboranes and examined them for enantioselective aldol reactions. One of these *B,B*-dibromo-2-isocaranylborane appears to be a very promising reagent.^{13,14} We are continuing the investigations on the scope of this reagent. We are also examining these and several new R^*BX_2 reagents for asymmetric aldolizations.

Acknowledgements

The financial assistance from the U.S. Office of Naval Research is gratefully acknowledged.

References

- For reviews on the aldol reactions, see: (a) Mukaiyama, T. *Org. React.* **1982**, 28, 203. (b) Evans, D. A.; Nelson, J. V.; Taber, T. R.; *Top. Stereochem.* **1982**, 13, 1. (c) Heathcock, C. H. in *Asymmetric Synthesis*, Morrison, J. D. Ed., Academic Press, Orlando, FL, 1983, Vol.3, Chapter 2. (d) Heathcock, C. H. in *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I. Eds.; Pergamon: Oxford, 1991, Vol. 2. For reactions of achiral boron enolates, see: (e) Fenzl, W.; Koster, R. *Liebigs Ann. Chem.* **1975**, 1322. (f) Masamune, S.; Mori, S.; Van Horn, D.; Brooks, D. W. *Tetrahedron Lett.* **1989**, 1665. (g) Brown, H. C.; Dhar, R. K.; Bakshi, R. K.; Pandiarajan, P. K.; Singaram, B. *J. Am. Chem. Soc.* **1989**, 111, 3441. (h) Brown, H. C.; Ganesan, K.; Dhar, R. K. *J. Org. Chem.* **1992**, 57, 3767. (i) *idem.*, *ibid.* **1993**, 58, 147. (j) Ganesan, K.; Brown, H. C. *J. Org. Chem.* **1993**, 58, 7162.
- (a) Hamana, H.; Sasakura, K.; Sugasawa, T. *Chem. Lett.* **1984**, 1729. (b) Evans, D. A.; Calter, M. A. *Tetrahedron Lett.* **1993**, 34, 6871. (c) Luke, G. P.; Morris, J. *J. Org. Chem.* **1995**, 60, 3013.
- Ramachandran, P. V.; Xu, W. C.; Brown, H. C. *Tetrahedron Lett.* **1997**, 38, 769.

4. (a) Evans, D. A.; Nelson, J. V.; Taber, T. R. in *Topics in Stereochemistry*; Wiley-Interscience, New York, 1982. Vol. 13, p.1. (b) Bernardi, A.; Gennari, C.; Goodman, J. M.; Paterson, I. *Tetrahedron: Asym.* **1995**, *6*, 2613.
5. Meyers, A. I.; Yamamoto, Y. *J. Am. Chem. Soc.* **1981**, *103*, 4278. (b) Paterson, I.; Goodman, J. M.; Lister, M. A.; Schumann, R. C.; McLure, C. K.; Norcross, R. D. *Tetrahedron* **1990**, *46*, 4663.
6. Fringuelli, F.; Piermatti, O.; Pizzo, F. *J. Org. Chem.* **1995**, *60*, 7006.
7. Gennari, C.; Hewkin, C. T.; Molinari, F.; Bernardi, A.; Comotti, A.; Goodman, J. M.; Paterson, I. *J. Org. Chem.* **1992**, *57*, 5173.
8. (a) Soundararajan, R.; Matteson, D. S. *J. Org. Chem.* **1990**, *55*, 2274; *Organometallics* **1995**, *14*, 4157. For other methods of preparing R*BCl₂ reagents, see: Brown, H. C.; Ramachandran, P. V.; Chandrasekharan, J. *Heteroatom Chem.* **1995**, *7*, 117.
9. Brown, H. C.; Ramachandran, P. V. In *Advances in Asymmetric Synthesis* Vol, 1., Hassner, A. Ed. JAI Press: Greenwich, CT, 1995, pp 147–210.
10. Dale, J. A.; Dull, D. L.; Mosher, H. S. *J. Org. Chem.* **1969**, *34*, 2543.
11. Brown, H. C.; Ramachandran, P. V.; Teodorovic', A. V.; Swaminathan, S. *Tetrahedron Lett.* **1991**, *32*, 6691.
12. Brown, H. C.; Randad, R. S.; Bhat, K. S.; Zaidlewicz, M.; Racherla, U. S. *J. Am. Chem. Soc.* **1990**, *112*, 2389.
13. All of the operations were carried out under nitrogen.¹⁴ An oven-dried, 50 mL round-bottom flask equipped with a side-arm, magnetic stirring bar, and a connecting tube was cooled to rt in a stream of nitrogen. 2-IcrBBR₂ (1.9 g, 6 mmol) and CH₂Cl₂ (30 mL) were added via a cannula to the flask, followed by the addition of 3-pentanone (0.43 g, 5 mmol). The flask was cooled to -78°C and *i*-Pr₂EtN (0.87 g, 6.5 mmol) was added using a syringe. The enolborinate was generated instantaneously with concurrent formation of *i*-Pr₂EtN·HBr. The ¹¹B NMR spectrum of an aliquot showed a singlet at δ 41. The mixture was stirred at this temperature for 1 h. PhCHO was added to this mixture and the reaction was followed using ¹¹B NMR spectroscopy of aliquots. Upon completion of the reaction (¹¹B NMR δ 31) methanol (8 mL) was added to the mixture, and oxidized with alkaline H₂O₂. The crude product was extracted with CH₂Cl₂, concentrated, chromatographed through silica (hexane:ethyl acetate, 10:1), and the solvents were removed to obtain (0.89 g) 92% of (1*R*,2*R*)-2-methyl-3-oxo-1-phenylpentanol. The structure was confirmed by ¹H and ¹³C NMR spectroscopy. The *syn/anti* ratio was determined from the ¹H NMR spectrum. The % ee and configuration were determined by analysis of the ¹⁹F NMR spectrum of the MTPA ester.
14. Brown, H. C.; Kramer, G. W.; Levy, A. B.; Midland, M. M. *Organic Syntheses via Boranes* Wiley-Interscience: New York, **1975**; Chapter 9.

(Received in USA 27 February 1997)