



Diastereoselective synthesis of pyridyl substituted thiazolidin-4-ones. New ligands for the Cu(I) catalyzed asymmetric conjugate addition of diethylzinc to enones

André H. M. de Vries, Robert P. Hof, Danny Staal, Richard M. Kellogg*[†] and
Ben L. Feringa*^{**}

Department of Organic and Molecular Inorganic Chemistry, Groningen Center for Catalysis and Synthesis,
University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

Abstract: Several novel thiazolidin-4-ones have been synthesized from chiral non-racemic α -mercapto acids by a three component reaction in high yields and with diastereomeric excesses up to 91%. After recrystallization, the thiazolidinones were obtained diastereomerically pure and were employed as chiral ligands in the copper catalyzed conjugate addition of diethylzinc to cyclohexenone and chalcone as model substrates. Enantioselectivities up to 62% were achieved. © 1997 Elsevier Science Ltd

Introduction

Chiral non-racemic β -amino alcohols are popular ligands in asymmetric catalysis.¹ The use of the corresponding β -amino thiols, however, has been limited due to the lack, until recently, of simple synthetic transformations to these interesting molecules. Building on the synthesis of enantiomerically pure β -amino thiols, -sulfides, and -disulfides² and a recently improved synthesis of both enantiomers of thiolactic acid, starting from a cheap building block, ethyl (*S*)-lactate,³ we describe herein the synthesis and application of novel chiral thiazolidinones.

Sulfur containing compounds have been used as chiral ligands for stoichiometric⁴ and catalytic enantioselective conjugate addition reactions of Grignard reagents to enones.^{5–7} Furthermore, enantiomerically pure thiols and sulfides are successful catalysts in the enantioselective addition of diethylzinc to benzaldehyde.⁸ Because sulfides and pyridines have a pronounced affinity for Cu(I)⁹ and chiral copper catalysts have been used successfully in the enantioselective conjugate addition of dialkylzinc to enones,^{10,16b} we envisioned that combination of these elements in the form of chiral thiazolidinones also might lead to ligands suited for this asymmetric transformation.

Results and discussion


Synthesis of thiazolidin-4-ones 2

Although thiazolidinones have been the subject of extensive study and diverse biological activities have been found to be associated with this class of heterocycles,¹¹ the elegant cyclocondensation of a mercapto acid, an amine and an aldehyde under conditions of azeotropic removal of water to give thiazolidin-4-ones is relatively unexplored.¹² The 2-pyridyl-thiazolidin-4-ones **2** are obtained in quantitative yield just by mixing α -mercapto acid, aniline, and 2-pyridinecarboxaldehyde in appropriate amounts (Scheme 1). A mixture of *cis*- and *trans*-diastereomers is produced; the former predominates (*vide infra*). No racemization of the α -mercapto acid was observed. In contrast to reactions whereby chiral amines are used (no diastereoselectivity),^{12b} this synthesis using chiral α -mercapto acids proceeds with diastereomeric excesses (d.e., determined by ¹H NMR) varying from

* Corresponding author. E-mail: Feringa@chem.rug.nl

[†] Kellogg@chem.rug.nl

Table 1. Enantioselective 1,4-addition of diethylzinc to cyclohexenone catalysed by CuOTf and **2**^a

			
entry	ligand	e.e. of 4 (%) ^b	abs. conf. ^c
1	2a	47	<i>R</i>
2	2b	39	<i>R</i>
3	2c	49	<i>R</i>
4	2d	62	<i>R</i>

^aReactions at 1 mmol scale at -10°C in 5 ml of toluene. Reaction time 2–6 h. Conversion to the 1,4-product >95% (based on GC analysis). Isolated yields >70%. ^bEnantiomeric excess of **4** was determined by derivatization with enantiomerically pure 1,2-diphenylethylene diamine. ^cComparison of the specific rotation of **4** with known data gave the absolute configuration.¹⁷

General procedure for the synthesis of thiazolidinones **2**

A mixture of the (*R*)- α -mercapto acid (10 mmol), the aldehyde (10 mmol) and aniline (0.93 g, 10 mmol) was azeotropically refluxed in 100 ml benzene in a Dean–Stark apparatus overnight. After cooling, the solution was evaporated to dryness to give a quantitative yield of crude **2** as a mixture of diastereomers. The diastereomeric excess of crude **2** was determined by 200 MHz ¹H-NMR (the absorptions of the thioacetal proton were used to determine the ratio). Diastereomerically pure material (*cis*) was obtained after recrystallization as indicated.

(2*S*,5*R*)-5-Methyl-3-phenyl-2-pyridin-2-yl-thiazolidin-4-one *cis*-**2a**

From **1a**, aniline and 2-pyridinecarboxaldehyde crude **2a** (d.e. 91%) was obtained. This material was recrystallized from EtOH to give diastereomerically pure *cis*-**2a** (1.84 g, 6.81 mmol, 68%) as glistening needles; mp 204.8–205.2°C; $[\alpha]_{\text{D}} +287.0$ (*c* 1.03, CHCl₃); ¹H NMR δ 1.65 (d, *J*=7.0 Hz, 3H), 4.31 (q, *J*=7.0 Hz, 1H), 6.03 (s, 1H), 7.12–7.31 (m, 8H), 7.59–7.68 (m, 1H), 8.56 (d, *J*=4.4 Hz, 1H); ¹³C NMR δ 18.22 (q), 41.28 (d), 63.95 (d), 120.08 (d), 123.21 (d), 124.52 (d), 126.61 (d), 129.03 (d), 137.15 (d), 149.97 (d), 159.31 (s); Anal. Calcd. for C₁₅H₁₄N₂OS: C, 66.64; H, 5.22; N, 10.36; S, 11.86. Found: C, 66.64; H, 5.14; N, 10.21; S, 11.71.

(2*S*,5*R*)-5-Methyl-3-phenyl-2-(6-methyl-pyridin-2-yl)-thiazolidin-4-one *cis*-**2b**

From **1a**, aniline and 6-methyl-2-pyridinecarboxaldehyde crude **2b** (d.e. 51%) was obtained. This material was recrystallized from ether/hexane to give diastereomerically pure *cis*-**2b** (1.54 g, 5.4 mmol, 54%); mp 110.5–112.3°C; $[\alpha]_{\text{D}} +161.5$ (*c* 1.20, CHCl₃); ¹H NMR δ 1.64 (d, *J*=7.1 Hz, 3H), 2.53 (s, 3H), 4.30 (q, *J*=7.1 Hz, 1H), 5.98 (s, 1H), 6.97–7.55 (m, 8H); ¹³C NMR δ 17.99 (q), 24.21 (q), 41.13 (d), 63.97 (d), 116.67 (d), 122.80 (d), 124.43 (d), 126.50 (d), 128.99 (d), 137.31 (d), 138.13 (s), 158.87 (s); Anal. Calcd. for C₁₆H₁₆N₂OS: C, 67.58; H, 5.67; N, 9.85; S, 1.27. Found: C, 67.36; H, 5.69; N, 9.81; S, 11.18.

(2*S*,5*R*)-5-Methyl-3-phenyl-2-quinolin-2-yl-thiazolidin-4-one *cis*-**2c**

From **1a**, aniline and 2-quinolinecarboxaldehyde crude **2c** (d.e. 24%) was obtained. This material was recrystallized four times from EtOH to give *cis*-**2c** (697 mg, 2.17 mmol, 22%, d.e. 94%) as glistening plates; mp 179.8–180.5°C; $[\alpha]_{\text{D}} +327.9$ (*c* 1.20, CHCl₃); ¹H NMR δ 1.70 (d, *J*=7.1 Hz, 3H), 4.33 (dq, *J*=7.1 Hz and *J*=1.2 Hz, 1H), 6.25 (d, *J*=1.2 Hz, 1H), 7.08–8.17 (m, 11H); ¹³C NMR δ 18.59 (q), 41.44 (d), 64.46 (d), 117.66 (d), 124.62 (d), 126.67 (d), 126.93 (d), 127.57 (d), 129.05 (d), 129.30 (d), 130.01 (d), 137.82 (d), 137.98 (s), 147.43 (s), 159.33 (s), 174.03 (s).

(2S,5R)-5-Isopropyl-3-phenyl-2-pyridin-2-yl-thiazolidin-4-one cis-2d

Starting from 9.5 mmol (*R*)-2-mercapto-3-methylbutanoic acid **1b**,¹⁴ aniline and 2-pyridine-carboxaldehyde crude **2d** (d.e. 85%) was obtained. This material was recrystallized from EtOH to give diastereomerically pure *cis-2d* (1.09 g, 3.66 mmol, 39%) as a white powder; mp 119.6–120.9°C; $[\alpha]_D^{25} +292.2$ (c 0.51, CHCl₃); ¹H-NMR δ 1.08 (dd, *J*=7.0 and *J*=6.6 Hz, 6H), 2.62 (m, 1H), 4.30 (s, 1H), 6.09 (s, 1H), 7.12–7.27 (m, 7H), 7.61 (m, 1H), 8.50 (m, 1H); ¹³C-NMR δ 20.96 (q), 30.99 (d), 55.09 (d), 64.83 (d), 120.62 (d), 123.13 (d), 124.81 (d), 126.61 (d), 128.91 (d), 137.03 (d), 149.52 (d), CO and quaternary C not observed.

Conjugate addition of diethylzinc to cyclohexenone 3 and chalcone using catalytic amounts of (CuOTf)₂·benzene and chiral thiazolidinones 2

This procedure is typical for all conjugate addition reactions. A solution of (CuOTf)₂·benzene (5 mol%) and of chiral thiazolidinone (11 mol%) in 5 ml of toluene and 2 ml of CH₂Cl₂ was stirred at ambient temperature for 1 h under argon. A clear solution was formed. Substrate was added (1.0–2.0 mmol), the mixture was cooled to –20°C and diethylzinc in toluene (1.1 M, 1.5 equivalent) was added. Stirring was continued at –10°C for 2–6 h. An aliquot of the solution (0.1 ml) was taken and quenched with 1 ml of aqueous 1 N HCl. After extraction with 1 ml of diethyl ether the conversion was determined by GC analysis.¹⁸ In all cases complete conversion was achieved with regioselectivities for the 1,4-product of >95%. The mixture was poured into 25 ml of aqueous 1 N HCl and extracted with diethyl ether (3×20 ml). The combined organic layers were washed with brine (25 ml), dried (MgSO₄), filtered and evaporated to give the crude 1,4-product. After purification by column chromatography (SiO₂, hexane:diethyl ether 5:1) the e.e.'s were determined.¹⁸

References

1. a) *Catalytic Asymmetric Synthesis*; Ojima, I., Ed.; VCH Publishers: New York, 1993. b) Brunner, H.; Zettlmeier, W. *Handbook of Enantioselective Catalysis*; VCH Publishers: New York, 1993. c) Noyori, R. *Asymmetric Catalysis in Organic Chemistry*; Wiley: New York, 1994. See for example also de Vries, A.H.M.; Feringa, B.L. *Tetrahedron: Asymmetry* **1997**, *8*, 1377 and Ref. 18.
2. Poelert, M.A.; Hof, R.P.; Peper, N.C.M.W.; Kellogg, R.M. *Heterocycles* **1994**, *37*, 461.
3. Hof, R.P.; Kellogg, R.M. *J. Chem. Soc., Perkin Trans. I* **1995**, 1247.
4. a) Leyendecker, F.; Laucher, D. *Tetrahedron Lett.* **1983**, *24*, 3517. b) Leyendecker, F.; Laucher, D. *Nouv. J. Chim.* **1985**, *9*, 13.
5. a) Lambert, F.; Knotter, D.M.; Janssen, M.D.; van Klaveren, M.; Boersma, J.; van Koten, G. *Tetrahedron: Asymmetry* **1991**, *2*, 1097. b) van Klaveren, M.; Lambert, F.; Eijkelkamp, D.J.F.M.; Grove, D.M.; van Koten, G. *Tetrahedron Lett.* **1994**, *35*, 6135. c) Zhou, Q.-L.; Pfaltz, A. *Tetrahedron Lett.* **1993**, *34*, 7725. d) Zhou, Q.-L.; Pfaltz, A. *Tetrahedron* **1994**, *50*, 4467. e) Spescha, M.; Rihs, G. *Helv. Chim. Acta* **1993**, *76*, 1219.
6. For a recent review on catalytic enantioselective conjugate addition reactions, see: Feringa, B.L.; de Vries, A.H.M. in *Advances in Catalytic Processes, Vol 1: Asymmetric Chemical Transformations*; Doyle, M.P., Ed.; JAI Press, Connecticut, 1995, p. 151.
7. The use of a chiral bidentate phosphine as ligand in a CuI catalyzed addition of Grignard reagents to cyclic enones (e.e. up to 92%) has been reported recently: M. Kanai, K. Tomioka, *Tetrahedron Lett.* **1995**, *36*, 4275.
8. a) Hof, R.P.; Poelert, M.A.; Peper, N.C.M.W.; Kellogg, R.M. *Tetrahedron: Asymmetry* **1994**, *5*, 31. b) Fitzpatrick, K.; Hulst, R.; Kellogg, R.M. *Tetrahedron: Asymmetry* **1995**, *6*, 1861.
9. Hathaway, B.J. In *Comprehensive Coordination Chemistry*; Wilkinson, G.; Gillard, R.D.; McLeverly, J.A., Eds.; Pergamon: Oxford, **1987**, Vol. 5, Chapter 53.
10. de Vries, A.H.M.; Meetsma, A.; Feringa, B.L. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 2374.
11. Singh, S.P.; Parmar, S.S.; Raman, K.; Stenberg, V.I. *Chem. Rev.* **1981**, *81*, 175–203.
12. a) Johnson, M.R.; Fazio, M.J.; Ward, D.L.; Sousa, L.R. *J. Org. Chem.* **1983**, *48*, 495, and references therein. b) Botteghi, C.; Schionato, A.; Chelucci, G.; Brunner, H.; Kürzinger, A.; Obermann, U.

- J. Organomet. Chem.* **1989**, 370, 17. c) Tanabe, Y.; Yamamoto, H.; Murakami, M.; Yanagi, K.; Kubota, Y.; Okumura, H.; Sanmitsu, Y.; Suzukamo, G. *J. Chem. Soc., Perkin Trans I* **1995**, 935.
13. Seebach, D.; Naef, R.; Calderari, G. *Tetrahedron* **1984**, 40, 1313.
14. Strijtveen, B.M.; Kellogg, R.M. *J. Org. Chem.* **1986**, 51, 3664.
15. Alexakis, A.; Frutos, J.C.; Mangeney, P. *Tetrahedron: Asymmetry* **1993**, 4, 2431.
16. a) Very recently a β -amino disulfide, derived from proline, has been used as chiral ligand in the nickel catalyzed conjugate addition of diethylzinc to chalcone (e.e. <50%); no results with other enones, for example cyclohexenone, are described: Gibson, C.L. *Tetrahedron: Asymmetry* **1996**, 7, 3357. b) Except for our own results (reference 10), only one other example of enantioselectivity in a copper catalyzed addition of diethylzinc to cyclohexenone has been reported (e.e. 30%), see: Alexakis, A.; Frutos, J.; Mangeney, P. *Tetrahedron: Asymmetry* **1993**, 4, 2427.
17. Posner, G.H.; Frye, L.L. *Isr. J. Chem.* **1984**, 24, 88.
18. See de Vries, A.H.M.; Imbos, R.; Feringa, B.L. *Tetrahedron: Asymmetry* **1997**, 8, 1467.

(Received in UK 19 February 1997; accepted 25 March 1997)