



Pergamon

Tetrahedron: Asymmetry 9 (1998) 2493–2498

TETRAHEDRON:
ASYMMETRY

A facile stereodivergent synthesis of *threo*- and *erythro*-*N,N*-dibenzyl sphingosines from (*S*)-*N,N*-dibenzyl-*O*-TBDMS-serinal

José M. Andrés and Rafael Pedrosa *

Departamento de Química Orgánica, Facultad de Ciencias, Universidad de Valladolid, Doctor Mergelina s/n, 47011-Valladolid, Spain

Received 21 May 1998; accepted 10 June 1998

Abstract

The (*L*)-*threo*-*N,N*-dibenzyl sphingosine was prepared in two steps from the serinal derivative **1** by diastereoselective alkenylation with pentadec-1-enyl(ethyl)zinc and deblocking. (*D*)-*erythro*-*N,N*-Dibenzyl sphingosine was also prepared in two steps from **1** by a highly diastereoselective alkynylation with pentadecynyl magnesium bromide and subsequent stereoselective LAH reduction. © 1998 Elsevier Science Ltd. All rights reserved.

1. Introduction

The fact that sphingosines or their derivatives exert important biological activities¹ has attracted a lot of synthetic attention.² Some of the most interesting methods start from serine because this α -amino acid bears a part of the chiral moiety present in the final product.³ Starting from these derivatives two stereochemical aspects must be solved: the implantation of the *trans* double bond⁴ and the control of the stereochemistry at C-3.⁵

Our recent report on the stereoselective *syn*-addition of diethylzinc to α -*N,N*-dibenzylaminoaldehydes⁶ prompted us to prepare the (*S*)-serinal derivative **1** as a starting chiron capable of giving different stereoisomers by tuning the nature of the organometallic used as the nucleophile,⁷ and now we report here on the transformation of **1** into both *threo*- and *erythro*-*N,N*-dibenzyl sphingosines *syn*-**3** and *anti*-**3** in two steps.

* Corresponding author.

2. Results and discussion

(*S*)-*N,N*-Dibenzyl-*O*-TBDMS serinal **1** was prepared from (*L*)-*N,N*-dibenzyl methyl serinate in 74% overall yield. Protection of the alcohol as TBDMS ether followed by reduction of the ester with NaBH₄–LiCl in THF–ethanol gave the corresponding monoprotected aminodiols which were submitted to Swern oxidation affording the serinal derivative **1**.⁷

The synthesis of (2*S*,3*S*,4*E*)-2-(*N,N*-dibenzyl)-4-octadecene-1,3-diol (*L*-*threo*-sphingosine) *syn*-**3** was achieved by *syn*-addition of pentadec-1-enyl(ethyl)zinc to **1** (Scheme 1). The mixed zinc derivative was prepared as previously described⁸ by transmetalation with diethylzinc of the hydroboration product of 1-pentadecyne with dicyclohexylborane. The reaction of **1** with two equivalents of the zinc derivative in toluene–heptane at 0°C for 4 h gave the protected sphingosines **2** in 57% yield as a mixture (4:1) of epimers where the major diastereomer was *syn*-**2** as demonstrated by ¹H NMR analysis after separation of the mixture by flash chromatography (silica gel, toluene:hexane=3:1). The final *threo*-*N,N*-dibenzyl sphingosine (*syn*-**3**) was obtained in 78% yield from *syn*-**2** by treatment with TBAF in THF at 0°C for 12 h.

An alternative method allowed the preparation of (*D*)-*erythro*-*N,N*-dibenzyl sphingosine (*anti*-**3**) from the serinal derivative **1**. The reaction of **1** with pentadecynyl magnesium bromide⁹ at 0°C for 45 min in diethyl ether yielded *anti*-**4** as a single diastereomer in 82% yield after purification by flash chromatography.

Treatment of *anti*-**4** with LAH in THF at 0°C quantitatively removed the TBDMS protecting group leading to the propargyl amino diol *anti*-**5** in 80% yield after purification. However, when a mixture of *anti*-**4** and LAH was refluxed in THF for 6 h, a concomitant deprotection and reduction of the triple bond led to a mixture of (*D*)-*erythro*-*N,N*-dibenzyl sphingosine (*anti*-**3** and *anti*-**5**). Separation by flash chromatography allowed *anti*-**3** to be isolated in 62% yield and *anti*-**5** to be recovered in 17% yield, which was available for recycling.^{3c}


The stereochemistry of the final products, *syn*-**3** and *anti*-**3**, was determined by ¹H NMR spectroscopy. To this end, *syn*-**3** and *anti*-**3** were transformed into the dioxane derivatives *cis*-**6** and *trans*-**6** respectively by reaction with 2,2-dimethoxypropane (acetone, cat. TsOH, 8 h, reflux). The coupling constant¹⁰ (^{1,3}J=4.1 Hz) between H-4 (m, δ=4.43) and H-5 (m, δ=2.39) in *cis*-**6** showed a *cis* relationship of these protons, and consequently for the substituents at C-4 and C-5. However, the *trans* relationship for H-4 (dd, δ=4.36) and H-5 (m, δ=2.82) was deduced from their coupling constant¹⁰ (^{1,3}J=9.8 Hz) in *trans*-**6**.

The synthesis of *N,N*-dibenzyl-sphingosines presented here has several advantages. The stereochemistry at C-2 is introduced directly from *L*-serine. This stereocenter is then used to induce the stereochemistry at C-3. All four stereoisomeric *N,N*-dibenzyl sphingosines can be prepared in two steps and high yield by a single synthetic strategy depending only on the choice of starting amino aldehyde (*D* or *L*) and the alkylating agent.

3. Experimental

3.1. General

The reactions were carried out in oven-dried glassware, under an argon atmosphere, and using anhydrous solvents. Diethylzinc, as a 1 M solution in heptane, and borane–methyl sulfide complex were purchased from Aldrich. The ¹H NMR (300 MHz) and ¹³C NMR (75 MHz) spectra were registered on a Bruker AC 300 or Bruker AMX 300, using TMS as the internal standard. IR spectra were recorded on



Scheme 1.

a Philips PU 9706 spectrometer, as a film or by KBr dispersion. Optical rotations were measured on a Perkin-Elmer 241 polarimeter in a 1 dm cell.

3.2. Reaction of **1** with pentadec-1-enyl(ethyl)zinc

Cyclohexene (0.41 ml, 4 mmol) was added under Ar at 0°C to a stirred 1 M solution of borane–methyl sulfide complex (0.20 ml, 2 mmol) in toluene (2 ml). After 3 h at 0°C, 1-pentadecyne (417 mg, 0.53 ml, 2 mmol) was added, and the mixture was stirred at r.t. for 1 h. Then, the solution was cooled to –78°C and Et₂Zn 1 M solution in heptane (2.1 ml, 2.1 mmol) was added to it. The mixture was then placed in an ice-bath and the chiral aldehyde **1** (383 mg, 1 mmol) in toluene (4 ml) was added. The mixture was stirred for 4 h, after which it was treated with saturated aq. NH₄Cl solution (30 ml) to quench the reaction. The mixture was then extracted with ether (2×20 ml) and the combined extracts were washed with brine, dried (Na₂SO₄) and evaporated under reduced pressure. Purification of the residue by flash-

chromatography (silica gel, toluene:hexane=3:1) afforded *syn-2* (274 mg, 0.46 mmol; 46%) and *anti-2* (66 mg, 0.11 mmol; 11%).

3.3. (2S,3S,4E)-2-(N,N-Dibenzylamino)-1-O-(tert-butyldimethylsilyl)-4-octadecene-1,3-diol (*syn-2*)

Colorless oil. $[\alpha]_D^{23}=+45.4$ (c 1.15, CHCl₃). IR (film): 3400, 2910, 1450, 1250, 1090, 970, 835, 750, 700 cm⁻¹. ¹H NMR δ: 0.09 and 0.11 (2s, 6H); 0.88 (t, 3H, J=7.0 Hz); 0.96 (s, 9H); 1.25 (br s, 22H); 1.98 (m, 2H); 2.63 (m, 1H); 3.68 (d, 2H, J=13.2 Hz); 3.78 (dd, 1H, J₁=11.2 Hz, J₂=6.8 Hz); 3.86 (dd, 1H, J₁=11.2 Hz, J₂=3.1 Hz); 3.97 (m, 1H); 4.00 (d, 2H, J=13.2 Hz); 4.35 (br s, 1H); 5.18 (dd, 1H, J₁=15.3 Hz, J₂=8.0 Hz); 5.68 (dt, 1H, J₁=15.3 Hz, J₂=6.8 Hz); 7.20–7.40 (m, 10H). ¹³C NMR δ: -5.7; -5.6; 14.11; 8.1; 22.7; 25.9; 29.1; 29.2; 29.3; 29.5; 29.6; 29.7; 31.9; 32.3; 54.5; 59.6; 63.3; 68.6; 127.1; 128.4; 129.1; 129.9; 135.3; 139.2.

3.4. (2S,3R,4E)-2-(N,N-Dibenzylamino)-1-O-(tert-butyldimethylsilyl)-4-octadecene-1,3-diol (*anti-2*)

Colorless oil. $[\alpha]_D^{23}=-5.8$ (c 0.65, CHCl₃). IR (film): 3400, 2910, 2840, 1450, 1250, 1070, 830, 740, 690 cm⁻¹. ¹H NMR δ: 0.09 and 0.11 (2s, 6H); 0.88 (t, 3H, J=7.0 Hz); 0.92 (s, 9H); 1.25 (br s, 22H); 2.05 (m, 2H); 2.85 (m, 1H); 3.66 (d, 2H, J=13.6 Hz); 3.88 (d, 2H, J=13.6 Hz); 3.90 (dd, 1H, J₁=10.4 Hz, J₂=6.1 Hz); 4.00 (dd, 1H, J₁=10.4 Hz, J₂=5.8 Hz); 4.26 (m, 1H); 5.48 (dd, 1H, J₁=15.3 Hz, J₂=6.8 Hz); 5.72 (dt, 1H, J₁=15.3 Hz, J₂=6.6 Hz); 7.20–7.40 (m, 10H). ¹³C NMR δ: -5.6; 14.1; 18.1; 22.7; 25.9; 27.5; 29.2; 29.4; 29.7; 31.9; 32.4; 55.3; 61.7; 72.3; 127.0; 128.2; 128.9; 131.3; 132.4; 140.0. MS (m/z, %): 593.75 (M⁺, 0.63), 430 (33), 354 (100), 262 (15), 222 (17), 181 (34), 132 (29), 91 (77).

3.5. (2S,3S,4E)-2-(N,N-Dibenzylamino)-4-octadecene-1,3-diol (*syn-3*)

Tetrabutylammonium fluoride (142 mg, 0.45 mmol) in THF (0.5 ml) was slowly added to a solution of *syn-2* (178 mg, 0.3 mmol) in THF (3 ml) at 0°C. The mixture was stirred overnight at 0°C, and the reaction was quenched by the addition of water (3 ml). The aqueous phase was extracted with ether (3×10 ml), and the combined organic extracts were washed with brine, dried (Na₂SO₄), concentrated and flash-chromatographed (ethyl acetate:hexane=1:2) to yield *syn-3* (112 mg, 0.23 mmol; 78%) as a pure compound.

Colorless solid, m.p. 60–61°C (from hexane). $[\alpha]_D^{23}=+19.4$ (c 1, CHCl₃). IR (film): 3370, 2900, 2830, 1440, 955, 750, 690 cm⁻¹. ¹H NMR δ: 0.88 (t, 3H, J=7.0 Hz); 1.24 (br s, 22H); 1.99 (m, 2H); 2.73 (m, 1H); 3.48 (br s, 2H); 3.70 (m, 2H); 3.72 (d, 2H, J=13.1 Hz); 3.95 (d, 2H, J=13.1 Hz); 4.07 (m, 1H); 5.30 (dd, 1H, J₁=15.3 Hz, J₂=8.1 Hz); 5.73 (dt, 1H, J₁=15.3 Hz, J₂=6.8 Hz); 7.20–7.35 (m, 10H). ¹³C NMR δ: 14.1; 22.6; 29.0; 29.2; 29.3; 29.4; 29.5; 29.6; 31.9; 32.2; 54.2; 59.2; 63.2; 71.0; 127.2; 128.4; 129.2; 130.2; 135.4; 139.0. MS (m/z, %): 479.75 (M⁺, 0.79), 240 (100), 181 (43), 148 (14), 91 (80), 70 (20). Anal. calcd for C₃₂H₄₉NO₂: C, 80.11; H, 10.29; N, 2.92. Found: C, 80.06; H, 10.18; N, 3.25.

3.6. Reaction of **1** with pentadec-1-ynyl magnesium bromide

1-Pentadecyne (375 mg, 1.8 mmol) was dissolved in 2 ml of ether and added to a previously prepared solution of 1 M EtMgBr in ether (2 ml, 2 mmol) after which the solution was refluxed for 2.5 h. Then, the reagent was placed in an ice-bath and aminoaldehyde **1** (517 mg, 1.35 mmol) in ether (6 ml) was added dropwise. After 45 min at 0°C, the reaction was complete and was quenched with saturated NH₄Cl solution (10 ml). The mixture was extracted with ether (2×10 ml), the combined organic layers were

washed with brine, dried over anhydrous Na_2SO_4 and the solvents evaporated under vacuum. After flash-chromatography (silica gel, hexane:ethyl acetate=15:1) *anti-4* (657 mg, 1.11 mmol; 82%) was obtained as a pure compound.

3.7. (2S,3R)-2-(N,N-Dibenzylamino)-1-O-(tert-butyldimethylsilyl)-4-octadecyne-1,3-diol (*anti-4*)

Colorless oil. $[\alpha]_{\text{D}}^{23} = +5.3$ (c 1.1, CHCl_3). IR (film): 3420, 2920, 2840, 2220 (weak), 1450, 1250, 1090, 830, 770, 740, 690 cm^{-1} . ^1H NMR δ : 0.11 and 0.12 (2s, 6H); 0.88 (t, 3H, $J=6.9$ Hz); 0.95 (s, 9H); 1.25 (br s, 20H); 1.43 (m, 2H); 2.17 (td, 2H, $J_1=6.9$ Hz, $J_2=2.0$ Hz); 3.04 (m, 1H); 3.68 (d, 2H, $J=13.1$ Hz); 3.83 (dd, 1H, $J_1=10.4$ Hz, $J_2=6.4$ Hz); 3.90 (m, 1H); 4.11 (dd, 1H, $J_1=10.4$ Hz, $J_2=6.9$ Hz); 4.20 (d, 2H, $J=13.1$ Hz); 4.35 (brs, 1H); 7.20–7.35 (m, 10H). ^{13}C NMR δ : -5.51; 14.1; 18.1; 18.8; 22.7; 25.9; 28.5; 28.9; 29.1; 29.3; 29.5; 29.6; 31.9; 55.3; 60.2; 61.3; 80.0; 86.6; 127.1; 128.3; 129.2; 139.6. MS (m/z , %): 591.30 (M^+ , 0.09), 416 (7), 354 (100), 91 (45). Anal. calcd for $\text{C}_{38}\text{H}_{61}\text{NO}_2\text{Si}$: C, 77.10; H, 10.39; N, 2.37. Found: C, 76.54; H, 10.18; N, 2.58.

3.8. LAH reduction of *anti-4*

To a stirred solution of 93 mg (2.45 mmol, 3.5 equiv.) of LiAlH_4 in 6 ml of THF at 0°C was added 415 mg (0.7 mmol, 1 equiv.) of *anti-4* in 4 ml of THF. After refluxing for 6 h, the reaction was cooled and quenched with 0.1 ml of H_2O , 0.1 ml of 15% NaOH and 0.3 ml of H_2O . The solids were removed by filtration and washed with ether. The combined organic extracts were dried over anhydrous MgSO_4 and evaporated to yield 334 mg of crude product (a 77:23 mixture of *anti-3* and *anti-5* by ^1H NMR analysis). Flash-chromatography (silica gel, hexane:ethyl acetate=4:1) gave 206 mg of *anti-3* (62%) and 53 mg of *anti-5* (17%).

3.9. (2S,3R)-2-(N,N-Dibenzylamino)-4-octadecyne-1,3-diol (*anti-5*)

Colorless oil. $[\alpha]_{\text{D}}^{23} = -38.1$ (c 1, CHCl_3). IR (film): 3350, 2890, 2830, 2220, 1450, 1020, 900, 730, 690 cm^{-1} . ^1H NMR δ : 0.89 (t, 3H, $J=7.0$ Hz); 1.25 (m, 20H); 1.48 (m, 2H); 2.22 (td, 2H, $J_1=7.1$ Hz, $J_2=2.0$ Hz); 3.04 (m, 1H); 3.79 (d, 2H, $J=13.3$ Hz); 3.85 (dd, 1H, $J_1=11.0$ Hz, $J_2=6.0$ Hz); 3.94 (dd, 1H, $J_1=11.0$ Hz, $J_2=7.7$ Hz); 4.02 (d, 2H, $J=13.3$ Hz); 4.52 (dt, 1H, $J_1=6.0$ Hz, $J_2=2.0$ Hz); 7.20–7.40 (m, 10H). ^{13}C NMR δ : 14.1; 18.8; 22.6; 28.5; 29.0; 29.1; 29.3; 29.5; 29.6; 31.9; 54.6; 59.6; 60.6; 62.0; 79.9; 87.4; 127.2; 128.4; 129.1; 139.2. MS (m/z , %): 477.30 (M^+ , 0.50), 446 (2), 282 (2), 240 (100), 181 (6), 91 (99).

3.10. (2S,3R,4E)-2-(N,N-Dibenzylamino)-4-octadecene-1,3-diol (*anti-3*)

Colorless oil. $[\alpha]_{\text{D}}^{23} = -47.0$ (c 1.1, CHCl_3). IR (film): 3360, 2920, 2850, 1450, 1025, 970, 745, 695 cm^{-1} . ^1H NMR δ : 0.88 (t, 3H, $J=6.9$ Hz); 1.26 (m, 22H); 2.50 (br s, 2H); 2.79 (m, 1H); 3.71 (d, 2H, $J=13.5$ Hz); 3.76 (m, 1H); 3.78 (d, 2H, $J=13.5$ Hz); 3.87 (dd, 1H, $J_1=11.0$ Hz, $J_2=7.0$ Hz); 4.37 (t, 1H, $J=6.8$ Hz); 5.48 (dd, 1H, $J_1=15.4$ Hz, $J_2=7.2$ Hz); 5.71 (dt, 1H, $J_1=15.4$ Hz, $J_2=6.6$ Hz); 7.20–7.40 (m, 10H). ^{13}C NMR δ : 29.1; 29.3; 29.5; 29.7; 31.9; 32.3; 54.4; 59.2; 62.3; 72.3; 127.1; 128.3; 128.9; 131.8 (C-5); 133.2; 139.5.

3.11. (2S,3S,4E)-2-(N,N-Dibenzylamino)-1,3-O-isopropylidene-4-octadecene-1,3-diol (cis-6)

Colorless oil. $^1\text{H NMR}$ δ : 0.81 (t, 3H, $J=6.2$ Hz); 1.19 (m, 22H); 1.35 (s, 3H); 1.38 (s, 3H); 2.11 (m, 2H); 2.39 (m, 1H); 3.55 (d, 2H, $J=14.4$ Hz); 3.85 (dd, 1H, $J_1=12.8$ Hz, $J_2=3.7$ Hz); 4.22 (m, 3H); 4.43 (m, 1H); 5.61 (dt, 1H, $J_1=15.5$ Hz, $J_2=6.5$ Hz); 5.83 (dd, 1H, $J=15.5$ Hz, $J=6.4$ Hz); 7.10–7.40 (m, 10H).

3.12. (2S,3R,4E)-2-(N,N-Dibenzylamino)-1,3-O-isopropylidene-4-octadecene-1,3-diol (trans-6)

Colorless oil. $^1\text{H NMR}$ δ : 0.89 (t, 3H, $J=6.9$ Hz); 1.28 (m, 22H); 1.35 (s, 3H); 1.46 (s, 3H); 2.15 (m, 2H); 2.82 (m, 1H); 3.64 (d, 2H, $J=13.9$ Hz); 3.86 (dd, 1H, $J_1=11.8$ Hz, $J_2=5.7$ Hz); 3.89 (d, 2H, $J=13.9$ Hz); 3.96 (dd, 1H, $J_1=11.8$ Hz, $J_2=8.3$ Hz); 4.36 (dd, 1H, $J_1=9.8$ Hz, $J_2=7.8$ Hz); 5.41 (dd, 1H, $J_1=15.4$ Hz, $J_2=7.8$ Hz); 5.87 (dt, 1H, $J_1=15.4$ Hz, $J_2=6.6$ Hz); 7.20–7.40 (m, 10H).

Acknowledgements

We thank the Spanish DGICYT (Project PB95-707) and the Junta de Castilla y León (Project VA42/97) for the financial support of this work.

References

- (a) Hannun, Y. A.; Loomis, C. R.; Merrill Jr., A. H.; Bell, R. M. *J. Biol. Chem.*, **1986**, 261, 12604. (b) Hannun, Y. A.; Ben, R. M.; Merrill, A. H. *Science*, **1989**, 243, 500. (c) Hori, T.; Sugita, M. *Progr. Lipid Res.*, **1993**, 32, 25.
- From sugars: (a) Gigg, R.; Conant, R. *J. Chem. Soc. Perkin Trans. 1*, **1977**, 2006. (b) Schmidt, R. R.; Zimmermann, P. *Tetrahedron Lett.*, **1986**, 27, 481. (c) Kumar, P.; Schmidt, R. R. *Synthesis*, **1998**, 33. Sharpless monoepoxidation of an enynol: (d) Julina, R.; Herzig, T.; Bennet, B.; Vasella, A. *Helv. Chim. Acta.*, **1986**, 27, 481. Asymmetric aldol reaction: (e) Kobayashi, S.; Hayashi, T.; Kawasuji, T. *Tetrahedron Lett.*, **1994**, 35, 9573. (f) Solladié-Cavallo, A.; Koessler, J. L. *J. Org. Chem.*, **1994**, 59, 3240. (g) Enders, D.; Whitehouse, D. L.; Runsink, J. *Chem. Eur. J.*, **1995**, 1, 382. (h) Spanu, P.; Rasso, G.; Pinna, L.; Battistini, L.; Casiraghi, G. *Tetrahedron: Asymmetry*, **1997**, 8, 3237. From other chiral pools: (i) Katsumura, S.; Yamamoto, N.; Fukuda, E.; Iwama, S. *Chem. Lett.*, **1995**, 5, 393. (j) Murakami, T.; Hato, M. *J. Chem. Soc., Perkin Trans. 1*, **1996**, 823. (k) Tuch, A.; Sanière, M.; Le Merrer, Y.; Depezay, J. C. *Tetrahedron: Asymmetry*, **1996**, 7, 897.
- (a) Newman, H. *J. Am. Chem. Soc.*, **1973**, 95, 4098. (b) Tkaczuk, P.; Thornton, E. R. *J. Org. Chem.*, **1981**, 46, 4393. (c) Mori, K.; Funaki, Y. *Tetrahedron*, **1985**, 41, 2379. (d) Boutin, R. H.; Rapoport, H. *J. Org. Chem.*, **1986**, 51, 5320. (e) Nimkar, S.; Menaldino, D.; Merrill, A. H.; Liotta, D. *Tetrahedron Lett.*, **1988**, 29, 2307. (f) Garner, P.; Park, J. M.; Malecki, E. *J. Org. Chem.*, **1988**, 53, 4395. (g) Radunz, H. E.; Devant, R. M.; Eiermann, V. *Liebigs Ann. Chem.*, **1988**, 11, 1103. (h) Herold, P. *Helv. Chim. Acta*, **1988**, 71, 354. (i) Dondoni, A.; Fantin, G.; Fogagnolo, M.; Pedrini, P. *J. Org. Chem.*, **1990**, 55, 1439. (j) Polt, R.; Peterson, M. A.; De Young, L. *J. Org. Chem.*, **1992**, 57, 5469.
- For recent solutions see: (a) Dondoni, A.; Perrone, D.; Turturici, E. *J. Chem. Soc., Perkin Trans. 1*, **1997**, 2389. (b) Li, Y.-L.; Wu, Y. L. *Liebigs Ann.*, **1996**, 2079 and references cited therein.
- Soai, K.; Takahashi, K. *J. Chem. Soc., Perkin Trans. 1*, **1994**, 1257.
- Andrés, J. M.; Barrio, R.; Martínez, M. A.; Pedrosa, R.; Pérez-Encabo, A. *J. Org. Chem.*, **1996**, 61, 4210.
- (a) Andrés, J. M.; Pedrosa, R. *Tetrahedron*, **1998**, 54, 5607. (b) Laib, T.; Chastanet, J.; Zhu, J. *J. Org. Chem.*, **1998**, 63, 1709.
- (a) Srebnik, M. *Tetrahedron Lett.*, **1991**, 32, 2449. (b) Oppolzer, W.; Radinov, R. N. *Helv. Chim. Acta*, **1992**, 75, 170.
- 1-Pentadecyne was prepared in 75% yield by alkylation of lithium acetylide ethylenediamine complex with 1-bromotridecane in DMSO at room temperature by analogy with that described by Burgsthaler, A. W.; Weigel, L. O.; Sanders, M. E.; Shaefer, C. G. *J. Org. Chem.*, **1977**, 42, 566.
- (a) Hafner, A.; Duthaler, R. O.; Marti, R.; Rihs, G.; Rothe-Streit, P.; Schwarzenbach, F. *J. Am. Chem. Soc.*, **1992**, 114, 2321. (b) Villard, R.; Fotiadu, F.; Buono, G. *Tetrahedron: Asymmetry*, **1998**, 9, 607.