


Contents

COMMUNICATIONS

Asymmetric Negishi reaction for sterically hindered couplings: synthesis of chiral binaphthalenes

pp 2593–2595


Miroslav Genov, Beatriz Fuentes, Pablo Espinet* and Beatriz Pelaz



Two complementary, diversity-driven asymmetric syntheses of a 2,2-disubstituted piperidine NK₁ antagonist

pp 2596–2598


Dong Xiao,* Cheng Wang, Anandan Palani, Gregory Reichard, Robert Aslanian, Neng-Yang Shih and Alexei Buevich




Resolution of 3-methyl-3-phospholene 1-oxides by molecular complex formation with TADDOL derivatives

pp 2599–2602

Tibor Novák,* József Schindler, Viktória Ujj, Mátyás Czugler, Elemér Fogassy and György Keglevich



Shravan K. Srisailam and Antonio Togni*




ARTICLES

Stereoselective synthesis of the polyketide chain of nagahamide A


pp 2609–2616

Debendra K. Mohapatra,* Siddhartha Ray Chaudhuri, Gokarneswar Sahoo and Mukund K. Gurjar*

Enantioselective diethylzinc addition to the exocyclic C=N double bond of some 4-arylideneamino-3-mercapto-6-methyl-4*H*-1,2,4-triazin-5-one derivatives

pp 2617–2624

Ashraf A. El-Shehawy




Chiral 4-(1-arylpropyl)amino-3-mercapto-6-methyl-4*H*-1,2,4-triazin-5-ones **4** were synthesized easily through the enantioselective diethylzinc addition to the exocyclic C=N double bond of 4-arylideneamino-3-mercapto-6-methyl-4*H*-1,2,4-triazin-5-ones **2**. Enantiomeric excess of up to 92% was obtained using (1*S*,2*R*)-*N*-methyl-*N*-benzylnorephedrine **3b** as chiral ligand and TIPSCl as activator.

Intermolecular one-pot cyclization of formyl-pyrroles of amino acid esters with norephedrine: stereoselective routes to new tricyclic pyrrole-pyrazine-oxazole fused structures

pp 2625–2631


Ayhan S. Demir,* N. Tuna Subasi and Ertan Sahin



A short and efficient synthesis of (*R*)-(-)-sporochnol A

pp 2632–2636

Ramón Alibés, Félix Busqué,* Gisela G. Bardají, Pedro de March, Marta Figueredo and Josep Font




A short and efficient synthesis of (*R*)-(-)-sporochnol A in only five steps and 9% overall yield has been developed.

Stereoselective synthesis of δ -heteroaryl substituted β -hydroxy- γ,δ -unsaturated α -amino acids

pp 2637–2641


Giuseppe Cremonesi, Piero Dalla Croce, Francesco Fontana and Concetta La Rosa*



Chiral NMR discrimination of pyrrolidines using (18-crown-6)-2,3,11,12-tetracarboxylic acid


pp 2642–2648

Ann E. Lovely and Thomas J. Wenzel*



Increasing the enantioselectivity of cyclopentanone monooxygenase (CPMO): profile of new CPMO mutants pp 2649–2653


Christopher M. Clouthier and Margaret M. Kayser*



Synthesis of a Schiff's base chiral ligand with a trifluoromethyl carbinol moiety

pp 2654–2658


Yasser Samir Sokeirik, Masaaki Omote, Kazuyuki Sato, Itsumaro Kumadaki and Akira Ando*



Asymmetric addition of trimethylsilyl cyanide to aldehydes promoted by chiral polymeric vanadium(V) salen complex as an efficient and recyclable catalyst

pp 2659–2666


Noor-ul H. Khan,* Santosh Agrawal, Rukhsana I. Kureshy, Sayed H. R. Abdi, Vishal J. Mayani and Raksh V. Jasra



Multicomponent diversity and enzymatic enantioselectivity as a route towards both enantiomers of α -amino acids—a model study

pp 2667–2671


Wiktor Szymanski and Ryszard Ostaszewski*



Complete reversal of enantioselection using oxazoline-containing Schiff base ligands derived from L-serine in enantioselective addition of diketene to aldehydes

pp 2672–2677


Changhu Chu, Koji Morishita, Takanori Tanaka and Masahiko Hayashi*



Synthesis of chiral hydroxylated cyclopentanones and cyclopentanes

pp 2678–2683


Allan Niidu, Anne Paju, Margus Eek, Aleksander-Mati Müürisepp, Tõnis Pehk and Margus Lopp*



Straightforward chemo-enzymatic synthesis of new aminocyclitols, analogues of valiolamine and their evaluation as glycosidase inhibitors

pp 2684–2688


Lahssen El Bliidi, Mustapha Ahbala, Jean Bolte and Marielle Lemaire*



 $\text{Rh}_2[(R)\text{-}(+)\text{-MTPA}]_4$ as an NMR auxiliary for the enantiodifferentiation of chiral secondary and tertiary phosphine–borane complexes

pp 2689–2696


Jens Mattiza, Dieter Albert, Marek Stankevič, Kamil Dziuba, Anna Szmigielska, K. Michał Pietrusiewicz* and Helmut Duddeck*



An approach to the synthesis and assignment of the absolute configuration of all enantiomers of ethyl hydroxy(phenyl)methane(P-phenyl)phosphinate

pp 2697–2701


Paulina Majewska,* Paweł Kafarski, Barbara Lejczak, Iwona Bryndal and Tadeusz Lis



Isolation synthesis and glycosidase inhibition profile of 3-*epi*-casuarine

pp 2702–2712


Jeroen Van Ameijde, Graeme Horne, Mark R. Wormald, Raymond A. Dwek, Robert J. Nash, Paul Wyn Jones, Emma L. Evinson and George W. J. Fleet*



Lipase-catalyzed resolution of β -hydroxy selenides

pp 2713–2721

Michelangelo Gruttadauria,* Paolo Lo Meo, Serena Riela, Francesca D'Anna and Renato Noto

**OTHER CONTENTS**

Corrigendum

p 2722

Stereochemistry abstracts

pp A575–A603

Instructions to contributors

pp I–IV

Cumulative author index

pp V–XI

*Corresponding author

Available online at www.sciencedirect.com

Indexed/Abstracted in: Beilstein, BIOSIS Previews, Chemical Abstracts, Current Contents: Physical, Chemical and Earth Sciences, Derwent Biotechnology Abstracts, Derwent Drug File, Ei Compendex, EMBASE/Excerpta Medica, PASCAL, Research Alert, Science Citation Index, SciSearch. Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®



ELSEVIER

ISSN 0957-4166