

Corrigendum

Corrigendum to “New efficient synthesis of Taniaphos ligands: application in ruthenium- and rhodium-catalyzed enantioselective hydrogenations”


[*Tetrahedron: Asymmetry* 15 (2004) 91]

Katja Tappe and Paul Knochel*


Department of Chemistry, Ludwig-Maximilians-University, Butenandtstr. 5-13, 81377 München, Germany

Available online 15 April 2008

Following a remark by Fukuzawa et al.¹ we confirm that the stereochemistry of Taniaphos (**19**, Scheme 6) is not (*R,S*_{Fe}), as mentioned in the original communication, but (*R,R*_{Fe}), as it is indicated by the X-ray structure of the [Rh(nbd)(**2a**)]BF₄ complex, included in the previous manuscript.² Thus, the ferrocene derivatives **19** and **1** (Scheme 1) possess (*R,R*_{Fe}) configuration.



Scheme 1. (Revised).



Scheme 6. (Revised).

References

1. Fukuzawa, S.; Yamamoto, M.; Hosaka, M.; Kikuchi, S. *Eur. J. Org. Chem.* **2007**, 5540.
2. Ireland, T.; Grossheimann, G.; Wieser-Junesse, C.; Knochel, P. *Angew. Chem., Int. Ed.* **1999**, 38, 3212.

DOI of original article: 10.1016/j.tetasy.2003.11.004

* Corresponding author. Tel.: +49 0 89 2180 77681; fax: +49 0 89 2180 77680; e-mail: paul.knochel@cup.uni-muenchen.de