

CONFORMATIONAL EFFECTS IN THE OPENING OF  
CIS/TRANS 1,4-DIALKYL SUBSTITUTED 1,2-  
CYCLOHEXENE EPOXIDES WITH ACETIC ACID


John C. Leffingwell<sup>1</sup> and E. Earl Royals

Department of Chemistry






Emory University, Atlanta, Georgia

(Received 23 August 1965)



In our continuing studies on the reactions of  
terpene oxides<sup>2</sup> we wish to report the following cases  
of stereospecific epoxide openings in the p-menthane  
series.



The peracid epoxidation of limonene (I), menthene-1  
(II) and menthene-3 (III) leads, in each case, to  
approximately a 1:1 mixture of the corresponding cis  
and trans epoxides<sup>3</sup> as would be expected from the  
recent findings of Rickborn and Lwo<sup>3d</sup>. Although  
the cis/trans epoxide mixtures are separable by dis-  
tillation only with difficulty, they may be separated  
conveniently on a 15 ft. 15% Carbowax 20M on chro-  
sorb W GLPC column<sup>4</sup>. In each of the three cases cited,  
reaction of the cis epoxide (IVa,b,c) with a sodium





acetate buffered solution of acetic acid leads exclusively to a single product containing trans tertiary acetoxy and secondary hydroxyl groups and trans 1,4-alkyl groups (VII a,b,c), as would be expected by an extension of the Fürst-Plattner rule<sup>5</sup> to monocyclic systems<sup>3a</sup>, assuming that  $R_2$  "fixes" the conformation of the cyclohexane ring system. Similarly, reaction of the



trans epoxides (Va,b,c) leads predominantly<sup>6</sup> to epoxide opening resulting from cleavage of the "equatorial"-C-O

linkage in the transition state (VIII) to afford the diaxial products (IXa,b,c) with secondary acetoxy and tertiary hydroxyl groups. The proof of structure in each case follows from pyrolysis of the acetate to yield known products<sup>2,3a</sup>. For example, ( $\pm$ )-cis-3-menthene epoxide (IVc) affords a hydroxyacetate (VIIc) which pyrolyzes at 390° to a mixture of ( $\pm$ )-trans-3-menthene-5-ol (X)<sup>7a</sup> and ( $\pm$ )-menthone-isomenthone (XI) in




a product ratio, 7:3. The ( $\pm$ )-trans-3-menthene epoxide affords a crystalline ( $\pm$ )-hydroxyacetate (IXc), m.p. 88.5-89.5°, which pyrolyzed at 450° to ( $\pm$ )-trans-2-

menthene-4-ol<sup>7b</sup> (XII). In each case, the infrared spectra of the pyrolysis products were superposable on those of authentic samples<sup>7a,b</sup>. Further structure proof of (X) came from hydrogenation (PtO<sub>2</sub> - ethyl acetate) to a mixture of 88% ( $\pm$ )-isomenthol (XIII) and 12% ( $\pm$ )-neomenthol (XIV). Hydrogenation of (XII) afforded trans-p-menthan-4-ol (XV).

The high degree of stereospecificity in these epoxide openings leads us to propose the following:

- (1) The appreciable amount of abnormal, "diequatorial", opening observed with nucleophilic reagents on monocyclic cis-1,4-dialkyl cyclohexene epoxides (vide infra) does not necessarily arise from reaction of the less stable conformational isomer<sup>8</sup> (as is observed in conformationally mobile cyclohexene epoxide systems<sup>9</sup>) but rather, perhaps, from the innate difficulty of nucleophilic displacements at tertiary centers<sup>10</sup>.
- (2) The 4-alkyl substituent determines the conformation in which the cis- and trans-<sup>6</sup> 1,4-alkyl substituted cyclohexene-1,2-epoxides react with acidic reagents (i.e., HOAc) via a normal, "diaxial", opening (Fürst-Plattner Rule<sup>5</sup>).


Thus the lithium aluminum hydride (LAH) reduction of (+)-cis-1,2-limonene epoxide (IVa)<sup>3a, 11a</sup> affords a product ratio of 15% cis- $\beta$ -terpineol (XVI) via "effective diequatorial opening" and 85% (+)-neodihydrocarveol (XVII) via "normal diaxial opening". With other nucleophilic reagents, i.e., dimethylamine, the percentage of "diequatorial" opening is even greater (30-40%)<sup>11b</sup>.



There are two ways in which such diequatorial openings may occur; (a) through diaxial opening of the less stable epoxide conformer with subsequent inversion (process "A") or (b) via diaxial opening of a "twist" transition state (process "B"). Inasmuch as there should be a considerable steric (shielding) effect exhibited by the "axial" 4-alkyl group on the incoming nucleophile in process "A", in addition to the already energetically unfavorable epoxide conformation, we tend to favor

process "B".

In the opening of the cis-dialkyl epoxide (VIc) with acetic acid, although  $R_1$  is isopropyl and  $R_2$  is methyl, only a single product is obtained (governed by diaxial opening with  $R_2$  equatorial). Therefore, even though the 1,4-dialkyl substituted cyclohexene epoxide has been postulated<sup>12</sup> to assume a "nearly chair" conformation in the reaction transition state, the 1-alkyl substituent plays little or no part in determining the conformational preference of the transition state<sup>13</sup>. It is probable, therefore, that the transition states for the reactions of the cis/trans-1,4-dialkyl cyclohexene epoxides as illustrated above (VI and VIII) are a bit exaggerated, since to develop a true "chair" conformation the C-O bond must effectively be broken (or at the least very stretched). Transition states such as (VIId) and (VIIIId) might be better representations. In such transition states, as the protonated oxirane starts to assume the status of an axial hydroxyl there is still effective overlap between the p-orbital of oxygen and the vacant p-orbital of the "incipient carbonium ion". Since  $R_1$  in such a transition state for the cis epoxide (VI) can be neither axial nor equatorial it should not have any effect on conformational preference (as indeed is the case). Such overlap would be less favorable for a



diequatorial opening (which is not observed for the cis epoxides of type IV). The small percentage (ca. 10%) of an "effective diequatorial opening" observed for the trans epoxides (Va and Vb) may occur by either a process of type "A" or "B" (vide supra). Although the results presented do not distinguish between these two mechanisms it is again feasible that process "B" may function inasmuch as the normal diaxial opening of a "twist" conformer of (V) would be enhanced by formation of an incipient tertiary positive carbon center.

In any event, the stereospecific opening of 4-alkyl substituted cyclohexene epoxides, coupled with the convenient pyrolysis of the resultant hydroxyacetates affords an excellent method for the preparation of specific allylic alcohols.

Acknowledgements. The authors wish to thank Dr. Gunther Ohloff and Dr. J. Gresham for valuable comments on various parts of this work and especially Dr. Bruce Rickborn who contributed insight into the discussion of the possible transition states.

REFERENCES

- (1). Present address, R. J. Reynolds Tobacco Co., Research Center, Winston-Salem, North Carolina.
- (2). E. E. Royals and J. C. Leffingwell, J. Am. Chem. Soc., 86, 2067(1964); J. Org. Chem., 29, 2098(1964); in press.
- (3). (a) E. E. Royals and J. C. Leffingwell, in press; (b) A. B. Booth, U. S. Pat. 2,945,068 (1960); (c) W. F. Newhall, J. Org. Chem., 29, 185(1964); (d) B. Rickborn and S.-Y Lwo, J. Org. Chem., 30, 2212(1965).
- (4). J. C. Leffingwell, unpublished results; Ref. 3c, footnote 5.
- (5). A. Furst and P. A. Plattner, Abstracts of Papers, 12th International Congress of Pure and Applied Chemistry, New York, 1951, p. 409; see also, D. H. R. Barton, J. Chem. Soc., 1953, 1027.
- (6). In the case of (Va,b) ca. 10% of the abnormal product arising from an effective "diequatorial" opening was observed; see Ref. 3a.
- (7). (a) D. Malcolm and J. Read, J. Chem. Soc., 1939, 1037; (b) G. Ohloff and G. Uhde, Helv. Chim. Acta., 48, 10(1965).
- (8). S. J. Angyal, Chem. and Ind., 1954, 1230; Quart. Revs., 11, 212(1957); see also R. C. Cookson, Chem. and Ind., 1954, 223, 1512.
- (9). For examples see Ref. 3a,c; W. F. Newhall, J. Org. Chem., 24, 1637(1959); 29, 185(1964); H. Kuczyński and A. Zabza, Roczniki Chem., 37, 733(1963).
- (10). E. L. Eliel, Stereochemistry of Carbon Compounds, McGraw Hill Co., New York, 1962, p. 290.
- (11). (a) E. Kuczyński and K. Piatkowski, Roczniki Chem., 33, 299, 311(1959); (b) H. Kuczyński and A. Zabza, Bull. Acad. Polon. Sci., Ser. sci. chem., 9, 551(1961); Roczniki Chem., 37, 773 (1963).
- (12). R. E. Parker and N. S. Isaacs, Chem. Revs., 59, 742(1959).



- (13). The term 1-alkyl and 4-alkyl substituent refers to the systematic naming of cyclohexane systems in general. Therefore, in each of the cases cited here, the alkyl substituent substituted on the epoxide bearing carbon is the 1-alkyl substituent. The fact that the isopropyl group in the 3-menthene epoxides is on the 4-carbon atom according to nomenclature in the p-menthane systems thus should not confuse the reader in the interpretation of the general effect.