

SOME NOVEL PERFLUOROALKANEDIOIC ACID DERIVATIVES AND  $\alpha,\omega$ -DI-IODOPERFLUOROALKANES


V. C. R. McLoughlin

Materials Department, Royal Aircraft Establishment, Farnborough, Hampshire.

(Received in UK 12 July 1968; accepted for publication 16 August 1968)

Perfluoroalkanedioic acids,  $\text{HO}_2\text{C}(\text{CF}_2)_n\text{CO}_2\text{H}$  (I), and  $\alpha,\omega$ -di-iodoperfluoroalkanes,  $\text{I}(\text{CF}_2)_n\text{I}$  (II), where  $n = 1,3,5,7$ , etc. were hitherto unknown for values of  $n \geq 5$ . In contrast, the corresponding compounds where  $n = 2,4,6,8$ , etc. are available by several routes (1,2,3,4). It has now been found possible to derive the compounds where  $n$  is odd from the acid chlorides III, which are readily available (5).

Fluorocarbons containing terminal difluoromethyl groups can be oxidised to fluoro-carboxylic acids by reaction with mixtures of nitrogen dioxide and chlorine in the gas phase (6). This technique has now been applied to the acid chlorides III when  $n+1 = 4,8$ , and 16,



and the crude oxidation products (IV) were converted into either dimethyl perfluoroalkanedioates (V) or perfluoroalkanedioic acids (I); the acids were then converted into perfluoroalkanedioyl dichlorides (VI). The overall yields of V or VI were generally between 30 and 50%, the major by-products being derivatives of the perfluoroalkanedioic acids  $\text{HO}_2\text{C}(\text{CF}_2)_{n-1}\text{CO}_2\text{H}$ , obtained in 10 to 15% yields. Some physical constants of the esters V (characterised by elemental analysis and mass spectroscopy) are shown in the Table.

Fluoroalkanyl chlorides react with potassium iodide at  $200^\circ$  in a sealed vessel to give fluoroalkyl iodides (7). It has now been found that the passage of fluoroalkanyl chlorides through a glass tube packed with crystalline potassium iodide (dried at  $200^\circ$ ) at temperatures between  $270$  and  $375^\circ$ , gives fluoroalkyl iodides and carbon monoxide. When a perfluoroalkanedioyl dichloride of the type VI is treated in this manner, the product consists of an  $\alpha,\omega$ -di-iodoperfluoroalkane (II), an  $\omega$ -iodoperfluoroalkanyl chloride (VII), and, at the lower temperatures of operation, the starting material (VI). The ratio of the

products can be controlled by the rate of addition of the acid chloride (VI), the temperature of the tube, and by recycling the mixture. On several occasions, VI,  $n = 3$  (ca 500g) was so treated giving II,  $n = 3$  (50 - 60%) and VII,  $n = 3$  (5 - 20%), the latter isolated as the free acid. On a smaller scale, VI,  $n = 7$  containing VI,  $n = 6$ , and VI,  $n = 9$  containing VI,  $n = 8$  were converted into the corresponding mixtures of di-iodides II. Some physical constants of these products (characterised by elemental analysis) are given in the Table below.

| Compound  | Boiling point   | Melting point | Refractive index, $n_D^{20}$ |
|---|-----------------|---------------|------------------------------|
| $\text{MeO}_2\text{C}(\text{CF}_2)_6\text{CO}_2\text{Me}$ | 236° a          |               | 1.3415                       |
| $\text{MeO}_2\text{C}(\text{CF}_2)_7\text{CO}_2\text{Me}$ | 249°            |               | 1.3400                       |
| $\text{MeO}_2\text{C}(\text{CF}_2)_8\text{CO}_2\text{Me}$ | 261° b          |               | 1.3394                       |
| $\text{MeO}_2\text{C}(\text{CF}_2)_9\text{CO}_2\text{Me}$ | 275°            | 31-2°         | 1.3314 c                     |
| $\text{I}(\text{CF}_2)_3\text{CO}_2\text{H}$              | 107-8°/35 mm Hg | 31-4°         |                              |
| $\text{I}(\text{CF}_2)_6\text{I}$                         |                 | 25°           |                              |
| $\text{I}(\text{CF}_2)_7\text{I}$                         |                 | 51-2°         |                              |
| $\text{I}(\text{CF}_2)_8\text{I}$                         |                 | 75-6°         |                              |
| $\text{I}(\text{CF}_2)_9\text{I}$                         |                 | 94-5          |                              |

a 126-7°/25 mm Hg (2)

b 144-6°/25 mm Hg (2)

c at 36°

## REFERENCES

1. I. L. Knunyants, Chih-Yuan Li, and V. V. Shokina, Izvest.Akad.Nauk S.S.S.R., Otdel.Khim. Nauk, 1462(1961); Chem.Abs., 56, 302(1962).
2. R. B. Ward, J.Org.Chem., 30, 3009 (1965).
3. Y. K. Khim and O. R. Pierce, ibid. 33, 442 (1968).
4. R. N. Haszeldine, Nature, 167, 139 (1951).
5. R. M. Joyce, Jr., U.S. 2,559,628 (1951); K. L. Berry, U.S. 2,559,629 (1951); O. H. Bullitt, Jr., U.S. 2,559,630 (1951)(Chem.Abs. 46, 3063-4 (1952)).
6. W. A. Severson and T. J. Brice, J.Amer.Chem.Soc., 80, 2313 (1958).
7. C. G. Krespan, J.Org.Chem., 23, 2016 (1958).