


ELECTROPHILIC AROMATIC SUBSTITUTION ·  
ELECTRON TRANSFER ROUTES IN SIDE CHAIN SUBSTITUTION

by Jay K Kochi

Department of Chemistry, Indiana University, Bloomington, Indiana 47401

(Received in USA 19 September 1974; received in UK for publication 26 November 1974)


In a continuing series of interesting reports, Baciocchi, Illuminati and coworkers have described the side chain substitution of alkyl-aromatics under a variety of electrophilic conditions<sup>1</sup> For example, hexamethylbenzene is converted to pentamethylbenzyl chloride by chlorine in acetic acid in the dark<sup>2</sup> The scope of these non-conventional substitution reactions comprise a variety of electrophilic reactions including halogenation, (chlorine, bromine and iodine), nitration and thiocyanation which proceed generally on polyalkylbenzenes and other electron-rich aromatics such as thiophenes, etc<sup>3-4</sup>



atics such as thiophenes, etc<sup>3-4</sup>



The heterolytic mechanism in Scheme I which they propose for side chain substitution under electrophilic conditions include first, the formation of a  $\sigma$ -adduct I between the electrophile and arene in eq 2, followed by either a stepwise or synchronous  $\alpha$ -proton loss and rearrangement (eq 3), e.g.,<sup>2</sup>

Scheme I



It has recently been shown, however, that aromatic substitution, either in the nuclear or side chain positions, may proceed by electron-transfer processes<sup>5</sup> The electron transfer steps



in Scheme II have been effected either electrochemically or by metal oxidants such as cobalt(III), manganese(III), cerium(IV) capable of effecting one-electron changes <sup>5,6</sup> The formation of Scheme II



the aromatic cation-radical II is the key to the electron transfer mechanism, since side chain substitution follows directly from the loss of an  $\alpha$ -proton as shown in eq 5 Two factors are important in facilitating the electron transfer process in Scheme II a relatively low oxidation potential for the aromatic compound and the availability of a one-electron oxidant We feel that both conditions are fulfilled in a number of the side chain substitution reactions reported by Illuminati et al In particular, many of the electrophiles available for aromatic substitution are also capable of performing as one-electron oxidants Thus, chlorine, bromine and iodine readily undergo one-electron reduction and are also highly effective electron acceptors in charge transfer complexes with a variety of arenes <sup>7</sup> Similarly, nitronium ions involved directly or indirectly in nitration have a reasonable high reduction potential relative to nitrogen dioxide <sup>8</sup>

We wish to show that chlorination of hexamethylbenzene under conditions described by Illuminati et al may proceed by an electron transfer mechanism Thus, acetic acid solutions of chlorine and hexamethylbenzene were mixed directly in the cavity of an electron spin resonance spectrometer The intense and well-resolved esr spectrum of hexamethylbenzene cation-radical shown in Figure 1 was readily obtained at room temperature Other arenes such as pentamethylbenzene and durene showed similar behavior The facile formation of the radical-cation is consistent with an electron-transfer mechanism described in Scheme III

Scheme III






Figure 1 Esr spectrum of the hexamethylbenzene radical-cation (central portion) obtained during the reaction of hexamethylbenzene and chlorine in acetic acid at 25 °C. Proton nmr field markers are in kHz.

The electron transfer mechanism in Scheme III involves readily accessible intermediates and avoids the postulation of cyclohexadienes and their rearrangement. A rigorous distinction between heterolytic and electron transfer pathways in Schemes I and III, respectively, is difficult to make since electrophilic addition (eq 2) and electron transfer (eq 7) have much the same electronic requirements.<sup>5b</sup> In some electrophilic substitutions, such as proton exchange,<sup>9</sup> acceptors are unavailable for an electron transfer process to occur, and the reaction must perforce take a heterolytic pathway.

#### References

- 1 E Bacciocchi, P Faggioni and G Illuminati, *Gazz Chim Ital* , 103, 1311 (1973)
- 2 E Bacciocchi, A Ciana, G. Illuminati and C Pasini, *J Amer Chem Soc* , 87, 3953 (1965)
- 3 (a) G Illuminati, L Mandolini and A Patara, *Tetrahedron Letters*, 4161 (1972)  
 (b) G Illuminati, L Mandolini, E M Arnett and R Smoyer, *J Chem Soc* , B, 2206 (1971)  
 (c) E Bacciocchi and G Illuminati, *Tetrahedron Letters*, 637 (1962)
- 4 (a) L J Andrews and R M Keefer, *J Amer Chem Soc* , 86, 4158 (1964)  
 (b) K Nakamura, *Bull Chem Soc* , Japan, 44, 133 (1971)  
 (c) H Suzuki, *Ibid* , 43, 879 (1970)  
 (d) R Astolfi, E Bacciocchi and G Illuminati, *La Chim e l'Ind* , 53, 1153 (1971)
- 5 (a) E I Heiba, R M Dessau and W J Koehl, *J Amer Chem Soc* , 91, 6830 (1969)  
 (b) J K Kochi, R T Tang and T Bernath, *Ibid* , 95, 7114 (1973).  
 (c) L Ebersson, *Ibid* , 89, 4669 (1967).

- 6 It is also possible with apparent 2-electron oxidants such as Tl(III) [ I H Elson and J K Kochi, J Amer Chem Soc , 95, 5061 (1973) ]
- 7 L J Andrews and R N Keefer, "Molecular Complexes in Organic Chemistry", Holden Day, San Francisco, Calif , 1964
- 8 F A Cotton and G Wilkinson "Advanced Inorganic Chemistry", Third Ed , Interscience Pub , 1972, p 357
- 9 (a) V G Shubin, A A Tabatskaya and V A Koptug, Zhur org Khim , 6, 2081 (1970)  
(b) C Eaborn and G J Wright, J Chem , Soc , B, 2262 (1971)