





2-PHENYLSULFONYLETHYL, A NEW PHOSPHATE PROTECTING GROUP: ITS APPLICATION
IN THE SYNTHESIS OF DODECATHYMYDYLIC ACID.







N. Balgobin, S. Josephson and J.B. Chattopadhyaya*

Department of Microbiology, Box 581, Biomedical Centre, Uppsala University, S-751 23 Uppsala,
Sweden.

Summary: 2-Phenylsulphonylethyl group has been employed for the first time to protect 3'-terminal phosphodiester to the triester level in the synthesis of dodecathymidylic acid using the phosphotriester approach.

2-Arylthioethyl group was first introduced by Narang¹ and then Khorana² attempted to use this phosphate protecting group in the phosphodiester approach. The novelties of this phosphate protecting group were considered to lie in the facts that (i) the oligodeoxynucleotides with this group would be sufficiently stable during the removable of acid and base labile protecting groups and (ii) yet, after the conversion of sulfide to the sulfoxide or sulfone oxidation state, it would be removable by base induced β -elimination of the phosphate without any side reactions. Narang used 2M NaOH to activate such a β -elimination of the sulfoxide¹ which resulted in the removal of N-acyl protecting groups. Khorana then improved on this observation by using 1M NaOH for the removal of arylsulfonyl derivative². In the phosphotriester approach³ with the aryl protecting group at the internucleotide junctions, any one of the above conditions is clearly very drastic beside the fact that attempts to oxidize 2-arylthioethyl group led to the oxidations of adenine and guanine residues. Thus 2-arylthioethyl group, as such, for phosphate protection can not be employed successfully. We, therefore, concentrated on exploring the stability of the 2-phenylsulfinylethyl- and 2-phenylsulphonylethyl phosphotriester derivatives (1) and (2) respectively⁴. The 2-phenylsulfinylethyl group in (1) could be removed very slowly with Et₃N (15 equiv.) in dry pyridine (t_{1/2} Ca. 26h at 20°C). However, the result of removal of 2-phenylsulphonylethyl group from (2) was very encouraging by the fact that it could be removed completely under 3h at 20°C using only 2 equiv. of Et₃N in dry pyridine (10ml/mmol) compared to at least 15 equiv. of Et₃N in pyridine for the removal of 2-cyanoethyl protecting group in (3)⁵. Furthermore, We observed that 2-phenylsulphonylethyl group in (2) was stable under the conditions which are normally employed in oligodeoxyribonucleotide synthesis^{5,6,7} despite the relatively mild conditions employable for its removal. Thus the 5'-O-(9-phenylxanthan-9-yl)-(Pixyl) group^{7,8} could be easily removed using 4-toluenesulfonic acid, H₂O in 2% ethanol-CHCl₃ to obtain the corresponding 5'-hydroxy compound (4) quantitatively and both (2) and (4) were reasonably stable during column chromatographic conditions over silica gel^{6,7}. These observations led us to synthesize dodecathymidylic acid using 2-phenylsulphonylethyl group





PSE = 2-Phenylsulphonylethyl

Px = 9-Phenylxanthen-9-yl

Ar = o-Chlorophenyl

to demonstrate its application in oligodeoxyribonucleotide synthesis following the phosphotriester approach^{3,4,9}. Thus, We condensed a slight excess of (5) with (4) in pyridine in presence of a large excess (5 equiv.) of 1-mesitylenesulfonyl-3-nitro-1,2,4-triazole (MS-NT)¹⁰ under a standard condition⁷ to obtain (6), $R_f=0.53$ ¹¹, in 83% yield (powder) after usual work-up and chromatography⁷. Similarly, We obtained (8) in 86% yield, $R_f=0.68$ ¹¹, by reacting (5) with (7) in pyridine soln. in presence of MS-NT. The (6) and (8) could be depixylated^{7,8} in quantitative yields to the corresponding 5'-hydroxy compounds (9) and (10) respectively by 4-toluenesulfonic acid, H₂O in 2% EtOH-CHCl₃ at 20°C. The 2-phenylsulfonylethyl group from (6) was then deprotected to obtain the phosphodiester triethylammonium salt (11) using Et₃N (2 equiv.) in dry pyridine (10ml/mmol) for 3h at 20°C. The deprotection of (6) to (11) was smooth, clean and quantitative. At the end of the reaction, the volatile matters were removed using a rotavapor. The residue was dissolved in small volume of CHCl₃ and then this solution was precipitated from a mixture of Et₂O-petether (30-50'), 1:1 v/v to remove 2-phenylsulfonylethylene completely. The dry powder of (11) was then used, in slight excess (1.1 equiv.), for further condensations with (9) and (10), using standard procedures⁷, to obtain (12) and (13), $R_f=0.48$ & 0.58 respectively¹¹, in 73.0 and 71.0% yields (powders) respectively. The comparison of isolated % yields of (12) and (13) adequately support that 2-phenylsulfonylethyl group at 3'-terminal phosphotriesters is, indeed, stable during the condensation reactions, work-up procedures and column chromatographic conditions which are normally necessary to synthesize and purify a fully protected oligodeoxyribonucleotide fragment. The 2-phenylsulfonylethyl group from (12) was then deprotected to the phosphodiester salt (14) using an identical stoichiometry of Et₃N (i.e 2 equiv.) in pyridine to that of deprotection of (6) to (11) and the reaction was complete within 3h at 20°C. An excess of (14) was then coupled to (15) in a usual fashion to obtain the fully protected octamer (16), $R_f=0.47$ ¹¹, in 74.0% yields (powder) which was then depixylated^{7,8} to obtain the 5'-hydroxy octamer (17) quantitatively. Finally, a condensation of (14) with (17), again under a standard condition⁷, gave the desired dodecathymidine undecaphosphate (18), $R_f=0.38$ ¹¹, in 69.0% yield (powder) after work-up and chromatography (8% EtOH-CHCl₃)^{5,6,7}. The (18) was then deprotected following a literature procedure^{8,10}. The HPLC elution pattern of the crude dodecathymidylic acid through a Permaphase AAX column at 60°C (linear gradient: 0.01M KH₂PO₄, 0.0M KCl to 0.05M KH₂PO₄ and 0.7M KCl, pH 4.45) is shown in Figure 1 and the main peak was found to contain more than 95% of the desired material. An aliquot of this material was ³²P-labelled with γ -³²P-ATP and kinase¹². This ³²P-labelled thymidylic acid was partially digested with *Crotalus adamantus* snake venom phosphodiesterase and then it was electrophoretically shown to contain the desired number of fragments of differently charged species expected from dodecathymidylic acid as shown in Figure 2.

Thus it is clearly demonstrated that 2-phenylsulfonylethyl group can be used for the protection of 3'-phosphodiester function and We are currently exploiting this observation to synthesize oligodeoxyribonucleotide of mixed sequences in our laboratory.

Acknowledgements: The authors thank Professor L. Philipson for encouragements and Swedish Board for Technical Development for generous financial assistance. One of us (J.B.C) specially thank Professor C.B. Reese for many serious discussions during J.B.C's postdoctoral years in C.B.R's laboratory.




Figure 1




Figure 2

References and Notes

1. R.H. Wightman, S.A. Narang and K. Itakura, *Can. J. Chem.* **50**, 456 (1972)
S.A. Narang, O.S. Bhanot, J. Goodchild, R.H. Wightman and S.K. Dheer, *J. Am. Chem. Soc.* **94**, 6183 (1972).
2. K.L. Agarwal, M. Fridkin, E. Jay and H.G. Khorana, *J. Am. Chem. Soc.* **95**, 2020 (1973).
3. C.B. Reese, *Tetrahedron*, **34**, 3143 (1978),
R. Arentzen and C.B. Reese, *J.C.S. Perkin I*, 445 (1977)
4. The phosphotriesters (1) and (2) were synthesized using 2-phenylsulphinyethanol (for (1)) and 2-phenylsulphonyethanol (for (2)) following basically a literature procedure which was used for the preparation of (3), N. Katagiri, K. Itakura and S.A. Narang in *J. Am. Chem. Soc.* **97**, 7332 (1975). 2-Phenylsulphinyethanol and 2-Phenylsulphonyethanol were prepared according to the procedure described by H.S. Schultz, H.B.F. Freyermuth and S.R. Buc in *J. Org. Chem.* **28**, 1141 (1963). Appropriate PMR (CDCl_3) were obtained in support of the structures (1), (2) and (3).
5. R.W. Adamiak, M.Z. Braciszewska, E. Bala, K. Grazeskowiak, R. Kierzek, A. Krazewski, W.T. Markiewicz and M. Wiewiorowski, *Nucleic Acid Res.* **3**, 3397 (1976).
A.K. Sood and S.A. Narang, *Nucleic Acid Res.* **4**, 2757 (1977)
6. W.L. Sung, M.H. Hsiung, R. Brosseau, J. Michniewicz, R. Wu and S.A. Narang, *Nucleic Acid Res.* **7**, 2199 (1979)
W. Pfeleiderer, *Les Colloques de l'INSERM Nucleosides, Nucleotides* **81**, 177 (1978)
J.B. Chattopadhyaya and C.B. Reese, *Nucleic Acid Res.* **8**, 2039 (1980)
7. N. Balgobin, S. Josephson and J.B. Chattopadhyaya, *Acta Chem. Scand. Ser. B*, (in press)
8. J.B. Chattopadhyaya and C.B. Reese, *Chem. Comm.* 639 (1978)
9. T.C. Catlin and F. Crammer, *J. Org. Chem.* **38**, 245 (1973)
10. C.B. Reese, R.C. Titmus and L. Yau, *Tetrahedron Lett.* 2727 (1978)
11. Merck Silica gel 60 F₂₅₄ pre-coated plates have been used for TLC in 10% MeOH- CHCl_3 and Merck kieselgel H was used for short column chromatography using mixtures of EtOH- CHCl_3 as the mobile phase.
12. E. Jay, R. Bambara, R. Padmanabhan and R. Wu, *Nucleic Acid Res.* **1**, 331 (1974)