

SIMPLE REGIOSELECTIVE SYNTHESIS OF TRANS-7a-METHYLHYDRIND-4-EN-1-ONE,
A KEY INTERMEDIATE FOR STEROID TOTAL SYNTHESIS¹


Michael E. Jung^{*2} and Kim M. Halweg³
Department of Chemistry, University of California,
Los Angeles, California 90024

Abstract: Intramolecular Diels-Alder cycloaddition and hydrolysis of the ketals 11 and 12, derived from the trienone 6, afforded mainly the trans-hydrindenone 8a.

The last few years have witnessed a great upsurge in interest in the development of new methods for the total synthesis of steroids.⁴ Most of the new routes have been designed to produce A-ring aromatic steroids such as estrone,⁵ although a few have been aimed at the more medicinally interesting corticosteroid group, e.g., cortisone 1.⁶ We report here a very simple method for the stereoselective preparation of the key trans C,D-ring intermediate 8a for the total synthesis of steroids.


For some time, we have envisioned the approach shown retrosynthetically in Scheme I, namely the preparation of cortisone 1 by standard methods from the dione 2 which could itself be produced from a Diels-Alder reaction of the diene 3⁷ with the hydrindenedione 4.⁸ This key enedione might be prepared by any of several methods, one of the simplest and most intriguing being the intramolecular Diels-Alder addition of either of the two acyclic trienones, 5 or 6, followed by oxidation to the ketone. Thus, it was decided to investigate this expedient route, namely the cyclization of 6 and its derivatives.

Sutherland⁹ reported several years ago that heating of 6 in benzene at 190°C for 13 hours afforded an 85% yield of a mixture of three products in a ratio of 30:4:3. The major product, isolated by preparative gas chromatography (GC), was shown to be the cis hydrindenone 7a, while the other two products were not identified. This result seemed somewhat peculiar in view of our recent work on related systems leading to the total synthesis of coronafacic acid,¹⁰ in which significant amounts of the trans hydrindenone were also produced. Therefore, Sutherland's experiment was repeated under various sets of conditions with somewhat different results. Heating of a solution of 6 in benzene (190°C, 13h) or toluene (170°C, 24h) produced a 99% yield of a mixture of the two cyclized products, 7a and 8a. The ratio of the two products was calculated to be approximately 70:30 favoring the cis by careful integration of the two methyl peaks in the 200 MHz ¹H NMR spectrum. In the cis compound 7a, the methyl appears as a singlet at 1.042δ⁹ while in the trans isomer 8a, the methyl is a small doublet at 0.893δ (W-coupling of 0.7Hz to the proton at the ring juncture). To show that these structural assignments were indeed correct, pure samples of each adduct were isolated by preparative GC or, more easily, by HPLC (Waters 500) using EtOAc/hexane and hydrogenated to the known cis (methyl



at 1.041 δ) and *trans* (methyl at 0.873 δ) hydrindanones¹¹ (yield of hydrogenation, 81%).

Examination of molecular models or three-dimensional drawings (Scheme II) helps one to understand why such high temperatures are required for this cyclization (negligible cyclization in toluene or chloroform solution at temperatures ranging from 25°C to 130°C, even in the presence of acids and/or metal ions) when compared to corresponding cycloadditions in the 1-octalone series (cyclization at 0°C or 25°C, in the presence of acids and/or metal ions).¹² In the *endo* transition state for cyclization 9a, the carbonyl group does not overlap well with the olefin and thus this enone is not as good a dienophile as it would be in an intermolecular reaction. One would also surmise that this non-planarity of the enone system causes the favorable "secondary overlap" leading to the normal stability of the *endo* transition state to be weakened here. Since the enone is "non-conjugated" in the transition state for cyclization, the system resembles a simple triene and thus requires very high temperatures for cyclization. It is interesting to point out that the addition of one methylene unit to the chain allows the carbonyl to overlap effectively with the olefin in the transition state for cycloaddition so that the cycloaddition occurs at 0°C and gives complete *endo* stereospecificity in the 1-octalone formed.¹²



Scheme II

REFERENCES AND NOTES

1. Presented at the meeting of the French Chemical Society, Paris, March 1981.
2. Camille and Henry Dreyfus Teacher-Scholar, 1978-1983; Alfred P. Sloan Foundation Fellow, 1979-1981.
3. University of California Regents Fellow, 1979-1980.
4. R. L. Funk and K. P. C. Vollhardt, Chem. Soc. Rev., 9, 41 (1980).
5. In addition to the references given in ref. 4 above, for more recent examples see: S. Djuric, T. Sarkar, and P. Magnus, J. Am. Chem. Soc., 102, 6885 (1980); Y. Ito, M. Nakatsuka, and T. Saegusa, ibid., 103, 476 (1980); 102, 863 (1980) and references therein.
6. An ingenious intramolecular Diels-Alder approach to cortisone has been described though it is still unpublished. G. Stork, 13th Sheffield Stereochemistry Conference, December 1979. For a discussion, see: Chem. Brit., 270 (1980).
7. For the use of dienes very similar to 3 in Diels-Alder reactions, see: N. L. Goldman, Diss. Abst., 20, 886 (1959); S. P. Tanis and K. Nakanishi, J. Am. Chem. Soc., 101, 4398 (1979).
8. Addition of 3 and 4 should produce the stereochemistry (approach of diene from less hindered side and Alder's endo rule) and regiochemistry (Houk's molecular orbital coefficient analysis) shown in 2 as the major product.
9. J. K. Sutherland, J. Chem. Soc. Perkin Trans. 1, 1559 (1975).
10. M. E. Jung and K. M. Halweg, Tetrahedron Lett., in press.
11. P. T. Lansbury, P. C. Briggs, T. R. Demmin, and G. E. DuBois, J. Am. Chem. Soc., 93, 1311 (1971).
12. a) O. P. Vig, I. R. Trehan, and R. Kumar, Ind. J. Chem., 15B, 319 (1977); O. P. Vig, I. R. Trehan, N. Malik, and R. Kumar, ibid., 16B, 449 (1978).
 b) J.-L. Gras and M. Bertrand, Tetrahedron Lett., 4549 (1979).
 c) D. F. Taber and B. P. Gunn, J. Am. Chem. Soc., 101, 3992 (1979).
13. The tetramethylethylene ketal corresponding to 11 could be formed in modest yield from 11 and pinacol in benzene with acid catalysis. Its intramolecular cycloaddition required three days of heating at 170°C and produced after acidic hydrolysis, an approximately 1:3 mixture of 7a and 8a in 97% yield. It should be pointed out that Prof. Paul Helquist has observed similar stereoselectivity in the internal cyclization of another ketal of 6; see accompanying paper.
14. An additional proposal for increasing the stereoselectivity involves having a bulky substituent at the 7-position of the 1,6,8-trien-3-one, as in cyclizations leading to coronafacic acid.^{10,15} For example, intramolecular cycloaddition of the compound corresponding to 11 in which the C7 and C8 hydrogens have been replaced by a tetramethylene unit (i.e., a 1,2-alkylidene methylene cyclohexane) would be expected to afford more of the trans isomer due to increased steric interaction between the ketal functionality and the methylene group attached to C7. This is being tested currently.
15. A. Ichihara, R. Kimura, S. Yamada, and S. Sakamura, J. Am. Chem. Soc., 102, 6353 (1980).

(Received in USA 18 June 1981)