

DIENAMINES AS DIELS-ALDER DIENES. A NOVEL BENZANNULATION SEQUENCE.


Roger L. Snowden* and Manfred Wüst

Firmenich SA, Research Laboratories, 1211 Geneva 8, Switzerland

Abstract: *Diels-Alder* reaction between (E)-dienamines 1 - 7 and methyl propiolate followed by elimination of Me₂NH from the intermediate cycloadducts III affords the aromatic methyl esters 8 - 14 in good yield.

The preceding Letter¹ describes an efficient sequence for the cyclohexannulation of cycloalkanones I using, as the key step, a *Diels-Alder* reaction between (E)-dienamines II and methyl acrylate. We now report an analogous benzannulation sequence, I → IV², which involves the regioselective [4+2]cycloaddition of II (R'=H) with methyl propiolate and subsequent elimination of Me₂NH from the intermediate cycloadduct III³ (*cf. Scheme*). Accordingly, reaction of dienamines

Scheme




1 - 7 with methyl propiolate (1.5 mole equiv.) in toluene (10% solution) at 25⁰ or 50⁰ (*cf. Table*) followed by treatment with silica gel at 60⁰ during 2 h afforded the methyl esters 8 - 14 in good yield. This one-pot procedure thus allows the efficient construction of aromatic carboxylic esters IV from (E)-dienamines II with complete regiochemical control⁴.

References and Notes

- [1] See preceding paper.
- [2] For examples of previous benzannulation methods, see: J.B.Dickenson & W.Reusch, *Syn.Comm.* 13(4), 303 (1983); M.A.Tius & A.Thurkauf, *J.Org.Chem.* 48, 3839 (1983); D.L.Boger & M.D.Mullican, *Tetrahedron Lett.* 24, 4939 (1983); K.Takaki, M.Ohsugi, M.Okada, M.Yasumura & K.Negoro, *J.Chem.Soc. Perkin I*, 741 (1984); G.Singh, H.Ila & H.Junjappa, *Tetrahedron Lett.* 25, 5095 (1984); R.K.Dieker & Y.J.Lin, *ibid* 26, 39 (1985) and references cited therein.
- [3] For an analogous *Diels-Alder* reaction of acyclic dienamines, see: S.Tanimoto, Y.Matsumura, T.Sugimoto & M.Okano, *Tetrahedron Lett.*, 2899 (1977).
- [4] The influence of the dimethylamino group on the rate and regioselectivity of the *Diels-Alder*

Table




Entry	b) DIENE AMINE	D.-A. React. Cond.	PRODUCT ^{c)}	Yield ^{d)} %	IVa m.p.(°C)
1	<u>1</u> (n=1)	25°/5h	<u>8</u> ⁸ (n=1)	89	180 - 182 ⁰⁵
2	<u>2</u> (n=2)	25°/24h	<u>9</u> ⁶ (n=2)	76	152 - 153 ⁰⁶
3	<u>3</u> (n=3)	25°/5h	<u>10</u> (n=3)	78	174 - 175 ⁰⁷
4	<u>4</u> (n=8) e)	50°/24h	<u>11</u> (n=8) (m.p. 113-5 ⁰)	52	181 - 182 ⁰
5	<u>5</u>	50°/5h	<u>12</u>	81	203 - 205 ⁰
6	<u>6</u>	50°/24h	<u>13</u>	83	144 - 146 ⁰
7	<u>7</u>	25°/24h	<u>14</u> (m.p. 31-3 ⁰)	82	179 - 181 ⁰

a) The presumed stereochemistry of III is consistent with the ¹H-NMR(360 MHz, CDCl₃) spectral data, e.g. entry 7: δ1.06, 1.15 (2 s, CH₃); 1.00 - 2.10 (6 H); 2.24 (s, N(CH₃)₂); 2.88 (m, H-C(8a)); 3.76 (s, CO₂CH₃); 4.21 (dd, J = 4, 4 Hz, H-C(3)); 5.54 (d, J = 4 Hz, H-C(4)); 6.82 (d, J = 4 Hz, H-C(1)); b) for the preparations of 1, 2, 3, 6 and 7 cf. reference 1; 4 and 5 were prepared from cyclododecanone and 4-t butylcyclohexanone resp. (cf. Method A¹); c) all new compounds have been fully characterised spectroscopically e.g. 14: IR(film): 1718, 1284, 1196, 762 cm⁻¹; UV(EtOH): λ_{max}. 242 (ε 13,700); ¹H-NMR(CDCl₃): δ1.30 (s, 6 H); 1.60 - 1.90 (4 H); 2.81 (t, J = 7 Hz, 2 H); 3.89 (s, 3 H); 7.38 (d, J = 8 Hz, H-C(4)); 7.74 (br.s, H-C(1)); 7.78 (br.d, J = 8 Hz, H-C(3)); MS: 218 (19, M⁺), 203 (100), 171 (25), 144 (15), 129 (18); d) yields refer to chromatographically pure, distilled products; e) 1:1 mixture of E/Z-1'-cyclododeceny1 double bond isomers

reactions of 1 and 2 is demonstrated by comparison with the dienes i and ii, see:

J.H.Markgraf, E.W.Greeno, M.D.Miller, W.J.Zaks & G.A.Lee, *Tetrahedron Lett.* 24, 241 (1983).



[5] J.Lindner, F.Schmitt & B.Zaubauer, *Monatsh.* 72, 216 (1939).

[6] M.S.Newman & H.V.Zahm, *J.Am.Chem.Soc.* 65, 1097 (1943).

[7] R.H.Wightman, D.E.Laycock & H.W.Avdovich, *J.Org.Chem.* 43, 2167 (1978).

[8] I.M.Hunsberger, D.Lednicer, H.Girtowsky, D.Bunker & P.Taussig, *J.Am.Chem.Soc.* 2466 (1955).

(Received in France 5 December 1985)